

Patient Information Patient's name: xxx

gender

male

age

44

ECOG-PS

0

Child-Pugh

A

Extrahepatic metastasis

none

CNLC

IA

Laboratory test indicators

Indicator Name	value	unit	Normal range
Alanine aminotransferase (ALT)	37.9	U/L	9-50
Activated partial thromboplastin time (APTT)	24.9	sec	22.7-36.5
Alpha-fetoprotein (AFP)	31.17 ↑	ng/mL	<20

MRI Imaging Report - Preoperative Imaging Examination - MRI Imaging. 2

[Impressions from Radiography]

Liver cirrhosis,
hepatocellular carcinoma, hepatocellular carcinoma,
gallbladder polyps? Small stones?
No enhancing lesions in the spleen, benign vascular tumor,
left retroperitoneal nodule, hemangioma?

[Imaging Findings]

The liver is of regular shape with irregular edges, and the proportions of each lobe are normal. ...
A T2WI high signal nodule, about 6 mm in diameter, with significant enhancement, is seen subcutaneously on the left posterior abdominal wall (SE9, IM26).

Model recommendation scheme


optimal: Surgical resection
Candidates: Surgical resection, ablation therapy, liver transplantation



* Scores are based on predictions from the IDEAL model.

Expected Survival

Median survival	186 months
1-year survival rate	95.6%
3-year survival rate	87.4%
5-year survival rate	80%



Model reasoning process

The patient is a 44-year-old male diagnosed with hepatocellular carcinoma (NOS), pathological grade I, with a tumor size of 12mm × 20mm located in the S3 region of the liver. Imaging findings suggest a high probability of hepatocellular carcinoma against a background of cirrhosis, with no extrahepatic metastasis, no portal vein tumor thrombus, and a Child-Pugh score of [missing information].APS rating:0,noneRelocation to distant placesnoneTumor emboli, number of tumors1Maximum tumor diameter12mm. According to the decision tree, the current PS score is...0child-pugh level isAlmagingnoneTumor thrombus, andnonelt showed distant metastasis, number of tumors1The sample has a maximum diameter of 12mm and is less than 5cm; therefore, this patient belongs to CNLC stage Ia. Based on the guideline decision tree, recommended treatment options include:Surgical resection,Ablation therapyas well asLiver transplantDetailed analysis of each potential treatment option: 1. Surgical resection: According to guidelines, surgical resection is the preferred treatment for CNLC stage Ia, Ib, and IIa hepatocellular carcinoma (Level 1 evidence, Recommendation A). The patient's tumor size is 12 mm, the Child-Pugh score is A, and there is no portal hypertension or extrahepatic metastasis, meeting the indications for surgical resection. Furthermore, imaging studies do not indicate tumor invasion of important blood vessels or bile ducts, therefore surgical resection is technically feasible. 2. Ablation therapy: According to guidelines, ablation therapy is mainly applicable to CNLC stage Ia and some stage Ib hepatocellular carcinomas (i.e., a single tumor ≤5 cm in diameter; or 2–3 tumors, with the largest diameter ≤3 cm). The patient's tumor size is 12 mm, meeting the indications for ablation therapy, and no contraindications have been violated (e.g., the tumor is not near the porta hepatis or first-order bile ducts). Ablation therapy has similar efficacy to surgical resection for small hepatocellular carcinomas, especially when the tumor diameter is