

Patient Information Patient's name: xxx

gender

male

age

44

ECOG-PS

0

Child-Pugh

A

Extrahepatic metastasis

none

CNLC

IA

Laboratory test indicators

Indicator Name	value	unit	Normal range
Alanine aminotransferase (ALT)	37.9	U/L	9-50
Activated partial thromboplastin time (APTT)	24.9	sec	22.7-36.5
Alpha-fetoprotein (AFP)	31.17 ↑	ng/mL	<20

MRI Imaging Report - Preoperative Imaging Examination - MRI Imaging. 2

[Impressions from Radiography]

Liver cirrhosis,
hepatocellular carcinoma, hepatocellular carcinoma,
gallbladder polyps? Small stones?
No enhancing lesions in the spleen, benign vascular tumor,
left retroperitoneal nodule, hemangioma?

[Imaging Findings]

The liver is of regular shape with irregular edges, and the proportions of each lobe are normal. ...
A T2WI high signal nodule, about 6 mm in diameter, with significant enhancement, is seen subcutaneously on the left posterior abdominal wall (SE9, IM26).

Model recommendation scheme

optimal: Surgical resection


Candidates: Surgical resection, ablation therapy, liver transplantation



* Scores are based on predictions from the IDEAL model.

Expected Survival

Median survival	186 months
1-year survival rate	95.6%
3-year survival rate	87.4%
5-year survival rate	80%



Model reasoning process

The patient is a 44-year-old male diagnosed with hepatocellular carcinoma (NOS), pathological grade I, tumor size 12mm × 20mm, located in the S3 region of the liver. Imaging findings suggest a high probability of hepatocellular carcinoma against a background of cirrhosis, with no extrahepatic metastasis, no portal vein tumor thrombus, Child-Pugh score **A** , PS score **0** , **no** distant metastasis, **no** tumor thrombus, **1** tumor , and a maximum tumor diameter of **12mm** .

According to the decision tree, with a current PS score of **0** , Child-Pugh grade **A** , **no** tumor thrombus on imaging , **no** distant metastasis, **1** tumor , and a maximum diameter of 12mm (less than 5cm), this patient is classified as CNLC stage Ia. Based on the guideline decision tree, recommended treatment options include: **surgical resection** , **ablation therapy** , and **liver transplantation** .

Detailed analysis of each potential option

: 1. Surgical resection: According to the guideline, surgical resection is the preferred treatment for CNLC stage Ia, Ib, and IIa (Level of evidence 1, Recommendation A). The patient's tumor size is 12mm, the Child-Pugh score is A, and there is no portal hypertension or extrahepatic metastasis, meeting the indications for surgical resection. Furthermore, imaging studies do not indicate tumor invasion of major blood vessels or bile ducts; therefore, surgical resection is technically feasible.

2. Ablation therapy: According to guidelines, ablation therapy is mainly suitable for CNLC stage Ia and some