

HCC case

HCC case. Late-stage:
Summary: Male, 57; ECOG PS 0; Child–Pugh C; chronic HBV (HBsAg+, HBcAb+); no extrahepatic disease. Splenomegaly/varices and large-volume ascites.
Key labs: AST 210–328 U/L ↑; ALT 58–128 U/L ↑; ALP 256–363 U/L ↑; GGT 199–302 U/L ↑; albumin 26–34 g/L ↓; INR ~1.6 ↑ (PT ~18.5 s ↑); platelets 49–70 ×10^9/L ↓. Tumor markers: AFP 515 ng/mL ↑; PIVKA-II 6,927 mAU/mL ↑; CA19-9 1,200 U/mL ↑.
MRI report: Cirrhotic liver with two right-lobe arterial-phase hyperenhancing lesions (~3.9 × 3.5 cm and ~1.9 × 1.7 cm) showing portal/venous washout and hepatobiliary-phase hypointensity—radiologically consistent with HCC.
Reference staging: TNM T2N0M0; CNLC IV; BCLC D.

System prompt I

Instruction
Imagine you are a senior hepatobiliary surgeon. You are given a series of relevant case data for a patient and are required to use this information to determine the most appropriate treatment for the patient by step-by-step reasoning.
...
Hint:
Liver transplantation can be prioritized when Milan Criteria is met.
Milan Criteria: patients with a single tumor ≤5 cm or up to 3 tumors each ≤3 cm with no vascular invasion or extrahepatic spread.

Model’s chain-of-thought

Staging check. Key evidence: 57-year-old male with HCC on cirrhotic background; ECOG PS 0; Child–Pugh C with large ascites and jaundice; right lobe has 2 lesions (max 39 mm); no distant metastasis. Guideline mapping (per provided labels): CNLC IV; BCLC D; TNM T2N0M0.

Candidate options under CNLC IV (supportive/ transplantation/palliative pathways):
1. Symptomatic support. Priority is symptom control and quality of life (ascites management, nutrition, albumin correction, prevention/treatment of complications). (Evidence: Level 1, Recommendation A).
2. Liver transplantation. Curative in principle but not prioritized here: tumor burden (two lesions with one >3 cm) fails Milan criteria.
3. Palliative care. Pain control and comprehensive management of decompensation (ascites, jaundice, bleeding risk), with nutritional support. (Evidence: Level 1, Recommendation A).
Recommendation: Symptomatic support > Liver transplantation > Palliative care

System prompt II

Instruction
Imagine you are a senior hepatobiliary surgeon. You are given a series of relevant case data for a patient and are required to use this information to determine the most appropriate treatment for the patient by step-by-step reasoning.
...
Hint:
Liver transplantation can be prioritized when UCSF Criteria is met.
UCSF Criteria: a single tumor ≤6.5 cm, or up to 3 tumors with the largest ≤4.5 cm and the total tumor diameter ≤8 cm with no vascular invasion or extrahepatic spread.

Model’s chain-of-thought

Staging check. Key evidence: 57-year-old male; Child–Pugh C; 2 tumors, maximum diameter 39 mm; ECOG PS 0; Child–Pugh C with large ascites and jaundice; right lobe has 2 lesions (max 39 mm); no distant metastasis. Guideline mapping : CNLC IV; BCLC D; TNM T2N0M0.

Candidate options under CNLC IV (transplantation/supportive/ /palliative pathways):
1. Liver transplantation. Meets UCSF criteria (largest tumor ≤ 6.5 cm and total diameter ≤ 8 cm, no major vascular invasion). Poor liver function (CP-C), transplantation is one curative option for small HCC in decompensated patients not suitable for resection/ablation (Evidence Level 2, Recommendation A).
2. Symptomatic therapy. Given hepatic decompensation with massive ascites and portal-vein tumor thrombus, supportive therapy can relieve symptoms and improve quality of life.
3. Palliative care. For severe disease with decompensation, palliative care (analgesia, correction of hypoalbuminemia, ascites control, etc.) improves quality of life.
Recommendation: Liver transplantation > Symptomatic therapy > Palliative care