

HCC case

HCC case. Late-stage:
Summary: Male, 57; ECOG PS 0; Child–Pugh C; chronic HBV (HBsAg+, HBcAb+); no extrahepatic disease. Splenomegaly/varices and large-volume ascites.
Key labs: AST 210–328 U/L ↑; ALT 58–128 U/L ↑; ALP 256–363 U/L ↑; GGT 199–302 U/L ↑; albumin 26–34 g/L ↓; INR ~1.6 ↑ (PT ~18.5 s ↑); platelets 49–70 ×10⁹/L ↓. Tumor markers: AFP 515 ng/mL ↑; PIVKA-II 6,927 mAU/mL ↑; CA19-9 1,200 U/mL ↑.
MRI report: Cirrhotic liver with two right-lobe arterial-phase hyperenhancing lesions (~3.9 × 3.5 cm and ~1.9 × 1.7 cm) showing portal/venous washout and hepatobiliary-phase hypointensity—radiologically consistent with HCC.
Reference staging: TNM T2N0M0; CNLC IV; BCLC D.



System prompt I

Instruction
Imagine you are a senior hepatobiliary surgeon. You are given a series of relevant case data for a patient and are required to use this information to determine the most appropriate treatment for the patient by step-by-step reasoning.
...
Hint:
Liver transplantation can be prioritized when Milan Criteria is met.
Milan Criteria: patients with a single tumor ≤5 cm or up to 3 tumors each ≤3 cm with no vascular invasion or extrahepatic spread.



Model's chain of thought

Staging check. Key evidence: 57-year-old male with HCC on cirrhotic background; ECOG PS 0; Child–Pugh C with large ascites and jaundice; right lobe has 2 lesions (max 39 mm); no distant metastasis. Guideline mapping (per provided labels): **CNLC IV; BCLC D; TNM T2N0M0.**

Candidate options under CNLC IV (supportive/ **transplantation**/palliative pathways):

- Symptomatic support.** Priority is symptom control and quality of life (ascites management, nutrition, albumin correction, prevention/treatment of complications). (*Evidence: Level 1, Recommendation A*).
- Liver transplantation.** Curative in principle but **not prioritized here: tumor burden (two lesions with one >3 cm) fails Milan criteria.**
- Palliative care.** Pain control and comprehensive management of decompensation (ascites, jaundice, bleeding risk), with nutritional support. (*Evidence: Level 1, Recommendation A*).

Recommendation: Symptomatic support > Liver transplantation > Palliative care



System prompt II

Instruction
Imagine you are a senior hepatobiliary surgeon. You are given a series of relevant case data for a patient and are required to use this information to determine the most appropriate treatment for the patient by step-by-step reasoning.
...
Hint:
Liver transplantation can be prioritized when UCSF Criteria is met.
UCSF Criteria: a single tumor ≤6.5 cm, or up to 3 tumors with the largest ≤4.5 cm and the total tumor diameter ≤8 cm with no vascular invasion or extrahepatic spread.



Model's chain of thought

Staging check. Key evidence: 57-year-old male; Child–Pugh C; 2 tumors, maximum diameter 39 mm; ECOG PS 0; Child–Pugh C with large ascites and jaundice; right lobe has 2 lesions (max 39 mm); no distant metastasis. Guideline mapping : **CNLC IV; BCLC D; TNM T2N0M0.**

Candidate options under CNLC IV (**transplantation**/supportive/ /palliative pathways):

- Liver transplantation.** Meets UCSF criteria (largest tumor ≤ 6.5 cm and total diameter ≤ 8 cm, no major vascular invasion). Poor liver function (CP-C), transplantation is one curative option for small HCC in decompensated patients not suitable for resection/ablation (*Evidence Level 2, Recommendation A*).
- Symptomatic therapy.** Given hepatic decompensation with massive ascites and portal-vein tumor thrombus, supportive therapy can relieve symptoms and improve quality of life.
- Palliative care.** For severe disease with decompensation, palliative care (analgesia, correction of hypoalbuminemia, ascites control, etc.) improves quality of life.

Recommendation: Liver transplantation > Symptomatic therapy > Palliative care