

## Getting / cleaning data 2

---

# Tidy data

---

All of the material in this section comes directly from Hadley Wickham's [paper on tidy data](#). You will need to read this paper to prepare for the quiz on this section.

# Characteristics of tidy data

Characteristics of tidy data are:

1. Each variable forms a column.
2. Each observation forms a row.
3. Each type of observational unit forms a table.

Getting your data into a “tidy” format makes it easier to model and plot. By taking the time to tidy your data at the start of an analysis, you will save yourself time, and make it easier to plan out, later steps.

# Five common problems

Here are five common problems that Hadley Wickham has identified that keep data from being tidy:

1. Column headers are values, not variable names.
2. Multiple variables are stored in one column.
3. Variables are stored in both rows and columns.
4. Multiple types of observational units are stored in the same table.
5. A single observational unit is stored in multiple tables.

In the following slides, I'll give examples of each of these problems.

## Five common problems

(1.) Column headers are values, not variable names.

religion	<\$10k	\$10-20k	\$20-30k	\$30-40k	\$40-50k	\$50-75k
Agnostic	27	34	60	81	76	137
Atheist	12	27	37	52	35	70
Buddhist	27	21	30	34	33	58
Catholic	418	617	732	670	638	1116
Don't know/refused	15	14	15	11	10	35
Evangelical Prot	575	869	1064	982	881	1486
Hindu	1	9	7	9	11	34
Historically Black Prot	228	244	236	238	197	223
Jehovah's Witness	20	27	24	24	21	30
Jewish	19	19	25	25	30	95

## Five common problems

Solution:

religion	income	freq
Agnostic	<\$10k	27
Agnostic	\$10-20k	34
Agnostic	\$20-30k	60
Agnostic	\$30-40k	81
Agnostic	\$40-50k	76
Agnostic	\$50-75k	137
Agnostic	\$75-100k	122
Agnostic	\$100-150k	109
Agnostic	>150k	84
Agnostic	Don't know/refused	96

## Five common problems

(2.) Multiple variables are stored in one column.

country	year	column	cases
AD	2000	m014	0
AD	2000	m1524	0
AD	2000	m2534	1
AD	2000	m3544	0
AD	2000	m4554	0
AD	2000	m5564	0
AD	2000	m65	0
AE	2000	m014	2
AE	2000	m1524	4
AE	2000	m2534	4
AE	2000	m3544	6
AE	2000	m4554	5
AE	2000	m5564	12
AE	2000	m65	10
AE	2000	f014	3



## Five common problems

Solution:

country	year	sex	age	cases
AD	2000	m	0-14	0
AD	2000	m	15-24	0
AD	2000	m	25-34	1
AD	2000	m	35-44	0
AD	2000	m	45-54	0
AD	2000	m	55-64	0
AD	2000	m	65+	0
AE	2000	m	0-14	2
AE	2000	m	15-24	4
AE	2000	m	25-34	4
AE	2000	m	35-44	6
AE	2000	m	45-54	5
AE	2000	m	55-64	12
AE	2000	m	65+	10
AE	2000	f	0-14	3

## Five common problems

(3.) Variables are stored in both rows and columns.

id	year	month	element	d1	d2	d3	d4	d5	d6	d7	d8
MX17004	2010	1	tmax	—	—	—	—	—	—	—	—
MX17004	2010	1	tmin	—	—	—	—	—	—	—	—
MX17004	2010	2	tmax	—	27.3	24.1	—	—	—	—	—
MX17004	2010	2	tmin	—	14.4	14.4	—	—	—	—	—
MX17004	2010	3	tmax	—	—	—	—	32.1	—	—	—
MX17004	2010	3	tmin	—	—	—	—	14.2	—	—	—
MX17004	2010	4	tmax	—	—	—	—	—	—	—	—
MX17004	2010	4	tmin	—	—	—	—	—	—	—	—
MX17004	2010	5	tmax	—	—	—	—	—	—	—	—
MX17004	2010	5	tmin	—	—	—	—	—	—	—	—

## Five common problems

Solution:

id	date	element	value
MX17004	2010-01-30	tmax	27.8
MX17004	2010-01-30	tmin	14.5
MX17004	2010-02-02	tmax	27.3
MX17004	2010-02-02	tmin	14.4
MX17004	2010-02-03	tmax	24.1
MX17004	2010-02-03	tmin	14.4
MX17004	2010-02-11	tmax	29.7
MX17004	2010-02-11	tmin	13.4
MX17004	2010-02-23	tmax	29.9
MX17004	2010-02-23	tmin	10.7

id	date	tmax	tmin
MX17004	2010-01-30	27.8	14.5
MX17004	2010-02-02	27.3	14.4
MX17004	2010-02-03	24.1	14.4
MX17004	2010-02-11	29.7	13.4
MX17004	2010-02-23	29.9	10.7
MX17004	2010-03-05	32.1	14.2
MX17004	2010-03-10	34.5	16.8
MX17004	2010-03-16	31.1	17.6
MX17004	2010-04-27	36.3	16.7
MX17004	2010-05-27	33.2	18.2

## Five common problems

(4.) Multiple types of observational units are stored in the same table.

year	artist	time	track	date	week	rank
2000	2 Pac	4:22	Baby Don't Cry	2000-02-26	1	87
2000	2 Pac	4:22	Baby Don't Cry	2000-03-04	2	82
2000	2 Pac	4:22	Baby Don't Cry	2000-03-11	3	72
2000	2 Pac	4:22	Baby Don't Cry	2000-03-18	4	77
2000	2 Pac	4:22	Baby Don't Cry	2000-03-25	5	87
2000	2 Pac	4:22	Baby Don't Cry	2000-04-01	6	94
2000	2 Pac	4:22	Baby Don't Cry	2000-04-08	7	99
2000	2Ge+her	3:15	The Hardest Part Of ...	2000-09-02	1	91
2000	2Ge+her	3:15	The Hardest Part Of ...	2000-09-09	2	87
2000	2Ge+her	3:15	The Hardest Part Of ...	2000-09-16	3	92
2000	3 Doors Down	3:53	Kryptonite	2000-04-08	1	81
2000	3 Doors Down	3:53	Kryptonite	2000-04-15	2	70
2000	3 Doors Down	3:53	Kryptonite	2000-04-22	3	68
2000	3 Doors Down	3:53	Kryptonite	2000-04-29	4	67
2000	3 Doors Down	3:53	Kryptonite	2000-05-06	5	66

## Five common problems

Solution:

id	artist	track	time	id	date	rank
1	2 Pac	Baby Don't Cry	4:22	1	2000-02-26	87
2	2Ge+her	The Hardest Part Of ...	3:15	1	2000-03-04	82
3	3 Doors Down	Kryptonite	3:53	1	2000-03-11	72
4	3 Doors Down	Loser	4:24	1	2000-03-18	77
5	504 Boyz	Wobble Wobble	3:35	1	2000-03-25	87
6	98~0	Give Me Just One Nig...	3:24	1	2000-04-01	94
7	A*Teens	Dancing Queen	3:44	1	2000-04-08	99
8	Aaliyah	I Don't Wanna	4:15	2	2000-09-02	91
9	Aaliyah	Try Again	4:03	2	2000-09-09	87
10	Adams, Yolanda	Open My Heart	5:30	2	2000-09-16	92
11	Adkins, Trace	More	3:05	3	2000-04-08	81
12	Aguilera, Christina	Come On Over Baby	3:38	3	2000-04-15	70
13	Aguilera, Christina	I Turn To You	4:00	3	2000-04-22	68
14	Aguilera, Christina	What A Girl Wants	3:18	3	2000-04-29	67
15	Alice Deejay	Better Off Alone	6:50	3	2000-05-06	66

## Five common problems

(5.) A single observational unit is stored in multiple tables.

Example: exposure and outcome data stored in different files:

- File 1: Daily mortality counts
- File 2: Daily air pollution measurements