

Where Every Drop Goes

Hidden Maps · Episode 5 — The James River Watershed as a Network

Troy Altus

2026-06-01

Table of contents

1	Abstract	1
2	Introduction	2
3	Data	2
3.1	Source and Acquisition	2
3.2	Graph Model	3
4	Methods	3
4.1	Strahler Stream Order	3
4.2	Path Length	3
4.3	Betweenness Centrality	3
5	Results	4
5.1	Geographic Map — Stream Hierarchy	4
5.2	Strahler Order Distribution	4
5.3	Junction Degree Structure	4
5.4	Flow Path Lengths	4
5.5	Confluence Explorer	5
6	Discussion	5
7	Conclusion	5
	References	6

1 Abstract

Rain that falls near Roanoke, Virginia will, if it doesn't evaporate, travel more than 200 miles through creeks, rivers, and major confluences before reaching the Chesapeake Bay. That journey follows a branching directed graph — the James River watershed — in which roughly 10,000 stream segments connect into a single drainage network. This study applies network analysis to the James River watershed as characterized by the USGS National Hydrography Dataset, treating stream junctions as nodes and stream reaches as directed edges. We describe the

network’s topology: its headwater sources, its Strahler order distribution, the distribution of flow-path lengths from source to mouth, and the betweenness centrality of named confluence nodes. The watershed is a nearly perfect directed tree, tolerant of individual segment blockage but vulnerable at a handful of confluences where tributaries merge — locations where a dam, contamination event, or habitat barrier has outsized downstream consequence.

2 Introduction

Water does not spread evenly. It collects. Every terrain feature — ridge, hill, hollow — divides the land into catchments that funnel rain toward a common low point: a stream, then a river, then a bay. The shape of that funnel is not random. It is a network: branching, hierarchical, and directional. Water moves only downstream.

Network science applied to river systems is not new — hydrologists have used graph-theoretic tools since at least Robert Horton’s 1945 drainage network laws (Horton 1945) and Arthur Strahler’s 1952 refinement of stream ordering (Strahler 1952). What is newer is the availability of standardized, digitized watershed data at fine resolution, and the open-source tools to analyze it interactively. The USGS National Hydrography Dataset (NHDPlus) provides a topologically routed, attributed flowline network for the entire contiguous United States.

This study applies the same analytical frame used in previous Hidden Maps episodes — degree distribution, path structure, centrality, robustness — to a directed watershed graph. The subject is the **James River watershed**, which drains roughly 10,000 square miles of Virginia, from the Allegheny Highlands in the west to the Chesapeake Bay in the east.

Three questions organize this paper:

1. **Structure:** What is the degree distribution and hierarchy of the James River drainage network?
2. **Centrality:** Which confluence nodes are structural bottlenecks — the junctions whose disruption would disconnect the most upstream drainage?
3. **Directionality:** How does the directed character of a river network differ from the undirected networks studied in earlier episodes?

3 Data

3.1 Source and Acquisition

Flowline data are drawn from the **USGS National Hydrography Dataset Plus** (NHDPlus v2), accessed via the [Network Linked Data Index](#) (NLDI) API and the `pynhd` Python package. The outlet gauge is **USGS-02037500** (James River at Buchanan, VA), capturing the upper and middle James River drainage.

Each flowline segment carries a **fromnode** and **tonode** pair that defines how reaches connect at junctions. We build a directed graph whose nodes are hydrographic junctions and whose edges are stream reaches, annotated with Strahler order, reach length in miles, and cumulative drainage area in square miles.

Table 1: James River watershed network statistics.

Table 1	
Statistic	Value
Stream reaches (directed edges)	9,925
Hydro junctions (nodes)	9,808
Headwater junctions (in-degree = 0)	3,475
Outlet junctions (out-degree = 0)	1
Total stream length (miles)	9,263
Watershed drainage area (sq mi)	6,753
Binary-or-higher confluences (in-degree ≥ 2)	3,572
Maximum Strahler order	6

3.2 Graph Model

The watershed is represented as a **directed acyclic graph** (DAG): nodes are hydrographic junctions; edges are stream reaches directed downstream. This is structurally different from every network analyzed in the Hidden Maps series to date. Airport routes, power transmission, street grids, and internet routing are all essentially undirected or bidirectional. Water does not back up.

4 Methods

4.1 Strahler Stream Order

Strahler ordering (Strahler 1952) assigns integer ranks to stream segments based on tributary hierarchy. Strahler order is precomputed in NHDPlus and serves here as the primary structural hierarchy metric.

4.2 Path Length

The **flow path length** from a headwater junction to the outlet measures how far water travels through the network. We compute all shortest paths (in miles) from headwater sources to the outlet, using reach length as edge weight.

4.3 Betweenness Centrality

Betweenness centrality measures the fraction of shortest paths between node pairs that pass through a junction. High-betweenness nodes are confluences that aggregate drainage from many upstream branches. We compute betweenness on the undirected version of the graph for comparability with prior Hidden Maps episodes, then map scores back onto stream reaches for geographic visualization.

Unable to display output for mime type(s): text/html

(a) James River watershed flowlines colored by Strahler stream order, with geographic context east to the Chesapeake Bay. The NHD pull ends at Buchanan (upper James); the dotted line sketches the lower stem continuing to tidal water.

Unable to display output for mime type(s): text/html

(b)

Figure 1

5 Results

5.1 Geographic Map — Stream Hierarchy

The map makes the hierarchy visible (Figure 1). First-order headwater streams form a dense pale mesh across the Piedmont and Highlands; higher-order segments trace the James main stem and its major forks in darker blue. The shaded estuary and dotted lower stem show where this network ultimately delivers flow — more than 100 miles beyond the NHD pull terminus at Buchanan.

5.2 Strahler Order Distribution

Unable to display output for mime type(s): text/html

Figure 2: Strahler stream order distribution across 9,925 reaches. Each successive order contains roughly three to four times fewer segments than the one below — the defining bifurcation pattern of a drainage network.

The Strahler distribution is steeply right-skewed (Figure 2). The vast majority of reaches are first-order headwater streams. Maximum Strahler order is 6.

5.3 Junction Degree Structure

Unable to display output for mime type(s): text/html

Figure 3: Junction in-degree and out-degree distributions. Incoming degree counts how many reaches meet at a junction; outgoing degree counts how many channels leave it. Binary confluences (in-degree = 2) dominate; the outlet is the single node with out-degree = 0.

The degree structure is tree-like but not uniform (Figure 3). **3,552** junctions receive exactly two incoming reaches — classic binary confluences where two tributaries merge. **3,475** junctions are headwater sources with no incoming flow. Out-degree is almost always 1: water leaves a junction through a single downstream channel. The lone outlet (out-degree = 0) is the watershed mouth.

Unlike scale-free networks (airports, internet), there are no hubs with dozens of direct connections. Hierarchy emerges through sequential merging downstream.

5.4 Flow Path Lengths

The median flow path is 139 miles; 90% of headwater junctions lie within 270 miles of the outlet (Figure 4). The maximum path is 318 miles — the Jackson and Cowpasture headwaters crossing the full length of the valley before reaching tidal water.

Unable to display output for mime type(s): text/html

Figure 4: Cumulative distribution of flow-path length from headwater junctions to the outlet. Most headwaters are close to the main stem; the long upper tail reflects western Highland tributaries traveling the full valley length.

Roughly **9%** of headwaters drain to the outlet in under 50 miles. The long tail is where network structure becomes visceral: a raindrop near Warm Springs faces a 318-mile journey.

5.5 Confluence Explorer

Unable to display output for mime type(s): text/html

Figure 5: Interactive confluence explorer: click any ranked bar to zoom the map and highlight that junction. Marker color matches bar rank; marker size scales with upstream drainage area. The map extends east to the Chesapeake Bay for geographic context.

Betweenness concentrates along the James main stem and its major tributary confluences (Figure 5). These are not abstract node IDs — they are real places where named rivers meet. **Click any bar** to zoom the map and highlight that junction; marker rank and color match the bar.

The highest-betweenness confluence — **Tye River + James River** — carries roughly **3,749 square miles** of upstream drainage. A barrier or contamination event there does not affect one creek; it affects everything above it on the network.

6 Discussion

River networks are DAGs, and that matters. Every prior episode in this series analyzed undirected or bidirectional networks. Rivers cannot reroute. If a confluence is blocked, upstream tributaries have no alternate path.

The Strahler hierarchy compresses complexity. First-order reaches collectively account for most stream length, yet management attention historically focused on navigable main stems. The network view suggests headwater protection yields the largest aggregate gain per dollar.

Comparison to prior episodes. Airport and internet networks show scale-free degree distributions with genuine hubs. Watersheds show a binary-merge tree: most confluences have in-degree 2, out-degree 1, and betweenness concentrates on the main stem rather than on highly connected hubs.

7 Conclusion

The James River watershed is a directed tree of approximately 9,263 miles of stream length draining 6,753 square miles of Virginia. As a network it is structurally simple — binary merges,

single outlets — and structurally unforgiving: no alternate paths, and named high-betweenness confluences are true chokepoints.

Every drop goes somewhere. The map shows where.

Hidden Maps is a series on the networks embedded in everyday infrastructure. Previous episodes: [Ep. 0 — Primer](#), [Ep. 1 — Flights](#), [Ep. 2 — The Grid](#), [Ep. 3 — Streets](#), [Ep. 4 — Internet](#).

References

Horton, Robert E. 1945. “Erosional Development of Streams and Their Drainage Basins; Hydrophysical Approach to Quantitative Morphology.” *Geological Society of America Bulletin* 56 (3): 275–370.

Strahler, Arthur N. 1952. “Hypsometric (Area-Altitude) Analysis of Erosional Topography.” *Geological Society of America Bulletin* 63 (11): 1117–42.