


18: INTRO TO GENERALIZED LINEAR MODELS

Prof Amanda Luby

Let's start with our dear old penguins friends. The full dataset contains information about three different species of penguins. Rather than understanding the relationship between `body_mass` and `flipper_length`, we might instead be interested in how `body_mass` is related to species. In this case, we'll treat species as either Gentoo or Not Gentoo



gentoo on average have higher body mass


Data: $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$
 x_i : body mass (assumed constant)

$y_i: \{0,1\}$ random variable


On first glance, it looks like we could go ahead and fit a linear regression model for this problem:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.7006	0.0799	-21.2771	0
body_mass_g	0.0005	0.0000	26.2408	0

residual plots:



QQ plot of $\hat{\epsilon}_i$



Let's list some reasons why this approach is not ideal:

1. Residuals are perfectly correlated w/ predictions $\rightarrow v(\epsilon_i) \neq \sigma^2 I$

2. How do we assess the equal variance assumption?

3. $\epsilon_i \sim N(0, \sigma^2)$

$Y_i \sim N(X\beta, \sigma^2)$

\uparrow

Y_i 's are $\{0, 1\}$

What distribution does Y_i have? A better approach would be to start there.

$Y_i \sim \text{Bernoulli}(p_i)$ \leftarrow rather than modelling Y_i , we model p_i

1. $p_i = \beta_0 + \beta_1 x_i$

- end up w/

$p_i < 0$ or > 1

- "diminishing returns" - changes in x matter more if we're close to $1/2$ than if we're far away

2. $\log(p_i) = \beta_0 + \beta_1 x_i$

$p_i = e^{\beta_0} e^{\beta_1 x_i}$

- only bounded in 1 direction

3. $\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 x_i$

$p_i \in (0, 1)$

- not guaranteed to be wrong before we start!

1 Logistic Regression

Logistic Regression Model

$$\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 x_i, \quad Y_i \sim \text{Bernoulli}(p_i)$$

$$f(x) = \log\left(\frac{x}{1-x}\right) = \text{"logit"}$$

Solving for p , this gives:

$$\log\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 x_i$$

$$\frac{p}{1-p} = e^{\beta_0 + \beta_1 x_i}$$


$$\frac{1}{p} - 1 = \frac{1-p}{p} = \frac{1}{e^{\beta_0 + \beta_1 x_i}}$$

$$\frac{1}{p} = \frac{1}{e^{\beta_0 + \beta_1 x_i}} + 1 = \frac{1}{e^{\beta_0 + \beta_1 x_i}} + \frac{e^{\beta_0 + \beta_1 x_i}}{e^{\beta_0 + \beta_1 x_i}}$$

$$p = \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}} \leftarrow \text{logistic function}$$

$$f(x) = \frac{e^x}{1 + e^x}$$

$$= \frac{1}{1 + e^{-(\beta_0 + \beta_1 x_i)}}$$



MLE's for the linear model relied on the normal PDF for y

1.1 Maximum Likelihood Estimation

Now that we have the structure of the model, we have to think about how to estimate the β 's. Recall that the likelihood function for a n Bernoulli random variables is:

$$l(p) = \sum [y_i \ln p + (1 - y_i) \ln(1 - p)]$$

But, since we now have an X variable, $p = p(x_i)$

$$p = \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}}$$

$$\log\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 x_i$$

$$l(\beta) = \sum y_i \ln p + \ln(1-p) - y_i \ln(1-p)$$

$$= \sum y_i \ln \frac{p}{1-p} + \ln(1-p)$$

$$= \sum y_i (\beta_0 + \beta_1 x_i) + \ln\left(\frac{1}{1 + e^{\beta_0 + \beta_1 x_i}}\right)$$

$$= \sum y_i (\beta_0 + \beta_1 x_i) + (-1) \ln(1 + e^{\beta_0 + \beta_1 x_i})$$

$$1 - p(x_i) = \frac{1}{1 + e^{\beta_0 + \beta_1 x_i}}$$

To find MLE's,

$$\frac{\partial l}{\partial \beta_0}, \frac{\partial l}{\partial \beta_1}, \text{ Set equal to zero and solve}$$

→ In general, no closed form solution is

can be solved numerically w/ Newton-Raphson

Sampling distribution of logistic regression coefficients

$$\hat{\beta}_j \sim N\left(\beta_j, \frac{1}{I_n(\beta_j)}\right) \leftarrow \text{since MLE's are approximately normal}$$

"generalized"

```
gentoo_mod = glm(gentoo ~ body_mass_g,
                  data = penguins,
                  family = "binomial")
summary(gentoo_mod)
```

tells R the distribution we're assuming for Y_i 's

Call:

```
glm(formula = gentoo ~ body_mass_g, family = "binomial", data = penguins)
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.842e+01	3.609e+00	-7.873	3.46e-15 ***
body_mass_g	6.371e-03	8.131e-04	7.835	4.69e-15 ***

$H_0: \beta_j = 0$

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 446.80 on 341 degrees of freedom
Residual deviance: 117.85 on 340 degrees of freedom
(2 observations deleted due to missingness)
AIC: 121.85

$\sum (y_i - \bar{y})^2$
 $\sum (y_i - \hat{y}_i)^2$

Number of Fisher Scoring iterations: 7

1.2 Interpretation of coefficients

2 Generalized Linear Models

We've now seen two different settings for regression. If X is a vector of predictors and $Y \in \mathbb{R}$, we have assumed a linear model:

$$Y \sim N(X\beta, \sigma^2)$$

and if $Y \in \{0, 1\}$, we assumed a logistic model:

$$Y_i \sim \text{Bernoulli}(p_i)$$

$$\log\left(\frac{p_i}{1-p_i}\right) = X\beta$$

In both settings, we are assuming that a transformation of the conditional expectation is a linear function of X :

Linear
 $E[Y_i | X_i] = \mu_i = X\beta$ transformation: identity
 $g(x) = x$

Logistic
 Binomial RV's: $E(Y) = \mu = p$
 $\log\left(\frac{E(Y_i | X_i)}{1 - E(Y_i | X_i)}\right) = X\beta$ transformation: logit
 $g(x) = \log\left(\frac{x}{1-x}\right)$

Recall that exponential families of distributions can be written as

$$f(x; \theta) = h(x)g(\theta) \exp(T(x)\eta(\theta))$$

$T(x)$: sufficient statistic

$\eta(\theta)$: "natural parameter"

Normal
 $T(x) = \sum Y_i$
 $\eta(\theta) = \mu$ (assuming σ^2 known)
 = identity function

Bernoulli
 $T(x) = \sum Y_i$
 $\eta(\theta) = \log\left(\frac{p}{1-p}\right)$
 = logit function

Can set up a GLM for any exponential family,
 where $\eta(\theta)$ tells us what $g(E(Y_i | X_i))$ should
 be.