

18: The Normal Distribution

Stat 120 | Spring 2026

Prof Amanda Luby

Suppose weights of newborn babies in one community are normally distributed with a mean of 7.7 pounds and a standard deviation of 1.25 pounds

1. Use the 95% rule to sketch a graph of this normal density curve. Include a scale with at least three values on the horizontal axis.

2. Suppose I wanted to know the percent of newborns weighing less than 4.5 pounds.

- (a) Use StatKey to answer this question
- (b) Use the `pnorm` function to answer this question.

Answer: _____

3. Suppose I wanted to know the 15th percentile of the distribution.

- (a) Use StatKey to answer this question
- (b) Use the `qnorm` function to answer this question.

Answer: _____

4. APM Research Lab ran a [survey of likely Minnesota voters](#) between Sept 16-18 of 2024. Here's an excerpt from their methodology report:

The margin for error, according to standards customarily used by statisticians, is no more than ± 3.5 percentage points. This means that there is a 95% probability that the “true” figure would fall within that range if all voters were surveyed. The margin for error is higher for any subgroup, such as a gender or age grouping.

They reported 48.4% in support of the Harris/Walz ticket, and 43.3% in support of the Trump/Vance ticket.

(a) What is the standard error for each of these estimates?

(b) Build a 95% confidence interval for one of these proportions

(dc Use the normal distribution to build a **99%** confidence interval for one of these proportions

The report also says:

Overall margin of error = 3.5 percentage points (and therefore approximately 7 percentage points for the difference between two data points)

(e) Use this information to build a 90% confidence interval for the *difference in proportions* between the Harris/Walz supporters and the Trump/Vance supporters.

5. True or False?

- (a) The central limit theorem says that the population distribution is approximately normal if the population is big enough.
- (b) The central limit theorem says that the sampling distribution for the mean of a random sample is approximately normal if the sample is big enough.
- (c) The central limit theorem says that the sampling distribution for the mean is only normal if the population distribution is normal.
- (d) The central limit theorem says that any statistic is normally distributed if the sample size is big enough.
- (e) The central limit theorem applies to proportions and means

5. Using `pnorm` or `qnorm`, find the following quantities.

- (a) $P(Z < .25)$
- (b) $P(Z > 1.5)$
- (c) $P(-.5 < Z < 1.5)$
- (d) z^* where $P(Z < z^*) = .33$
- (e) z^* where $P(Z > z^*) = .33$
- (f) $P(X < 10)$ where $X \sim N(15, 22)$
- (g) $P(X > 40)$ where $X \sim (15, 22)$

i When you're done...

Please make sure that 1 person submits your group answers to gradescope, and includes all folks on the submission