

NOTES 22: INFERENCE FOR >2 PROPORTIONS

Stat 120 | Spring 2026

Prof Amanda Luby

What happens when we have *more* than two categories within a variable? (so response is not Yes/No or True/False)

Example: Some people think that children who are the older ones in their class at school naturally perform better in sports and that these children then get more coaching and encouragement. Could that make a difference in who makes it to the professional level in sports? Below is the birth month of 1478 major league players born since 1975, along with the national birth percentage across the same years. In the US, the cutoff for starting kindergarten is typically Sept 1, meaning children born in Sept, Oct, Nov are usually the oldest in their class, while Jun, Jul, Aug are usually the youngest.

Possible research questions:

- Are MLB players more likely to be born in Sept, Oct, Nov than the general US population?
- Are MLB players more likely to be born in Sept, Oct, Nov than Jun, July, Aug?
- Is the proportion of MLB players born in Aug lower than the proportion born in Sept?
- Does the birth month of MLB players follow the same distribution as the general US population?

Month	1	2	3	4	5	6	7	8	9	10	11	12
Obs	137	121	116	121	126	114	102	165	134	115	105	122
Birth %	8%	7%	8%	8%	8%	8%	9%	9%	9%	9%	8%	9%
Exp	118.2	103.4	118.2	118.2	118.2	118.2	133.0	133.0	133.0	133.0	118.2	133.0

How did I find the Expected Count?

Let's walk through an appropriate hypothesis test for this data

(1) Hypotheses

(2) Conditions/Assumptions


- Independence
- Expected count ≥ 5
- Counts

(3) Test statistic:

$$\chi^2 = \sum_{\text{all cells}} \frac{(Obs - Exp)^2}{Exp}$$

(4) p-value:

The χ^2 distribution looks like:



For χ^2 tests, we **always** calculate the p value as

$$P(\chi^2_{n-1} > 26.48) = \text{pchisq}(26.48, df = 11, \text{lower.tail} = \text{FALSE}) = .0055$$

This is a χ^2 **Goodness of Fit** test.

1 Two Categorical Variables

When we have two categorical variables, each that could have 2 or more levels, the only thing that changes is how we compute the *expected* counts within each cell.

Example: We are assessing the efficacy of three different drugs. If the type of drug does *not* matter, what is the probability of the condition improving for each drug? How does that translate to an expected count?

	Condition Improved	Did Not Improve	
Drug A			70
Drug B			100
Drug C			150
	180	140	320

(1) Hypotheses

(2) Conditions/Assumptions

- Independent observations
- Expected Counts ≥ 5
- Counts

(3) Test Statistic

(4) P-value

This is a χ^2 **Test for Association**

2 Group Problems

- Below is a sample of sauces. Does Taco Bell distribute sauce packets with equal probability?



- In order to assess whether habitat conditions are related to the sunlight choices a lizard makes for resting, Western fence lizard (*Sceloporus occidentalis*) were observed across three different micro-habitats. The results are shown below:

site	sunlight			Total
	sun	partial	shade	
desert	16	32	71	119
mountain	56	36	15	107
valley	42	40	24	106
Total	114	108	110	332

- If the site and the amount of sunlight are independent, what proportion of lizards (total) would be expected in the sun? In partial sun? In shade?
- Given the proportions of each sunlight condition, how many lizards *of each type* would you expect to see in the sun? in the partial sun? in the shade?

	sun	partial	shade	Total
desert				119
mountain				107

	sun	partial	shade	Total
valley				106
Total	114	108	110	332

- c. Compare the observed (original data) and expected (part b.) tables. From a first glance, does it seem as though the habitat and choice of sunlight may be associated?
- d. Perform a χ^2 test to formally test for an association between habitat and choice of sunlight. Write out the appropriate hypotheses, check conditions, compute the test statistic, find the p-value, and interpret your result in context.