23: Overdispersion and Quasibinomial Logistic Regression

Stat 230 - Spring 2026

```
library(patchwork)
library(ggformula)
```

1 Data overview

Orobanche is a genus of parasitic plants without chlorophyll that grow on the roots of flowering plants. In an experiment, seeds from two varieties, Orobanche aegyptiaca 75 and Orobanche aegyptiaca 73, where brushed with extract from either a cucumber root or a bean root. Researchers recorded the number of seeds that eventually germinated.

Run the code chunk below to load the data:

```
orobanche <- read.csv("https://aluby.github.io/stat230/data/orabanche.csv")
```

The variables in the data set are:

- y: count of seeds germinated
- n: number of seeds
- `variety`: either oa75 or oa73
- `root`: either cucumber or bean

2 Model from last time

Last class we fit the following model:

$$\text{logit}(\pi_i) = \beta_0 + \beta_1 \text{variety} + \beta_2 \text{root} + \beta_3 \text{variety} \cdot \text{root}$$

```r
oro_glm <- glm(y/n ~ variety + root + variety:root,
               weights = n,
               data = orobanche,
               family = "binomial")

summary(oro_glm)
```

```
Call:
glm(formula = y/n ~ variety + root + variety:root, family = "binomial",
    data = orobanche, weights = n)

Coefficients:
                         Estimate Std. Error z value Pr(>|z|)
(Intercept)               -0.4122     0.1842  -2.238   0.0252 *
varietyoa75               -0.1459     0.2232  -0.654   0.5132
rootcucumber               0.5401     0.2498   2.162   0.0306 *
varietyoa75:rootcucumber   0.7781     0.3064   2.539   0.0111 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

    Null deviance: 98.719  on 20  degrees of freedom
Residual deviance: 33.278  on 17  degrees of freedom
AIC: 117.87

Number of Fisher Scoring iterations: 4
```

1. Perform an overall goodness of fit test. What do you conclude?

2. We should also check the *validity* of the model, specifically whether linearity is satsified and if there are any outliers. The plots below show the pearson residual plots. Do you see any evidence of nonlinearity or extreme outliers?

```r
aug_interaction <- broom::augment(oro_glm, type.residuals = "pearson", type.predict =


p1 <- gf_point(.resid ~ .fitted, data = aug_interaction, xlab = "Linear predictor", y
```

```
  gf_hline(yintercept = 0, linetype = 2)

p2 <- gf_point(.resid ~ variety, data = aug_interaction, xlab = "variety", ylab = "Pe
  gf_hline(yintercept = 0, linetype = 2)


p3 <- gf_point(.resid ~ root, data = aug_interaction, xlab = "root", ylab = "Pearson
  gf_hline(yintercept = 0, linetype = 2)

p1 + p2 + p3
```



3 Checking and correcting for overdispersion

1. Perform the ad hoc test of overdispersion. What do you conclude?

```
x2 <- sum(residuals(oro_glm, type = "pearson")^2)
df <- ___
1 - pchisq(___, ____)
```

2. What is the ad-hoc estimate of $\widehat{\psi}$? Use this to calculate $\text{SE}_{quasil}(\hat{\beta}_2)$. (*Note:* do this by hand, without fitting another model)

3. Construct a 95% confidence interval for β_2 using the adjusted standard error. (*Note:* do this by hand, without fitting another model)

4. Now, confirm your answer by fitting a `glm()` model with `family = "quasibinomial"` and finding the confidence interval

```
oro_qglm <- glm(___ ~ ___,
                weights = ___,
                data = ___,
                family = ___)
```

6. What does your result from Q4 imply about the necessity of the interaction term? Is this the same or different than you conclusion about the interaction term using the binomial model?

---

4 Function quick reference

The following table summarizes the functions discussed above:

Function	Purpose
`glm()`	Fit generalized linear models, including quasibinomial logistic regression models.