
Day 09: Intro to Multiple Linear Regression

Note: the first section recaps what was in the lecture slides. You should fill in the second section in your groups using the R activity.

Capital Bikeshare is a bike sharing service in the Washington, D.C. area. To best serve its registered members, the company must understand the demand for its service. We have data on the number of rides taken on a random sample of n days, along with:

- Weekend/holiday/etc.
- Day of the measurement
- Weather information

We want to predict the number of `rides` based on `temp_actual` and `windspeed`. Both are correlated with `rides`:



Neither predictor alone explains a large amount of the variation in ridership, and fitting two separate models would give us two different predictions for the same day, with no way to combine them into one. Instead, we want to build a single model that summarizes the relationship between `temp_actual` and `rides` *after accounting for windspeed* and vice-versa.

Multiple Linear Regression (MLR)

$$Y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + \varepsilon_i, \quad \varepsilon_i \overset{\text{iid}}{\sim} N(0, \sigma^2)$$

This model requires:

1. $\mu\{Y \mid x_1, x_2, \ldots, x_p\}$ is a _____ function of the predictors
2. The error terms are _____
3. The error terms follow a Normal distribution with mean 0 and variance σ^2

Fitting the model: least squares estimation chooses the coefficients to minimize

$$\text{SSE} = \sum_{i=1}^{n} \left(Y_i - \hat{Y}_i\right)^2, \qquad \text{then} \quad \hat{\sigma} = \sqrt{\frac{\text{SSE}}{n-(p+1)}}$$

```
bikes_mlr <- lm(rides ~ temp_actual + windspeed, data = bikes)
coef(bikes_mlr)
```

```
(Intercept) temp_actual   windspeed
-1525.53822    87.01789   -38.62817
```

$\hat{\beta}_0$ is the expected value of the response when all predictors are 0.

Example: the expected number of rides is about -1525.5 when temperature = 0 and windspeed = 0 – not meaningful in this dataset!


$\hat{\beta}_j$ is the expected change in the response for a 1-unit increase in x_j, when all other predictors are held constant.

Example:

- We expect 87.02 *additional* rides for every 1 degree increase in actual temperature, holding windspeed constant.
- We expect 38.63 *fewer* rides for every 1 MPH increase in windspeed, holding temperature constant.

Geometric interpretation: the MLR ``line'' is a plane

With two predictors, the fitted model is no longer a line but a **plane** through 3-dimensional space:



Takeaways

- EDA for more than one predictor variable
- Write the theoretical MLR model
- Write the fitted MLR model
- Interpret the MLR coefficients

MLR Activity

To complete this handout, open the R activity (linked on moodle)

Data

- In an effort to understand the important aspects of a satisfactory supervisor, clerical employees at a large financial organization were asked to rate their immediate supervisor.
- The survey questions were designed to measure overall performance of the supervisor, as well as six additional characteristics.
- Employees were asked to rate the following statements on a scale from 0 to 100 (0 = "completely disagree", 100 = "completely agree")

Variable	Description
`rating`	Overall rating of supervisor performance
`complaints`	Score for "Your supervisor handles employee complaints appropriately."
`privileges`	Score for "Your supervisor allows special privileges."
`learn`	Score for "Your supervisor provides opportunities to learn new things."
`raises`	Score for "Your supervisor bases raises on performance."
`critical`	Score for "Your supervisor is too critical of poor performance."
`advance`	Score for "I am not satisfied with the rate I am advancing in the company?"

Q1: Before looking at the data, how do you think each predictor variable will be related to `rating`? What do you think will have the strongest relationship? The weakest?

Q2: Use the pairs plot to answer the following

- Which predictor variables appear to be most closely related to `overall`?

- Which predictor variables appear to be most closely related to *each other*?

Q3: SLR. Choose what you think will be the best *single* predictor and fit an SLR model. Interpret the intercept and slope coefficient and report the R^2 value.

Q4: MLR. Use all predictors and fit an MLR model. Interpret the intercept and two of the slope coefficients and report the R^2 value.

Q5: Use the `predict` function to predict the overall manager score for a manager with scores of 50% for all predictors

Q7: What if the manager from the previous question decides to try to improve her `critical` score by 10 points (so 40% instead of 50%) and her `complaints` score by 10 points (so 60% instead of 50%). How does that change the prediction?

Function quick reference

Function	Purpose
`lm(formula, data)`	Fit a MLR model
`predict(model, newdata, interval, level, se.fit)`	Make predictions and calculate intervals
`data |> select(vars) |> ggpairs()`	Make pairs plot for selected variables