

# Group Work 07

```
library(bayesrules) # R package for our textbook
library(tidyverse) # Collection of packages for tidying and plotting data
library(janitor) # Helper functions like tabyl
library(rstan) # for fitting models
library(rstanarm) # for fitting standard regression models
library(broom.mixed) # for tidy() function
library(bayesplot) # helpful plotting functions
library(tidybayes) # helpful for wrangling Bayesian model output
```

## Note

I've given you most of the R code you need for today to help you get through the questions more quickly and focus on the concepts. Make sure you understand what all the code is doing, enough so that you can replicate it on your own in the future.

In code chunks where you need to fill in the blank, remember to remove `#! eval: false` to get your document to render correctly

The data for today is an expansion of the `weather_perth` data that you worked with last week, called `weather_WU`. We'll continue to model the daily 3pm temperature, starting with using the 9am temperature and later adding more predictors.

```
data("weather_WU")
```

We'll use the following Bayesian regression model:

$$Y_i | \beta_0, \beta_1, \sigma \sim N(\mu_i, \sigma^2)$$

$$\mu_i = \beta_0 + \beta_1 X_i$$

$$\beta_{0c} \sim N(30, 10^2)$$

$$\beta_1 \sim N(0, 2.5^2)$$

$$\sigma \sim \text{Exp}(0.1)$$



Tip

Your R code from last week may be helpful here!

## 1 Simulate the posterior

- (a) Since you worked with (essentially) this data last week, let's begin by simulating the posterior.

```
# Simulate the posterior
```

```
model_1 <- stan_glm(...)
```

- (b) To ensure that we can “trust” the MCMC posterior simulation results, create trace plots and density plots of the multiple chains.

```
# your code here
```

- (c) How wrong is the model? To answer this question, perform a posterior predictive check using `pp_check()`. Does the underlying structure of our Normal regression model appear to be reasonable?

```
## 50 simulated posterior model lines
```

```
pp_check(model_1)
```

## 2 Fix the model

The `pp_check()` revealed a structural issue with our model. (a) Identify what you think the issue is. Support your guess by constructing a relevant visualization of the raw weather data. (b) Build and simulate a new model which fixes the issue. Store this as `model_2`. NOTE: Do not take `temp9am` out of the model and use the same priors. (c) Perform a `pp_check()` to confirm that `model_2` is structurally better than `model_1`.

Feel free to chat with me if you're unsure what to do!

```
# your R code here
```

### 3 Posterior predictions

We can assess the quality of predictions using both visual and numerical summaries.

- (a) Plot the posterior predictive models for all data points. Leave code comments for any arguments you're unsure about

```
# First obtain posterior predictive models
# for each data point in weather:
set.seed(84735)
predictions_2 <- posterior_predict(model_2, newdata = weather_WU)

# Plot the posterior predictive intervals
ppc_intervals_grouped(
  weather$temp3pm, yrep = predictions_2,
  x = weather_WU$temp9am, group = weather_WU$location,
  prob = 0.5, prob_outer = 0.95,
  facet_args = list(scales = "fixed")) +
  labs(x = "temp9am", y = "temp3pm")
```

- (b) Evaluate the accuracy of the posterior predictions. Interpret each of the four reported numbers and make sure you can match them to the plot in part a.

```
set.seed(84735)
prediction_summary(model = model_2, data = weather)
```

### 4 Prediction accuracy and Cross Validation


- (a) In terms of MAE, how accurate is our model at predicting the `temp3pm` values in our original sample?
- (b) In terms of MAE, how accurate is our model at predicting `temp3pm` values on NEW days?
- (c) When communicating the posterior prediction accuracy for our model, which number should you report?

## 5 Model comparison

Using a sample of data on just Wollongong, let's model `temp3pm` by `day_of_year`, 1 through 365:

```
# Obtain the sample data
wollongong <- weather_WU |>
  filter(location == "Wollongong")

# Plot temp3pm vs day_of_year
ggplot(wollongong, aes(x = day_of_year, y = temp3pm)) +
  geom_point()
```



Consider 3 polynomial models of this relationship:

- `mod1` has no predictors:  $\mu = \beta_0$
- `mod2` assumes a quadratic relationship and has 2 predictors:  $\mu = \beta_0 + \beta_1 x + \beta_2 x^2$
- `mod3` assumes a 12th order polynomial relationship:  $\mu = \beta_0 + \beta_1 x + \beta_2 x^2 + \dots + \beta_{12} x^{12}$

Simulate the posteriors for these 3 models:

```
mod_1 <- stan_glm(
  temp3pm ~ 1, data = wollongong,
  family = gaussian,
  prior_intercept = normal(30, 10),
  chains = 4, iter = 5000*2, seed = 84735, refresh = 0)
```

```

mod_2 <- stan_glm(
  temp3pm ~ poly(day_of_year, 2), data = wollongong,
  family = gaussian,
  prior_intercept = normal(30, 10),
  chains = 4, iter = 5000*2, seed = 84735, refresh = 0)

mod_3 <- stan_glm(
  temp3pm ~ poly(day_of_year, 12), data = wollongong,
  family = gaussian,
  prior_intercept = normal(30, 10),
  chains = 4, iter = 5000*2, seed = 84735, refresh = 0)

```

Plot 200 posterior lines:

```

wollongong %>%
  add_fitted_draws(mod_1, n = 200) %>%
  ggplot(aes(x = day_of_year, y = temp3pm)) +
    geom_line(aes(y = .value, group = .draw), alpha = 0.15) +
    geom_point(data = wollongong)

wollongong %>%
  add_fitted_draws(mod_2, n = 200) %>%
  ggplot(aes(x = day_of_year, y = temp3pm)) +
    geom_line(aes(y = .value, group = .draw), alpha = 0.15) +
    geom_point(data = wollongong)

wollongong %>%
  add_fitted_draws(mod_3, n = 200) %>%
  ggplot(aes(x = day_of_year, y = temp3pm)) +
    geom_line(aes(y = .value, group = .draw), alpha = 0.15) +
    geom_point(data = wollongong)

```

## 6 Which is the best model?

- Examine the `pp_check()` for each model. Which model is the most wrong? The least?
- Which model do you think will have the best “in-sample” prediction error, that is the best predictions of the data points in our `wollongong` sample? Test your guess with a `prediction_summary()` of each model.
- Which model do you think will have the best cross-validated prediction error, that is the best predictions of new data points? Test your guess with a `prediction_summary_cv()`\$cv of each model.