

Homework 1: Due 24/03/2025 before 23.55

Lecturer: Aurelien Lucchi

Problem 1 (Convex Sets, 10 points):

a. Consider the set

(3 Pts)

$$S = \{(x, y) \in \mathbb{R}^2 \mid x \geq 0, x^2 + y^2 \leq 4x\}.$$

Show whether S is convex or not. (Hint: notice that $x^2 + y^2 \leq 4x$ can be rewritten as a circle shifted from the origin.)

b. Define

(4 Pts)

$$T = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid (x_1, x_2) \in \text{conv}\{(0, 0), (2, 0), (0, 2)\}, x_3 \geq x_1 + x_2\}.$$

Prove that T is convex.

c. Let

(4 Pts)

$$U := \{(x, y) \in \mathbb{R}^2 : \min\{x^2 + y^2, (x - 1)^2 + y^2\} \leq 1\}.$$

Show that U is the union of two disks of radius 1, centered at $(0, 0)$ and $(1, 0)$. Check whether this union is convex or not.

Problem 2 (Function Regularity and the PL Inequality, 10 points):a. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

(4 Pts)

$$f(x) = \frac{x^4}{4} - x^2.$$

i) (*Smoothness*) Is f L -smooth on all of \mathbb{R} ? If yes, determine such an L . If no, explain which calculations imply that no global L can exist.

ii) (*PL Inequality*) The Polyak-Łojasiewicz (PL) condition says there is a $\mu > 0$ such that

$$\frac{1}{2} |f'(x)|^2 \geq \mu (f(x) - f^*) \quad \text{for all } x \in \mathbb{R},$$

where $f^* = \min_{x \in \mathbb{R}} f(x)$. Find f^* by direct calculation (use $f'(x) = 0$). Then, check whether you can identify a positive μ that satisfies the PL condition for every $x \in \mathbb{R}$. Make sure you include explicit computations of both sides to conclude whether such a μ exists or not.

b. (*Example of PL without Strong Convexity*)

(3 Pts)

Define $g : \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$g(\mathbf{x}) = \sum_{i=1}^n \max\{0, x_i\}^2.$$

(i) Show that g is *not* strongly convex: for example, any vector \mathbf{x} with all $x_i \leq 0$ has $g(\mathbf{x}) = 0$, so there is a whole “flat” (plateau) region where g is constant. This violates the strict curvature needed for strong convexity.

(ii) Verify that g *does* satisfy the PL inequality

$$\frac{1}{2} \|\nabla g(\mathbf{x})\|^2 \geq \mu (g(\mathbf{x}) - g^*),$$

for some $\mu > 0$, where $g^* = \min_{\mathbf{x}} g(\mathbf{x}) = 0$.

c. (*Linear Convergence via PL*)

(3 Pts)

Suppose that $g(\mathbf{x})$ in part (b) is also L -smooth (or pick another suitable g that is both PL and L -smooth). With the standard gradient descent update

$$\mathbf{x}^{k+1} = \mathbf{x}^k - \alpha \nabla g(\mathbf{x}^k),$$

demonstrate, by explicit calculation, that for a proper step size α there is a linear rate of convergence:

$$g(\mathbf{x}^k) - g^* \leq \rho^k (g(\mathbf{x}^0) - g^*), \quad \text{for some } \rho \in (0, 1).$$

Make sure to use both L -smoothness and the PL condition in your derivation.

Problem 3 (Convergence of Gradient Descent, 10 points):

In this problem, we assume that $f: \mathbb{R}^d \rightarrow \mathbb{R}$ is differentiable, L -smooth and μ -strongly convex. Let us define the Bregman divergence associated with the function f as follows

$$D_f(\mathbf{x}, \mathbf{y}) := f(\mathbf{x}) - f(\mathbf{y}) - \langle \nabla f(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle. \quad (1)$$

Bregman divergence is useful in deriving convergence guarantees of gradient descent with tighter dependencies on the problem-specific constants (e.g., L or μ). In particular, following (6) from Lecture 2: Gradient Descent the smoothness, the L -smoothness and μ -strong convexity imply

$$\frac{\mu}{2} \|\mathbf{x} - \mathbf{y}\|^2 \leq D_f(\mathbf{x}, \mathbf{y}) \leq \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|^2. \quad (2)$$

Moreover, a convex and L -smooth function satisfies

$$\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\|^2 \leq 2LD_f(\mathbf{x}, \mathbf{y}). \quad (3)$$

We will prove the following convergence rate for the gradient descent which is equivalent to Theorem 1 from Lecture 2: Gradient Descent using Bregman divergence

$$\|\mathbf{x}_k - \mathbf{x}^*\|^2 \leq (1 - \eta\mu)^k \|\mathbf{x}_0 - \mathbf{x}^*\|^2, \quad (4)$$

where the stepsize $\eta \leq \frac{1}{L}$.

a. Using (2) and definition of Bregman divergence (1) demonstrate that the following inequality holds (2 Pts)

$$\frac{\mu}{2} \|\mathbf{x} - \mathbf{y}\|^2 + D_f(\mathbf{x}, \mathbf{y}) \leq \langle \nabla f(\mathbf{x}) - \nabla f(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle.$$

b. Demonstrate that the following equality holds (1 Pt)

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2 = \|\mathbf{x}_k - \mathbf{x}^*\|^2 - 2\eta \langle \nabla f(\mathbf{x}_k) - \nabla f(\mathbf{x}^*), \mathbf{x}_k - \mathbf{x}^* \rangle + \eta^2 \|\nabla f(\mathbf{x}_k) - \nabla f(\mathbf{x}^*)\|^2.$$

c. Using the result of 1. demonstrate that (2 Pts)

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2 \leq (1 - \eta\mu) \|\mathbf{x}_k - \mathbf{x}^*\|^2 - 2\eta D_f(\mathbf{x}_k, \mathbf{x}^*) + \eta^2 \|\nabla f(\mathbf{x}_k) - \nabla f(\mathbf{x}^*)\|^2.$$

d. Using (3) demonstrate that (2 Pts)

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2 \leq (1 - \eta\mu) \|\mathbf{x}_k - \mathbf{x}^*\|^2 - 2\eta(1 - \eta L) D_f(\mathbf{x}_k, \mathbf{x}^*).$$

e. Using the stepsize restriction derive the recursion (2 Pts)

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2 \leq (1 - \eta\mu) \|\mathbf{x}_k - \mathbf{x}^*\|^2.$$

f. Unrolling the recursion obtain the final convergence result (1 Pt)

$$\|\mathbf{x}_k - \mathbf{x}^*\|^2 \leq (1 - \eta\mu)^k \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$