

Exercise 3: Complexity Theory

Lecturer: Aurelien Lucchi

Teaching Assistants: Enea Monzio Compagnoni, Jim Zhao

Problem 1 (Barron norm of Gaussian measure):

Consider the density function of a Gaussian random variable, $f(x) = (2\pi\sigma^2)^{d/2} \exp(-\frac{\|x\|^2}{2\sigma^2})$. We will calculate the complexity measure $\int \|\widehat{\nabla} f(\omega)\| d\omega$ that appears in Barron's theorem discussed in class.

a) Use the following facts: (Why are they true?)

- For f defined as above, we have $|\hat{f}(\omega)| = \exp(-2\pi^2\sigma^2\|\omega\|^2)$;
- (Cauchy-Schwartz inequality in expected value) $\mathbb{E}[\|X\|] \leq (\mathbb{E}[\|X\|^2])^{\frac{1}{2}}$ where X is a random vector.

Show that $\int \|\omega\| |\hat{f}(\omega)| d\omega \leq Z \left(\int \|\omega\|^2 Z^{-1} \hat{f}(\omega) d\omega \right)^{1/2}$ where $Z = (2\pi\sigma^2)^{-d/2}$.

b) Using the fact that $\int g(\omega) Z^{-1} \hat{f}(\omega) d\omega$ is the expectation of the function $g(\omega)$ with respect to the density $\mathcal{N}(0, \frac{1}{4\pi^2\sigma^2})$, show that $\int \|\omega\| |\hat{f}(\omega)| d\omega \leq \sqrt{\frac{d}{4\pi^2\sigma^2}} \cdot (2\pi\sigma^2)^{-d/2}$.

c) Since $\int \|\widehat{\nabla} f(\omega)\| d\omega = 2\pi \int \|\omega\| \cdot |\hat{f}(\omega)| d\omega$, what do you conclude about the complexity measure $\int \|\widehat{\nabla} f(\omega)\| d\omega$ when d is very large?

Problem 2 (Maurey's lemma):

Let $X = \mathbb{E}V$ be given, with V a random vector supported on a subset S of the event space, and let $V_1, \dots, V_k$ be i.i.d. realization of V .

a) Show that

$$\mathbb{E}_{V_i} \left\| X - \frac{1}{k} \sum_{i=1}^k V_i \right\|^2 = \frac{1}{k} \mathbb{E}_V \|V - X\|^2.$$

b) Show that

$$\frac{1}{k} \mathbb{E}_V \|V - X\|^2 \leq \frac{\sup_{U \in S} \|U\|^2}{k}.$$

Problem 3 (Number of affine pieces in a ReLU network):

We will prove the lemma that bounds the number of affine pieces in a ReLU network stated in the lecture notes. Denote by $\delta_A(f)$ be the (minimum) number of affine pieces of a piece-wisely linear continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$. Show that

- a) $\delta_A(\sum_k a_k f_k + b_k) \leq \sum_k \delta_A(f_k)$, for any finite sequence of piece-wisely linear continuous functions f_k , and for any real sequences a_k and b_k .
- b) Let $g_i: \mathbb{R} \rightarrow \mathbb{R}$ denote the output of some node in layer i in a ReLU network with L layers of widths $(m_1, \dots, m_L)$ as a function of the input. Using induction on i , show that the number of affine pieces $\delta_A(g_i)$ satisfies

$$\delta_A(g_i) \leq 2^i \prod_{j < i} m_j.$$

- c) Recall that $g_L: \mathbb{R} \rightarrow \mathbb{R}$ is a ReLU network with L layers of widths $(m_1, \dots, m_L)$ such that $m = \sum_{i=1}^L m_i$. The number of affine pieces in g_L satisfies

$$\delta_A(g_L) \leq \left(\frac{2m}{L} \right)^L.$$

- d) What value of L maximizes the upper bound $\left(\frac{2m}{L} \right)^L$? Although taking this value of L gives a large value of $\delta_A(g_L)$, give a reason why in practice we usually pick a value much smaller than this.