

NMF in LM

1 Introduction

Localisation microscopy (LM) is a conceptually simple and accessible technique providing superresolution fluorescent images.^{2,4,6,11} The structure of the sample is reconstructed by localising individual fluorophores with precision surpassing the classical resolution limit.¹³ LM make use of the fluorophore transition between bright (ON) and dark (OFF) states. Standard LM techniques (fPALM,² STORM¹⁵) control ON-OFF transition by photo-switching of the fluorophores as they require only one source to be ON within a diffraction limited area at a time. This is achieved by driving the sample into a state where only a small subset of the fluorophores are in ON state. The superresolution image is composed from the repetitive localisation of the different subsets of individual fluorophores. An optimal number of the ON sources in each acquisition frame must be experimentally estimated as a small density lead to a long acquisition time whereas a high density of the ON fluorophores results in overlapping sources that cannot be localised.¹⁶

Quantum dots (QD) are an order of magnitude brighter compared to the organic dyes used in the standard LM.^{5,14} Under a continuous excitation the QDs exhibit a stochastic blinking between ON and OFF states.^{7,12,17} An excellent photo-stability, low cytotoxicity and unique spectral properties¹⁴ make QDs very attractive for biological research. However, the stochastic blinking makes QDs impractical for the standard LM as the rate of ON-OFF transition is difficult to control. This results in highly overlapping sources that cannot be localised with standard LM techniques.

Some methods exploiting the blinking behaviour of the QD have been proposed. Maximum a posteriori fitting of the positions and the intensities of known individual point spread functions (PSFs) into a movie of blinking QD (QD data) has been proposed in.¹ Inde-

pendent component analysis (ICA) of the QD data was suggested in¹⁰ and a resolution improvement by a statistical analysis of the intensity fluctuation (SOFI) has been demonstrated.⁷

2 NMF

Non-negative matrix factorisation (NMF)⁸ is a natural model for QD data. It decomposes spatio-temporal data - a movie of the blinking QDs - into a spatial component - images of the individual sources - and a temporal component - their fluctuating intensities. Unlike ICA, NMF imposes a natural non-negativity constraints on the images of the individual sources (point spread function PSF) and their intensities. Moreover, we used a NMF algorithm⁹ that maximizes the likelihood of model for data corrupted with Poisson noise which makes it a method of choice compared to ICA.

NMF does not put any constraints on the shape or blinking behavior of the individual sources (apart from being non-negative). Therefore NMF can separate images overlapping fluorophores each having different shape and blinking behavior. This can, for example, arise in a 3D sample where sources can be in different focal depth. This results in variety of point spread function

3 Model comparison

NMF requires a prior knowledge about the number of sources (K) to be separated. Principal component analysis (PCA) can be used as a simple method for dimensionality estimation. However, for noisy data the estimation of K is difficult.

4 Out of focus PSF

5 Notes

Fitting of the multiple PSF into the STORM data is in.³

Supplementary materials

Spatial temporal data (movie) of blinking quantum dots can be regarded as a $N \times T$ data matrix $\mathbf{D}(x, t)$ where N is a number of pixels and T is a number of time frames in the movie. Each frame in the movie is transformed into

a column of the matrix \mathbf{D} by concatenating columns of the 2D image into a $N \times 1$ vector. Non-negative matrix factorisation (NMF) makes an approximative decomposition

$$\mathbf{D} \approx \mathbf{W}\mathbf{H}, \quad (1)$$

where the $N \times T$ matrix \mathbf{D} is expressed as a multiplication of the $N \times K$ matrix \mathbf{W} and $K \times T$ matrix \mathbf{H} subject to the non-negativity constraints on the entries $w_{xk} \geq 0$ and $h_{kt} \geq 0$. Each column \mathbf{w}_k of the matrix \mathbf{W} ($N \times 1$ vector) then represents a k th image of one source and each row \mathbf{h}_t^T of the matrix \mathbf{H} ($1 \times T$ vector) represents the time profile of the k th source blinking.

The NMF algorithm⁹ makes the decomposition such that the likelihood function of the model is maximised under assumption of the Poisson noise. The approximative factorisation(1) can then be written as

$$\mathbb{E}[\mathbf{D}] = \mathbf{W}\mathbf{H},$$

where $\mathbb{E}[\cdot]$ denotes expectation value of the noisy data.

References

- [1] Paul Harrington, Jonas Anderson, Bernd Rieger, Diane Lidke, and Keith A Lidke. Poster: A Bayesian Approach to Fluorescence Intermittency Based Localization Microscopy. *Supplement of Biophysical Journal*, 96:20–20, 2008. 1
- [2] Samuel T Hess, Thanu P K Girirajan, and Michael D Mason. Ultra-high resolution imaging by fluorescence photoactivation localization microscopy. *Biophysical journal*, 91(11):4258–72, December 2006. 1
- [3] Seamus J Holden, Stephan Uphoff, and Achillefs N Kapanidis. DAOSTORM: an algorithm for high- density super-resolution microscopy. *Nature methods*, 8(4):279–80, April 2011. 1
- [4] Bo Huang, Wenqin Wang, Mark Bates, and Xiaowei Zhuang. Three-dimensional super-resolution imaging by stochastic optical reconstruction microscopy. *Science (New York, N.Y.)*, 319(5864):810–3, February 2008. 1
- [5] Jyoti K Jaiswal and Sanford M Simon. Potentials and pitfalls of fluorescent quantum dots for biological imaging. *Trends in cell biology*, 14(9):497–504, September 2004. 1
- [6] Sara a Jones, Sang-Hee Shim, Jiang He, and Xiaowei Zhuang. Fast, three-dimensional super-resolution imaging of live cells. *Nature methods*, 8(6):499–505, June 2011. 1
- [7] M. Kuno, D. P. Fromm, H. F. Hamann, A. Gallagher, and D. J. Nesbitt. "On"/"off" fluorescence intermittency of single semiconductor quantum dots. *The Journal of Chemical Physics*, 115(2):1028, 2001. 1
- [8] D D Lee and H S Seung. Learning the parts of objects by non-negative matrix factorization. *Nature*, 401(6755):788–91, October 1999. 1
- [9] D.D. Lee and H.S. Seung. Algorithms for non-negative matrix factorization. *Advances in neural information processing systems*, 13, 2001. 1, 2
- [10] Keith a. Lidke, Bernd Rieger, Thomas M. Jovin, and Rainer Heintzmann. Superresolution by localization of quantum dots using blinking statistics. *Optics Express*, 13(18):7052, 2005. 1
- [11] S. Linde, R. Kasper, M. Heilemann, and M. Sauer. Photoswitching microscopy with standard fluorophores. *Applied Physics B*, 93(4):725–731, October 2008. 1
- [12] M Nirmal, B O Dabbousi, M G Bawendi, J J Macklin, J K Trautman, T D Harris, and L E Brus. Fluorescence intermittency in single cadmium selenide nanocrystals. *Nature*, 1996. 1
- [13] Raimund J Ober, Sripad Ram, and E Sally Ward. Localization accuracy in single-molecule microscopy. *Biophysical journal*, 86(2):1185–200, February 2004. 1
- [14] Ute Resch-Genger, Markus Grabolle, Sara Cavaliere-Jaricot, Roland Nitschke, and Thomas Nann. Quantum dots versus organic dyes as fluorescent labels. *Nature methods*, 5(9):763–75, September 2008. 1
- [15] M.J. Rust, Mark Bates, and Xiaowei Zhuang. Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). *Nature methods*, 3(10):793–796, 2006. 1
- [16] Alexander R Small. Theoretical limits on errors and acquisition rates in localizing switchable fluorophores. *Biophysical journal*, 96(2):L16–8, January 2009. 1
- [17] Fernando D. Stefani, Jacob P. Hoogenboom, and Eli Barkai. Beyond quantum jumps: Blinking nanoscale light emitters. *Physics Today*, 62(2):34, 2009. 1