


Figure 16 Evaluation of patients with a working diagnosis of MINOCA. ACh, acetylcholine; CMRI, cardiac magnetic resonance imaging; CT, computed tomography; CTPA, computed tomography pulmonary angiogram; ECG, electrocardiogram; IVUS, intravascular ultrasound; LV, left ventricular; LVeDP, left ventricular end-diastolic pressure; MINOCA, myocardial infarction with non-obstructive coronary arteries; NSTE-ACS, Non-ST elevation acute coronary syndrome; NTpro BNP, N terminal pro brain natriuretic peptide; OCT, optical coherence tomography; STEMI, ST-elevation myocardial infarction; UA, unstable angina. Patients presenting with STEMI present directly to catheter lab as per the current standard of care pathway **(1)**. In this context, when non-obstructive coronary arteries are identified then further assessment should be considered. When patients are subsequently admitted to the ward then investigations as shown in **(2)** should be considered. Patients presenting with NSTE-ACS or UA are often stabilized on the ward **(2)** prior to transfer to the cath lab **(1)**. In this context the order in which the investigations are carried out will vary depending on the location these patients are managed during first contact. MINOCA patients require follow-up review **(3)** and may require repeat assessment using echocardiography and magnetic resonance imaging, depending on the initial findings. [a]Options for adjunctive tests. Patients will not require all investigations but instead the appropriate tests should be selected based on their presentation and clinical course. [b]Examples of potential blood tests include: full blood count, renal profile, troponin, C-reactive protein, D-dimer, NT-pro BNP. [c]A CT scan of the brain should be considered if a cranial pathology (i.e. intracranial bleed) is suspected that might have resulted in ST elevation.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

If the underlying cause of MINOCA is not established using functional coronary angiography, then non-invasive imaging (i.e. echocardiography, CMR, CT) is recommended, as clinically appropriate. CMR is one of the key diagnostic tools to determine the underlying cause of MINOCA.[540–544] CMR can identify the underlying cause in up to 87% of patients with a working diagnosis of MINOCA and should be performed as soon as possible after presentation in these patients to maximize its diagnostic yield, ideally during the index admission.[545]

Diagnosis of the underlying cause of MINOCA will enable the appropriate treatment to be initiated based on the final diagnosis. Secondary prevention therapies should be considered for those with evidence of coronary atherosclerotic disease and to control risk factors. The management of takotsubo syndrome is not informed by any prospective RCTs, and treatment is largely supportive and empiric.[546,547] The treatment of patients with myocarditis has been covered by previous ESC documents.[548,549] Ischemia with non-obstructive coronary arteries (INOCA) has also been described in the context of CCS.[550,551] Additional information about MINOCA is provided in the Supplementary data online, including *Table S13*.

Recommendation Table 13 — Recommendations for myocardial infarction with non-obstructive coronary arteries

Recommendations	Class[a]	Level[b]
In patients with a working diagnosis of MINOCA, CMR imaging is recommended after invasive angiography if the final diagnosis is not clear.[544,545]	I	B
Management of MINOCA according to the final established underlying diagnosis is recommended, consistent with the appropriate disease-specific guidelines.[546,550,552]	I	B
In all patients with an initial working diagnosis of MINOCA, it is recommended to follow a diagnostic algorithm to determine the underlying final diagnosis.	I	C

CMR, cardiac magnetic resonance; MINOCA, myocardial infarction with non-obstructive coronary arteries.
[a]Class of recommendation.
[b]Level of evidence.

© ESC 2023

12. Special situations

12.1. Type 2 myocardial infarction and acute myocardial injury

Pathological processes other than atherothrombosis commonly underlie the presentation of patients with acute chest pain with troponin elevation. These include Type 2 MI and myocardial injury as defined in the fourth universal definition of MI.[1] Type 2 MI is an ischaemic myocardial injury in the context of a mismatch between oxygen supply and demand that is not related to acute coronary atherothrombosis. This may occur in the context of atherosclerosis and an oxygen supply/demand imbalance, with an oxygen supply/demand imbalance alone, secondary to vasospasm or coronary microvascular dysfunction, or secondary to non-atherosclerotic coronary dissection. These causes of Type 2 MI can be divided into those with underlying coronary (e.g. coronary embolus, dissection, spasm, microvascular dysfunction) or non-coronary mechanisms (supply demand mismatch due to hypoxia, hypotension, anaemia, tachycardia, bradycardia).[1] Type 2 MI is common and associated with a prognosis similar to Type 1 MI.[12]

Myocardial injury is characterized by myocyte necrosis and troponin elevation due to mechanisms other than myocardial ischaemia and can be acute (e.g. sepsis, myocarditis, takotsubo) or chronic (e.g. HF, cardiomyopathies, severe valve heart disease). Myocardial injury is increasingly appreciated in the era of hs-cTn assays, which are not specific for MI. In patients who have elevated hs-cTn values and do not have evidence of acute myocardial ischaemia, a diagnosis of myocardial injury can be made. It is important to recognize that this diagnosis can change if subsequent investigations indicate that the patient meets the criteria for MI.

Despite some common risk factors, the pathophysiology of Type 2 MI is different to that of Type 1 MI. Therefore, the natural history and appropriate management strategy of these two conditions also differs in some important respects. Type 2 and Type 1 MI require diagnostic distinction, which is best achieved by following an algorithmic approach.[1,553] Once patients with suspected Type 2 MI and myocardial injury have been stabilized and any precipitating illnesses have been treated, targeted echocardiography and/or coronary angiography (invasive or CCTA) can be used to identify contributory (and prognostically important) cardiac conditions and to guide appropriate long-term cardiovascular treatments.[12] Due to the lack of robust scientific evidence investigating management strategies and the wide range of precipitating causes, there are currently no specific recommended pharmacological interventions for patients with Type 2 MI. Therefore, management should instead focus on identifying and treating any precipitating conditions (e.g. anaemia, hypoxia) alongside strict control of CV risk factors.

12.2. Complications

12.2.1. Heart failure

Acute HF may occur as a complication of ACS. Acute HF as a result of ACS significantly increases the risk of other in-hospital complications, including worsening of renal function, respiratory failure, pneumonia, and death. *De novo* acute HF complicating ACS should be distinguished from pre-existing HF exacerbated by ACS.[554–556] This can be challenging and the presence of acute HF may impede the straightforward diagnosis of ACS. Patients with ACS and acute HF are more likely to present with resting dyspnoea and clinical signs/symptoms of fluid overload. In some clinical scenarios, increased troponin levels in patients with acute HF may reflect myocardial injury due to HF rather than myocardial necrosis due to ischaemia.

Patients with ACS complicated by acute HF require urgent and coordinated management of both conditions. The management of acute HF should follow current recommendations included in the ESC Guidelines on HF and ancillary documents.[557–559] The use of diuretics, vasodilators, inotropic agents, and vasopressors should be considered according to the established algorithms. Mechanical circulatory support may also be considered in selected cases. Invasive respiratory support and/or renal replacement therapy may be required in some circumstances.[557–559]

Patients presenting with acute HF (including patients with CS) complicating ACS require immediate ICA.[250,394,396] These patients should also undergo emergency echocardiography/chest ultrasonography to gather information about LV and RV function, regional wall motion abnormalities, valvular function, and possible mechanical complications.[250,557,560] In patients with ACS, CS may occur as a result of

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

extensive ischaemia due to MVD, acute severe mitral regurgitation, and mechanical complications. Patients with ACS and CS should be transferred as soon as possible to a PCI centre where immediate coronary angiography, and PCI of the IRA if needed, can be performed.[404,505] In patients with CS complicating ACS in whom the coronary anatomy is not suitable for PCI, emergency CABG is recommended. Management of MVD in this context is detailed in *Section 10*.

The clinical benefit of percutaneous MCS devices and/or VA-ECMO in the context of ACS remains unclear.[402,561] Micro-axial MCS devices have not been associated with lower 30-day mortality in comparison to IABP in observational studies.[400] In a large retrospective registry of 48 306 patients (>80% ACS) undergoing PCI with MCS, micro-axial MCS support was associated with higher mortality and bleeding rates in comparison to IABP.[562] Similar results were observed in another propensity-matched registry analysis confined to patients with CS, where micro-axial MCS support was also associated with more complications and higher mortality than IABP.[563] In the IABP-SHOCK II trial, the routine use of IABP in patients with ACS and CS did not reduce 30-day, 1-year, or 6-year mortality.[399,405,407] Based on these data, a benefit of LVAD in patients with ACS has not been demonstrated, and given that observational data have suggested that this may be associated with harm, caution is advised in this regard until further RCT evidence is available.

12.2.2. Mechanical complications

Mechanical complications may occur in the first days following MI, most commonly in patients presenting with STEMI. The incidence of mechanical complications has fallen significantly in the era of PPCI.[564] A recent large epidemiological investigation including almost 9 million ACS patients reported an overall prevalence of mechanical complications in 0.27% of STEMI cases and 0.06% of NSTEMI cases, with in-hospital mortality rates of 42.4% and 18%, respectively.[564] Mechanical complications are life-threatening and therefore require prompt identification and management (Supplementary data online, *Table S14*). Sudden hypotension, the recurrence of chest pain, new cardiac murmurs suggestive of acute mitral regurgitation or a ventricular septal defect, pulmonary congestion, or jugular vein distension should raise suspicion of a mechanical complication. Immediate echocardiographic assessment is indicated when mechanical complications are suspected.

The use of temporary MCS for mechanical complications, either to improve pre-operative clinical/haemodynamic status or prophylactically, represents a new trend in management. However, this approach requires more data and evidence in order to determine if it provides a clinical benefit.[565–568] Surgery is currently regarded as the treatment of choice for patients with ACS and mechanical complications, although percutaneous strategies are occasionally used in selected candidates with a prohibitive risk profile or contraindications to a surgical approach.[569–572] A multidisciplinary approach to the management of these patients is of paramount importance, and should apply to all stages of care, from the initial stabilization of the patient to discussion and application of the therapeutic strategy, including palliative care.[573,574] Patients with ACS-related mechanical complications should be considered for IABP while awaiting surgery.

12.2.3. Left ventricular thrombus

While the incidence of LV thrombus following AMI has declined due to advances in reperfusion and antithrombotic therapies, it remains relatively common, particularly following anterior STEMI, where it can be present in >9% of patients according to a large meta-analysis.[575,576]

Echocardiography remains the first-line imaging test for the detection of LV thrombus. In patients where the apex is not well visualized on regular echocardiography, contrast echocardiography may be considered for improved image quality. CMR is the gold standard imaging modality for the diagnosis and assessment of LV thrombi. Contemporary CMR data report LV thrombi in up to 6.3% of all STEMI patients and in 12.2% of those with anterior STEMI, suggesting that the incidence of LV thrombi may be underestimated with echocardiography.[577] Patients with LV thrombi that were not evident on echocardiography but were detected by CMR appear to have similar clinical outcomes to patients with LV thrombi that were evident on echocardiography.[578] Therefore, CMR should be considered in patients with equivocal echocardiographic images or in patients considered to be at a particularly high risk of LV thrombus.

The timing of imaging for LV thrombus may also be relevant, given that the identification of LV thrombus has been reported to increase in the first 2 weeks post-MI.[579] While more contemporary data are required, these data suggest that a high proportion of LV thrombi may develop following hospital discharge, indicating that delayed imaging at 2 weeks in high-risk patients may be of value.

Once an LV thrombus has been diagnosed, OAC therapy (warfarin or NOAC) should be considered for 3–6 months, guided by repeated echocardiography or CMR and with consideration of bleeding risk and the need for concomitant antiplatelet therapy.[580,581] However, there are a lack of prospective randomized data on the optimal anticoagulation regimen, anticoagulation duration, and the combination of oral anticoagulation with antiplatelet agents in patients with LV thrombus following MI.[581] The choice of therapy should be tailored to the patient's clinical status and the results of follow-up investigations.

12.2.4. Post-acute coronary syndrome pericarditis

Pericardial complications that may develop after an AMI include early infarct-associated pericarditis (occurring from a few hours to 4 days after AMI, mostly transient), late pericarditis or post-cardiac injury (Dressler) syndrome (typically occurring 1–2 weeks after AMI), and pericardial effusion.[548,582] This topic is discussed further in the Supplementary data online.

12.2.5. Arrhythmias

12.2.5.1. Atrial fibrillation

Atrial fibrillation is the most frequent supraventricular arrhythmia in patients with ACS.[583] AF may be pre-existing, first time detected, or of new onset during ACS management. Patients with AF have a greater number of comorbidities compared with patients without AF and are at higher risk of complications.[584] In most cases, AF is well tolerated and no specific treatment is required, apart from anticoagulation.[585] Prompt treatment is required for AF causing acute haemodynamic instability, with electrical cardioversion being the preferred approach. Adequate rate control can be achieved by administration of beta-blockers depending on the presence of HF and low ejection fraction. For patients with depressed LVEF, amiodarone or digoxin could be used (preferably amiodarone). In cases of hypotension, digoxin is preferred over amiodarone or beta-blockers. Patients with AF and risk factors for thrombo-embolism should be adequately treated with chronic oral anticoagulation.[585] ACS patients with documented AF of any

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

length have worse short- and long-term prognoses when compared with patients in sinus rhythm.[584,586] There is some evidence to suggest that transient, self-terminating AF during STEMI may be a predictor of an increased risk of stroke during long-term follow-up.[584,587]

12.2.5.2. Ventricular arrhythmias

With the widespread increased uptake of emergency reperfusion therapies for patients with STEMI, the incidence of malignant arrhythmias (ventricular tachycardia [VT] and ventricular fibrillation [VF]) has significantly declined. Nevertheless, 6–8% of patients with STEMI develop haemodynamically significant VT or VF.[588] The typical arrhythmia presentation is unstable, frequently polymorphic, and relatively fast VT, often degenerating into VF. Urgent reperfusion is most important as ischaemia is often the trigger for these arrhythmias. Early administration of i.v. or oral beta-blockers reduces the incidence of malignant arrhythmias.[163,164,169,589] Beta-blockers or amiodarone are recommended if malignant arrhythmias occur and lidocaine may be considered if these are contraindicated.[163,164,169,589,590] The prognostic role of early VT/VF within the first 48 h of STEMI is still controversial. Several studies have suggested that patients with early VT/VF have increased 30-day mortality but no increase in long-term arrhythmic risk.[591–593] Another study has suggested that while malignant ventricular arrhythmias occurring at the time of reperfusion do not confer poor prognosis, sustained VT or VF occurring during ongoing ischaemia or late after reperfusion (>48 h) is associated with an increase in long-term mortality.[594] Sustained VT/VF late after reperfusion (>48 h) requires an evaluation for ICD implantation for secondary prevention of sudden cardiac death. Ventricular premature beats are very frequent during the first 24 h after reperfusion for STEMI and no specific therapy is required.

Primary prevention of sudden cardiac death with ICD implantation within 40 days after MI is generally not indicated. Patients should be re-evaluated for ICD implantation post-revascularization after a period of 6–12 weeks on evidence-based treatments, although patients with a pre-existing impaired LVEF may be considered for ICD implantation for primary prevention even within the early post-infarction period. Some patients may develop electrical storm and/or incessant VT despite complete revascularization and treatment with anti-arrhythmic drugs. Overdrive stimulation may help to control this situation; however, recurrence of VT/VF upon cessation of stimulation is frequent and catheter ablation of such triggers appears to be the preferred treatment option in centres with that expertise. Successful radiofrequency ablation has been shown to abolish recurrent VT/VF.[595]

Non-sustained monomorphic VT is the most common form of ventricular arrhythmia in the early phase of ACS, and usually does not require anti-arrhythmic treatment. Accelerated idioventricular rhythm at reperfusion is frequent and does not require intervention given its benign nature.[596]

12.2.6. Bleeding

Bleeding is associated with a poor prognosis in ACS patients.[231,597,598] The mechanisms by which bleeding increases the risk of death are complex and multifactorial.[599] While intracranial or massive haemorrhage directly threatens life through fatal brain damage or sudden cardiocirculatory collapse, other less severe forms of haemorrhage may increase the risk of death through indirect mechanisms. Blood transfusion may increase systemic inflammation and represents one of the possible links between bleeding and subsequent mortality.[600] Bleeding is also a major driver of unplanned DAPT discontinuation and the interruption of other medication (e.g. statins, beta-blockers).[601,602]

12.2.6.1. Management of bleeding

See Supplementary data online, *Section 12.1.3.1*.

Recommendation Table 14 — Recommendations for acute coronary syndrome complications

Recommendations	Class[a]	Level[b]
Heart failure		
IABP should be considered in patients with haemodynamic instability/cardiogenic shock due to ACS-related mechanical complications.	IIa	C
LV thrombus		
CMR imaging should be considered in patients with equivocal echocardiographic images or in cases of high clinical suspicion of LV thrombus.[577,578]	IIa	C
Oral anticoagulant therapy (VKA or NOAC) should be considered for 3–6 months in patients with confirmed LV thrombus.[603]	IIa	C
Following an acute anterior MI, a contrast echocardiogram may be considered for the detection of LV thrombus if the apex is not well visualized on echocardiography.[604]	IIb	C
Atrial fibrillation		
Intravenous beta-blockers are recommended when rate control is needed in the absence of acute HF or hypotension.[605]	I	C
Intravenous amiodarone is recommended when rate control is needed in the presence of acute HF and no hypotension.[606]	I	C
Immediate electrical cardioversion is recommended in patients with ACS and haemodynamic instability and when adequate rate control cannot be achieved promptly with pharmacological agents.	I	C
Intravenous amiodarone is recommended to facilitate electrical cardioversion and/or decrease risk for early recurrence of AF after electrical cardioversion in unstable patients with recent-onset AF.[607,608]	I	C
In patients with documented *de novo* AF during the acute phase of ACS, long-term oral anticoagulation should be considered depending on the CHA_2DS_2-VASc score, after taking the HAS-BLED score and the need for concomitant antiplatelet therapy into consideration. NOACs are the preferred drugs.[583,584,587]	IIa	C
Ventricular arrythmias		
ICD therapy is recommended to reduce sudden cardiac death in patients with symptomatic HF (NYHA Class II–III) and LVEF ≤35% despite optimal medical therapy for >3 months and at least 6 weeks after MI who are expected to survive for at least 1 year with good functional status.[434,609,610]	I	A

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Intravenous beta-blocker and/or amiodarone treatment is recommended for patients with polymorphic VT and/or VF unless contraindicated.[611–614]	I	B
Prompt and complete revascularization is recommended to treat myocardial ischaemia that may be present in patients with recurrent VT and/or VF.[368,388]	I	C
Transvenous catheter pacing termination and/or overdrive pacing should be considered if VT cannot be controlled by repeated electrical cardioversion.	IIa	C
Radiofrequency catheter ablation at a specialized ablation centre followed by ICD implantation should be considered in patients with recurrent VT, VF, or electrical storm despite complete revascularization and optimal medical therapy.	IIa	C
Treatment of recurrent VT with haemodynamic relevance (despite repeated electrical cardioversion) with lidocaine may be considered if beta-blockers, amiodarone, and overdrive stimulation are not effective/applicable.[615]	IIb	C
In patients with recurrent life-threatening ventricular arrhythmias, sedation or general anaesthesia to reduce sympathetic drive may be considered.[616]	IIb	C
ICD implantation or the temporary use of a wearable cardioverter defibrillator may be considered <40 days after MI in selected patients (incomplete revascularization, pre-existing LVEF dysfunction, occurrence of arrhythmias >48 h after STEMI onset, polymorphic VT or VF).	IIb	C
Treatment of asymptomatic and haemodynamically irrelevant ventricular arrhythmias with anti-arrhythmic drugs is not recommended.	III	C
Bradyarrhythmias		
In cases of sinus bradycardia with haemodynamic intolerance or high-degree AV block without stable escape rhythm:		
• i.v. positive chronotropic medication (adrenaline, vasopressin, and/or atropine) is recommended.[617,618]	I	C
• temporary pacing is recommended in cases of failure to respond to atropine.	I	C
• urgent angiography with a view to revascularization is recommended if the patient has not received previous reperfusion therapy.	I	C
Implantation of a permanent pacemaker is recommended when high-degree AV block does not resolve within a waiting period of at least 5 days after MI.	I	C
In selected patients with high-degree AV block in the context of an anterior wall MI and acute HF, early device implantation (CRT-D/CRT-P) may be considered.[619,620]	IIb	C
Pacing is not recommended if high-degree AV block resolves after revascularization or spontaneously.[620–622]	III	B

Continued

© ESC 2023

ACS, acute coronary syndrome; AF, atrial fibrillation; AV, atrioventricular; CHA_2DS_2-VASc, Congestive heart failure, Hypertension, Age ≥75 (doubled), Diabetes, previous Stroke/transient ischaemic attack/thrombo-embolism (doubled), Vascular disease, Age: 65–74, Sex (female); CMR, cardiac magnetic resonance; CRT-D/CRT-P, cardiac resynchronization therapy—defibrillator/pacemaker; HAS-BLED, Hypertension, Abnormal liver/renal function, Stroke history, Bleeding history or predisposition, Labile INR, Elderly, Drug/alcohol usage; HF, heart failure; IABP, intra-aortic balloon pump; ICD, implantable cardioverter defibrillator; LV, left ventricular; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NOAC, non-vitamin K antagonist oral anticoagulant; NYHA, New York Heart Association; STEMI, ST-elevation myocardial infarction; VF, ventricular fibrillation; VKA, vitamin K antagonist; VT, ventricular tachycardia.
[a]Class of recommendation.
[b]Level of evidence.

12.3. Comorbid conditions

12.3.1. Patients at high bleeding risk and with blood disorders (anaemia and thrombocytopaenia)

Anaemia is more prevalent in elderly/frail ACS patients and in patients with multimorbidity (i.e. HF, chronic kidney disease [CKD], diabetes mellitus, cancer, and autoimmune diseases). In some cases, severe anaemia may precipitate Type 2 MI. Persistent or worsening anaemia in patients with ACS is associated with an increased risk of recurrent ischaemic events, death, and major bleeding.[623–625] According to the ARC-HBR, haemoglobin <11 g/dL at the time of PCI constitutes a major criterion for HBR, whereas haemoglobin between 11 and 13 g/dL (12 g/dL for women) is a minor criterion.

There is no established strategy for treating anaemia in patients with ACS. The efficacy and safety of blood transfusion in this clinical scenario remains unknown. In the majority of studies investigating different transfusion protocols, a liberal blood transfusion strategy has been defined as any red blood cell transfusion at a haemoglobin level <9–10 g/dL, while a restrictive blood transfusion strategy has been defined as any transfusion at a haemoglobin level <7–8 g/dL. Observational data suggest that a liberal blood transfusion strategy may be associated with an increase in all-cause mortality.[626–630] The open-label Restrictive and Liberal Transfusion Strategies in Patients With Acute Myocardial Infarction (REALITY) trial enrolled 668 ACS patients who were randomized to management with a restrictive (triggered by haemoglobin ≤8) or a liberal (triggered by haemoglobin ≤10) transfusion strategy.[631] The composite outcome (all-cause death, stroke, recurrent MI, or emergency revascularization) at 30 days occurred in a comparable number of patients in both arms (11% vs. 14%, RR 0.79, with a one-sided 97.5% CI of 0.00–1.19), meeting the pre-specified non-inferiority criterion. All components of the composite endpoint were numerically higher in the liberal transfusion strategy arm. The trial was not powered to detect superiority of the restrictive strategy, and the CI included what may be a clinically important harm. The pre-specified 1-year follow-up of the REALITY trial yielded contradictory conclusions to the 30-day outcomes: at 1 year, the restrictive transfusion strategy (vs. a liberal

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

approach) did not achieve non-inferiority in terms of MACE. In addition, a *post-hoc* analysis of MACE between day 30 and 1 year demonstrated an increased risk in the restrictive transfusion strategy group.[632] Therefore, no formal recommendation as to the optimal transfusion strategy (liberal vs. restrictive) in patients with ACS can be made at present.

Although there are several classifications to grade the severity of thrombocytopaenia, clinically relevant thrombocytopaenia can be defined as a platelet count <100 000/μL or a relative drop in platelet count of 50% from baseline in the context of ACS. Thrombocytopaenia increases the risk of death, major bleeding events, and life-threatening thrombotic events.[633,634] The ARC-HBR criteria define a platelet count <100 000/μL as a major criterion for HBR. Management of GP IIb/IIIa inhibitor- and heparin-induced thrombocytopaenia is discussed in the Supplementary data online.

12.3.2. Chronic kidney disease

Moderate to severe CKD (stages III–V) is present in more than 30% of ACS patients.[635] Patients with ACS and concomitant CKD receive less interventional and pharmacological treatment and have a worse prognosis than patients with normal kidney function.[636–638] Likely contributing factors to this worse prognosis include a larger number of comorbidities and an increased risk of in-hospital complications, including serious bleeding complications.[639] Although evidence from RCTs is lacking, data from observational and registry-based studies indicate that ACS patients with moderate to severe CKD have a better prognosis with early revascularization than with medical therapy alone.[640,641]

The type and dose of antithrombotic agent (see Supplementary data online, *Table S15*) and the amount of contrast agent should be considered based on kidney function.[635,642] In relation to supplementary i.v. hydration during and after revascularization, the evidence around choice, timing, and duration of treatment is somewhat conflicting.[643] Taking the clinical circumstances and patient characteristics into consideration, i.v. hydration should be considered as part of the management of ACS patients with a low eGFR undergoing invasive management to minimize the risk of contrast-induced nephropathy.[250,635,642,644,645] For recommendations on long-term treatment in patients with ACS and concomitant CKD, please refer to the 2021 ESC Guidelines on cardiovascular disease prevention.[646]

12.3.3. Diabetes mellitus

ACS patients with diabetes mellitus (DM) may more commonly present with non-specific symptoms, which can lead to delays in both diagnosis and access to treatment.[647,648] Both treatment in the acute phase and risk factor management post-ACS is poorer in patients with DM and these patients tend to have more advanced CAD at diagnosis. These factors likely contribute to the worse long-term prognosis associated with ACS in patients with DM, particularly in patients requiring insulin treatment.[649–651]

All patients with ACS, regardless of a history of DM, should have their glycaemic status evaluated during hospitalization. Given that the ACS itself may give rise to hyperglycaemia due to catecholamine-induced stress, a diagnosis of DM made during hospitalization should be subsequently confirmed. While several studies have shown the benefits of managing hyperglycaemia (>11.0 mmol/L or 200 mg/dL) in hospitalized ACS patients, the risk of hypoglycaemia-related events when using intensive insulin therapy should not be neglected.[652–654]

Glucose lowering is important in order to prevent microvascular complications in patients with DM. However, recent trial evidence has shown that the reduction in the risk of new ACS events, HF, and renal impairment with glucose-lowering medications like sodium–glucose co-transporter 2 (SGLT2) inhibitors or glucagon-like peptide-1 receptor agonists (GLP-1-RA) is independent of baseline glycosylated haemoglobin (HbA1c) levels.[655–657] This should be taken into consideration when choosing glucose-lowering therapy for patients with DM and concomitant CAD. For further details, please refer to the 2023 ESC Guidelines on diabetes and cardiovascular diseases and the 2021 ESC Guidelines on cardiovascular disease prevention.[646,658]

12.3.4. Older adults with frailty and multimorbidity

12.3.4.1. The older person

Older adults represent an increasing proportion of ACS patients. One of the major predictors of adverse outcomes following ACS is age, but patients aged ≥75 years are often excluded from or under-represented in clinical trials.[659,660] Older age is associated with frailty, multimorbidity, and a greater risk of both ischaemic and bleeding events in patients with ACS.[661] Hs-cTn assays have an excellent diagnostic performance in the older person, but the specificity of the test is lower than in younger patients, and elevated cTn levels are more commonly associated with conditions other than ACS in older patients.[662]

There are limited data on the optimal management of older adults with ACS.[663] A small RCT enrolling older patients (≥80 years) with NSTE-ACS reported the superiority of an invasive vs. a conservative strategy in the reduction of the composite of MI, need for urgent revascularization, stroke, and death. No treatment effect was shown for all-cause death and the benefit associated with the invasive strategy was diluted with increasing age.[664] In the absence of robust clinical trial evidence, decisions regarding how to manage older patients should be individualized based on patient characteristics (i.e. ischaemic and bleeding risks, estimated life expectancy, comorbidities, the need for non-cardiac surgery, quality of life, frailty, cognitive and functional impairment, patient values and preferences, and the estimated risks and benefits of an invasive strategy).

In the context of STEMI, PPCI has drastically improved outcomes for all ages. However, data are limited in the 'very old' cohort, with lack of formal assessment of frailty or comorbidity.[665] In the context of CS and cardiac arrest, age is an independent predictor of mortality following PCI.[666,667] In the absence of robust RCT data, PPCI should be considered for all patients with STEMI. When PPCI cannot be performed in a timely manner, fibrinolysis may be a reasonable strategy in these patients. For details regarding pharmacotherapy in older patients, please see the Supplementary data online.

12.3.4.2. Frailty and multimorbidity

Geriatric syndromes (i.e. frailty and multimorbidity) are associated with adverse outcomes in older patients with ACS.[668,669] Frailty is a syndrome characterized by reduced biological reserve, leading to a failure of homeostatic mechanisms following stressor events, including ACS. There is a lack of consensus on which frailty assessment tool is optimal in older patients with CV disease.[670,671]

Frail patients with NSTE-ACS less frequently receive ACS pharmacotherapies and invasive assessment, have more complex coronary disease, have longer durations of hospital stay, and are at higher risk of death.[672] Specifically, frail patients are reported to have a higher rate of a composite of all-cause mortality, MI, stroke, unplanned

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

revascularization, and major bleeding.[673] Frail older adults with NSTE-ACS have poor health-related quality of life (HRQoL) at baseline. Invasive management appears to be associated with modest improvements in HRQoL through to 1 year follow-up in these patients. This improvement in HRQoL is most marked in frail and pre-frail patients, who receive a proportionally larger benefit than robust patients.[674] In older adults with NSTE-ACS referred for coronary angiography, the presence of multimorbidity is associated with an increased risk of long-term adverse CV events, driven by a higher risk of all-cause mortality.[675] Undiagnosed cognitive impairment is also common in older patients with NSTE-ACS undergoing ICA, and these patients are more likely to experience MACE at 1 year.[676]

In the absence of robust RCT data to inform healthcare professionals about the management of frail patients presenting with ACS, it is recommended to adapt a holistic approach to individualize interventional and pharmacological treatments after careful evaluation of risks vs. benefits. To aid in decision-making, the routine assessment of frailty (e.g. Rockwood Frailty Score) and comorbidity (e.g. Charlson index) in ACS patients is recommended. Following risk stratification using frailty assessment and evaluation of the comorbidity burden, it may be reasonable to offer optimal medical therapy plus an invasive strategy to frail patients at high risk of future CV events and low risk of complications, and to offer optimal medical therapy alone to those who are deemed to be at low risk of future events with a high risk of developing procedural complications. For those patients for whom any form of treatment might be futile, then a palliative end-of-life care approach should be considered.

12.3.5. Pregnancy

Acute coronary syndrome diagnostic criteria are the same for pregnant and non-pregnant patients.[677] Pregnant women with STEMI should not be managed differently to non-pregnant women. Given the high mortality associated with STEMI in pregnancy, PPCI is the preferred reperfusion therapy.[678] The management plan for pregnant women with ACS should be determined by a multidisciplinary team consisting of cardiologists, obstetricians, anaesthesiologists, and neonatologists, and these patients should be treated in an intensive care unit that can provide maternal monitoring and obstetric care.[678,679] ACS treatment should not be delayed for delivery. Delivery should be ideally postponed for at least 2 weeks post-ACS as there is increased risk of maternal mortality during this time.[678] It has been demonstrated that SCAD is the most common cause of AMI in pregnancy, and this tends to occur mainly in the late pregnancy or early post-partum periods.[680,681] Further details are provided in the Supplementary data online.

12.3.6. Drug abuse

Acute coronary syndrome in the setting of drug abuse is covered in the Supplementary data online.

12.3.7. Patients with cancer

The four most common types of cancer in patients with ACS are prostate, breast, colon, and lung.[682] Patients with a history of cancer should be treated like all other ACS patients, but the management of ACS patients with active cancer has some specific issues that need to be taken into consideration. Outcomes vary across types of cancer and the balance between the ischaemic and bleeding risks should be considered on an individual basis.

The percentage of ACS patients with a current diagnosis of cancer is rising, and currently constitutes ~3% of patients in large observational studies.[683] Patients with active cancer presenting with ACS pose important challenges as there are significant gaps in scientific knowledge. Therefore, recommendations based on solid evidence are scarce. Patients with active cancer presenting with ACS tend to be older, with a larger number of comorbidities and more extensive CAD. These patients often have concomitant haematologic and coagulation abnormalities that may present a challenge with respect to both the use of antithrombotic therapy and the performing of PCI.[684] Observational studies have reported that ACS in patients with cancer is associated with increased risk of major CV events, bleeding, and cardiac and non-cardiac mortality.[682,683,685,686] As per the ARC-HBR criteria, patients with active cancer diagnosed in the past 12 months are considered as HBR.

The diagnosis of ACS in patients with cancer should be based on the same principles as in patients without cancer. The management of ACS in patients with cancer can be challenging because of frailty, increased bleeding risk, thrombocytopaenia, and increased thrombotic risk.[687] Temporary interruption of cancer treatment and an urgent multidisciplinary approach is recommended.[688] Cancer patients with ACS have been reported to less frequently undergo invasive management; however, invasive management (and PCI with DES if needed) is recommended in ACS patients with cancer, as long as the prognosis is >6 months or, irrespective of the prognosis, if the patient is unstable.[689] Retrospective data have reported both a lower use of invasive management in cancer patients with STEMI, and better outcomes in patients who do undergo invasive management.[682,686,689] Invasive management in patients with advanced cancer or life expectancy <6 months has been reported to not demonstrate a mortality benefit compared with a conservative approach and therefore a conservative strategy should be considered in these patients.[690] When the coronary anatomy is not amenable for PCI, CABG surgery can be considered after a multidisciplinary team discussion and where the cancer prognosis is >12 months. Given that they are considered to be HBR, the preferred $P2Y_{12}$ inhibitor for ACS patients with active cancer is clopidogrel.[687] Potential drug–drug interactions with cancer therapies should be checked when using ticagrelor or clopidogrel, since some pharmacokinetic-based drug–drug interactions via CYP450 may occur.

When acute ischaemia is provoked by cancer therapy, alternative cancer therapies should be considered after a multidisciplinary team discussion. Some specific cancer treatments can have cardiotoxic vascular effects that can lead to ACS (Supplementary data online, *Table S16*). Following ACS, a review of the cancer medications is recommended, and any cancer drug associated with thrombosis and MI should be stopped. Cancer therapies that are not associated with MI can be restarted once revascularization (when indicated) has been completed and the patient is stabilized on ACS medical therapy without complications. Additional information can be found in the Supplementary data online, including Supplementary data online, *Table S16* and in the 2022 ESC Guidelines on cardio-oncology.[684]

12.3.8. Coronavirus disease (COVID-19)

A section on the impact of Coronavirus disease (COVID-19) on ACS management is presented in the Supplementary data online.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Recommendation Table 15 — Recommendations for acute coronary syndrome comorbid conditions

Recommendations	Class[a]	Level[b]
Chronic kidney disease		
The use of low- or iso-osmolar contrast media (at the lowest possible volume) is recommended for invasive strategies.[691–693]	I	A
It is recommended to assess kidney function using eGFR in all patients with ACS.	I	C
It is recommended to apply the same diagnostic and therapeutic strategies in patients with CKD (dose adjustment may be necessary) as in patients with normal kidney function.	I	C
Hydration during and after angiography should be considered in patients at risk of contrast-induced nephropathy, especially in patients with acute kidney injury and/or CKD with eGFR <30 mL/min/1.73 m^2.[694–697]	IIa	B
Diabetes		
It is recommended to base the choice of long-term glucose-lowering treatment on the presence of comorbidities, including heart failure, CKD, and obesity.[698–704]	I	A
It is recommended to assess glycaemic status at initial evaluation in all patients with ACS.[705–707]	I	B
It is recommended to frequently monitor blood glucose levels in patients with known diabetes mellitus or hyperglycaemia (defined as glucose levels ≥11.1 mmol/L or ≥200 mg/dL).	I	C
Glucose-lowering therapy should be considered in patients with ACS with persistent hyperglycaemia, while episodes of hypoglycaemia should be avoided.[708,709]	IIa	C
Older adults		
It is recommended to apply the same diagnostic and treatment strategies in older patients as in younger patients.[662,664,665,710,711]	I	B
It is recommended to adapt the choice and dosage of antithrombotic agent, as well as of secondary prevention medications, to renal function, co-medications, comorbidities, frailty, cognitive function, and specific contraindications.[363,712]	I	B
For frail older patients with comorbidities, a holistic approach is recommended to individualize interventional and pharmacological treatments after careful evaluation of the risks and benefits.[668,673,676]	I	B
Patients with cancer		
An invasive strategy is recommended in cancer patients presenting with high-risk ACS with expected survival ≥6 months.[682,689,690]	I	B
A temporary interruption of cancer therapy is recommended in patients in whom the cancer therapy is suspected to be a contributing cause of ACS[c].[713,714]	I	C
A conservative non-invasive strategy should be considered in ACS patients with poor cancer prognosis[d] (i.e. with expected survival <6 months) and/or very high bleeding risk.[690]	IIa	C
Aspirin is not recommended in cancer patients with a platelet count <10 000/μL.[715]	III	C
Clopidogrel is not recommended in cancer patients with a platelet count <30 000/μL.	III	C
In ACS patients with cancer and <50 000/μL platelet count, prasugrel or ticagrelor are not recommended.	III	C

Continued

ACS, acute coronary syndrome; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate.
[a]Class of recommendation.
[b]Level of evidence.
[c]Anticancer therapies associated with high risk of ACS (very common [>10%]) include: capecitabine, paclitaxel, cisplatin, carfilzomib, bevacizumab, ramucirumab, aflibercept, axitinib, sorafenib, pazopanib, cabozantinib, lenvatinib, ponatinib, and erlotinib.
[d]Related to advanced cancer stage and/or severe irreversible non-CV comorbidities.

13. Long-term treatment

Secondary prevention after ACS is central to increase quality of life and to decrease morbidity and mortality. This should start as early as possible after the index event.[716–718] The topic is covered in detail in the 2019 CCS Guidelines and the 2021 Prevention Guidelines.[195,646] Optimal medical therapy and treatment targets are well defined and are summarized in *Figure 17*. A figure aimed at educating patients on improving their 'heart health' after an ACS event is provided in the Supplementary data online, *Figure S5*.

© ESC 2023

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026




Figure 17 Long-term management after acute coronary syndrome. ACS, acute coronary syndrome; HbA1c, glycosylated haemoglobin; LDL-C, low-density lipoprotein cholesterol; OPD, outpatient department. [a]See Recommendation Table 16 for other pharmacological treatments after ACS. [b]For patients ≥70 years of age the systolic target should be <140 mmHg and down to 130 mmHg if tolerated.[646] [c]For patients with diabetes mellitus.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

13.1. Cardiac rehabilitation

13.1.1. Comprehensive cardiac rehabilitation

Secondary prevention is most effectively provided through cardiac rehabilitation (CR).[716,717] All ACS patients should participate in a comprehensive CR programme, which should start as early as possible after the ACS event.[716,717,719] CR may be performed in inpatient or outpatient settings, taking age, frailty, results of prognostic risk stratification, and comorbidities into account.[716] Comprehensive CR is a multidisciplinary intervention, supervised and performed by a team and usually co-ordinated by a cardiologist.[716] The core components of CR include patient assessment, management and control of CV risk factors, physical activity counselling, prescription of exercise training, dietary advice, tobacco counselling, patient education, psychosocial management, and vocational support.[716] Several studies have found that CR programmes after atherosclerotic cardiovascular disease (ASCVD) events or revascularization reduce CV hospitalizations, MI, CV mortality and, in some studies, all-cause mortality.[720–725] Despite proven benefits, the rates of referral to, participation in, and implementation of CR programmes are low.[726–730] Another identified issue is that many patients adopt healthier lifestyles during CR but relapse to pre-morbid habits when returning to everyday life.[731] Therefore, there is an unmet need for complementary pathways to the classical centre-based CR model. In addition to alternatives to CR, there is also a need for stronger endorsement of CR by physicians, cardiologists, and healthcare professionals.[732,733] It is also important to initiate and establish a strong partnership between patients and healthcare professionals as early as possible.[732–734]

13.1.2. Digital health

Telerehabilitation may be an effective strategy to maintain a healthy lifestyle over time and can support or even partially replace conventional, centre-based CR.[729] Telerehabilitation means rehabilitation from a distance, covering all CR core components, including telecoaching, social interaction, telemonitoring, and e-learning.[735,736] Studies in patients with CAD have shown that telerehabilitation can be equivalent to traditional CR in terms of achieving functional improvement, managing risk factors, and increasing patient well-being.[737–741] Few data are available about the effect of telerehabilitation on recurrent events.[742] Nevertheless, in a meta-analysis no significant difference was found between mortality following telehealth interventions and centre-based supervised CR.[743] Also, most trials have only focused on one of the CR core components—exercise training and/or physical activity.[742] Therefore, more research on the impact of telerehabilitation on outcomes is still needed, as are investigations into health and digital literacy in CR.

13.1.3. Adherence and persistence

Promotion of both adherence (the extent to which a patient adheres to a prescribed treatment or lifestyle advice) and persistence (the length of time between initiation and discontinuation of a prescribed treatment or lifestyle advice) are key in preventing recurrent CV events after ACS. Adherence to medication has been shown to be sub-optimal, ranging from 50% in primary prevention to 66% in secondary prevention. It is estimated that 9% of ASCVD events in Europe occur as a result of sub-optimal medication adherence.[646] Contributors to sub-optimal adherence and persistence are multidimensional and include: polypharmacy, drug regimen complexity, the doctor–patient relationship, a lack of patient-centred care and disease acceptance, concern regarding side effects, cognitive ability, mental and physical disorders, financial aspects, living alone, and depression.[646,744–749] Polypills, which include guideline-recommended treatments for secondary prevention, have been shown to increase adherence in post-ACS patients and may improve therapeutic targets.[750–752] The Secondary Prevention of Cardiovascular Disease in the Elderly (SECURE) study is the only RCT testing the impact of a strategy based on a polypill (containing aspirin, ramipril, and atorvastatin) vs. usual care on hard outcomes in ACS patients. The polypill strategy was associated with a significant reduction in major CV events, driven by a significant 33% reduction in CV mortality.[753] The use of technology to improve medication adherence is also generating interest: mobile phone applications and mobile health (mHealth) tools may improve medication adherence, but clinical trials of sufficient size and duration are needed.[754–756] Finally, it is important to recognize that adherence has complex underlying psychological drivers, and therefore a whole-systems approach is mandatory. This should include the education of health professionals, the use of patient-reported outcomes and experience measures, patient education, and patient-centred care.[734,757,758]

13.2. Lifestyle management

Lifestyle management is one of the cornerstones of comprehensive CR.[716] While most of the evidence regarding the benefits of a healthy lifestyle on prognosis comes from primary prevention, studies in secondary prevention settings indicate similar beneficial effects.[716,724,759–763]

13.2.1. Tobacco

Tobacco abstinence is associated with a reduced risk of re-infarction (30–40%) and death (35–45%) after ACS.[763–765] Measures to promote cessation of smoking are therefore a priority after ACS. Interventions for smoking cessation should begin during hospitalization using a combination of behavioural interventions, pharmacotherapy, and counselling.[18,766] Many patients continue or resume smoking after ACS, in particular patients with depression and environmental exposures.[646] During encounters with smokers, the 'very brief advice' evidence-based intervention should be used to facilitate dialogue between the patient and healthcare worker.[646] Drug interventions, including nicotine-replacement therapy (NRT), bupropion and varenicline, should be considered along with behavioural support. All forms of NRT are effective, and the anti-depressant bupropion aids in long-term smoking cessation with similar efficacy to NRT.[646,766] Varenicline is the most effective medical treatment to support smoking cessation and is safe to use in ACS patients.[767–770] An average weight gain of 5 kg can be expected when a person quits smoking, but it is important to recognize that the CV risk from continued smoking outweighs the CV risk from gaining weight.[646]

E-cigarettes have been used to help smokers quit, but evidence on their impact on successful smoking cessation is insufficient, particularly with regard to whether using e-cigarettes actually helps the person remain tobacco free. While e-cigarettes do contain nicotine, they do not contain as many tobacco chemicals as cigarettes. Caution should be given with respect to the use of e-cigarettes, as current evidence suggests they are harmful to CV health by increasing arterial stiffness, heart rate and blood pressure, and by causing endothelial dysfunction.[771]

13.2.2. Nutrition and alcohol

A healthy diet and eating habits influence CV risk. Adopting a Mediterranean-style diet can help reduce CV risk in all individuals, including persons at high CV risk and patients with ASCVD.[761,762,772]

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Supplementary data online, *Table S17* summarizes the characteristics of a healthy diet that should be adhered to. For further details on nutrition, please refer to the 2021 ESC Guidelines on cardiovascular disease prevention.[646]

With regard to alcohol consumption, recent data suggest that alcohol abstainers have the lowest risk of CVD outcomes, that any amount of alcohol uniformly increases blood pressure and body mass index, and that a weekly consumption of >100 g of alcohol is associated with decreased life expectancy.[773–775] Accordingly, it is recommended to restrict alcohol consumption to a maximum of 100 g per week (same limit for men and women).[646]

13.2.3. Physical activity and exercise

Based on extensive data from the general population, sedentary behaviour, defined as time spent sitting or lying with low energy expenditure, while awake, is an independent risk factor for all-cause mortality.[776,777] According to recommendations from the World Health Organization, adults with chronic conditions should limit their amount of sedentary time, replacing it with physical activity of any intensity (including light intensity).[646,778] General physical activity recommendations include a combination of regular aerobic physical activity and resistance exercise throughout the week, which also forms the basis of recommendations for patients post-ACS.[646,778] However, it is important to recognize that daily physical activity does not replace participation in exercise-based CR. With support from multiple randomized trials, exercise training is a pivotal part of comprehensive CR and participation in exercise-based CR should be offered to all patients after ACS.[779] Cardiorespiratory fitness is a strong predictor of future prognosis both in the general population and in post-ACS patients.[780]

13.2.4. Psychological considerations

There is a two-fold risk of anxiety and mood disorders in patients with heart disease. Depression, anxiety, and psychological stress are associated with worse outcomes. Psychological and pharmacological interventions can have a beneficial effect and should be considered for ACS patients with depression, anxiety, and stress.[781] It is recommended that all patients have their mental well-being assessed using validated tools before discharge, with consideration of onward psychological referral when appropriate.[782] For further details, please refer to the 2021 ESC Guidelines on cardiovascular disease prevention.[646]

13.2.5. Resumption of activities

Information on the resumption of activities, sexual activity, and environmental factors is presented in the Supplementary data online, *Section 13.1.2*.

13.3. Pharmacological treatment

13.3.1. Antithrombotic therapy

Recommendations for antithrombotic therapy are included in *Section 6*.

13.3.2. Lipid-lowering therapy

Dyslipidaemia should be managed according to the current dyslipidaemia guidelines, with a combination of lifestyle and pharmacological interventions.[783] Trials have consistently demonstrated that lower low-density lipoprotein-cholesterol (LDL-C) levels after ACS are associated with lower CV event rates.[784] The current treatment goal for secondary prevention is to lower LDL-C to <1.4 mmol/L (<55 mg/dL) and to achieve a ≥50% LDL-C reduction from baseline. For patients who experience a second CV event within 2 years (not necessarily of the same type as the first event), an LDL-C goal of <1.0 mmol/L (<40 mg/dL) appears to confer additional benefit.[783,785,786]

After an ACS event, lipid-lowering treatment should be initiated as early as possible, both for prognostic benefit and to increase patient adherence after discharge. It is recommended that a high-intensity statin (e.g. atorvastatin or rosuvastatin) is initiated as early as possible after hospital admission, preferably before planned PCI, and prescribed up to the highest tolerated dose in order to reach the LDL-C goals.[783,787] The intensity of statin therapy should be increased in patients who were receiving low- or moderate-intensity statin treatment before the ACS event. In IMPROVE-IT (Improved Reduction of Outcomes: Vytorin Efficacy International Trial), ezetimibe treatment early after ACS (within 10 days) was added on top of prior statin therapy or initiated concomitantly in statin-naïve patients (two-thirds of patients) and compared with statin monotherapy.[788] Treatment with ezetimibe was shown to be safe and provided long-term benefits for CV outcomes. As such, if patients are on a maximally tolerated statin dose, or have no prior statin treatment, and have LDL-C levels which indicate it is unlikely that targets will be reached with statin therapy alone, initiating ezetimibe in addition to a statin (or statin plus ezetimibe combination treatment) may be considered during the ACS hospitalization.[783,788] In the ODYSSEY OUTCOMES (Evaluation of Cardiovascular Outcomes After an Acute Coronary Syndrome During Treatment With Alirocumab) trial, treatment with the proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor alirocumab was initiated as early as 1 month after ACS.[786] Treatment with PCSK9 inhibitors has been shown to be safe and effective in lowering LDL-C in patients hospitalized with ACS.[789–791] Recent data have also shown improvements in plaque phenotype and plaque regression in ACS patients treated with PCSK9 inhibitors.[792,793] Combined with the data from trials on the long-term benefits of PCSK9 inhibitors and observational data on the importance of lowering LDL-C early after ACS, PCSK9 inhibitor treatment should be initiated during ACS hospitalization in patients who were not at their LDL-C goal despite being on statin and ezetimibe treatment before admission.[785,786,794–796]

In all cases, lipid levels should be re-evaluated 4–6 weeks after each treatment or dose adjustment to determine whether treatment goals have been achieved and to check for any safety issues; the therapeutic regimen can then be adapted accordingly. If the LDL-C goals are not achieved with the maximum tolerated dose of a statin alone after 4–6 weeks following ACS, adding ezetimibe is recommended.[783,788] Initiation of PCSK9 inhibitor treatment is recommended in patients who do not reach their LDL-C goal despite maximum tolerated statin and ezetimibe therapy.[783,785,786] Finally, icosapent ethyl, at a dose of 2 g b.i.d., can be used in combination with a statin in patients with ACS and triglyceride levels of 1.5–5.6 mmol/L (135–499 mg/dL) despite statin treatment.[783,797] An algorithm for lipid-lowering management in ACS patients is outlined in *Figure 18*.

For a detailed description of the different lipid-lowering drug classes and respective trial data, please refer to the Supplementary data online.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026




Figure 18 Lipid-lowering therapy in ACS patients. ACS, acute coronary syndrome; LDL-C, low-density lipoprotein cholesterol; LLT, lipid-lowering therapy; PCSK9i, proprotein convertase subtilisin/kexin type 9 inhibitor. [a]Consider LDL-C <1.0 mmol/L if recurrent event.

13.3.3. Beta-blockers

The clinical benefit of beta-blockers after ACS in patients with reduced LVEF is supported by evidence from contemporary trials.[557,798–800] However, the evidence for prescribing beta-blockers after uncomplicated ACS in patients with LVEF >40% is less well established. With the exception of the CAPRICORN (CArvedilol Post-infaRct survIval COntRolled evaluatioN) trial, which only recruited patients with LVEF ≤40%, all large RCTs testing the benefits of post-MI beta-blocker maintenance were performed in the pre-reperfusion era.[801] Pooled data demonstrated that post-MI beta-blocker therapy reduced the

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

risk of death by >20%. These trials mostly enrolled patients with STEMI, making the evidence for their benefit in NSTEMI less robust. In addition, since these trials were performed, the clinical scenario has changed dramatically, with improvements in invasive strategies and associated pharmacotherapy resulting in an improved prognosis for patients with ACS.[718] Modern observational studies and meta-analyses of these trials have yielded mixed results, with some studies suggesting a benefit of beta-blocker therapy irrespective of LVEF, and others reaching the opposite conclusion.[557,800,802–804]

There is only one small, open-label trial, CAPITAL-RCT (Carvedilol Post-Intervention Long-Term Administration in Large-scale Randomized Controlled Trial), that randomized 801 STEMI patients with successful PPCI and preserved LVEF to carvedilol or control.[805] During a 3-year follow-up, the incidence of a composite of all-cause death, MI, hospitalization for HF, and hospitalization for ACS was not significantly different between the two groups. However, the trial was underpowered and therefore this scientific question remains open. There are four ongoing pragmatic prospective large-scale RCTs in Europe randomizing ACS patients without reduced LVEF to beta-blocker or control: REBOOT-CNIC (TREatment With Beta-blockers After myOcardial Infarction withOut Reduced Ejection fracTion), 8468 ACS patients with LVEF >40%; REDUCE-SWEDEHEART (Evaluation of Decreased Usage of Betablockers After Myocardial Infarction in the SWEDEHEART Registry), 5000 ACS patients with LVEF ≥50% (NCT03278509); BETAMI (BEtablocker Treatment After Acute Myocardial Infarction in Patients Without Reduced Left Ventricular Systolic Function), 10 000 ACS patients with LVEF >40%; and DANBLOCK (Danish Trial of Beta Blocker Treatment After Myocardial Infarction Without Reduced Ejection Fraction), 3570 ACS patients with LVEF >40%.[806–808]

The duration of beta-blocker therapy after uncomplicated ACS is also another controversial topic. There are some observational studies suggesting that the clinical benefit of beta-blocker therapy is restricted to the first year after the index ACS event, but the non-randomized nature of the studies limits their conclusions.[809] There are two ongoing large-scale RCTs testing the impact of beta-blocker withdrawal after 6–12 months following uncomplicated ACS in patients with preserved LVEF: AβYSS (Beta Blocker Interruption After Uncomplicated Myocardial Infarction; NCT03498066) and SMART-DECISION (Long-term Beta-blocker Therapy After Acute Myocardial Infarction; NCT04769362).[810]

13.3.4. Nitrates and calcium channel blockers

Intravenous nitrates may be useful during the acute phase in STEMI patients with hypertension or HF, provided there is no hypotension or RV infarction. In the ISIS-4 (Fourth International Study of Infarct Survival) trial, oral nitrates had no survival benefit in MI patients.[811] Their use is therefore restricted to the control of residual angina, as recommended in the 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes.[195] Calcium channel blocker use was not associated with prognostic benefit in a systematic review including 28 trials.[812] Calcium channel blocker use can be considered in the context of residual angina and for blood pressure control as recommended in the 2021 ESC Guidelines on CVD prevention and the 2019 ESC Guidelines for the diagnosis and management of CCS.[195,646]

13.3.5. Renin–angiotensin–aldosterone system inhibitors

Angiotensin-converting enzyme (ACE) inhibitors have been demonstrated to improve outcomes in post-MI patients with additional conditions, such as clinical HF and/or LVEF ≤40%, diabetes, CKD, and/or hypertension.[813–817] A systematic overview of (old) trials of ACE inhibition early in STEMI showed that their use is associated with a small but significant reduction in 30-day mortality, especially in anterior MIs.[818]

In the VALsartan In Acute myocardial iNfarcTion (VALIANT) trial, valsartan was found to be non-inferior to captopril in patients with a recent MI plus HF and/or LVEF ≤40%.[819]

There is established evidence that patients with heart failure with reduced ejection fraction (HFrEF), regardless of aetiology, benefit from ACE inhibitors.[820–823] Angiotensin receptor/neprilysin inhibitors (ARNI) have been shown to be superior to ACE inhibitors in patients with established HF (of different aetiologies) and LVEF ≤40%.[824] However, in the more recent PARADISE-MI (Prospective ARNI vs ACE Inhibitor Trial to Determine Superiority in Reducing Heart Failure Events After MI), a dedicated study in patients with recent ACS (1–7 days) complicated by HF and/or LVEF ≤40%, an ARNI combination (sacubitril plus valsartan) was not associated with a significantly lower incidence of death from CV causes or incident HF in comparison to the active comparator ramipril.[825]

In general, ACE inhibitors (or sacubitril plus valsartan as a replacement for them) are recommended for patients with established HFrEF regardless of the aetiology.[557] These agents may be considered for patients with HF with mildly reduced ejection fraction.[557] Patients who tolerate neither ACE inhibitors nor ARNI are recommended to be treated with an angiotensin receptor blocker.

In the Eplerenone Post-AMI Heart failure Efficacy and SUrvival Study (EPHESUS), the mineralocorticoid receptor antagonist (MRA) eplerenone was associated with reduced mortality and CV hospitalizations in patients with a recent MI and LV dysfunction with symptoms of either HF or diabetes.[826] The Double-Blind, Randomized, Placebo-Controlled Trial Evaluating The Safety And Efficacy Of Early Treatment With Eplerenone In Patients With Acute Myocardial Infarction (REMINDER) trial randomized 1012 patients with acute STEMI without HF to eplerenone or placebo within 24 h of symptom onset.[827] The primary endpoint was the composite of CV mortality, rehospitalization, or extended initial hospital stay due to diagnosis of HF, sustained VT or VF, ejection fraction ≤40%, or elevated BNP/NT-pro BNP at 1 month or more after randomization. Eplerenone was associated with a significant reduction in the primary composite endpoint, although this difference was primarily driven by BNP levels.[827]

13.3.6. Medications for diabetes

13.3.6.1. Sodium–glucose co-transporter 2 inhibitors

Pharmacological blockade of SGLT2 induces glycosuria with lowering of plasma glucose levels, improving glycaemic control without hypoglycaemia, and leading to reductions in weight and blood pressure.[828] In patients with type 2 diabetes and established ASCVD, three trials (with empagliflozin, canagliflozin, and dapagliflozin) have demonstrated significant CV benefits.[656,829,830] In a meta-analysis of these three trials, MACE were reduced by 11%, with no clear effect on stroke or MI. This benefit was only seen in patients with established ASCVD.[698] The benefits of SGLT2 inhibitors may relate more to cardio-renal haemodynamic effects than to atherosclerosis.[646] Further recommendations for patients with diabetes can be found in the current ESC Guidelines on diabetes, pre-diabetes,and cardiovascular diseases.[831]

In patients with HF regardless of their LVEF, dapagliflozin and empagliflozin have been shown to significantly reduce the risk of worsening HF or CV death, both in the presence or absence of type 2

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

diabetes.[702,703,832,833] In the EMMY (EMpagliflozin in patients with acute MYocardial infarction) trial, empagliflozin led to a significant improvement in NT-pro BNP reduction over 26 weeks post-MI, accompanied by a significant improvement in echocardiographic functional and structural parameters.[834] Ongoing outcome trials in ACS populations will be useful to better define the role of these agents in the absence of HF.[835]

13.3.6.2. Glucagon-like peptide-1 receptor agonists

In a systematic review and meta-analysis of seven trials (56 004 patients with type 2 diabetes) testing different GLP1-RAs, their use was associated with reductions in the incidence of MACE, CV death, all-cause mortality, MI, and stroke.[699]

13.3.7. Proton pump inhibitors

Proton pump inhibitors (PPIs) reduce the risk of upper gastroduodenal bleeding in patients treated with antiplatelet agents.[287,836,837] Therapy with a PPI is indicated for patients receiving any antithrombotic regimen who are at high risk of gastrointestinal bleeding (for details see *Section 8.2.2.3*, Bleeding risk assessment, in the Supplementary data online).

PPIs that inhibit CYP2C19, particularly omeprazole and esomeprazole, may reduce the pharmacodynamic response to clopidogrel, though there is no strong evidence that this results in an increased risk of ischaemic events or stent thrombosis in clinical trials and propensity score-matched studies.[287,288,838–842] Importantly, no interaction between the concomitant use of PPIs and aspirin, prasugrel or ticagrelor has been observed.

13.3.8. Vaccination

An annual influenza vaccination in patients with stable ASCVD appears to be associated with reduced incidence of MI, an improved prognosis in patients with HF, and decreased CV risk in adults aged 65 years and older.[843,844] In addition, influenza vaccination given early after an MI or in high-risk CAD has been shown to result in a lower risk of all-cause death and CV death at 12 months.[845–847] Therefore, influenza vaccination is recommended for all ACS patients and should be given preferentially during index hospitalization during influenza season for those not protected by a seasonal influenza vaccination.

13.3.9. Anti-inflammatory drugs

Inflammation plays a central role in the pathogenesis of atherosclerosis and acute coronary events. Several recent trials have tested the role of the anti-inflammatory agent colchicine in acute and chronic coronary syndromes.[848,849] In the Colchicine Cardiovascular Outcomes Trial (COLCOT), which enrolled 4745 patients with a recent ACS event, low-dose colchicine (0.5 mg daily) was associated with a significant reduction of the primary composite endpoint (CV death, resuscitated cardiac arrest, MI, stroke, or urgent revascularization) in comparison to placebo.[850] Of note, pneumonia was more frequent in the colchicine group. The Low-dose Colchicine trial-2 (LoDoCo2) enrolled 5522 patients with CCS (84% of whom had prior ACS) who were randomized to colchicine (0.5 mg daily) or placebo.[851] The primary endpoint (composite of CV death, MI, stroke, or ischaemia-driven coronary revascularization) rate was significantly lower in the colchicine group; however, the incidence of non-CV death was higher in the colchicine group. The benefits of colchicine in reducing CV events have been shown to be consistent irrespective of history and timing of prior ACS.[852]

13.3.10. Hormone replacement therapy

For further information on hormone replacement therapy in patients with ACS, please see the Supplementary data online.

Recommendation Table 16 — Recommendations for long-term management

Recommendations	Class[a]	Level[b]
Cardiac rehabilitation		
It is recommended that all ACS patients participate in a medically supervised, structured, comprehensive, multidisciplinary exercise-based cardiac rehabilitation and prevention programme.[721–724,853,854]	I	A
Lifestyle management		
It is recommended that ACS patients adopt a healthy lifestyle, including: • stopping all smoking of tobacco • healthy diet (Mediterranean style) • alcohol restriction • regular aerobic physical activity and resistance exercise • reduced sedentary time[724,761,763,772,773,776,777,855–858]	I	B
In smokers, offering follow-up support, nicotine replacement therapy, varenicline or bupropion, individually or in combination, should be considered.[859–864]	IIa	A
Pharmacological treatment		
Lipid-lowering therapy		
It is recommended that high-dose statin therapy is initiated or continued as early as possible, regardless of initial LDL-C values.[787,865–867]	I	A
It is recommended to aim to achieve an LDL-C level of <1.4 mmol/L (<55 mg/dL) and to reduce LDL-C by ≥50% from baseline.[868,869]	I	A
If the LDL-C goal is not achieved despite maximally tolerated statin therapy after 4–6 weeks, the addition of ezetimibe is recommended.[788]	I	B
If the LDL-C goal is not achieved despite maximally tolerated statin therapy and ezetimibe after 4–6 weeks, the addition of a PCSK9 inhibitor is recommended.[785,786,795,796]	I	A
It is recommended to intensify lipid-lowering therapy[c] during the index ACS hospitalization for patients who were on lipid-lowering therapy before admission.	I	C
For patients with a recurrent atherothrombotic event (recurrence within 2 years of first ACS episode) while taking maximally tolerated statin-based therapy, an LDL-C goal of <1.0 mmol/L (<40 mg/dL) may be considered.[785,786]	IIb	B
Combination therapy with high-dose statin plus ezetimibe may be considered during index hospitalization.[788]	IIb	B

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Beta-blockers		
Beta-blockers are recommended in ACS patients with LVEF ≤40% regardless of HF symptoms.[801,870–872]	I	A
Routine beta-blockers for all ACS patients regardless of LVEF should be considered.[798,873–878]	IIa	B
RAAS system inhibitors		
Angiotensin-converting enzyme (ACE) inhibitors[d] are recommended in ACS patients with HF symptoms, LVEF ≤40%, diabetes, hypertension, and/or CKD.[195,813–817,879]	I	A
Mineralocorticoid receptor antagonists are recommended in ACS patients with an LVEF ≤40% and HF or diabetes.[826,880]	I	A
Routine ACE inhibitors for all ACS patients regardless of LVEF should be considered.[816,817]	IIa	A
Adherence to medication		
A polypill should be considered as an option to improve adherence and outcomes in secondary prevention after ACS.[753]	IIa	B
Imaging		
In patients with pre-discharge LVEF ≤40%, repeat evaluation of the LVEF 6–12 weeks after an ACS (and after complete revascularization and the institution of optimal medical therapy) is recommended to assess the potential need for sudden cardiac death primary prevention ICD implantation.	I	C
Cardiac magnetic resonance imaging should be considered as an adjunctive imaging modality in order to assess the potential need for primary prevention ICD implantation.	IIa	C
Vaccination		
Influenza vaccination is recommended for all ACS patients.[843,845–847]	I	A
Anti-inflammatory drugs		
Low-dose colchicine (0.5 mg once daily) may be considered, particularly if other risk factors are insufficiently controlled or if recurrent cardiovascular disease events occur under optimal therapy.[850,851]	IIb	A

Continued

ACS, acute coronary syndrome; CKD, chronic kidney disease; HF, heart failure; ICD, implantable cardioverter defibrillator; LDL-C, low-density lipoprotein-cholesterol; LVEF, left ventricular ejection fraction; PCSK9, proprotein convertase subtilisin/kexin type 9; RAAS, renin–angiotensin–aldosterone system.
[a]Class of recommendation.
[b]Level of evidence.
[c]Increase statin potency/dose if the patient was on low-potency/low-dose statin, add ezetimibe if the patient was only on statins at highest tolerated dose, or add PCSK9 inhibitor if the patient was on statins plus ezetimibe.
[d]Angiotensin receptor blockers in cases of intolerance.

© ESC 2023

14. Patient perspectives

14.1. Patient-centred care

The management of patients with ACS should not only consider the best available evidence with regard to clinical management strategies, but also should be mindful of the provision of care that is respectful of and responsive to individual patient preferences, needs, and values, ensuring that these values are included in clinical decision-making.[881]

Patient-centred care should be guided by ethical values when considering a patient's physical, emotional, and psychological needs. Adopting a person-centred care approach after an ACS event improves patient outcomes and enhances quality of life.[882] Patients who are regarded as equal partners in their ACS medical management are more likely to actively engage and participate in their own healthcare.[883]

Educating and involving patients in their care should be seen as a continuous process. Engaging and educating the patient is a key component of ACS care and should take place throughout their patient journey, from admission to hospital discharge and cardiac rehabilitation (*Figure 19*).

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026



Figure 19 A person-centred approach to the ACS journey. ACS, acute coronary syndrome.

14.2. Shared decision-making

Shared decision-making is a process, during which the patient and a healthcare professional work together to make an informed decision about the patient's care.[884] During this process, information is provided, comprehension checked, and the patient is given an opportunity to ask questions in order to equip them with the tools needed to make an informed decision.

Using a shared decision-making approach during the consent process allows the patient's preferences to be established.[884] Discovery of the patient's concerns, goals, preferences, and values should be a central component of this process. The use of validated decision aids and audio-visual tools may also be helpful to facilitate informed consent and promote patient involvement.[884–887]

14.3. Informed consent

Informed consent should include the components listed in Supplementary data online, *Table S18*.[885,888] Informed consent is an opportunity to educate patients about the proposed procedure, the associated risks and benefits, and any available alternative interventions or treatments.[886,887] Assessment of the patient's understanding of the information given to them during the informed consent process using the 'teach back' technique should be considered (Supplementary data online, *Figure S6*).[885,889–891] The teach back method assesses understanding by asking patients to state in their own words what they need to know or do about their health.

Informed consent is an ethical and legal obligation for medical practitioners and is required before any invasive procedure. The information

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

should be provided in a simple and clear format. In patients undergoing emergency invasive angiography, a shortened informed consent process is appropriate. If a shortened informed consent process has been used, it is important that there is contact with the patient and/or family member after the intervention when the patient is physically and psychologically stable or following the death of the patient.[892] Further information can be found in the Supplementary data online.

14.4. Research participation and consent in the acute setting

With unstable ACS patients, it is often challenging to obtain their consent for emergency procedures—and even more challenging to enrol in clinical trials due to a number of factors, including the necessity for prompt clinical care, increased pain and stress levels, and impairment of consciousness. Where clinical trials are conducted, patient involvement in enrolment decisions is paramount, if possible.[893,894] A short witnessed verbal consent, followed by written consent after the acute phase, has been shown to be less stressful and more positively received than written consent in the acute setting.[894] The research and consent process must follow the ethical and legal requirements in the relevant country. Further information can be found in the Supplementary data online.

14.5. Patient satisfaction and expectations

Focusing healthcare around the needs and preferences of patients has the potential to improve clinical outcomes, quality of care and patient satisfaction, while decreasing healthcare costs and health disparities.[881] Patient perception of care is built on interpersonal interactions, the quality of clinical communication, delivery of care, and the administrative management of care. ACS patient expectations are summarized in *Figure 20* and further information can be found in the Supplementary data online, *Table S19*.




Figure 20 Acute coronary syndrome patient expectations. ACS, acute coronary syndrome.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

14.6. Patient-reported outcome measures and patient-reported experience measures

Understanding and measuring patient expectations and health outcomes using patient-reported outcome measures (PROMs) and patient-reported experience measures (PREMs) is central to improving patient satisfaction and delivering patient-centred care.[895] The quality of care for ACS patients should be measured during the patient's journey from initial presentation until discharge. Further information can be found in the Supplementary data online. Further information on PROMs and PREMs is also provided in the Supplementary data online.

14.7. Preparation for discharge

Many ACS patients may not be fully aware of what has happened to them and how to best manage their healthcare after discharge, leading to them both wanting and needing more information upon discharge.[896] Cognitive impairment can occur as a complication of ACS and some patients may have difficulty with instructions for care when transitioning towards discharge home.[897] Therefore, discharge information should be provided in both verbal and written formats and should include a discharge letter outlining the key components of the evidence-based discharge plan (Supplementary data online, *Table S20*).[898–901] Some important messages aimed at patients on how to improve their heart health after ACS are demonstrated in Supplementary data online, *Figure S5*. Moreover, following an ACS event, anxiety and depression are frequently encountered and confer an increased risk of non-adherence to medications and lifestyle changes, subsequent MACE, and death.[902–904] Non-adherence also generally increases over time, which has additional impact on clinical outcomes.[905] Assessing and identifying these patients and intervening with onward psychological referral is recommended.[858] Further information can be found in the Supplementary data online. A summary of patient concerns and educational needs throughout their ACS journey is also provided in Supplementary data online, *Figure S7*.

Recommendation Table 17 — Recommendations for patient perspectives in acute coronary syndrome care

Recommendations	Class[a]	Level[b]
Patient-centred care is recommended by assessing and adhering to individual patient preferences, needs and beliefs, ensuring that patient values are used to inform all clinical decisions.[744,881,906,907]	I	B
It is recommended to include ACS patients in decision-making (as much as their condition allows) and to inform them about the risk of adverse events, radiation exposure, and alternative options. Decision aids can be used to facilitate the discussion.[908,909]	I	B
It is recommended to assess symptoms using methods that help patients to describe their experience.[910]	I	C
Use of the 'teach back' technique for decision support during the securing of informed consent should be considered.[885,889–891]	IIa	B
Patient discharge information should be provided in both written and verbal formats prior to discharge. Adequate preparation and education for patient discharge using the teach back technique and/or motivational interviewing, giving information in chunks, and checking for understanding should be considered.[885,896,911]	IIa	B
Assessment of mental well-being using a validated tool and onward psychological referral when appropriate should be considered.[903,904,912,913]	IIa	B

© ESC 2023

ACS, acute coronary syndrome.
[a]Class of recommendation.
[b]Level of evidence.

15. Key messages

Epidemiology of ACS

Acute coronary syndromes encompass a spectrum of conditions that include patients with a recent change in clinical symptoms or signs, with or without changes on 12-lead ECG and with or without acute elevations in cardiac troponin concentrations. ACS are commonly classified based on ECG at presentation and the presence or absence of troponin elevation into UA, NSTEMI, or STEMI. The incidence of STEMI is decreasing whereas the incidence of NSTEMI is increasing. While there are some sex differences in the epidemiology of ACS, women and men receive equal benefit from invasive and non-invasive management strategies and, in general, should be managed similarly.

Diagnostic tools (ECG, troponin, and non-invasive imaging)

Chest pain/discomfort is the most common symptom initiating the ACS diagnostic and therapeutic pathway. High-sensitivity troponin measurements and rapid 'rule-in' and 'rule-out' algorithms should be used in patients with suspected NSTE-ACS. MI is not the only condition resulting in cardiomyocyte injury and cardiac troponin elevation, and other conditions should also be considered in the differential diagnosis. Non-invasive imaging can be useful to increase diagnostic accuracy and optimize risk assessment.

STEMI management networks

Co-ordination between EMS and hospitals with common written protocols is central to the management of STEMI. EMS should transfer patients immediately to 24/7 high-volume PCI centres regardless of the initial treatment strategy (PPCI or pre-hospital fibrinolysis). EMS should always alert the PCI centre immediately after selection of the reperfusion strategy, and patient transfer to the PCI centre should bypass the ED.

Invasive strategy and reperfusion therapy

An invasive strategy is recommended for patients with ACS. Invasive strategies are time sensitive. For STEMI and very high-risk NSTE-ACS, an immediate invasive strategy is recommended. For patients with NSTE-ACS an inpatient invasive strategy is recommended; in NSTE-ACS patients with high-risk characteristics, an early invasive strategy (<24 h) should be considered. If timely (within 120 min from time of diagnosis) PPCI cannot be performed in patients with STEMI, fibrinolytic therapy is indicated within 12 h of symptom onset in patients without contraindications.

Antithrombotic therapy

Antithrombotic therapy is indicated in all ACS patients, regardless of the management strategy. This consists of both antiplatelet and anticoagulant therapy. Aspirin is recommended for all ACS patients at an initial loading dose and a longer-term maintenance dose. In addition to aspirin, a $P2Y_{12}$ receptor inhibitor is recommended, and should be maintained over 12

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

months unless there are concerns regarding HBR. Regarding $P2Y_{12}$ receptor inhibitor choice, prasugrel and ticagrelor are recommended in preference to clopidogrel, and prasugrel should be considered in preference to ticagrelor for ACS patients who undergo PCI. Pre-treatment (i.e. treatment with a $P2Y_{12}$ receptor inhibitor prior to coronary angiography) in patients with NSTE-ACS is not recommended routinely but may be considered for patients with STEMI undergoing PPCI. Parenteral anticoagulation is recommended for all patients at the time of diagnosis. Discontinuation of parenteral anticoagulation should be considered immediately after the invasive procedure. Some patients with ACS will also have an indication for long-term OAC, most commonly AF. In these patients, TAT for up to 1 week, followed by DAT using a NOAC at the recommended dose for stroke prevention and a single oral antiplatelet agent (preferably clopidogrel), is recommended as the default strategy.

ACS with unstable presentation

A PPCI strategy is recommended in patients with resuscitated cardiac arrest and an ECG with persistent ST elevation (or ST elevation equivalents), whereas routine immediate angiography is not recommended in patients with an ECG without persistent ST elevation (or equivalents). Temperature control (i.e. continuous monitoring of core temperature and active prevention of fever [i.e. >37.7°C]) is recommended in patients with OHCA who remain unresponsive after ROSC. In patients with CS complicating ACS, emergency coronary angiography is recommended, whereas the routine use of IABP in ACS patients with CS and no mechanical complications is not.

Early care

Following reperfusion, it is recommended to admit high-risk ACS patients, including all STEMI patients, to a CCU/ICCU. ECG monitoring for arrhythmias and ST-segment changes is recommended for at least 24 h after symptom onset in all high-risk patients with ACS. It is recommended that all hospitals participating in the care of high-risk ACS patients have an ICCU/CCU equipped to provide all required aspects of care including treatment of ischaemia, severe HF, arrhythmias, and common comorbidities. It is also recommended that the LVEF is determined before hospital discharge in all patients with ACS. Discharge of high-risk ACS patients within 48–72 h should be considered in selected patients if early rehabilitation and adequate follow-up are arranged.

Technical aspects during PPCI

Routine radial access and use of DES are the standard of care during PCI for ACS. Intravascular imaging should be considered to guide PCI and may be considered in patients with ambiguous culprit lesions. Routine thrombus aspiration is not recommended. CABG should be considered in patients with an occluded IRA when PCI is not feasible or unsuccessful and there is a large area of myocardium in jeopardy. In patients presenting with SCAD, PCI is recommended only for patients with symptoms and signs of ongoing myocardial ischaemia, a large area of myocardium in jeopardy, and reduced antegrade flow.

Management of patients with MVD

For patients with MVD, it is recommended to base the revascularization strategy (IRA PCI, multivessel PCI/CABG) on the patient's clinical status and comorbidities, as well as their disease complexity, according to the principles of management of myocardial revascularization. For patients with MVD presenting with CS, IRA-only PCI during the index procedure is recommended. For patients with STEMI undergoing PPCI, complete revascularization is recommended either during the index PCI or within 45 days. In patients presenting with NSTE-ACS and MVD, complete revascularization should be considered, preferably during the index procedure. For patients with STEMI, it is recommended that decisions regarding PCI of non-IRA are based on angiographic severity, whereas for patients with NSTE-ACS, functional invasive evaluation of non-IRA severity during the index procedure may be considered.

MINOCA

The term MINOCA refers to the situation where patients present with symptoms suggestive of ACS and demonstrate troponin elevation and non-obstructive coronary arteries at the time of coronary angiography, i.e. coronary artery stenosis <50% in any major epicardial vessel. MINOCA is best considered as a working diagnosis that encompasses a heterogenous group of underlying causes (both cardiac and extra-cardiac) and is found in 1–14% of patients with ACS. In all patients with an initial working diagnosis of MINOCA, it is recommended to follow a diagnostic algorithm to determine the underlying cause. CMR imaging is a key diagnostic tool in patients with a working diagnosis of MINOCA.

Special patient subsets

Chronic kidney disease: moderate to severe CKD is present in >30% of ACS patients. These patients receive less interventional and pharmacological treatment and have a worse prognosis in comparison to patients with normal kidney function. It is recommended to apply the same diagnostic and therapeutic strategies in patients with CKD (dose adjustment may be necessary) as for patients with normal kidney function.

Older adults: in general, older adults should undergo the same diagnostic and treatment strategies, including invasive angiography and revascularization, as younger patients.

Patients with cancer: management of ACS in patients with cancer can be challenging for several reasons, including frailty, increased bleeding risk, thrombocytopaenia, and increased thrombotic risk. An invasive strategy is recommended in cancer patients presenting with high-risk ACS with expected survival ≥6 months. A conservative non-invasive strategy should be considered in ACS patients with poor cancer prognosis (with expected survival <6 months) and/or very high bleeding risk.

Long-term treatment

Secondary prevention after ACS should be offered to every patient and should start as early as possible after the index event. This includes cardiac rehabilitation, lifestyle management, and pharmacological treatment, and has been shown to both increase quality of life and decrease morbidity and mortality.

Patient perspectives

Some of the key first steps in the timely diagnosis and treatment of ACS are reliant on a comprehensive assessment of symptoms. An incomplete history or poorly elicited symptoms can result in delay or misdiagnosis. Patient-centred care is recommended as a critical tenet of routine clinical management and involves consideration of a patient's physical, emotional, and psychological needs.

The provision of care that is respectful of, and responsive to, individual patient preferences, needs and values, is important in the management of patients with ACS. It is recommended, as much as possible, to include ACS patients in decision-making. Preparing for discharge begins on admission. Educating and informing the patient using the teach back method and educationally appropriate material should be integrated into the patient care pathway.

Quality indicators

Acute coronary syndrome QIs aim to audit practice and improve clinical outcomes in real-life patients by demonstrating the gap between optimal guideline-based treatment and actual care of ACS patients. Subsequent measures to improve QI attainment can be implemented based on the local, regional, and global assessment of QIs.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

16. Gaps in evidence

Table 8 Gaps in evidence

	Section	Gaps in evidence	Research recommendations to address these gaps
3	**Triage and diagnosis**	• Observe zone: how can we improve the guidance for and management of patients assigned to the observe zone of the 0 h/1 h and 0 h/2 h ESC algorithms to improve their poor outcome? • No testing rule: what is the added value of biomarkers other than hs-cTn for rapid rule-out of NSTE-ACS compared with usual care? • There is insufficient evidence to set sex-specific thresholds for troponin levels. • The role of non-invasive anatomy (e.g. CCTA) or functional imaging (e.g. stress testing strategies) for low-risk NSTE-ACS patients should be further evaluated.	• Observe zone: prospectively evaluate changes in the 0 h/1 h and 0 h/2 h ESC algorithms to improve the outcomes of patients assigned to the observe zone. • No testing rule: randomization of patients to strategies with and without new biomarkers to evaluate whether their use improves clinical outcomes. • Prospectively evaluate the impact of using sex-specific cut-offs on the diagnosis, treatment, and outcomes of patients presenting to the ED with suspected ACS. • Adequately powered RCTs testing whether non-invasive imaging improves clinical outcomes in patients presenting with NSTE-ACS.
4	**Initial measures for patients presenting with suspected STEMI | Initial treatment**	• The impact of early i.v. beta-blockers on clinical outcomes in patients with a working diagnosis of STEMI remains unclear. • Infarct size and microvascular obstruction are the main determinants of long-term prognosis. Interventions which serve to limit infarct size are needed.	• Patients randomized to i.v. beta-blockers (ideally metoprolol) or placebo before PPCI, with hard endpoints evaluated. • Translate cardio-protective therapies from experimental to clinical setting by executing adequately powered trials.
5	**Acute-phase management of patients with NSTE-ACS**	• The comparison of routine or selective invasive assessment in low-risk NSTE-ACS has not been adequately evaluated. • The optimal timing of invasive angiography in high-risk NSTE-ACS patients remains uncertain.	• Low-risk patients should be randomized to routine or selective invasive strategy. • RCTs testing different time intervals to perform angiography within the 72 h window after the initial presentation.
6	**Antithrombotic therapy**	• Whether pre-treatment with oral $P2Y_{12}$ receptor inhibitors prior to ICA improves clinical outcomes in NSTE-ACS patients is uncertain. • Whether platelet function testing or genetic testing to guide de-escalation of oral $P2Y_{12}$ receptor inhibitors after the first month of therapy following PCI improves clinical management and outcomes remains unclear. • The optimal long-term antithrombotic regimen in NSTE-ACS patients who have undergone PCI is unknown. • After stopping DAPT, a head-to-head comparison based on superiority between aspirin monotherapy and clopidogrel monotherapy is required.	• Randomize patients to pre-treatment with oral $P2Y_{12}$ receptor inhibitors or no pre-treatment, prior to ICA. • Randomize ACS patients to prasugrel or ticagrelor, both without pre-treatment. • A strategy based on platelet function testing or genetic testing should be prospectively tested in patients who may benefit from de-escalating antithrombotic therapy. • RCTs evaluating the benefit-risk balance for ischaemic bleeding events for different periods of antithrombotic duration. • A head-to-head randomized comparison testing for superiority is needed to compare aspirin vs. clopidogrel monotherapy after DAPT.
7	**Acute coronary syndrome with unstable presentation**	• The role of percutaneous MCS devices in patients presenting with ACS and CS remains unclear.	• Randomized comparisons between standard of care and percutaneous MCS devices in ACS with CS.
8	**Management of acute coronary syndrome during hospitalization**	• Clinical improvement through the use of risk stratification based on risk prediction models.	• Patients randomized to a particular intervention or to usual care based on validated risk prediction models.
9	**Technical aspects of invasive strategies**	• Does intravascular imaging-guided revascularization strategy improve clinical outcomes in patients with ACS? • Does intracoronary physiology assessment of myocardial reperfusion after PPCI improve risk stratification and/or stratified medicine for limiting microvascular dysfunction and reperfusion injury/MVO post ACS?	• RCTs evaluating the efficacy of an intravascular imaging-guided revascularization strategy to improve meaningful clinical outcomes in patients with ACS. • Prospectively evaluate whether intracoronary physiology assessment of myocardial reperfusion better stratifies patient risk.

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

		• In ACS patients with an IRA that is unsuitable for stent implantation, does drug-coated balloon treatment of the IRA improve clinical outcomes? • Microvascular obstruction associated with PPCI represents an unmet clinical need in patients with ACS. Development of therapies for the prevention and treatment of MVO is urgently needed. • Does early implementation of MCS in the management of high-risk ACS patients improve clinical outcomes? • Does intracoronary hypothermia reduce infarct size and improve clinical outcomes in STEMI patients undergoing PPCI? • What is the optimal antiplatelet strategy in patients presenting with SCAD? Specific gaps in knowledge surround antithrombotic treatment in the acute and post-ACS periods, including the optimal combination and duration of treatment.	• Patients with an IRA unsuitable for stent implantation randomized to drug-coated balloon treatment or usual care to evaluate clinical outcomes. • Pre-clinical and clinical research is needed to evaluate cardio-protective therapies aimed at reducing microvascular obstruction. • RCTs evaluating the benefit of using MCS in high-risk patients. • Randomized trials are needed to demonstrate both whether intracoronary hypothermia reduces myocardial infarct size, and if this translates into clinical improvement. • RCTs evaluating several antiplatelet strategies in patients with SCAD with the aim of determining which results in the greatest clinical benefit.
10	**Management of patients with multivessel disease**	• Does complete revascularization of NSTE-ACS with multivessel CAD improve clinical outcomes vs. culprit-only PCI? • Does management of non-infarct-related CAD with intravascular imaging guidance to identify rupture-prone atherosclerotic plaque improve clinical outcomes? • Does FFR-guided management improve clinical outcomes vs. standard angiography-guided management in NSTE-ACS? • What is the optimal timing of coronary revascularization (immediate vs. index hospitalization vs. staged) for non-IRA revascularization in STEMI and NSTE-ACS? • Does intensive medical therapy improve outcomes in patients with MVD compared with standard secondary prevention? • The clinical utility of hybrid coronary revascularization in ACS patients with multivessel CAD is uncertain.	• Patients with NSTE-ACS and MVD randomized to complete vs. culprit-only PCI. • RCTs testing whether the use of intravascular imaging to guide the management of non-infarct-related lesions improves clinical outcomes. • Patients randomized to FFR-guided management vs. standard angiography-guided management in NSTE-ACS. • Three-arm study comparing the clinical benefits of immediate, in-hospital and staged coronary revascularization strategies. • Patients with MVD randomized to intensive secondary prevention vs. usual care to evaluate whether the former strategy improves clinical outcomes. • RCTs assessing the clinical benefit of hybrid revascularization.
12	**Special situations**	• How to better differentiate Type 2 from Type 1 MI before invasive assessment. • The optimal management strategy in older adults with NSTE-ACS is not known. • The optimal management strategy in older frail, comorbid adults with NSTE-ACS is not known. • The optimal management strategy in older frail, comorbid adults with STEMI is not known. • Optimal antiplatelet therapy and its duration to manage ACS in pregnant patients are not known. • The optimal management strategy for pregnant women with NSTE-ACS is not known. • There is a need to further evaluate the contribution of social determinants of health.	• Prospective evaluation of diagnostic strategies aimed at better classifying patients according to their type of MI (Type 1 vs. Type 2). • Further studies recruiting older adults should be conducted to evaluate whether the current standard of care also benefits this subset of patients. • Older frail, comorbid patients should not be systematically excluded from RCTs. • Prospective data are needed to better understand which antiplatelet therapy regimen is best for pregnant women. • Observational data are needed in patients with ACS to evaluate the real impact of social determinants of health on clinical outcomes. Randomized interventions aimed at reducing social inequalities are needed to evaluate how to reduce this gap.
13	**Long-term treatment**	• To evaluate the uptake, safety, and outcomes for alternative forms of cardiac rehabilitation, with a focus on telemedicine and eHealth. • How to improve referral for and uptake of CR, especially for groups with low participation, including women, older persons, and ethnic minorities.	• Remote cardiac rehabilitation methods need randomized data to evaluate their true potential. • Further monitoring is needed to increase the participation of historically under-represented patients in CR.

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

		• The role of personalized medicine in the short- and long-term treatment of ACS needs to be further studied. • How to address additional risk from non-traditional risk factors, e.g. cardio-obstetrics, cardio-oncology, and inflammatory conditions, needs further attention. • Inflammation as a treatment target in patients with atherosclerosis still needs unravelling, as well as the use of biomarkers of inflammation (high-sensivity C-reactive protein, interleukins 1 and 6) to guide treatment of residual risk. • The role of lipoprotein (a) in guiding treatment and as an independent treatment target needs to be studied further. • The added cardio-protective role of beta-blockers in post-ACS patients without reduced LVEF on otherwise optimal medical therapy needs to be clarified. • The added cardio-protective role of ACE inhibitors/ARBs in post-ACS patients without reduced LVEF on otherwise optimal medical therapy needs to be clarified. • The future role of new treatment options, using mRNA- and siRNA-based therapies targeting lipid metabolism and inflammation, needs to be explored. • It has to be determined whether SGLT2 inhibitors—in the specific group of patients with ACS without heart failure or diabetes—improve clinical outcomes, regardless of diabetes status.	• Patients randomized to personalized strategies vs. usual care are needed to determine the role of precision medicine in ACS. • Prospective cohorts are needed to evaluate non-traditional risk factors and residual risk. • RCTs testing whether management based on the use of biomarkers of inflammation improves clinical outcomes. • RCTs testing whether lipoprotein (a) measurement to guide medical management further improves clinical outcomes. • Patients randomized to beta-blocker and no beta-blocker use to evaluate treatment efficacy in patients with ACS and LVEF >40%. • RCTs evaluating the benefit of using ACE inhibitors/ARBs vs. placebo on top of standard care in ACS patients with LVEF >40%. • Randomized data are needed to evaluate the role of mRNA- and siRNA-based therapies in the current context of lipid management and lipid targets. • ACS patients without HF or diabetes should be randomized to SGLT2 inhibitors vs. standard of care.
14	**Patient perspectives**	• The feasibility of performing short witnessed verbal consent followed by written consent after the acute phase needs further evaluation. • There is a need to assess the contribution of social determinants of health on ACS incidence and prognosis. • The use of validated patient-reported outcome and experience measures in evidence-based medicine should be increased. • Quality of life is a relevant outcome not captured in most trials. • Use of validated decision aids and audio-visual tools can be helpful to make informed choices that consider patients' values and preferences and promote patient involvement.	• Studies comparing verbal vs. written consent to evaluate safety endpoints and any ethical concerns. • The influence of social determinants of health on clinical outcomes should be evaluated, as well as those interventions aimed at reducing social inequalities. • PROMs/PREMs should have a more prominent role in RCTs evaluating patients with ACS. • Include quality of life as a prominent outcome in clinical trials. • Testing the use of validated decision aids and audio-visual tools to improve decisions around informed choices.
19	**Quality indicators**	• There is a lack of implementation studies evaluating whether prospectively monitoring and reporting ESC QIs for ACS improve clinical outcomes.	• Implementation studies evaluating a quality of care programme based on the evaluation of ESC QIs for ACS.
	General	• Patients included in clinical trials represent a relatively small proportion of real-life patients.	• Conduct clinical trials that enrol more representative patient populations (e.g. pragmatic clinical trials).

Ongoing trials addressing some of these gaps in evidence are presented in the Supplementary data online.

ACE, angiotensin-converting enzyme; ACS, acute coronary syndrome; ARB, angiotensin receptor blocker; CAD, coronary artery disease; CCTA, coronary computed tomography angiography; CR, cardiac rehabilitation; CS, cardiogenic shock; DAPT, dual antiplatelet therapy; ED, emergency department; ESC, European Society of Cardiology; FFR, fractional flow reserve; HF, heart failure; ICA, invasive coronary angiography; IRA, infarct-related artery; i.v., intravenous; LVEF, left ventricular ejection fraction; MCS, mechanical circulatory support; MI, myocardial infarction; MINOCA, myocardial infarction with non-obstructive coronary arteries; mRNA, messenger ribonucleic acid; MVD, multivessel disease; MVO, microvascular obstruction; NSTE-ACS, non-ST-elevation acute coronary syndrome; PCI, percutaneous coronary intervention; PPCI, primary percutaneous coronary intervention; PREM, patient-reported experience measure; PROM, patient-reported outcome measure; QI, quality indicator; RCT, randomized controlled trial; SCAD, spontaneous coronary artery dissection; SGLT2, sodium-glucose co-transporter 2; siRNA, small interfering ribodeoxynucleic acid; STEMI, ST-elevation myocardial infarction.

17. Sex differences

There are currently no data supporting the differential management of ACS based on sex. However, several studies have reported that women presenting with ACS are treated differently than men.[914–918] This includes being less likely than men to receive ICA, timely revascularization, CR, and secondary prevention medications.[914–918]

Healthcare providers and policymakers should be conscious of this potential gender bias in the management of ACS and make a concerted effort to ensure that women with ACS receive evidence-based care.

In order to ensure the generalizability of the findings yielded by RCTs, patient recruitment should be reflective of real-world populations from different socioeconomic backgrounds.[919] Several studies have reported that a disproportionately low proportion of women are recruited to CV trials.[920–922] Alongside historic under-representation of other subsets of patients, including older patients and ethnic minorities, this suggests an underlying recruitment bias.[923] Increased representation of female patients in future clinical trials is required to better inform the optimal management of women with ACS.[924]

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

© ESC 2023

18. 'What to do' and 'What not to do' messages from the Guidelines

Table 9 'What to do' and 'What not to do'

Recommendations for clinical and diagnostic tools for patients with suspected acute coronary syndrome	Class[a]	Level[b]
It is recommended that patients with suspected STEMI are immediately triaged for an emergency reperfusion strategy.	I	A
It is recommended to base the diagnosis and initial short-term risk stratification of ACS on a combination of clinical history, symptoms, vital signs, other physical findings, ECG, and hs-cTn.	I	B
Twelve-lead ECG recording and interpretation is recommended as soon as possible at the point of first medical contact, with a target of <10 min.	I	B
Continuous ECG monitoring and the availability of defibrillator capacity is recommended as soon as possible in all patients with suspected STEMI, in suspected ACS with other ECG changes or ongoing chest pain, and once the diagnosis of MI is made.	I	B
The use of additional ECG leads (V3R, V4R, and V7–V9) is recommended in cases of inferior STEMI or if total vessel occlusion is suspected and standard leads are inconclusive.	I	B
An additional 12-lead ECG is recommended in cases with recurrent symptoms or diagnostic uncertainty.	I	C
It is recommended to measure cardiac troponins with high-sensitivity assays immediately after presentation and to obtain the results within 60 min of blood sampling.	I	B
It is recommended to use an ESC algorithmic approach with serial hs-cTn measurements (0 h/1 h or 0 h/2 h) to rule in and rule out NSTEMI.	I	B
Additional testing after 3 h is recommended if the first two hs-cTn measurements of the 0 h/1 h algorithm are inconclusive and no alternative diagnoses explaining the condition have been made.	I	B
Recommendations for non-invasive imaging in the initial assessment of patients with suspected acute coronary syndrome		
Emergency TTE is recommended in patients with suspected ACS presenting with cardiogenic shock or suspected mechanical complications.	I	C
Routine, early coronary computed tomography angiography in patients with suspected ACS is not recommended.	III	B
Recommendations for the initial management of patients with acute coronary syndrome		
It is recommended that the pre-hospital management of patients with a working diagnosis of STEMI is based on regional networks designed to deliver reperfusion therapy expeditiously and effectively, with efforts made to make PPCI available to as many patients as possible.	I	B
It is recommended that PPCI-capable centres deliver a 24/7 service and are able to perform PPCI without delay.	I	B
It is recommended that patients transferred for PPCI bypass the emergency department and CCU/ICU and are transferred directly to the catheterization laboratory.	I	B
Oxygen is recommended in patients with hypoxaemia (SaO_2 <90%).	I	C
It is recommended that EMS transfer patients with suspected STEMI to a PCI-capable centre, bypassing non-PCI centres.	I	C
It is recommended that ambulance teams are trained and equipped to identify ECG patterns suggestive of acute coronary occlusion and to administer initial therapy, including defibrillation, and fibrinolysis when applicable.	I	C
It is recommended that all hospitals and EMS participating in the care of patients with suspected STEMI record and audit delay times and work together to achieve and maintain quality targets.	I	C
Routine oxygen is not recommended in patients with oxygen saturation >90%.	III	A
Recommendations for reperfusion therapy and timing of invasive strategy		
Recommendations for reperfusion therapy for patients with STEMI		
Reperfusion therapy is recommended in all patients with a working diagnosis of STEMI (persistent ST-segment elevation or equivalents) and symptoms of ischaemia of ≤12 h duration.	I	A
A PPCI strategy is recommended over fibrinolysis if the anticipated time from diagnosis to PCI is <120 min.	I	A
If timely PPCI (<120 min) cannot be performed in patients with a working diagnosis of STEMI, fibrinolytic therapy is recommended within 12 h of symptom onset in patients without contraindications.	I	A
Rescue PCI is recommended for failed fibrinolysis (i.e. ST-segment resolution <50% within 60–90 min of fibrinolytic administration) or in the presence of haemodynamic or electrical instability, worsening ischaemia, or persistent chest pain.	I	A

Continued

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

In patients with a working diagnosis of STEMI and a time from symptom onset >12 h, a PPCI strategy is recommended in the presence of ongoing symptoms suggestive of ischaemia, haemodynamic instability, or life-threatening arrhythmias.	I	C
Routine PCI of an occluded IRA is not recommended in STEMI patients presenting >48 h after symptom onset and without persistent symptoms.	III	A
Transfer/interventions after fibrinolysis		
Transfer to a PCI-capable centre is recommended in all patients immediately after fibrinolysis.	I	A
Emergency angiography and PCI of the IRA, if indicated is recommended in patients with new-onset or persistent heart failure/shock after fibrinolysis.	I	A
Angiography and PCI of the IRA, if indicated, is recommended between 2 and 24 h after successful fibrinolysis.	I	A
Invasive strategy in NSTE-ACS		
An invasive strategy during hospital admission is recommended in NSTE-ACS patients with high-risk criteria or with a high index of suspicion for unstable angina.	I	A
A selective invasive approach is recommended in patients without very high- or high-risk features and a low index of suspicion for NSTE-ACS.	I	A
An immediate invasive strategy is recommended in patients with a working diagnosis of NSTE-ACS and with at least one of the following very high-risk criteria: • Haemodynamic instability or cardiogenic shock • Recurrent or refractory chest pain despite medical treatment • In-hospital life-threatening arrhythmias • Mechanical complications of MI • Acute heart failure presumed secondary to ongoing myocardial ischaemia *Recurrent* dynamic ST-segment or T wave changes, particularly intermittent ST-segment elevation.	I	C
Recommendations for antiplatelet and anticoagulant therapy in acute coronary syndrome		
Antiplatelet therapy		
Aspirin is recommended for all patients without contraindications at an initial oral LD of 150–300 mg (or 75–250 mg i.v.) and an MD of 75–100 mg o.d. for long-term treatment.	I	A
In all ACS patients, a $P2Y_{12}$ receptor inhibitor is recommended in addition to aspirin, given as an initial oral LD followed by an MD for 12 months unless there is high bleeding risk.	I	A
A proton pump inhibitor in combination with dual antiplatelet therapy is recommended in patients at high risk of gastrointestinal bleeding.	I	A

Continued

Prasugrel is recommended in $P2Y_{12}$ receptor inhibitor-naïve patients proceeding to PCI (60 mg LD, 10 mg o.d. MD, 5 mg o.d. MD for patients aged ≥75 years or with a body weight <60 kg).	I	B
Ticagrelor is recommended irrespective of the treatment strategy (invasive or conservative) (180 mg LD, 90 mg twice a day MD).	I	B
Clopidogrel (300–600 mg LD, 75 mg o.d. MD) is recommended when prasugrel or ticagrelor are not available, cannot be tolerated, or are contraindicated.	I	C
If patients presenting with ACS stop DAPT to undergo CABG, it is recommended they resume DAPT after surgery for at least 12 months.	I	C
Pre-treatment with a glycoprotein IIb/IIIa antagonist is not recommended.	III	A
Routine pre-treatment with a $P2Y_{12}$ receptor inhibitor in NSTE-ACS patients in whom coronary anatomy is not known and early invasive management (<24 h) is planned is not recommended.	III	A
Anticoagulant therapy		
Parenteral anticoagulation is recommended for all patients with ACS at the time of diagnosis.	I	A
Routine use of a UFH bolus (weight-adjusted i.v. bolus during PCI of 70–100 IU/kg) is recommended in patients undergoing PCI.	I	C
Patients with STEMI		
Fondaparinux is not recommended in patients with STEMI undergoing PPCI.	III	B
Patients with NSTE-ACS		
For patients with NSTE-ACS in whom early invasive angiography (i.e. within 24 h) is not anticipated, fondaparinux is recommended.	I	B
Combining antiplatelets and OAC		
As the default strategy for patients with atrial fibrillation and CHA_2DS_2-VASc score ≥1 in men and ≥2 in women, after up to 1 week of triple antithrombotic therapy following the ACS event, dual antithrombotic therapy using a non-vitamin K antagonist oral anticoagulant at the recommended dose for stroke prevention and a single oral antiplatelet agent (preferably clopidogrel) for up to 12 months is recommended.	I	A
During PCI, a UFH bolus is recommended in any of the following circumstances: • if the patient is on a NOAC • if the INR is <2.5 in VKA-treated patients.	I	C
The use of ticagrelor or prasugrel as part of triple antithrombotic therapy is not recommended.	III	C

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Recommendations for alternative antithrombotic therapy regimens		
Discontinuation of antiplatelet treatment in patients treated with an oral anticoagulant is recommended after 12 months.	I	B
De-escalation of antiplatelet therapy in the first 30 days after ACS is not recommended.	III	B
Recommendations for fibrinolytic therapy		
When fibrinolysis is the reperfusion strategy, it is recommended to initiate this treatment as soon as possible after diagnosis in the pre-hospital setting (aim for target of <10 min to lytic bolus).	I	A
A fibrin-specific agent (i.e. tenecteplase, alteplase, or reteplase) is recommended.	I	B
Antiplatelet co-therapy with fibrinolysis		
Aspirin and clopidogrel are recommended.	I	A
Anticoagulation co-therapy with fibrinolysis		
Anticoagulation is recommended in patients treated with fibrinolysis until revascularization (if performed) or for the duration of hospital stay (up to 8 days).	I	A
Enoxaparin i.v. followed by subcutaneous is recommended as the preferred anticoagulant.	I	A
When enoxaparin is not available, unfractionated heparin is recommended as a weight-adjusted i.v. bolus, followed by infusion.	I	B
Recommendations for cardiac arrest and out-of-hospital cardiac arrest		
A PPCI strategy is recommended in patients with resuscitated cardiac arrest and an ECG with persistent ST-segment elevation (or equivalents).	I	B
Temperature control (i.e. continuous monitoring of core temperature and active prevention of fever [i.e. >37.7°C]) is recommended after either out-of-hospital or in-hospital cardiac arrest for adults who remain unresponsive after return of spontaneous circulation.	I	B
Routine immediate angiography after resuscitated cardiac arrest is not recommended in haemodynamically stable patients without persistent ST-segment elevation (or equivalents).	III	A
Systems of care		
It is recommended that healthcare systems implement strategies to facilitate transfer of all patients in whom ACS is suspected after resuscitated cardiac arrest directly to a hospital offering 24/7 PPCI via one specialized EMS.	I	C
Evaluation of neurological prognosis		
Evaluation of neurological prognosis (no earlier than 72 h after admission) is recommended in all comatose survivors after cardiac arrest.	I	C

Continued

Recommendations for cardiogenic shock		
Immediate coronary angiography and PCI of the IRA (if indicated) is recommended in patients with CS complicating ACS.	I	B
Emergency CABG is recommended for ACS-related CS if PCI of the IRA is not feasible/ unsuccessful.	I	B
In cases of haemodynamic instability, emergency surgical/catheter-based repair of mechanical complications of ACS is recommended, based on Heart Team discussion.	I	C
The routine use of an intra-aortic balloon pump in ACS patients with CS and without mechanical complications is not recommended.	III	B
Recommendations for in-hospital management		
It is recommended that all hospitals participating in the care of high-risk patients have an ICCU/CCU equipped to provide all required aspects of care, including treatment of ischaemia, severe heart failure, arrhythmias, and common comorbidities.	I	C
It is recommended that high-risk patients (including all STEMI patients and very high-risk NSTE-ACS patients) have ECG monitoring for a minimum of 24 h.	I	C
It is recommended that high-risk patients with successful reperfusion therapy and an uncomplicated clinical course (including all STEMI patients and very high-risk NSTE-ACS patients) are kept in the CCU/ ICCU for a minimum of 24 h whenever possible, after which they may be moved to a step-down monitored bed for an additional 24–48 h.	I	C
Recommendations for technical aspects of invasive strategies		
Radial access is recommended as the standard approach, unless there are over-riding procedural considerations.	I	A
PCI with stent deployment in the IRA during the index procedure is recommended for patients undergoing PPCI.	I	A
Drug-eluting stents are recommended in preference to bare metal stents in all cases.	I	A
In patients with spontaneous coronary artery dissection, PCI is recommended only for patients with symptoms and signs of ongoing myocardial ischaemia, a large area of myocardium in jeopardy, and reduced antegrade flow.	I	C
The routine use of thrombus aspiration is not recommended.	III	A
Recommendations for management of patients with multivessel disease		
It is recommended to base the revascularization strategy (IRA PCI, multivessel PCI/CABG) on the patient's clinical status and comorbidities, as well as their disease complexity, according to the principles of management of myocardial revascularization.	I	B

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Multivessel disease in ACS patients presenting in cardiogenic shock		
IRA-only PCI during the index procedure is recommended.	I	B
Multivessel disease in haemodynamically stable STEMI patients undergoing PPCI		
Complete revascularization is recommended either during the index PCI procedure or within 45 days.	I	A
It is recommended that PCI of the non-IRA is based on angiographic severity.	I	B
Invasive epicardial functional assessment of non-culprit segments of the IRA is not recommended during the index procedure.	III	C
Recommendations for myocardial infarction with non-obstructive coronary arteries		
In patients with a working diagnosis of MINOCA CMR imaging is recommended after invasive angiography if the final diagnosis is not clear	I	B
Management of MINOCA according to the final established underlying diagnosis is recommended, consistent with the appropriate disease-specific guidelines.	I	B
In all patients with an initial working diagnosis of MINOCA, it is recommended to follow a diagnostic algorithm to determine the underlying final diagnosis.	I	C
Recommendations for acute coronary syndrome complications		
Atrial fibrillation		
Intravenous beta-blockers are recommended when rate control is needed in the absence of acute HF or hypotension.	I	C
Intravenous amiodarone is recommended when rate control is needed in the presence of acute HF and no hypotension.	I	C
Immediate electrical cardioversion is recommended in patients with ACS and haemodynamic instability and when adequate rate control cannot be achieved promptly with pharmacological agents.	I	C
Intravenous amiodarone is recommended to facilitate electrical cardioversion and/or decrease risk of early recurrence of AF after electrical cardioversion in unstable patients with recent-onset AF.	I	C
Ventricular arrythmias		
Implantable cardioverter defibrillator use is recommended to reduce sudden cardiac death in patients with symptomatic HF (NYHA Class II–III) and LVEF ≤35% despite optimal medical therapy for >3 months and at least 6 weeks after MI who are expected to survive for at least 1 year with good functional status.	I	A
Intravenous beta-blocker and/or amiodarone treatment is recommended for patients with polymorphic VT and/or VF unless contraindicated.	I	B
Prompt and complete revascularization is recommended to treat myocardial ischaemia that may be present in patients with recurrent VT and/or VF.	I	C
Bradyarrhythmias		
In cases of sinus bradycardia with haemodynamic intolerance or high-degree AV block without stable escape rhythm:		
• i.v. positive chronotropic medication (adrenaline, vasopressin, and/or atropine) is recommended. • temporary pacing is recommended in cases of failure to respond to atropine. • urgent angiography with a view to revascularization is recommended if the patient has not received previous reperfusion therapy.	I	C
Implantation of a permanent pacemaker is recommended when high-degree AV block does not resolve within a waiting period of at least 5 days after MI.	I	C
Pacing is not recommended if high-degree AV block resolves after revascularization or spontaneously.	III	B
Treatment of asymptomatic and haemodynamically irrelevant ventricular arrhythmias with anti-arrhythmic drugs is not recommended.	III	C
Recommendations for acute coronary syndrome comorbid conditions		
Chronic kidney disease		
The use of low- or iso-osmolar contrast media (at the lowest possible volume) is recommended for invasive strategies.	I	A
It is recommended to assess kidney function using eGFR in all patients with ACS.	I	C
It is recommended to apply the same diagnostic and therapeutic strategies in patients with CKD (dose adjustment may be necessary) as in patients with normal kidney function.	I	C
Diabetes		
It is recommended to base the choice of long-term glucose-lowering treatment on the presence of comorbidities, including heart failure, CKD, and obesity.	I	A
It is recommended to assess glycaemic status at initial evaluation in all patients with ACS.	I	B
It is recommended to frequently monitor blood glucose levels in patients with known diabetes mellitus or hyperglycaemia (defined as glucose levels ≥11.1 mmol/L or ≥200 mg/dL).	I	C
Older adults		
It is recommended to apply the same diagnostic and treatment strategies in older patients as in younger patients.	I	B

Continued

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

It is recommended to adapt the choice and dosage of antithrombotic agent, as well as of secondary prevention medications, to renal function, co-medications, comorbidities, frailty, cognitive function, and specific contraindications.	I	B
For frail older patients with comorbidities, a holistic approach is recommended to individualize interventional and pharmacological treatments after careful evaluation of the risks and benefits.	I	B
An invasive strategy is recommended in cancer patients presenting with high-risk ACS with expected life survival ≥6 months.	I	B
A temporary interruption of cancer therapy is recommended in patients in whom the cancer therapy is suspected to be a contributing cause of ACS.	I	C
Aspirin is not recommended in cancer patients with a platelet count <10 000/μL.	III	C
Clopidogrel is not recommended in cancer patients with a platelet count <30 000/μL.	III	C
In ACS patients with cancer and <50 000/μL platelet count, prasugrel or ticagrelor are not recommended.	III	C
Recommendations for long-term management		
It is recommended that all ACS patients participate in a medically supervised, structured, comprehensive, multidisciplinary exercise-based cardiac rehabilitation and prevention programme.	I	A
It is recommended that ACS patients adopt a healthy lifestyle, including: • stopping all smoking of tobacco • healthy diet (Mediterranean style) • alcohol restriction • regular aerobic physical activity and resistance exercise • reduced sedentary time	I	B
Pharmacological treatment		
Lipid-lowering therapy		
It is recommended that high-dose statin therapy is initiated or continued as early as possible, regardless of initial LDL-C values.	I	A
It is recommended to aim to achieve an LDL-C level of <1.4 mmol/L (<55 mg/dL) and to reduce LDL-C by ≥50% from baseline.	I	A
If the LDL-C goal is not achieved despite maximally tolerated statin therapy and ezetimibe after 4–6 weeks, the addition of a PCSK9 inhibitor is recommended	I	A
If the LDL-C goal is not achieved despite maximally tolerated statin therapy after 4–6 weeks, the addition of ezetimibe is recommended.	I	B
It is recommended to intensify lipid-lowering therapy during the index ACS hospitalization for patients who were on lipid-lowering therapy before admission.	I	C
Beta-blockers		
Beta-blockers are recommended in ACS patients with LVEF ≤40% regardless of HF symptoms.	I	A
RAAS system inhibitors		
Angiotensin-converting enzyme inhibitors are recommended in ACS patients with HF symptoms, LVEF ≤40%, diabetes, hypertension, and/or CKD.	I	A
Mineralocorticoid receptor antagonists are recommended in ACS patients with an LVEF ≤40% and HF or diabetes.	I	A
Imaging		
In patients with pre-discharge LVEF ≤40%, repeat evaluation of the LVEF 6–12 weeks after an ACS (and after complete revascularization and the institution of optimal medical therapy) is recommended to assess the potential need for sudden cardiac death primary prevention ICD implantation.	I	C
Vaccination		
Influenza vaccination is recommended for all ACS patients.	I	A
Recommendations for patient perspectives in acute coronary syndrome care		
Patient-centred care is recommended by assessing and adhering to individual patient preferences, needs and beliefs, ensuring that patient values are used to inform all clinical decisions.	I	B
It is recommended to include ACS patients in decision-making (as much as their condition allows) and to inform them about the risk of adverse events, radiation exposure, and alternative options. Decision aids should be used to facilitate the discussion.	I	B
It is recommended to assess symptoms using methods that help patients to describe their experience.	I	C

ACS, acute coronary syndrome; AV, atrioventricular; CABG, coronary artery bypass grafting; CCU, cardiac care unit; CHA_2DS_2-VASc, Congestive heart failure, Hypertension, Age ≥75 years, Diabetes mellitus, Stroke or transient ischaemic attack, Vascular disease; CKD, chronic kidney disease; CMR, cardiac magnetic resonance; CS, cardiogenic shock; DAPT, dual antiplatelet therapy; ECG, electrocardiogram; ESC, European Society of Cardiology; HF, heart failure; hs-cTn, high-sensitivity cardiac troponin; ICU, intensive care unit; IRA, infarct-related artery; i.v., intravenous; LD, loading dose; LDL-C, low-density lipoprotein cholesterol; MD, maintenance dose; MINOCA, myocardial infarction with non-obstructive coronary arteries; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSTE-ACS, non-ST-elevation acute coronary syndrome; NSTEMI, non-ST-elevation myocardial infarction; NYHA, New York Heart Association; o.d., once daily; OAC, oral anticoagulant; PCI, percutaneous coronary intervention; PCSK9, proprotein convertase subtilisin/kexin type 9; PPCI, primary percutaneous coronary intervention; RAAS, renin–angiotensin–aldosterone system; STEMI, ST-elevation myocardial infarction; UFH, unfractionated heparin; VKA, vitamin K antagonist.

[a]Class of recommendation.

[b]Level of evidence.

© ESC 2023

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

19. Quality indicators

Quality indicators are tools that can be used to evaluate care quality, including structures, processes, and outcomes of care.[925] They may also serve as a mechanism for enhancing adherence to guideline recommendations, through associated quality improvement initiatives and the benchmarking of care providers.[926,927] As such, the role of QIs in improving care and outcomes for CVD is increasingly recognized by healthcare authorities, professional organizations, payers, and the public.[925]

The ESC understands the need for measuring and reporting quality and outcomes of CV care and has established methods for the development of the ESC QIs for the quantification of care and outcomes for CVDs.[925] To date, the ESC has developed QI suites for a number of CVDs in parallel with the writing of the ESC Clinical Practice Guidelines. Previous QIs for the management of AMI have been tested in numerous large registries.[928–933] A systematic review of these studies has shown that there is room for improvement in terms of levels of attainment of QIs.[934]

The ESC aims to harmonize its QIs for various CV conditions and integrate them with ESC registries.[935,936] This integrative approach provides 'real-world' data about the patterns and outcomes of care for CVD across Europe.

20. Supplementary data

Supplementary data are available at *European Heart Journal* online.

21. Data availability statement

No new data were generated or analysed in support of this research.

22. Author information

Author/Task Force Member Affiliations: Xavier Rossello, Cardiology Department, Hospital Universitari Son Espases, Palma de Mallorca, Spain, Clinical Research Department Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain, Health Research Institute of the Balearic Islands (IdiSBa), Universitat de les Illes Balears (UIB), Palma de Mallorca, Spain; **J.J. Coughlan**, Cardiovascular Research Institute, Mater Private Network, Dublin, Ireland; **Emanuele Barbato**, Clinical and Molecular Medicine, Sapienza University, Rome, Italy; **Colin Berry**, British Heart Foundation, Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, United Kingdom, Cardiology, NHS Golden Jubilee, Clydebank, United Kingdom, Cardiology, NHS Greater Glasgow and Clyde Health Board, Glasgow, United Kingdom; **Alaide Chieffo**, IRCCS San Raffaele Scientific Institute, IRCCS San Raffaele Scientific Institute, Milan, Italy, University Vita-Salute San Raffaele, Milan, Italy; **Marc J. Claeys**, Cardiology, Antwerp University Hospital, Edegem, Belgium; **Gheorghe-Andrei Dan**, Colentina University Hospital, Cardiology Dpt., "Carol Davila" University of Medicine Bucharest, Romania; **Marc R. Dweck**, British Heart Foundation Centre for Cardiovascular Sciences, Chancellors Building, Little France Crescent, Little France, Edinburgh, United Kingdom; **Mary Galbraith** (United Kingdom), ESC Patient Forum, Sophia Antipolis, France; **Martine Gilard**, Cardiology, INSERM UMR 1304 GETBO- Brest University, Brest, France; **Lynne Hinterbuchner**, Department of Cardiology, Clinic of Internal Medicine II, Paracelsus Medical University of Salzburg, Salzburg, Austria; **Ewa A. Jankowska**, Institute of Heart Diseases, Wroclaw Medical University, Wroclaw, Poland, Institute of Heart Diseases, University Hospital in Wroclaw, Wroclaw, Poland; **Peter Jüni**, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom; **Takeshi Kimura**, Department of Cardiology, Hirakata Kohsai Hospital, Osaka, Japan; **Vijay Kunadian**, Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom, Cardiothoracic Centre, Freeman Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, United Kingdom; **Margret Leosdottir**, Department of Cardiology, Skane University Hospital, Malmö, Sweden, Department of Clinical Sciences, Malmö, Lund University, Malmö, Sweden; **Roberto Lorusso**, Cardio-Thoracic Surgery, Maastricht University Medical Centre, Maastricht, Netherlands, Cardiovascular Research Institute Maastricht, Maastricht, Netherlands; **Roberto F.E. Pedretti**, Cardiovascular Department, IRCCS MultiMedica, Milan, Italy; **Angelos G. Rigopoulos**, Adult Cardiology, Mitera General Hospital-Hygeia Group, Athens, Greece; **Maria Rubini Gimenez**, Department of Internal Medicine and Cardiology, Heart Center Leipzig at the University of Leipzig, Leipzig, Germany, Cardiovascular Research Institute Basel, University Hospital Basel, Basel, Switzerland; **Holger Thiele**, Internal Medicine/Cardiology; Heart Center Leipzig at University of Leipzig, Leipzig, Germany, Leipzig Heart Science, Leipzig, Germany; **Pascal Vranckx**, Cardiology and Critical Care Medicine, Jessa Ziekenhuis, Hasselt, Belgium, Faculty of Medicine and Life Sciences, University of Hasselt, Hasselt, Belgium; **Sven Wassmann**, Cardiology, Cardiology Pasing, Munich, Germany; Faculty of Medicine, University of the Saarland, Homburg/Saar, Germany; **Nanette Kass Wenger**, Department of Medicine (Cardiology), Emory University School of Medicine, Atlanta, GA, United States of America, Consultant, Emory Heart and Vascular Center Emory University, School of Medicine, Atlanta, GA, United States of America, Founding Consultant, Emory Women's Heart Center, Emory University School of Medicine, Atlanta, GA, United States of America.

23. Appendix

ESC Scientific Document Group

Includes Document Reviewers and ESC National Cardiac Societies.

Document Reviewers: Sigrun Halvorsen (CPG Review Co-ordinator) (Norway), Stefan James (CPG Review Co-ordinator) (Sweden), Magdy Abdelhamid (Egypt), Victor Aboyans (France), Nina Ajmone Marsan (Netherlands), Sotiris Antoniou (United Kingdom), Riccardo Asteggiano (Italy), Maria Bäck (Sweden), Davide Capodanno (Italy), Ruben Casado-Arroyo (Belgium), Salvatore Cassese (Germany), Jelena Čelutkienė (Lithuania), Maja Cikes (Croatia), Jean-Philippe Collet (France), Gregory Ducrocq (France), Volkmar Falk (Germany), Laurent Fauchier (France), Tobias Geisler (Germany), Diana A. Gorog (United Kingdom), Lene Holmvang (Denmark), Tiny Jaarsma (Sweden), Hywel Wynne Jones (United Kingdom), Lars Køber (Denmark), Konstantinos C. Koskinas (Switzerland), Dipak Kotecha (United Kingdom), Konstantin A. Krychtiuk (Austria), Ulf Landmesser (Germany), George Lazaros (Greece), Basil S. Lewis (Israel), Bertil Lindahl (Sweden), Ales Linhart (Czech Republic), Maja-Lisa Løchen (Norway), Mamas A. Mamas

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

(United Kingdom), John William McEvoy (Ireland), Borislava Mihaylova (United Kingdom), Richard Mindham (United Kingdom), Christian Mueller (Switzerland), Lis Neubeck (United Kingdom), Josef Niebauer (Austria), Jens Cosedis Nielsen (Denmark), Alexander Niessner (Austria), Valeria Paradies (Netherlands), Agnes A. Pasquet (Belgium), Steffen E. Petersen (United Kingdom), Eva Prescott (Denmark), Amina Rakisheva (Kazakhstan), Bianca Rocca (Italy), Giuseppe M. C. Rosano (Italy), Leyla Elif Sade (United States of America/Türkiye), François Schiele (France), Jolanta M. Siller-Matula (Austria), Christian Sticherling (Switzerland), Robert F. Storey (United Kingdom), Matthias Thielmann (Germany), Christiaan Vrints (Belgium), Stephan Windecker (Switzerland), Rune Wiseth (Norway), and Adam Witkowski (Poland).

ESC National Cardiac Societies actively involved in the review process of the 2023 ESC Guidelines for the management of acute coronary syndromes:

Algeria: Algerian Society of Cardiology, Mohammed El Amine Bouzid; **Armenia:** Armenian Cardiologists Association, Hamlet Hayrapetyan; **Austria:** Austrian Society of Cardiology, Bernhard Metzler; **Belgium:** Belgian Society of Cardiology, Patrizio Lancellotti; **Bosnia and Herzegovina:** Association of Cardiologists of Bosnia and Herzegovina, Mugdim Bajrić; **Bulgaria:** Bulgarian Society of Cardiology, Kiril Karamfiloff; **Cyprus:** Cyprus Society of Cardiology, Andreas Mitsis; **Czechia:** Czech Society of Cardiology, Petr Ostadal; **Denmark:** Danish Society of Cardiology, Rikke Sørensen; **Egypt:** Egyptian Society of Cardiology, Tamer Elwasify; **Estonia:** Estonian Society of Cardiology, Toomas Marandi; **Finland:** Finnish Cardiac Society, Essi Ryödi; **France:** French Society of Cardiology, Jean-Philippe Collet; **Georgia:** Georgian Society of Cardiology, Archil Chukhrukidze; **Germany:** German Cardiac Society, Julinda Mehilli; **Greece:** Hellenic Society of Cardiology, Periklis Davlouros; **Hungary:** Hungarian Society of Cardiology, Dávid Becker; **Iceland:** Icelandic Society of Cardiology, Ingibjörg Jóna Guðmundsdóttir; **Ireland:** Irish Cardiac Society, James Crowley; **Israel:** Israel Heart Society, Yigal Abramowitz; **Italy:** Italian Federation of Cardiology, Ciro Indolfi; **Kazakhstan:** Association of Cardiologists of Kazakhstan, Orazbek Sakhov; **Kosovo (Republic of):** Kosovo Society of Cardiology, Shpend Elezi; **Kyrgyzstan:** Kyrgyz Society of Cardiology, Medet Beishenkulov; **Latvia:** Latvian Society of Cardiology, Andrejs Erglis; **Lebanon:** Lebanese Society of Cardiology, Nicolas Moussallem; **Libya:** Libyan Cardiac Society, Hisham Benlamin; **Lithuania:** Lithuanian Society of Cardiology, Olivija Dobilienė; **Luxembourg:** Luxembourg Society of Cardiology, Philippe Degrell; **Malta:** Maltese Cardiac Society, Matthew Mercieca Balbi; **Moldova (Republic of):** Moldavian Society of Cardiology, Aurel Grosu; **Morocco:** Moroccan Society of Cardiology, Zouhair Lakhal; **Netherlands**: Netherlands Society of Cardiology, Jurriën ten Berg; **North Macedonia**: The National Society of Cardiology of North Macedonia, Hristo Pejkov; **Norway:** Norwegian Society of Cardiology, Kristin Angel; **Poland:** Polish Cardiac Society, Adam Witkowski; **Portugal:** Portuguese Society of Cardiology, Manuel De Sousa Almeida; **Romania:** Romanian Society of Cardiology, Ovidiu Chioncel; **San Marino:** San Marino Society of Cardiology, Luca Bertelli; **Serbia:** Cardiology Society of Serbia, Sinisa Stojkovic; **Slovakia:** Slovak Society of Cardiology, Martin Studenčan; **Slovenia:** Slovenian Society of Cardiology, Peter Radšel; **Spain:** Spanish Society of Cardiology, Jose Luis Ferreiro; **Sweden:** Swedish Society of Cardiology, Annica Ravn-Fischer; **Switzerland:** Swiss Society of Cardiology, Lorenz Räber; **Syrian Arab Republic:** Syrian Cardiovascular Association, Mohammed Yassin Bani Marjeh; **Tunisia:** Tunisian Society of Cardiology and Cardiovascular Surgery, Majed Hassine; **Türkiye:** Turkish Society of Cardiology, Aylin Yildirir; **Ukraine:** Ukrainian Association of Cardiology, Alexander Parkhomenko; **United Kingdom of Great Britain and Northern Ireland:** British Cardiovascular Society, Adrian Paul Banning.

ESC Clinical Practice Guidelines (CPG) Committee: Eva Prescott (Chairperson) (Denmark), Stefan James (Co-Chairperson) (Sweden), Elena Arbelo (Spain), Colin Baigent (United Kingdom), Michael A. Borger (Germany), Sergio Buccheri (Sweden), Borja Ibanez (Spain), Lars Køber (Denmark), Konstantinos C. Koskinas (Switzerland), John William McEvoy (Ireland), Borislava Mihaylova (United Kingdom), Richard Mindham (United Kingdom), Lis Neubeck (United Kingdom), Jens Cosedis Nielsen (Denmark), Agnes A. Pasquet (Belgium), Amina Rakisheva (Kazakhstan), Bianca Rocca (Italy), Xavier Rossello (Spain), Ilonca Vaartjes (Netherlands), Christiaan Vrints (Belgium), Adam Witkowski (Poland), and Katja Zeppenfeld (Netherlands).

24. References

1. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, *et al.* Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019;**40**:237–269. https://doi.org/10.1093/eurheartj/ehy462
2. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;**392**:1736–1788. https://doi.org/10.1016/s0140-6736(18)32203-7
3. Timmis A, Vardas P, Townsend N, Torbica A, Katus H, De Smedt D, *et al.* European Society of Cardiology: cardiovascular disease statistics 2021. *Eur Heart J* 2022;**43**:716–799. https://doi.org/10.1093/eurheartj/ehab892
4. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, *et al.* 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;**39**:119–177. https://doi.org/10.1093/eurheartj/ehx393
5. Diercks DB, Peacock WF, Hiestand BC, Chen AY, Pollack CV, Kirk JD, *et al.* Frequency and consequences of recording an electrocardiogram >10 minutes after arrival in an emergency room in non-ST-segment elevation acute coronary syndromes (from the CRUSADE Initiative). *Am J Cardiol* 2006;**97**:437–442. https://doi.org/10.1016/j.amjcard.2005.09.073
6. Lopez-Sendon J, Coma-Canella I, Alcasena S, Seoane J, Gamallo C. Electrocardiographic findings in acute right ventricular infarction: sensitivity and specificity of electrocardiographic alterations in right precordial leads V4R, V3R, V1, V2, and V3. *J Am Coll Cardiol* 1985;**6**:1273–1279. https://doi.org/10.1016/s0735-1097(85)80213-8
7. Schmitt C, Lehmann G, Schmieder S, Karch M, Neumann F-J, Scho¨mig A. Diagnosis of acute myocardial infarction in angiographically documented occluded infarct vessel: limitations of ST-segment elevation in standard and extended ECG leads. *Chest* 2001;**120**:1540–1546. https://doi.org/10.1378/chest.120.5.1540
8. Kosuge M, Kimura K, Ishikawa T, Hongo Y, Shigemasa T, Sugiyama M, *et al.* Implications of the absence of ST-segment elevation in lead V4R in patients who have inferior wall acute myocardial infarction with right ventricular involvement. *Clin Cardiol* 2001;**24**:225–230. https://doi.org/10.1002/clc.4960240310
9. Yan AT, Yan RT, Kennelly BM, Anderson FA, Budaj A, López-Sendón J, *et al.* Relationship of ST elevation in lead aVR with angiographic findings and outcome in non-ST elevation acute coronary syndromes. *Am Heart J* 2007;**154**:71–78. https://doi.org/10.1016/j.ahj.2007.03.037
10. Hirano T, Tsuchiya K, Nishigaki K, Sou K, Kubota T, Ojio S, *et al.* Clinical features of emergency electrocardiography in patients with acute myocardial infarction caused by left main trunk obstruction. *Circ J* 2006;**70**:525–529. https://doi.org/10.1253/circj.70.525
11. Yamaji H, Iwasaki K, Kusachi S, Murakami T, Hirami R, Hamamoto H, *et al.* Prediction of acute left main coronary artery obstruction by 12-lead electrocardiography. ST segment elevation in lead aVR with less ST segment elevation in lead V(1). *J Am Coll Cardiol* 2001;**38**:1348–1354. doi: doi:10.1016/S0735-1097(01)01563-7
12. Chapman AR, Shah ASV, Lee KK, Anand A, Francis O, Adamson P, *et al.* Long-term outcomes in patients with type 2 myocardial infarction and myocardial injury.

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Circulation 2018;**137**:1236–1245. https://doi.org/10.1161/CIRCULATIONAHA.117.031806
13. Nestelberger T, Boeddinghaus J, Badertscher P, Twerenbold R, Wildi K, Breitenbücher D, *et al.* Effect of definition on incidence and prognosis of type 2 myocardial infarction. *J Am Coll Cardiol* 2017;**70**:1558–1568. https://doi.org/10.1016/j.jacc.2017.07.774
14. Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Renné T, Qaderi V, *et al.* Discrimination of patients with type 2 myocardial infarction. *Eur Heart J* 2017;**38**:3514–3520. https://doi.org/10.1093/eurheartj/ehx457
15. Shah ASV, Anand A, Strachan FE, Ferry AV, Lee KK, Chapman AR, *et al.* High-sensitivity troponin in the evaluation of patients with suspected acute coronary syndrome: a stepped-wedge, cluster-randomised controlled trial. *Lancet* 2018;**392**:919–928. https://doi.org/10.1016/S0140-6736(18)31923-8
16. Mandelzweig L, Battler A, Boyko V, Bueno H, Danchin N, Filippatos G, *et al.* The second Euro Heart Survey on acute coronary syndromes: characteristics, treatment, and outcome of patients with ACS in Europe and the Mediterranean Basin in 2004. *Eur Heart J* 2006;**27**:2285–2293. https://doi.org/10.1093/eurheartj/ehl196
17. Twerenbold R, Boeddinghaus J, Nestelberger T, Wildi K, Rubini Gimenez M, Badertscher P, *et al.* Clinical use of high-sensitivity cardiac troponin in patients with suspected myocardial infarction. *J Am Coll Cardiol* 2017;**70**:996–1012. https://doi.org/10.1016/j.jacc.2017.07.718
18. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, *et al.* 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2021;**42**:1289–1367. https://doi.org/10.1093/eurheartj/ehaa575
19. Rokos IC, French WJ, Koenig WJ, Stratton SJ, Nighswonger B, Strunk B, *et al.* Integration of pre-hospital electrocardiograms and ST-elevation myocardial infarction receiving center (SRC) networks: impact on door-to-balloon times across 10 independent regions. *JACC Cardiovasc Interv* 2009;**2**:339–346. https://doi.org/10.1016/j.jcin.2008.11.013
20. O'Doherty M, Tayler DI, Quinn E, Vincent R, Chamberlain DA. Five hundred patients with myocardial infarction monitored within one hour of symptoms. *Br Med J (Clin Res Ed)* 1983;**286**:1405–1408. https://doi.org/10.1136/bmj.286.6375.1405
21. Mehta RH, Starr AZ, Lopes RD, Hochman JS, Widimsky P, Pieper KS, *et al.* Incidence of and outcomes associated with ventricular tachycardia or fibrillation in patients undergoing primary percutaneous coronary intervention. *JAMA* 2009;**301**:1779–1789. https://doi.org/10.1001/jama.2009.600
22. Kulkarni AU, Brown R, Ayoubi M, Banka VS. Clinical use of posterior electrocardiographic leads: a prospective electrocardiographic analysis during coronary occlusion. *Am Heart J* 1996;**131**:736–741. https://doi.org/10.1016/s0002-8703(96)90280-x
23. Casas RE, Marriott HJL, Glancy DL. Value of leads V7-V9 in diagnosing posterior wall acute myocardial infarction and other causes of tall R waves in V1-V2. *Am J Cardiol* 1997;**80**:508–509. https://doi.org/10.1016/s0002-9149(97)00404-9
24. Zalenski RJ, Rydman RJ, Sloan EP, Hahn KH, Cooke D, Fagan J, *et al.* Value of posterior and right ventricular leads in comparison to the standard 12-lead electrocardiogram in evaluation of ST-segment elevation in suspected acute myocardial infarction. *Am J Cardiol* 1997;**79**:1579–1585. https://doi.org/10.1016/s0002-9149(97)00202-6
25. Reichlin T, Hochholzer W, Bassetti S, Steuer S, Stelzig C, Hartwiger S, *et al.* Early diagnosis of myocardial infarction with sensitive cardiac troponin assays. *N Engl J Med* 2009;**361**:858–867. https://doi.org/10.1056/NEJMoa0900428
26. Reichlin T, Twerenbold R, Reiter M, Steuer S, Bassetti S, Balmelli C, *et al.* Introduction of high-sensitivity troponin assays: impact on myocardial infarction incidence and prognosis. *Am J Med* 2012;**125**:1205–1213.e1. https://doi.org/10.1016/j.amjmed.2012.07.015
27. Neumann JT, Twerenbold R, Ojeda F, Sörensen NA, Chapman AR, Shah ASV, *et al.* Application of high-sensitivity troponin in suspected myocardial infarction. *N Engl J Med* 2019;**380**:2529–2540. https://doi.org/10.1056/NEJMoa1803377
28. Boeddinghaus J, Twerenbold R, Nestelberger T, Badertscher P, Wildi K, Puelacher C, *et al.* Clinical validation of a novel high-sensitivity cardiac troponin I assay for early diagnosis of acute myocardial infarction. *Clin Chem* 2018;**64**:1347–1360. https://doi.org/10.1373/clinchem.2018.286906
29. Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Schock A, Seddighizadeh P, *et al.* Evaluation of a new ultra-sensitivity troponin I assay in patients with suspected myocardial infarction. *Int J Cardiol* 2019;**283**:35–40. https://doi.org/10.1016/j.ijcard.2018.12.001
30. Neumann JT, Sorensen NA, Schwemer T, Ojeda F, Bourry R, Sciacca V, *et al.* Diagnosis of myocardial infarction using a high-sensitivity troponin I 1-hour algorithm. *JAMA Cardiol* 2016;**1**:397–404. https://doi.org/10.1001/jamacardio.2016.0695
31. Reichlin T, Schindler C, Drexler B, Twerenbold R, Reiter M, Zellweger C, *et al.* One-hour rule-out and rule-in of acute myocardial infarction using high-sensitivity cardiac troponin T. *Arch Intern Med* 2012;**172**:1211–1218. https://doi.org/10.1001/archinternmed.2012.3698
32. Boeddinghaus J, Twerenbold R, Nestelberger T, Koechlin L, Wussler D, Meier M, *et al.* Clinical use of a new high-sensitivity cardiac troponin I assay in patients with suspected myocardial infarction. *Clin Chem* 2019;**65**:1426–1436. https://doi.org/10.1373/clinchem.2019.304725
33. Boeddinghaus J, Lopez-Ayala P, Nestelberger T, Koechlin L, Münch T, Miro O, *et al.* Prospective validation of the ESC 0/1h-algorithm using high-sensitivity cardiac troponin I. *Am J Cardiol* 2021;**158**:152–153. https://doi.org/10.1016/j.amjcard.2021.08.007
34. Boeddinghaus J, Nestelberger T, Twerenbold R, Neumann JT, Lindahl B, Giannitsis E, *et al.* Impact of age on the performance of the ESC 0/1h-algorithms for early diagnosis of myocardial infarction. *Eur Heart J* 2018;**39**:3780–3794. https://doi.org/10.1093/eurheartj/ehy514
35. Twerenbold R, Badertscher P, Boeddinghaus J, Nestelberger T, Wildi K, Puelacher C, *et al.* 0/1-Hour triage algorithm for myocardial infarction in patients with renal dysfunction. *Circulation* 2018;**137**:436–451. https://doi.org/10.1161/CIRCULATIONAHA.117.028901
36. Boeddinghaus J, Reichlin T, Cullen L, Greenslade JH, Parsonage WA, Hammett C, *et al.* Two-hour algorithm for triage toward rule-out and rule-in of acute myocardial infarction by use of high-sensitivity cardiac troponin I. *Clin Chem* 2016;**62**:494–504. https://doi.org/10.1373/clinchem.2015.249508
37. Wildi K, Cullen L, Twerenbold R, Greenslade JH, Parsonage W, Boeddinghaus J, *et al.* Direct comparison of 2 rule-out strategies for acute myocardial infarction: 2-h accelerated diagnostic protocol vs 2-h algorithm. *Clin Chem* 2017;**63**:1227–1236. https://doi.org/10.1373/clinchem.2016.268359
38. Nestelberger T, Boeddinghaus J, Greenslade J, Parsonage WA, Than M, Wussler D, *et al.* Two-hour algorithm for rapid triage of suspected acute myocardial infarction using a high-sensitivity cardiac troponin I assay. *Clin Chem* 2019;**65**:1437–1447. https://doi.org/10.1373/clinchem.2019.305193
39. Koechlin L, Boeddinghaus J, Nestelberger T, Lopez-Ayala P, Wussler D, Shrestha S, *et al.* Performance of the ESC 0/2h-algorithm using high-sensitivity cardiac troponin I in the early diagnosis of myocardial infarction. *Am Heart J* 2021;**242**:132–137. https://doi.org/10.1016/j.ahj.2021.08.008
40. Wildi K, Nelles B, Twerenbold R, Rubini Giménez M, Reichlin T, Singeisen H, *et al.* Safety and efficacy of the 0 h/3 h protocol for rapid rule out of myocardial infarction. *Am Heart J* 2016;**181**:16–25. https://doi.org/10.1016/j.ahj.2016.07.013
41. Badertscher P, Boeddinghaus J, Twerenbold R, Nestelberger T, Wildi K, Wussler D, *et al.* Direct comparison of the 0/1h and 0/3h algorithms for early rule-out of acute myocardial infarction. *Circulation* 2018;**137**:2536–2538. https://doi.org/10.1161/CIRCULATIONAHA.118.034260
42. Chapman AR, Anand A, Boeddinghaus J, Ferry AV, Sandeman D, Adamson PD, *et al.* Comparison of the efficacy and safety of early rule-out pathways for acute myocardial infarction. *Circulation* 2017;**135**:1586–1596. https://doi.org/10.1161/CIRCULATIONAHA.116.025021
43. Chapman AR, Fujisawa T, Lee KK, Andrews JP, Anand A, Sandeman D, *et al.* Novel high-sensitivity cardiac troponin I assay in patients with suspected acute coronary syndrome. *Heart* 2019;**105**:616–622. https://doi.org/10.1136/heartjnl-2018-314093
44. Chew DP, Lambrakis K, Blyth A, Seshadri A, Edmonds MJR, Briffa T, *et al.* A randomized trial of a 1-hour troponin T protocol in suspected acute coronary syndromes: the rapid assessment of possible acute coronary syndrome in the emergency department with high-sensitivity troponin T study (RAPID-TnT). *Circulation* 2019;**140**:1543–1556. https://doi.org/10.1161/CIRCULATIONAHA.119.042891
45. Nestelberger T, Wildi K, Boeddinghaus J, Twerenbold R, Reichlin T, Giménez MR, *et al.* Characterization of the observe zone of the ESC 2015 high-sensitivity cardiac troponin 0h/1h-algorithm for the early diagnosis of acute myocardial infarction. *Int J Cardiol* 2016;**207**:238–245. https://doi.org/10.1016/j.ijcard.2016.01.112
46. Lopez-Ayala P, Nestelberger T, Boeddinghaus J, Koechlin L, Ratmann PD, Strebel I, *et al.* Novel criteria for the observe-zone of the ESC 0/1h-hs-cTnT algorithm. *Circulation* 2021;**144**:773–787. https://doi.org/10.1161/CIRCULATIONAHA.120.052982
47. Granger CB, Goldberg RJ, Dabbous O, Pieper KS, Eagle KA, Cannon CP, *et al.* Predictors of hospital mortality in the global registry of acute coronary events. *Arch Intern Med* 2003;**163**:2345–2353. https://doi.org/10.1001/archinte.163.19.2345
48. Fox KA, Fitzgerald G, Puymirat E, Huang W, Carruthers K, Simon T, *et al.* Should patients with acute coronary disease be stratified for management according to their risk? Derivation, external validation and outcomes using the updated GRACE risk score. *BMJ Open* 2014;**4**:e004425. https://doi.org/10.1136/bmjopen-2013-004425
49. Wenzl FA, Kraler S, Ambler G, Weston C, Herzog SA, Räber L, *et al.* Sex-specific evaluation and redevelopment of the GRACE score in non-ST-segment elevation acute coronary syndromes in populations from the UK and Switzerland: a multinational analysis with external cohort validation. *Lancet* 2022;**400**:744–756. https://doi.org/10.1016/s0140-6736(22)01483-0
50. Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. *Lancet* 1986;**1**:397–402.
51. Boersma E, Maas AC, Deckers JW, Simoons ML. Early thrombolytic treatment in acute myocardial infarction: reappraisal of the golden hour. *Lancet* 1996;**348**:771–775. https://doi.org/10.1016/s0140-6736(96)02514-7
52. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* 2003;**361**:13–20. https://doi.org/10.1016/s0140-6736(03)12113-7
53. Reichlin T, Twerenbold R, Maushart C, Reiter M, Moehring B, Schaub N, *et al.* Risk stratification in patients with unstable angina using absolute serial changes of 3 high-

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

sensitive troponin assays. *Am Heart J* 2013;**165**:371–378.e3. https://doi.org/10.1016/j.ahj.2012.11.010
54. Anand A, Lee KK, Chapman AR, Ferry AV, Adamson PD, Strachan FE, *et al.* High-sensitivity cardiac troponin on presentation to rule out myocardial infarction: a stepped-wedge cluster randomized controlled trial. *Circulation* 2021;**143**:2214–2224. https://doi.org/10.1161/CIRCULATIONAHA.120.052380
55. Mueller C. Biomarkers and acute coronary syndromes: an update. *Eur Heart J* 2014;**35**:552–556. https://doi.org/10.1093/eurheartj/eht530
56. Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, *et al.* 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;**37**:267–315. https://doi.org/10.1093/eurheartj/ehv320
57. Mueller C, Giannitsis E, Mockel M, Huber K, Mair J, Plebani M, *et al.* Rapid rule out of acute myocardial infarction: novel biomarker-based strategies. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:218–222. https://doi.org/10.1177/2048872616653229
58. Mockel M, Giannitsis E, Mueller C, Huber K, Jaffe AS, Mair J, *et al.* Rule-in of acute myocardial infarction: focus on troponin. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:212–217. https://doi.org/10.1177/2048872616653228
59. Roe MT, Harrington RA, Prosper DM, Pieper KS, Bhatt DL, Lincoff AM, *et al.* Clinical and therapeutic profile of patients presenting with acute coronary syndromes who do not have significant coronary artery disease. The Platelet Glycoprotein IIb/IIIa in Unstable Angina: Receptor Suppression Using Integrilin Therapy (PURSUIT) trial investigators. *Circulation* 2000;**102**:1101–1106. https://doi.org/10.1161/01.cir.102.10.1101
60. Westermann D, Neumann JT, Sorensen NA, Blankenberg S. High-sensitivity assays for troponin in patients with cardiac disease. *Nat Rev Cardiol* 2017;**14**:472–483. https://doi.org/10.1038/nrcardio.2017.48
61. Rubini Gimenez M, Twerenbold R, Reichlin T, Wildi K, Haaf P, Schaefer M, *et al.* Direct comparison of high-sensitivity-cardiac troponin I vs. T for the early diagnosis of acute myocardial infarction. *Eur Heart J* 2014;**35**:2303–2311. https://doi.org/10.1093/eurheartj/ehu188
62. Boeddinghaus J, Nestelberger T, Twerenbold R, Koechlin L, Meier M, Troester V, *et al.* High-sensitivity cardiac troponin I assay for early diagnosis of acute myocardial infarction. *Clin Chem* 2019;**65**:893–904. https://doi.org/10.1373/clinchem.2018.300061
63. Keller T, Zeller T, Peetz D, Tzikas S, Roth A, Czyz E, *et al.* Sensitive troponin I assay in early diagnosis of acute myocardial infarction. *N Engl J Med* 2009;**361**:868–877. https://doi.org/10.1056/NEJMoa0903515
64. Collinson PO, Saenger AK, Apple FS; IFCC C-CB. High sensitivity, contemporary and point-of-care cardiac troponin assays: educational aids developed by the IFCC Committee on Clinical Application of Cardiac Bio-Markers. *Clin Chem Lab Med* 2019;**57**:623–632. https://doi.org/10.1515/cclm-2018-1211
65. Camaro C, Aarts GWA, Adang EMM, van Hout R, Brok G, Hoare A, *et al.* Rule-out of non-ST-segment elevation acute coronary syndrome by a single, pre-hospital troponin measurement: a randomized trial. *Eur Heart J* 2023;**44**:1705–1714. https://doi.org/10.1093/eurheartj/ehad056
66. Pickering JW, Young JM, George PM, Watson AS, Aldous SJ, Troughton RW, *et al.* Validity of a novel point-of-care troponin assay for single-test rule-out of acute myocardial infarction. *JAMA Cardiol* 2018;**3**:1108–1112. https://doi.org/10.1001/jamacardio.2018.3368
67. Apple FS, Schulz K, Schmidt CW, van Domburg TSY, Fonville JM, de Theije FK. Determination of sex-specific 99th percentile upper reference limits for a point of care high sensitivity cardiac troponin I assay. *Clin Chem Lab Med* 2021;**59**:1574–1578. https://doi.org/10.1515/cclm-2021-0262
68. Sorensen NA, Neumann JT, Ojeda F, Giannitsis E, Spanuth E, Blankenberg S, *et al.* Diagnostic evaluation of a high-sensitivity troponin I point-of-care assay. *Clin Chem* 2019;**65**:1592–1601. https://doi.org/10.1373/clinchem.2019.307405
69. Azmy C, Guerard S, Bonnet X, Gabrielli F, Skalli W. EOS orthopaedic imaging system to study patellofemoral kinematics: assessment of uncertainty. *Orthop Traumatol Surg Res* 2010;**96**:28–36. https://doi.org/10.1016/j.rcot.2009.12.003
70. Chapman AR, Lee KK, McAllister DA, Cullen L, Greenslade JH, Parsonage W, *et al.* Association of high-sensitivity cardiac troponin I concentration with cardiac outcomes in patients with suspected acute coronary syndrome. *JAMA* 2017;**318**:1913–1924. https://doi.org/10.1001/jama.2017.17488
71. Hillinger P, Twerenbold R, Wildi K, Rubini Gimenez M, Jaeger C, Boeddinghaus J, *et al.* Gender-specific uncertainties in the diagnosis of acute coronary syndrome. *Clin Res Cardiol* 2017;**106**:28–37. https://doi.org/10.1007/s00392-016-1020-y
72. Miller-Hodges E, Anand A, Shah ASV, Chapman AR, Gallacher P, Lee KK, *et al.* High-sensitivity cardiac troponin and the risk stratification of patients with renal impairment presenting with suspected acute coronary syndrome. *Circulation* 2018;**137**:425–435. https://doi.org/10.1161/CIRCULATIONAHA.117.030320
73. Twerenbold R, Neumann JT, Sorensen NA, Ojeda F, Karakas M, Boeddinghaus J, *et al.* Prospective validation of the 0/1-h algorithm for early diagnosis of myocardial infarction. *J Am Coll Cardiol* 2018;**72**:620–632. https://doi.org/10.1016/j.jacc.2018.05.040
74. Rubini Gimenez M, Twerenbold R, Boeddinghaus J, Nestelberger T, Puelacher C, Hillinger P, *et al.* Clinical effect of sex-specific cutoff values of high-sensitivity cardiac troponin T in suspected myocardial infarction. *JAMA Cardiol* 2016;**1**:912–920. https://doi.org/10.1001/jamacardio.2016.2882
75. Mueller-Hennessen M, Lindahl B, Giannitsis E, Biener M, Vafaie M, deFilippi CR, *et al.* Diagnostic and prognostic implications using age- and gender-specific cut-offs for high-sensitivity cardiac troponin T-sub-analysis from the TRAPID-AMI study. *Int J Cardiol* 2016;**209**:26–33. https://doi.org/10.1016/j.ijcard.2016.01.213
76. Sorensen NA, Neumann JT, Ojeda F, Schäfer S, Magnussen C, Keller T, *et al.* Relations of sex to diagnosis and outcomes in acute coronary syndrome. *J Am Heart Assoc* 2018;**7**:e007297. https://doi.org/10.1161/JAHA.117.007297
77. Rubini Gimenez M, Badertscher P, Twerenbold R, Boeddinghaus J, Nestelberger T, Wussler D, *et al.* Impact of the US Food and Drug Administration-approved sex-specific cutoff values for high-sensitivity cardiac troponin T to diagnose myocardial infarction. *Circulation* 2018;**137**:1867–1869. https://doi.org/10.1161/circulationaha.117.031940
78. Lee KK, Ferry AV, Anand A, Strachan FE, Chapman AR, Kimenai DM, *et al.* Sex-specific thresholds of high-sensitivity troponin in patients with suspected acute coronary syndrome. *J Am Coll Cardiol* 2019;**74**:2032–2043. https://doi.org/10.1016/j.jacc.2019.07.082
79. Kimenai DM, Lindahl B, Jernberg T, Bekers O, Meex SJR, Eggers KM. Sex-specific effects of implementing a high-sensitivity troponin I assay in patients with suspected acute coronary syndrome: results from SWEDEHEART registry. *Sci Rep* 2020;**10**:15227. https://doi.org/10.1038/s41598-020-72204-2
80. Peacock WF, Baumann BM, Rivers EJ, Davis TE, Handy B, Jones CW, *et al.* Using sex-specific cutoffs for high-sensitivity cardiac troponin T to diagnose acute myocardial infarction. *Acad Emerg Med* 2021;**28**:463–466. https://doi.org/10.1111/acem.14098
81. Zhao Y, Izadnegahdar M, Lee MK, Kavsak PA, Singer J, Scheuermeyer F, *et al.* High-sensitivity cardiac troponin-optimizing the diagnosis of acute myocardial infarction/injury in women (CODE-MI): rationale and design for a multicenter, stepped-wedge, cluster-randomized trial. *Am Heart J* 2020;**229**:18–28. https://doi.org/10.1016/j.ahj.2020.06.013
82. Reichlin T, Irfan A, Twerenbold R, Reiter M, Hochholzer W, Burkhalter H, *et al.* Utility of absolute and relative changes in cardiac troponin concentrations in the early diagnosis of acute myocardial infarction. *Circulation* 2011;**124**:136–145. https://doi.org/10.1161/CIRCULATIONAHA.111.023937
83. Wildi K, Boeddinghaus J, Nestelberger T, Twerenbold R, Badertscher P, Wussler D, *et al.* Comparison of fourteen rule-out strategies for acute myocardial infarction. *Int J Cardiol* 2019;**283**:41–47. https://doi.org/10.1016/j.ijcard.2018.11.140
84. Ambavane A, Lindahl B, Giannitsis E, Roiz J, Mendivil J, Frankenstein L, *et al.* Economic evaluation of the one-hour rule-out and rule-in algorithm for acute myocardial infarction using the high-sensitivity cardiac troponin T assay in the emergency department. *PLoS One* 2017;**12**:e0187662. https://doi.org/10.1371/journal.pone.0187662
85. Boeddinghaus J, Nestelberger T, Twerenbold R, Wildi K, Badertscher P, Cupa J, *et al.* Direct comparison of 4 very early rule-out strategies for acute myocardial infarction using high-sensitivity cardiac troponin I. *Circulation* 2017;**135**:1597–1611. https://doi.org/10.1161/CIRCULATIONAHA.116.025661
86. Ljung L, Lindahl B, Eggers KM, Frick M, Linder R, Löfmark HB, *et al.* A rule-out strategy based on high-sensitivity troponin and HEART score reduces hospital admissions. *Ann Emerg Med* 2019;**73**:491–499. https://doi.org/10.1016/j.annemergmed.2018.11.039
87. Odqvist M, Andersson PO, Tygesen H, Eggers KM, Holzmann MJ. High-sensitivity troponins and outcomes after myocardial infarction. *J Am Coll Cardiol* 2018;**71**:2616–2624. https://doi.org/10.1016/j.jacc.2018.03.515
88. Twerenbold R, Jaeger C, Rubini Gimenez M, Wildi K, Reichlin T, Nestelberger T, *et al.* Impact of high-sensitivity cardiac troponin on use of coronary angiography, cardiac stress testing, and time to discharge in suspected acute myocardial infarction. *Eur Heart J* 2016;**37**:3324–3332. https://doi.org/10.1093/eurheartj/ehw232
89. Greenslade J, Cho E, Van Hise C, Hawkins T, Parsonage W, Ungerer J, *et al.* Evaluating rapid rule-out of acute myocardial infarction using a high-sensitivity cardiac troponin I assay at presentation. *Clin Chem* 2018;**64**:820–829. https://doi.org/10.1373/clinchem.2017.283887
90. Pickering JW, Than MP, Cullen L, Aldous S, ter Avest E, Body R, *et al.* Rapid rule-out of acute myocardial infarction with a single high-sensitivity cardiac troponin T measurement below the limit of detection: a collaborative meta-analysis. *Ann Intern Med* 2017;**166**:715–724. https://doi.org/10.7326/M16-2562
91. Shah AS, Anand A, Sandoval Y, Lee KK, Smith SW, Adamson PD, *et al.* High-sensitivity cardiac troponin I at presentation in patients with suspected acute coronary syndrome: a cohort study. *Lancet* 2015;**386**:2481–2488. https://doi.org/10.1016/S0140-6736(15)00391-8
92. Boeddinghaus J, Nestelberger T, Lopez-Ayala P, Koechlin L, Buechi M, Miro O, *et al.* Diagnostic performance of the European Society of Cardiology 0/1-h algorithms in late presenters. *J Am Coll Cardiol* 2021;**77**:1264–1267. https://doi.org/10.1016/j.jacc.2021.01.004
93. Mueller C, Giannitsis E, Christ M, Ordóñez-Llanos J, deFilippi C, McCord J, *et al.* Multicenter evaluation of a 0-hour/1-hour algorithm in the diagnosis of myocardial

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

infarction with high-sensitivity cardiac troponin T. *Ann Emerg Med* 2016;**68**:76–87.e4. https://doi.org/10.1016/j.annemergmed.2015.11.013
94. Stoyanov KM, Hund H, Biener M, Gandowitz J, Riedle C, Löhr J, *et al.* RAPID-CPU: a prospective study on implementation of the ESC 0/1-hour algorithm and safety of discharge after rule-out of myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2019;**9**:39–51. https://doi.org/10.1177/2048872619861911
95. Twerenbold R, Costabel JP, Nestelberger T, Campos R, Wussler D, Arbucci R, *et al.* Outcome of applying the ESC 0/1-hour algorithm in patients with suspected myocardial infarction. *J Am Coll Cardiol* 2019;**74**:483–494. https://doi.org/10.1016/j.jacc.2019.05.046
96. Nestelberger T, Boeddinghaus J, Wussler D, Twerenbold R, Badertscher P, Wildi K, *et al.* Predicting major adverse events in patients with acute myocardial infarction. *J Am Coll Cardiol* 2019;**74**:842–854. https://doi.org/10.1016/j.jacc.2019.06.025
97. Rubini Gimenez M, Wildi K, Wussler D, Koechlin L, Boeddinghaus J, Nestelberger T, *et al.* Early kinetics of cardiac troponin in suspected acute myocardial infarction. *Rev Esp Cardiol (Engl Ed)* 2021;**74**:502–509. https://doi.org/10.1016/j.rec.2020.04.008
98. Chiang CH, Chiang CH, Lee GH, Gi W-T, Wu Y-K, Huang S-S, *et al.* Safety and efficacy of the European Society of Cardiology 0/1-hour algorithm for diagnosis of myocardial infarction: systematic review and meta-analysis. *Heart* 2020;**106**:985–991. https://doi.org/10.1136/heartjnl-2019-316343
99. Vigen R, Kutscher P, Fernandez F, Yu A, Bertulfo B, Hashim IA, *et al.* Evaluation of a novel rule-out myocardial infarction protocol incorporating high-sensitivity troponin T in a US hospital. *Circulation* 2018;**138**:2061–2063. https://doi.org/10.1161/circulationaha.118.033861
100. Katus H, Ziegler A, Ekinci O, Giannitsis E, Stough WG, Achenbach S, *et al.* Early diagnosis of acute coronary syndrome. *Eur Heart J* 2017;**38**:3049–3055. https://doi.org/10.1093/eurheartj/ehx492
101. Koechlin L, Boeddinghaus J, Lopez-Ayala P, Nestelberger T, Wussler D, Mais F, *et al.* Diagnostic discrimination of a novel high-sensitivity cardiac troponin I assay and derivation/validation of an assay-specific 0/1h-algorithm. *Am Heart J* 2023;**255**:58–70. https://doi.org/10.1016/j.ahj.2022.10.007
102. Kaier TE, Twerenbold R, Puelacher C, Marjot J, Imambaccus N, Boeddinghaus J, *et al.* Direct comparison of cardiac myosin-binding protein C with cardiac troponins for the early diagnosis of acute myocardial infarction. *Circulation* 2017;**136**:1495–1508. https://doi.org/10.1161/CIRCULATIONAHA.117.028084
103. Boeddinghaus J, Reichlin T, Nestelberger T, Twerenbold R, Meili Y, Wildi K, *et al.* Early diagnosis of acute myocardial infarction in patients with mild elevations of cardiac troponin. *Clin Res Cardiol* 2017;**106**:457–467. https://doi.org/10.1007/s00392-016-1075-9
104. Hillinger P, Twerenbold R, Jaeger C, Wildi K, Reichlin T, Gimenez MR, *et al.* Optimizing early rule-out strategies for acute myocardial infarction: utility of 1-hour copeptin. *Clin Chem* 2015;**61**:1466–1474. https://doi.org/10.1373/clinchem.2015.242743
105. Keller T, Tzikas S, Zeller T, Czyz E, Lillpopp L, Ojeda FM, *et al.* Copeptin improves early diagnosis of acute myocardial infarction. *J Am Coll Cardiol* 2010;**55**:2096–2106. https://doi.org/10.1016/j.jacc.2010.01.029
106. Möckel M, Searle J, Hamm C, Slagman A, Blankenberg S, Huber K, *et al.* Early discharge using single cardiac troponin and copeptin testing in patients with suspected acute coronary syndrome (ACS): a randomized, controlled clinical process study. *Eur Heart J* 2015;**36**:369–376. https://doi.org/10.1093/eurheartj/ehu178
107. Mueller C, Mockel M, Giannitsis E, Huber K, Mair J, Plebani M, *et al.* Use of copeptin for rapid rule-out of acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:570–576. https://doi.org/10.1177/2048872617710791
108. Mueller-Hennessen M, Lindahl B, Giannitsis E, Vafaie M, Biener M, Haushofer AC, *et al.* Combined testing of copeptin and high-sensitivity cardiac troponin T at presentation in comparison to other algorithms for rapid rule-out of acute myocardial infarction. *Int J Cardiol* 2019;**276**:261–267. https://doi.org/10.1016/j.ijcard.2018.10.084
109. Reichlin T, Hochholzer W, Stelzig C, Laule K, Freidank H, Morgenthaler NG, *et al.* Incremental value of copeptin for rapid rule out of acute myocardial infarction. *J Am Coll Cardiol* 2009;**54**:60–68. https://doi.org/10.1016/j.jacc.2009.01.076
110. Stallone F, Schoenenberger AW, Puelacher C, Rubini Gimenez M, Walz B, Naduvilekoot Devasia A, *et al.* Incremental value of copeptin in suspected acute myocardial infarction very early after symptom onset. *Eur Heart J Acute Cardiovasc Care* 2016;**5**:407–415. https://doi.org/10.1177/2048872616641289
111. Vargas KG, Kassem M, Mueller C, Wojta J, Huber K. Copeptin for the early rule-out of non-ST-elevation myocardial infarction. *Int J Cardiol* 2016;**223**:797–804. https://doi.org/10.1016/j.ijcard.2016.08.304
112. Wildi K, Zellweger C, Twerenbold R, Jaeger C, Reichlin T, Haaf P, *et al.* Incremental value of copeptin to highly sensitive cardiac troponin I for rapid rule-out of myocardial infarction. *Int J Cardiol* 2015;**190**:170–176. https://doi.org/10.1016/j.ijcard.2015.04.133
113. Zellweger C, Wildi K, Twerenbold R, Reichlin T, Naduvilekoot A, Neuhaus JD, *et al.* Use of copeptin and high-sensitive cardiac troponin T for diagnosis and prognosis in patients with diabetes mellitus and suspected acute myocardial infarction. *Int J Cardiol* 2015;**190**:190–197. https://doi.org/10.1016/j.ijcard.2015.04.134
114. Restan IZ, Sanchez AY, Steiro OT, Lopez-Ayala P, Tjora HL, Langørgen J, *et al.* Adding stress biomarkers to high-sensitivity cardiac troponin for rapid non-ST-elevation myocardial infarction rule-out protocols. *Eur Heart J Acute Cardiovasc Care* 2022;**11**:201–212. https://doi.org/10.1093/ehjacc/zuab124
115. Dedic A, Lubbers MM, Schaap J, Lammers J, Lamfers EJ, Rensing BJ, *et al.* Coronary CT angiography for suspected ACS in the era of high-sensitivity troponins: randomized multicenter study. *J Am Coll Cardiol* 2016;**67**:16–26. https://doi.org/10.1016/j.jacc.2015.10.045
116. Hoffmann U, Truong QA, Schoenfeld DA, Chou ET, Woodard PK, Nagurney JT, *et al.* Coronary CT angiography versus standard evaluation in acute chest pain. *N Engl J Med* 2012;**367**:299–308. https://doi.org/10.1056/NEJMoa1201161
117. Gray AJ, Roobottom C, Smith JE, Goodacre S, Oatey K, O'Brien R, *et al.* Early computed tomography coronary angiography in patients with suspected acute coronary syndrome: randomised controlled trial. *BMJ* 2021;**374**:n2106. https://doi.org/10.1136/bmj.n2106
118. Lee KK, Bularga A, O'Brien R, Ferry AV, Doudesis D, Fujisawa T, *et al.* Troponin-guided coronary computed tomographic angiography after exclusion of myocardial infarction. *J Am Coll Cardiol* 2021;**78**:1407–1417. https://doi.org/10.1016/j.jacc.2021.07.055
119. Kofoed KF, Engstrom T, Sigvardsen PE, Linde JJ, Torp-Pedersen C, de Knegt M, *et al.* Prognostic value of coronary CT angiography in patients with non-ST-segment elevation acute coronary syndromes. *J Am Coll Cardiol* 2021;**77**:1044–1052. https://doi.org/10.1016/j.jacc.2020.12.037
120. Linde JJ, Hove JD, Sorgaard M, Kelbæk H, Jensen GB, Kühl JT, *et al.* Long-term clinical impact of coronary CT angiography in patients with recent acute-onset chest pain: the randomized controlled CATCH trial. *JACC Cardiovasc Imaging* 2015;**8**:1404–1413. https://doi.org/10.1016/j.jcmg.2015.07.015
121. Linde JJ, Kelbæk H, Hansen TF, Sigvardsen PE, Torp-Pedersen C, Bech J, *et al.* Coronary CT angiography in patients with non-ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol* 2020;**75**:453–463. https://doi.org/10.1016/j.jacc.2019.12.012
122. Samad Z, Hakeem A, Mahmood SS, Pieper K, Patel MR, Simel DL, *et al.* A meta-analysis and systematic review of computed tomography angiography as a diagnostic triage tool for patients with chest pain presenting to the emergency department. *J Nucl Cardiol* 2012;**19**:364–376. https://doi.org/10.1007/s12350-012-9520-2
123. Litt HI, Gatsonis C, Snyder B, Singh H, Miller CD, Entrikin DW, *et al.* CT angiography for safe discharge of patients with possible acute coronary syndromes. *N Engl J Med* 2012;**366**:1393–1403. https://doi.org/10.1056/NEJMoa1201163
124. Hulten E, Pickett C, Bittencourt MS, Villines TC, Petrillo S, Di Carli MF, *et al.* Outcomes after coronary computed tomography angiography in the emergency department: a systematic review and meta-analysis of randomized, controlled trials. *J Am Coll Cardiol* 2013;**61**:880–892. https://doi.org/10.1016/j.jacc.2012.11.061
125. Gaibazzi N, Reverberi C, Badano L. Usefulness of contrast stress-echocardiography or exercise-electrocardiography to predict long-term acute coronary syndromes in patients presenting with chest pain without electrocardiographic abnormalities or 12-hour troponin elevation. *Am J Cardiol* 2011;**107**:161–167. https://doi.org/10.1016/j.amjcard.2010.08.066
126. Lim SH, Anantharaman V, Sundram F, Chan ES-Y, Ang ES, Yo SL, *et al.* Stress myocardial perfusion imaging for the evaluation and triage of chest pain in the emergency department: a randomized controlled trial. *J Nucl Cardiol* 2013;**20**:1002–1012. https://doi.org/10.1007/s12350-013-9736-9
127. Nabi F, Kassi M, Muhyieddeen K, Chang SM, Xu J, Peterson LE, *et al.* Optimizing evaluation of patients with low-to-intermediate-risk acute chest pain: a randomized study comparing stress myocardial perfusion tomography incorporating stress-only imaging versus cardiac CT. *J Nucl Med* 2016;**57**:378–384. https://doi.org/10.2967/jnumed.115.166595
128. Jackson AM, Zhang R, Findlay I, Robertson K, Lindsay M, Morris T, *et al.* Healthcare disparities for women hospitalized with myocardial infarction and angina. *Eur Heart J Qual Care Clin Outcomes* 2020;**6**:156–165. https://doi.org/10.1093/ehjqcco/qcz040
129. Terkelsen CJ, Sørensen JT, Maeng M, Jensen LO, Tilsted HH, Trautner S, *et al.* System delay and mortality among patients with STEMI treated with primary percutaneous coronary intervention. *JAMA* 2010;**304**:763–771. https://doi.org/10.1001/jama.2010.1139
130. Jortveit J, Pripp AH, Halvorsen S. Outcomes after delayed primary percutaneous coronary intervention vs. pharmaco-invasive strategy in ST-segment elevation myocardial infarction in Norway. *Eur Heart J Cardiovasc Pharmacother* 2022;**8**:442–451. https://doi.org/10.1093/ehjcvp/pvab041
131. Larsen AI, Løland KH, Hovland S, Bleie Ø, Eek C, Fossum E, *et al.* Guideline-recommended time less than 90 minutes from ECG to primary percutaneous coronary intervention for ST-segment-elevation myocardial infarction is associated with major survival benefits, especially in octogenarians: a contemporary report in 11 226 patients from NORIC. *J Am Heart Assoc* 2022;**11**:e024849. https://doi.org/10.1161/jaha.122.024849
132. Fordyce CB, Al-Khalidi HR, Jollis JG, Roettig ML, Gu J, Bagai A, *et al.* Association of rapid care process implementation on reperfusion times across multiple ST-segment-elevation myocardial infarction networks. *Circ Cardiovasc Interv* 2017;**10:**e004061. https://doi.org/10.1161/circinterventions.116.004061
133. Stowens JC, Sonnad SS, Rosenbaum RA. Using EMS dispatch to trigger STEMI alerts decreases door-to-balloon times. *West J Emerg Med* 2015;**16**:472–480. https://doi.org/10.5811/westjem.2015.4.24248

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

134. Squire BT, Tamayo-Sarver JH, Rashi P, Koenig W, Niemann JT. Effect of prehospital cardiac catheterization lab activation on door-to-balloon time, mortality, and false-positive activation. *Prehosp Emerg Care* 2014;**18**:1–8. https://doi.org/10.3109/10903127.2013.836263
135. Shavadia JS, Roe MT, Chen AY, Lucas J, Fanaroff AC, Kochar A, *et al.* Association between cardiac catheterization laboratory pre-activation and reperfusion timing metrics and outcomes in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: a report from the ACTION registry. *JACC Cardiovasc Interv* 2018;**11**:1837–1847. https://doi.org/10.1016/j.jcin.2018.07.020
136. Kontos MC, Gunderson MR, Zegre-Hemsey JK, Lange DC, French WJ, Henry TD, *et al.* Prehospital activation of hospital resources (PreAct) ST-segment-elevation myocardial infarction (STEMI): a standardized approach to prehospital activation and direct to the catheterization laboratory for STEMI recommendations from the American Heart Association's mission: lifeline program. *J Am Heart Assoc* 2020;**9**:e011963. https://doi.org/10.1161/jaha.119.011963
137. Bagai A, Jollis JG, Dauerman HL, Peng SA, Rokos IC, Bates ER, *et al.* Emergency department bypass for ST-segment-elevation myocardial infarction patients identified with a prehospital electrocardiogram: a report from the American Heart Association Mission: Lifeline program. *Circulation* 2013;**128**:352–359. https://doi.org/10.1161/circulationaha.113.002339
138. Scholz KH, Friede T, Meyer T, Jacobshagen C, Lengenfelder B, Jung J, *et al.* Prognostic significance of emergency department bypass in stable and unstable patients with ST-segment elevation myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:34–44. https://doi.org/10.1177/2048872618813907
139. Meisel SR, Kleiner-Shochat M, Abu-Fanne R, Frimerman A, Danon A, Minha S, *et al.* Direct admission of patients with ST-segment-elevation myocardial infarction to the catheterization laboratory shortens pain-to-balloon and door-to-balloon time intervals but only the pain-to-balloon interval impacts short- and long-term mortality. *J Am Heart Assoc* 2021;**10**:e018343. https://doi.org/10.1161/jaha.120.018343
140. Wang TY, Nallamothu BK, Krumholz HM, Li S, Roe MT, Jollis JG, *et al.* Association of door-in to door-out time with reperfusion delays and outcomes among patients transferred for primary percutaneous coronary intervention. *JAMA* 2011;**305**:2540–2547. https://doi.org/10.1001/jama.2011.862
141. Huber K, De Caterina R, Kristensen SD, Verheugt FWA, Montalescot G, Maestro LB, *et al.* Pre-hospital reperfusion therapy: a strategy to improve therapeutic outcome in patients with ST-elevation myocardial infarction. *Eur Heart J* 2005;**26**:2063–2074. https://doi.org/10.1093/eurheartj/ehi413
142. Welsh RC, Chang W, Goldstein P, Adgey J, Granger CB, Verheugt FW, *et al.* Time to treatment and the impact of a physician on prehospital management of acute ST elevation myocardial infarction: insights from the ASSENT-3 PLUS trial. *Heart* 2005;**91**:1400–1406. https://doi.org/10.1136/hrt.2004.054510
143. Jollis JG, Al-Khalidi HR, Roettig ML, Berger PB, Corbett CC, Doerfler SM, *et al.* Impact of regionalization of ST-segment-elevation myocardial infarction care on treatment times and outcomes for emergency medical services-transported patients presenting to hospitals with percutaneous coronary intervention: mission: lifeline accelerator-2. *Circulation* 2018;**137**:376–387. https://doi.org/10.1161/circulationaha.117.032446
144. Fosbol EL, Granger CB, Jollis JG, Monk L, Lin L, Lytle BL, *et al.* The impact of a statewide pre-hospital STEMI strategy to bypass hospitals without percutaneous coronary intervention capability on treatment times. *Circulation* 2013;**127**:604–612. https://doi.org/10.1161/circulationaha.112.118463
145. Kalla K, Christ G, Karnik R, Malzer R, Norman G, Prachar H, *et al.* Implementation of guidelines improves the standard of care: the Viennese registry on reperfusion strategies in ST-elevation myocardial infarction (Vienna STEMI registry). *Circulation* 2006;**113**:2398–2405. https://doi.org/10.1161/circulationaha.105.586198
146. Henry TD, Sharkey SW, Burke MN, Chavez IJ, Graham KJ, Henry CR, *et al.* A regional system to provide timely access to percutaneous coronary intervention for ST-elevation myocardial infarction. *Circulation* 2007;**116**:721–728. https://doi.org/10.1161/circulationaha.107.694141
147. Le May MR, So DY, Dionne R, Glover CA, Froeschl MPV, Wells GA, *et al.* A citywide protocol for primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med* 2008;**358**:231–240. https://doi.org/10.1056/NEJMoa073102
148. Hofmann R, James SK, Jernberg T, Lindahl B, Erlinge D, Witt N, *et al.* Oxygen therapy in suspected acute myocardial infarction. *N Engl J Med* 2017;**377**:1240–1249. https://doi.org/10.1056/NEJMoa1706222
149. Stewart RAH, Jones P, Dicker B, Jiang Y, Smith T, Swain A, *et al.* High flow oxygen and risk of mortality in patients with a suspected acute coronary syndrome: pragmatic, cluster randomised, crossover trial. *BMJ* 2021;**372**:n355. https://doi.org/10.1136/bmj.n355
150. Henrikson CA, Howell EE, Bush DE, Miles JS, Meininger GR, Friedlander T, *et al.* Chest pain relief by nitroglycerin does not predict active coronary artery disease. *Ann Intern Med* 2003;**139**:979–986. https://doi.org/10.7326/0003-4819-139-12-200312160-00007
151. Charpentier S, Galinski M, Bounes V, Ricard-Hibon A, El-Khoury C, Elbaz M, *et al.* Nitrous oxide/oxygen plus acetaminophen versus morphine in ST elevation myocardial infarction: open-label, cluster-randomized, non-inferiority study. *Scand J Trauma Resusc Emerg Med* 2020;**28**:36. https://doi.org/10.1186/s13049-020-00731-y
152. Silvain J, Storey RF, Cayla G, Esteve J-B, Dillinger J-G, Rousseau H, *et al.* P2Y12 receptor inhibition and effect of morphine in patients undergoing primary PCI for ST-segment elevation myocardial infarction. The PRIVATE-ATLANTIC study. *Thromb Haemost* 2016;**116**:369–378. https://doi.org/10.1160/th15-12-0944
153. Parodi G, Bellandi B, Xanthopoulou I, Capranzano P, Capodanno D, Valenti R, *et al.* Morphine is associated with a delayed activity of oral antiplatelet agents in patients with ST-elevation acute myocardial infarction undergoing primary percutaneous coronary intervention. *Circ Cardiovasc Interv* 2015;**8**:e001593. https://doi.org/10.1161/circinterventions.114.001593
154. Saad M, Meyer-Saraei R, de Waha-Thiele S, Stiermaier T, Graf T, Fuernau G, *et al.* Impact of morphine treatment with and without metoclopramide coadministration on ticagrelor-induced platelet inhibition in acute myocardial infarction: the randomized MonAMI trial. *Circulation* 2020;**141**:1354–1356. https://doi.org/10.1161/circulationaha.119.042816
155. Stiermaier T, Schaefer P, Meyer-Saraei R, Saad M, de Waha-Thiele S, Pöss J, *et al.* Impact of morphine treatment with and without metoclopramide coadministration on myocardial and microvascular injury in acute myocardial infarction: insights from the randomized MonAMI trial. *J Am Heart Assoc* 2021;**10**:e018881. https://doi.org/10.1161/jaha.120.018881
156. Kubica J, Adamski P, Ostrowska M, Sikora J, Kubica JM, Sroka WD, *et al.* Morphine delays and attenuates ticagrelor exposure and action in patients with myocardial infarction: the randomized, double-blind, placebo-controlled IMPRESSION trial. *Eur Heart J* 2016;**37**:245–252. https://doi.org/10.1093/eurheartj/ehv547
157. Zhang Y, Wang N, Gu Q. Effects of morphine on P2Y(12) platelet inhibitors in patients with acute myocardial infarction: a meta-analysis. *Am J Emerg Med* 2021;**41**:219–228. https://doi.org/10.1016/j.ajem.2020.11.003
158. Furtado RHM, Nicolau JC, Guo J, Im K, White JA, Sabatine MS, *et al.* Morphine and cardiovascular outcomes among patients with non-ST-segment elevation acute coronary syndromes undergoing coronary angiography. *J Am Coll Cardiol* 2020;**75**:289–300. https://doi.org/10.1016/j.jacc.2019.11.035
159. Kubica A, Kosobucka A, Niezgoda P, Adamski P, Buszko K, Lesiak M, *et al.* ANalgesic Efficacy and safety of MOrphiNe versus methoxyflurane in patients with acute myocardial infarction: the rationale and design of the ANEMON-SIRIO 3 study: a multicentre, open-label, phase II, randomised clinical trial. *BMJ Open* 2021;**11**:e043330. https://doi.org/10.1136/bmjopen-2020-043330
160. Batchelor R, Liu DH, Bloom J, Noaman S, Chan W. Association of periprocedural intravenous morphine use on clinical outcomes in ST-elevation myocardial infarction (STEMI) treated by primary percutaneous coronary intervention: systematic review and meta-analysis. *Catheter Cardiovasc Interv* 2020;**96**:76–88. https://doi.org/10.1002/ccd.28561
161. Bonin M, Mewton N, Roubille F, Morel O, Cayla G, Angoulvant D, *et al.* Effect and safety of morphine use in acute anterior ST-segment elevation myocardial infarction. *J Am Heart Assoc* 2018;**7**:e006833. https://doi.org/10.1161/jaha.117.006833
162. Clemente-Moragón A, Gómez M, Villena-Gutiérrez R, Lalama DV, García-Prieto J, Martínez F, *et al.* Metoprolol exerts a non-class effect against ischaemia-reperfusion injury by abrogating exacerbated inflammation. *Eur Heart J* 2020;**41**:4425–4440. https://doi.org/10.1093/eurheartj/ehaa733
163. Ibanez B, Macaya C, Sánchez-Brunete V, Pizarro G, Fernández-Friera L, Mateos A, *et al.* Effect of early metoprolol on infarct size in ST-segment-elevation myocardial infarction patients undergoing primary percutaneous coronary intervention: the effect of metoprolol in cardioprotection during an acute myocardial infarction (METOCARD-CNIC) trial. *Circulation* 2013;**128**:1495–1503. https://doi.org/10.1161/circulationaha.113.003653
164. Roolvink V, Ibáñez B, Ottervanger JP, Pizarro G, van Royen N, Mateos A, *et al.* Early intravenous beta-blockers in patients with ST-segment elevation myocardial infarction before primary percutaneous coronary intervention. *J Am Coll Cardiol* 2016;**67**:2705–2715. https://doi.org/10.1016/j.jacc.2016.03.522
165. Pizarro G, Fernández-Friera L, Fuster V, Fernández-Jiménez R, García-Ruiz JM, García-Álvarez A, *et al.* Long-term benefit of early pre-reperfusion metoprolol administration in patients with acute myocardial infarction: results from the METOCARD-CNIC trial (effect of metoprolol in cardioprotection during an acute myocardial infarction). *J Am Coll Cardiol* 2014;**63**:2356–2362. https://doi.org/10.1016/j.jacc.2014.03.014
166. García-Ruiz JM, Fernández-Jiménez R, García-Alvarez A, Pizarro G, Galán-Arriola C, Fernández-Friera L, *et al.* Impact of the timing of metoprolol administration during STEMI on infarct size and ventricular function. *J Am Coll Cardiol* 2016;**67**:2093–2104. https://doi.org/10.1016/j.jacc.2016.02.050
167. Hoedemaker NP, Roolvink V, de Winter RJ, van Royen N, Fuster V, García-Ruiz JM, *et al.* Early intravenous beta-blockers in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a patient-pooled meta-analysis of randomized clinical trials. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:469–477. https://doi.org/10.1177/2048872619830609
168. Sterling LH, Filion KB, Atallah R, Reynier P, Eisenberg MJ. Intravenous beta-blockers in ST-segment elevation myocardial infarction: a systematic review and meta-analysis. *Int J Cardiol* 2017;**228**:295–302. https://doi.org/10.1016/j.ijcard.2016.11.133

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

169. Chatterjee S, Chaudhuri D, Vedanthan R, Fuster V, Ibanez B, Bangalore S, *et al.* Early intravenous beta-blockers in patients with acute coronary syndrome–a meta-analysis of randomized trials. *Int J Cardiol* 2013;**168**:915–921. https://doi.org/10.1016/j.ijcard.2012.10.050
170. Elgendy IY, Elgendy AY, Mahmoud AN, Mansoor H, Mojadidi MK, Bavry AA, *et al.* Intravenous β-blockers for patients undergoing primary percutaneous coronary intervention: a meta-analysis of randomized trials. *Int J Cardiol* 2016;**223**:891–897. https://doi.org/10.1016/j.ijcard.2016.08.293
171. García-Prieto J, Villena-Gutiérrez R, Gómez M, Bernardo E, Pun-García A, García-Lunar I, *et al.* Neutrophil stunning by metoprolol reduces infarct size. *Nat Commun* 2017;**8**:14780. https://doi.org/10.1038/ncomms14780
172. Stub D, Smith K, Bernard S, Nehme Z, Stephenson M, Bray JE, *et al.* Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation* 2015;**131**:2143–2150. https://doi.org/10.1161/circulationaha.114.014494
173. Widimsky P, Wijns W, Fajadet J, de Belder M, Knot J, Aaberge L, *et al.* Reperfusion therapy for ST elevation acute myocardial infarction in Europe: description of the current situation in 30 countries. *Eur Heart J* 2010;**31**:943–957. https://doi.org/10.1093/eurheartj/ehp492
174. Hall M, Laut K, Dondo TB, Alabas OA, Brogan RA, Gutacker N, *et al.* Patient and hospital determinants of primary percutaneous coronary intervention in England, 2003–2013. *Heart* 2016;**102**:313–319. https://doi.org/10.1136/heartjnl-2015-308616
175. Widimsky P, Fajadet J, Danchin N, Wijns W. "Stent 4 Life" targeting PCI at all who will benefit the most. A joint project between EAPCI, Euro-PCR, EUCOMED and the ESC Working Group on Acute Cardiac Care. *EuroIntervention* 2009;**4**:555–557. doi: doi:10.4244/EIJV4I5A94
176. Pinto DS, Kirtane AJ, Nallamothu BK, Murphy SA, Cohen DJ, Laham RJ, *et al.* Hospital delays in reperfusion for ST-elevation myocardial infarction: implications when selecting a reperfusion strategy. *Circulation* 2006;**114**:2019–2025. https://doi.org/10.1161/circulationaha.106.638353
177. Steg PG, Cambou JP, Goldstein P, Durand E, Sauval P, Kadri Z, *et al.* Bypassing the emergency room reduces delays and mortality in ST elevation myocardial infarction: the USIC 2000 registry. *Heart* 2006;**92**:1378–1383. https://doi.org/10.1136/hrt.2006.101972
178. Baran KW, Kamrowski KA, Westwater JJ, Tschida VH, Alexander CF, Beahrs MM, *et al.* Very rapid treatment of ST-segment-elevation myocardial infarction: utilizing prehospital electrocardiograms to bypass the emergency department. *Circ Cardiovasc Qual Outcomes* 2010;**3**:431–437. https://doi.org/10.1161/circoutcomes.110.942631
179. Huynh T, Perron S, O'Loughlin J, Joseph L, Labrecque M, Tu JV, *et al.* Comparison of primary percutaneous coronary intervention and fibrinolytic therapy in ST-segment-elevation myocardial infarction: Bayesian hierarchical meta-analyses of randomized controlled trials and observational studies. *Circulation* 2009;**119**:3101–3109. https://doi.org/10.1161/circulationaha.108.793745
180. Nallamothu BK, Bates ER. Percutaneous coronary intervention versus fibrinolytic therapy in acute myocardial infarction: is timing (almost) everything? *Am J Cardiol* 2003;**92**:824–826. https://doi.org/10.1016/s0002-9149(03)00891-9
181. Betriu A, Masotti M. Comparison of mortality rates in acute myocardial infarction treated by percutaneous coronary intervention versus fibrinolysis. *Am J Cardiol* 2005;**95**:100–101. https://doi.org/10.1016/j.amjcard.2004.08.069
182. Boersma E. Does time matter? A pooled analysis of randomized clinical trials comparing primary percutaneous coronary intervention and in-hospital fibrinolysis in acute myocardial infarction patients. *Eur Heart J* 2006;**27**:779–788. https://doi.org/10.1093/eurheartj/ehi810
183. Pinto DS, Frederick PD, Chakrabarti AK, Kirtane AJ, Ullman E, Dejam A, *et al.* Benefit of transferring ST-segment-elevation myocardial infarction patients for percutaneous coronary intervention compared with administration of onsite fibrinolytic declines as delays increase. *Circulation* 2011;**124**:2512–2521. https://doi.org/10.1161/circulationaha.111.018549
184. Armstrong PW, Gershlick AH, Goldstein P, Wilcox R, Danays T, Lambert Y, *et al.* Fibrinolysis or primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med* 2013;**368**:1379–1387. https://doi.org/10.1056/NEJMoa1301092
185. Gershlick AH, Stephens-Lloyd A, Hughes S, Abrams KR, Stevens SE, Uren NG, *et al.* Rescue angioplasty after failed thrombolytic therapy for acute myocardial infarction. *N Engl J Med* 2005;**353**:2758–2768. https://doi.org/10.1056/NEJMoa050849
186. Cantor WJ, Fitchett D, Borgundvaag B, Ducas J, Heffernan M, Cohen EA, *et al.* Routine early angioplasty after fibrinolysis for acute myocardial infarction. *N Engl J Med* 2009;**360**:2705–2718. https://doi.org/10.1056/NEJMoa0808276
187. Wijeysundera HC, Vijayaraghavan R, Nallamothu BK, Foody JoAnne M, Krumholz HM, Phillips CO, *et al.* Rescue angioplasty or repeat fibrinolysis after failed fibrinolytic therapy for ST-segment myocardial infarction: a meta-analysis of randomized trials. *J Am Coll Cardiol* 2007;**49**:422–430. https://doi.org/10.1016/j.jacc.2006.09.033
188. Sutton AG, Campbell PG, Graham R, Price DJA, Gray JC, Grech ED, *et al.* A randomized trial of rescue angioplasty versus a conservative approach for failed fibrinolysis in ST-segment elevation myocardial infarction: the Middlesbrough Early Revascularization to Limit INfarction (MERLIN) trial. *J Am Coll Cardiol* 2004;**44**:287–296. https://doi.org/10.1016/j.jacc.2003.12.059
189. Schömig A, Mehilli J, Antoniucci D, Ndrepepa G, Markwardt C, Di Pede F, *et al.* Mechanical reperfusion in patients with acute myocardial infarction presenting more than 12 hours from symptom onset: a randomized controlled trial. *JAMA* 2005;**293**:2865–2872. https://doi.org/10.1001/jama.293.23.2865
190. Ndrepepa G, Kastrati A, Mehilli J, Antoniucci D, Schömig A. Mechanical reperfusion and long-term mortality in patients with acute myocardial infarction presenting 12 to 48 hours from onset of symptoms. *JAMA* 2009;**301**:487–488. https://doi.org/10.1001/jama.2009.32
191. Bouisset F, Gerbaud E, Bataille V, Coste P, Puymirat E, Belle L, *et al.* Percutaneous myocardial revascularization in late-presenting patients with STEMI. *J Am Coll Cardiol* 2021;**78**:1291–1305. https://doi.org/10.1016/j.jacc.2021.07.039
192. Hochman JS, Lamas GA, Buller CE, Dzavik V, Reynolds HR, Abramsky SJ, *et al.* Coronary intervention for persistent occlusion after myocardial infarction. *N Engl J Med* 2006;**355**:2395–2407. https://doi.org/10.1056/NEJMoa066139
193. Menon V, Pearte CA, Buller CE, Steg PhG, Forman SA, White HD, *et al.* Lack of benefit from percutaneous intervention of persistently occluded infarct arteries after the acute phase of myocardial infarction is time independent: insights from Occluded Artery Trial. *Eur Heart J* 2009;**30**:183–191. https://doi.org/10.1093/eurheartj/ehn486
194. Ioannidis JPA, Katritsis DG. Percutaneous coronary intervention for late reperfusion after myocardial infarction in stable patients. *Am Heart J* 2007;**154**:1065–1071. https://doi.org/10.1016/j.ahj.2007.07.049
195. Knuuti J, Wijns W, Saraste A, Capodanno D, Barbato E, Funck-Brentano C, *et al.* 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J* 2020;**41**:407–477. https://doi.org/10.1093/eurheartj/ehz425
196. O'Donoghue M, Boden WE, Braunwald E, Cannon CP, Clayton TC, de Winter RJ, *et al.* Early invasive vs conservative treatment strategies in women and men with unstable angina and non-ST-segment elevation myocardial infarction: a meta-analysis. *JAMA* 2008;**300**:71–80. https://doi.org/10.1001/jama.300.1.71
197. Mehta SR, Cannon CP, Fox KAA, Wallentin L, Boden WE, Spacek R, *et al.* Routine vs selective invasive strategies in patients with acute coronary syndromes: a collaborative meta-analysis of randomized trials. *JAMA* 2005;**293**:2908–2917. https://doi.org/10.1001/jama.293.23.2908
198. Fox KA, Clayton TC, Damman P, Pocock SJ, de Winter RJ, Tijssen JGP, *et al.* Long-term outcome of a routine versus selective invasive strategy in patients with non-ST-segment elevation acute coronary syndrome a meta-analysis of individual patient data. *J Am Coll Cardiol* 2010;**55**:2435–2445. https://doi.org/10.1016/j.jacc.2010.03.007
199. Fanning JP, Nyong J, Scott IA, Aroney CN, Walters DL. Routine invasive strategies versus selective invasive strategies for unstable angina and non-ST elevation myocardial infarction in the stent era. *Cochrane Database Syst Rev* 2016;**2016**:CD004815. https://doi.org/10.1002/14651858.CD004815.pub4
200. Elgendy IY, Mahmoud AN, Wen X, Bavry AA. Meta-analysis of randomized trials of long-term all-cause mortality in patients with non-ST-elevation acute coronary syndrome managed with routine invasive versus selective invasive strategies. *Am J Cardiol* 2017;**119**:560–564. https://doi.org/10.1016/j.amjcard.2016.11.005
201. Navarese EP, Gurbel PA, Andreotti F, Tantry U, Jeong Y-H, Kozinski M, *et al.* Optimal timing of coronary invasive strategy in non-ST-segment elevation acute coronary syndromes: a systematic review and meta-analysis. *Ann Intern Med* 2013;**158**:261–270. https://doi.org/10.7326/0003-4819-158-4-201302190-00006
202. Jobs A, Mehta SR, Montalescot G, Vicaut E, van't Hof AWJ, Badings EA, *et al.* Optimal timing of an invasive strategy in patients with non-ST-elevation acute coronary syndrome: a meta-analysis of randomised trials. *Lancet* 2017;**390**:737–746. https://doi.org/10.1016/s0140-6736(17)31490-3
203. Kite TA, Kurmani SA, Bountziouka V, Cooper NJ, Lock ST, Gale CP, *et al.* Timing of invasive strategy in non-ST-elevation acute coronary syndrome: a meta-analysis of randomized controlled trials. *Eur Heart J* 2022;**43**:3148–3161. https://doi.org/10.1093/eurheartj/ehac213
204. Eggers KM, James SK, Jernberg T, Lindahl B. Timing of coronary angiography in patients with non-ST-elevation acute coronary syndrome: long-term clinical outcomes from the nationwide SWEDEHEART registry. *EuroIntervention* 2022;**18**:582–589. https://doi.org/10.4244/eij-d-21-00982
205. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet* 1994;**343**:311–322. https://doi.org/10.1016/S0140-6736(94)91161-4
206. Morrison LJ, Verbeek PR, McDonald AC, Sawadsky BV, Cook DJ. Mortality and prehospital thrombolysis for acute myocardial infarction: a meta-analysis. *JAMA* 2000;**283**:2686–2692. https://doi.org/10.1001/jama.283.20.2686
207. Steg PG, Bonnefoy E, Chabaud S, Lapostolle F, Dubien P-Y, Cristofini P, *et al.* Impact of time to treatment on mortality after prehospital fibrinolysis or primary angioplasty: data from the CAPTIM randomized clinical trial. *Circulation* 2003;**108**:2851–2856. https://doi.org/10.1161/01.Cir.0000103122.10021.F2
208. ASSENT-4 PCI Investigators. Primary versus tenecteplase-facilitated percutaneous coronary intervention in patients with ST-segment elevation acute myocardial infarction (ASSENT-4 PCI): randomised trial. *Lancet* 2006;**367**:569–578. https://doi.org/10.1016/s0140-6736(06)68147-6

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

209. Fazel R, Joseph TI, Sankardas MA, Pinto DS, Yeh RW, Kumbhani DJ, *et al.* Comparison of reperfusion strategies for ST-segment-elevation myocardial infarction: a multivariate network meta-analysis. *J Am Heart Assoc* 2020;**9**:e015186. https://doi.org/10.1161/jaha.119.015186
210. Sinnaeve PR, Armstrong PW, Gershlick AH, Goldstein P, Wilcox R, Lambert Y, *et al.* ST-segment-elevation myocardial infarction patients randomized to a pharmaco-invasive strategy or primary percutaneous coronary intervention: strategic Reperfusion Early After Myocardial Infarction (STREAM) 1-year mortality follow-up. *Circulation* 2014;**130**:1139–1145. https://doi.org/10.1161/circulationaha.114.009570
211. Arbel Y, Ko DT, Yan AT, Cantor WJ, Bagai A, Koh M, *et al.* Long-term follow-up of the trial of routine angioplasty and stenting after fibrinolysis to enhance reperfusion in acute myocardial infarction (TRANSFER-AMI). *Can J Cardiol* 2018;**34**:736–743. https://doi.org/10.1016/j.cjca.2018.02.005
212. Di Mario C, Dudek D, Piscione F, Mielecki W, Savonitto S, Murena E, *et al.* Immediate angioplasty versus standard therapy with rescue angioplasty after thrombolysis in the Combined Abciximab REteplase Stent Study in Acute Myocardial Infarction (CARESS-in-AMI): an open, prospective, randomised, multicentre trial. *Lancet* 2008;**371**:559–568. https://doi.org/10.1016/s0140-6736(08)60268-8
213. Bøhmer E, Hoffmann P, Abdelnoor M, Arnesen H, Halvorsen S. Efficacy and safety of immediate angioplasty versus ischemia-guided management after thrombolysis in acute myocardial infarction in areas with very long transfer distances results of the NORDISTEMI (NORwegian study on DIstrict treatment of ST-elevation myocardial infarction). *J Am Coll Cardiol* 2010;**55**:102–110. https://doi.org/10.1016/j.jacc.2009.08.007
214. Scheller B, Hennen B, Hammer B, Walle J, Hofer C, Hilpert V, *et al.* Beneficial effects of immediate stenting after thrombolysis in acute myocardial infarction. *J Am Coll Cardiol* 2003;**42**:634–641. https://doi.org/10.1016/s0735-1097(03)00763-0
215. Le May MR, Wells GA, Labinaz M, Davies RF, Turek M, Leddy D, *et al.* Combined angioplasty and pharmacological intervention versus thrombolysis alone in acute myocardial infarction (CAPITAL AMI study). *J Am Coll Cardiol* 2005;**46**:417–424. https://doi.org/10.1016/j.jacc.2005.04.042
216. Abdel-Qadir H, Yan AT, Tan M, Borgia F, Piscione F, Di Mario C, *et al.* Consistency of benefit from an early invasive strategy after fibrinolysis: a patient-level meta-analysis. *Heart* 2015;**101**:1554–1561. https://doi.org/10.1136/heartjnl-2015-307815
217. Madan M, Halvorsen S, Di Mario C, Tan M, Westerhout CM, Cantor WJ, *et al.* Relationship between time to invasive assessment and clinical outcomes of patients undergoing an early invasive strategy after fibrinolysis for ST-segment elevation myocardial infarction: a patient-level analysis of the randomized early routine invasive clinical trials. *JACC Cardiovasc Interv* 2015;**8**:166–174. https://doi.org/10.1016/j.jcin.2014.09.005
218. Andersen HR, Nielsen TT, Rasmussen K, Thuesen L, Kelbaek H, Thayssen P, *et al.* A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. *N Engl J Med* 2003;**349**:733–742. https://doi.org/10.1056/NEJMoa025142
219. Dalby M, Bouzamondo A, Lechat P, Montalescot G. Transfer for primary angioplasty versus immediate thrombolysis in acute myocardial infarction: a meta-analysis. *Circulation* 2003;**108**:1809–1814. https://doi.org/10.1161/01.Cir.0000091088.63921.8c
220. Gierlotka M, Gasior M, Wilczek K, Hawranek M, Szkodzinski J, Paczek P, *et al.* Reperfusion by primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction within 12 to 24 hours of the onset of symptoms (from a prospective national observational study [PL-ACS]). *Am J Cardiol* 2011;**107**:501–508. https://doi.org/10.1016/j.amjcard.2010.10.008
221. Busk M, Kaltoft A, Nielsen SS, Bottcher M, Rehling M, Thuesen L, *et al.* Infarct size and myocardial salvage after primary angioplasty in patients presenting with symptoms for<12 h vs. 12–72 h. *Eur Heart J* 2009;**30**:1322–1330. https://doi.org/10.1093/eurheartj/ehp113
222. Borgia F, Goodman SG, Halvorsen S, Cantor WJ, Piscione F, Le May MR, *et al.* Early routine percutaneous coronary intervention after fibrinolysis vs. standard therapy in ST-segment elevation myocardial infarction: a meta-analysis. *Eur Heart J* 2010;**31**:2156–2169. https://doi.org/10.1093/eurheartj/ehq204
223. D'Souza SP, Mamas MA, Fraser DG, Fath-Ordoubadi F. Routine early coronary angioplasty versus ischaemia-guided angioplasty after thrombolysis in acute ST-elevation myocardial infarction: a meta-analysis. *Eur Heart J* 2011;**32**:972–982. https://doi.org/10.1093/eurheartj/ehq398
224. Fernandez-Avilés F, Alonso JJ, Castro-Beiras A, Vázquez N, Blanco J, Alonso-Briales J, *et al.* Routine invasive strategy within 24 hours of thrombolysis versus ischaemia-guided conservative approach for acute myocardial infarction with ST-segment elevation (GRACIA-1): a randomised controlled trial. *Lancet* 2004;**364**:1045–1053. https://doi.org/10.1016/s0140-6736(04)17059-1
225. Hochman JS, Sleeper LA, White HD, Dzavik V, Wong SC, Menon V, *et al.* One-year survival following early revascularization for cardiogenic shock. *JAMA* 2001;**285**:190–192. https://doi.org/10.1001/jama.285.2.190
226. Lemkes JS, Janssens GN, van der Hoeven NW, van de Ven PM, Marques KMJ, Nap A, *et al.* Timing of revascularization in patients with transient ST-segment elevation myocardial infarction: a randomized clinical trial. *Eur Heart J* 2019;**40**:283–291. https://doi.org/10.1093/eurheartj/ehy651
227. Mehta SR, Granger CB, Boden WE, Steg PG, Bassand J-P, Faxon DP, *et al.* Early versus delayed invasive intervention in acute coronary syndromes. *N Engl J Med* 2009;**360**:2165–2175. https://doi.org/10.1056/NEJMoa0807986
228. Kofoed KF, Kelbæk H, Hansen PR, Torp-Pedersen C, Høfsten D, Kløvgaard L, *et al.* Early versus standard care invasive examination and treatment of patients with non-ST-segment elevation acute coronary syndrome. *Circulation* 2018;**138**:2741–2750. https://doi.org/10.1161/circulationaha.118.037152
229. Butt JH, Kofoed KF, Kelbæk H, Hansen PR, Torp-Pedersen C, Høfsten D, *et al.* Importance of risk assessment in timing of invasive coronary evaluation and treatment of patients with non-ST-segment-elevation acute coronary syndrome: insights from the VERDICT trial. *J Am Heart Assoc* 2021;**10**:e022333. https://doi.org/10.1161/jaha.121.022333
230. Barbarawi M, Kheiri B, Zayed Y, Barbarawi O, Chahine A, Haykal T, *et al.* Meta-analysis of optimal timing of coronary intervention in non-ST-elevation acute coronary syndrome. *Catheter Cardiovasc Interv* 2020;**95**:185–193. https://doi.org/10.1002/ccd.28280
231. Vranckx P, White HD, Huang Z, Mahaffey KW, Armstrong PW, Van de Werf F, *et al.* Validation of BARC bleeding criteria in patients with acute coronary syndromes: the TRACER trial. *J Am Coll Cardiol* 2016;**67**:2135–2144. https://doi.org/10.1016/j.jacc.2016.02.056
232. Ndrepepa G, Berger PB, Mehilli J, Seyfarth M, Neumann F-J, Schömig A, *et al.* Periprocedural bleeding and 1-year outcome after percutaneous coronary interventions: appropriateness of including bleeding as a component of a quadruple end point. *J Am Coll Cardiol* 2008;**51**:690–697. https://doi.org/10.1016/j.jacc.2007.10.040
233. Urban P, Mehran R, Colleran R, Angiolillo DJ, Byrne RA, Capodanno D, *et al.* Defining high bleeding risk in patients undergoing percutaneous coronary intervention: a consensus document from the Academic Research Consortium for High Bleeding Risk. *Eur Heart J* 2019;**40**:2632–2653. https://doi.org/10.1093/eurheartj/ehz372
234. Doomun D, Doomun I, Schukraft S, Arroyo D, Cook S, Huwyler T, *et al.* Ischemic and bleeding outcomes according to the academic research consortium high bleeding risk criteria in all comers treated by percutaneous coronary interventions. *Front Cardiovasc Med* 2021;**8**:620354. https://doi.org/10.3389/fcvm.2021.620354
235. Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy-I: prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *BMJ* 1994;**308**:81–106. https://doi.org/10.1136/bmj.308.6921.81
236. Valgimigli M, Bueno H, Byrne RA, Collet J-P, Costa F, Jeppsson A, *et al.* 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with EACTS: the Task Force for dual antiplatelet therapy in coronary artery disease of the European Society of Cardiology (ESC) and of the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2018;**39**:213–260. https://doi.org/10.1093/eurheartj/ehx419
237. Jones WS, Mulder H, Wruck LM, Pencina MJ, Kripalani S, Muñoz D, *et al.* Comparative effectiveness of aspirin dosing in cardiovascular disease. *N Engl J Med* 2021;**384**:1981–1990. https://doi.org/10.1056/NEJMoa2102137
238. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, *et al.* Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;**361**:1045–1057. https://doi.org/10.1056/NEJMoa0904327
239. Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, *et al.* Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2007;**357**:2001–2015. https://doi.org/10.1056/NEJMoa0706482
240. Aradi D, Kirtane A, Bonello L, Gurbel PA, Tantry US, Huber K, *et al.* Bleeding and stent thrombosis on P2Y12-inhibitors: collaborative analysis on the role of platelet reactivity for risk stratification after percutaneous coronary intervention. *Eur Heart J* 2015;**36**:1762–1771. https://doi.org/10.1093/eurheartj/ehv104
241. Aradi D, Storey RF, Komócsi A, Trenk D, Gulba D, Kiss RG, *et al.* Expert position paper on the role of platelet function testing in patients undergoing percutaneous coronary intervention. *Eur Heart J* 2014;**35**:209–215. https://doi.org/10.1093/eurheartj/eht375
242. Gimbel M, Qaderdan K, Willemsen L, Hermanides R, Bergmeijer T, de Vrey E, *et al.* Clopidogrel versus ticagrelor or prasugrel in patients aged 70 years or older with non-ST-elevation acute coronary syndrome (POPular AGE): the randomised, open-label, non-inferiority trial. *Lancet* 2020;**395**:1374–1381. https://doi.org/10.1016/s0140-6736(20)30325-1
243. Husted S, James S, Becker RC, Horrow J, Katus H, Storey RF, *et al.* Ticagrelor versus clopidogrel in elderly patients with acute coronary syndromes: a substudy from the prospective randomized PLATelet inhibition and patient Outcomes (PLATO) trial. *Circ Cardiovasc Qual Outcomes* 2012;**5**:680–688. https://doi.org/10.1161/circoutcomes.111.964395
244. Schüpke S, Neumann FJ, Menichelli M, Mayer K, Bernlochner I, Wöhrle J, *et al.* Ticagrelor or prasugrel in patients with acute coronary syndromes. *N Engl J Med* 2019;**381**:1524–1534. https://doi.org/10.1056/NEJMoa1908973
245. Montalescot G, van't Hof AW, Lapostolle F, Silvain J, Lassen JF, Bolognese L, *et al.* Prehospital ticagrelor in ST-segment elevation myocardial infarction. *N Engl J Med* 2014;**371**:1016–1027. https://doi.org/10.1056/NEJMoa1407024
246. Koul S, Smith JG, Götberg M, Omerovic E, Alfredsson J, Venetsanos D, *et al.* No benefit of ticagrelor pretreatment compared with treatment during percutaneous coronary

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

intervention in patients with ST-segment-elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Circ Cardiovasc Interv* 2018;**11**: e005528. https://doi.org/10.1161/circinterventions.117.005528
247. Montalescot G, Bolognese L, Dudek D, Goldstein P, Hamm C, Tanguay J-F, *et al.* Pretreatment with prasugrel in non-ST-segment elevation acute coronary syndromes. *N Engl J Med* 2013;**369**:999–1010. https://doi.org/10.1056/NEJMoa1308075
248. Tarantini G, Mojoli M, Varbella F, Caporale R, Rigattieri S, Andò G, *et al.* Timing of oral P2Y(12) inhibitor administration in patients with non-ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol* 2020;**76**:2450–2459. https://doi.org/10.1016/j.jacc.2020.08.053
249. Boersma E, Harrington RA, Moliterno DJ, White H, Théroux P, Van de Werf F, *et al.* Platelet glycoprotein IIb/IIIa inhibitors in acute coronary syndromes: a meta-analysis of all major randomised clinical trials. *Lancet* 2002;**359**:189–198. https://doi.org/10.1016/s0140-6736(02)07442-1
250. Neumann FJ, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, *et al.* 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019;**40**:87–165. https://doi.org/10.1093/eurheartj/ehy394
251. Harrington RA, Stone GW, McNulty S, White HD, Lincoff AM, Gibson CM, *et al.* Platelet inhibition with cangrelor in patients undergoing PCI. *N Engl J Med* 2009; **361**:2318–2329. https://doi.org/10.1056/NEJMoa0908628
252. Bhatt DL, Stone GW, Mahaffey KW, Gibson CM, Steg PG, Hamm CW, *et al.* Effect of platelet inhibition with cangrelor during PCI on ischemic events. *N Engl J Med* 2013; **368**:1303–1313. https://doi.org/10.1056/NEJMoa1300815
253. Bhatt DL, Lincoff AM, Gibson CM, Stone GW, McNulty S, Montalescot G, *et al.* Intravenous platelet blockade with cangrelor during PCI. *N Engl J Med* 2009;**361**: 2330–2341. https://doi.org/10.1056/NEJMoa0908629
254. Steg PG, Bhatt DL, Hamm CW, Stone GW, Gibson CM, Mahaffey KW, *et al.* Effect of cangrelor on periprocedural outcomes in percutaneous coronary interventions: a pooled analysis of patient-level data. *Lancet* 2013;**382**:1981–1992. https://doi.org/10.1016/s0140-6736(13)61615-3
255. Eikelboom JW, Anand SS, Malmberg K, Weitz JI, Ginsberg JS, Yusuf S. Unfractionated heparin and low-molecular-weight heparin in acute coronary syndrome without ST elevation: a meta-analysis. *Lancet* 2000;**355**:1936–1942. https://doi.org/10.1016/s0140-6736(00)02324-2
256. Ferguson JJ, Califf RM, Antman EM, Cohen M, Grines CL, Goodman S, *et al.* Enoxaparin vs unfractionated heparin in high-risk patients with non-ST-segment elevation acute coronary syndromes managed with an intended early invasive strategy: primary results of the SYNERGY randomized trial. *JAMA* 2004;**292**:45–54. https://doi.org/10.1001/jama.292.1.45
257. Cohen M, Mahaffey KW, Pieper K, Pollack CV, Antman EM, Hoekstra J, *et al.* A subgroup analysis of the impact of prerandomization antithrombin therapy on outcomes in the SYNERGY trial: enoxaparin versus unfractionated heparin in non-ST-segment elevation acute coronary syndromes. *J Am Coll Cardiol* 2006;**48**:1346–1354. https://doi.org/10.1016/j.jacc.2006.05.058
258. Montalescot G, Zeymer U, Silvain J, Boulanger B, Cohen M, Goldstein P, *et al.* Intravenous enoxaparin or unfractionated heparin in primary percutaneous coronary intervention for ST-elevation myocardial infarction: the international randomised open-label ATOLL trial. *Lancet* 2011;**378**:693–703. https://doi.org/10.1016/s0140-6736(11)60876-3
259. Li Y, Liang Z, Qin L, Wang M, Wang X, Zhang H, *et al.* Bivalirudin plus a high-dose infusion versus heparin monotherapy in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: a randomised trial. *Lancet* 2022;**400**:1847–1857. https://doi.org/10.1016/s0140-6736(22)01999-7
260. Yusuf S, Mehta SR, Chrolavicius S, Cohen M, Grines CL, Goodman S, *et al.* Effects of fondaparinux on mortality and reinfarction in patients with acute ST-segment elevation myocardial infarction: the OASIS-6 randomized trial. *JAMA* 2006;**295**: 1519–1530. https://doi.org/10.1001/jama.295.13.joc60038
261. Silvain J, Beygui F, Barthélémy O, Pollack C, Cohen M, Zeymer U, *et al.* Efficacy and safety of enoxaparin versus unfractionated heparin during percutaneous coronary intervention: systematic review and meta-analysis. *BMJ* 2012;**344**:e553. https://doi.org/10.1136/bmj.e553
262. The Fifth Organization to Assess Strategies in Acute Ischemic Syndromes Investigators. Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 2006;**354**:1464–1476. https://doi.org/10.1056/NEJMoa055443
263. Mehta SR, Yusuf S, Peters RJ, Bertrand ME, Lewis BS, Natarajan MK, *et al.* Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: the PCI-CURE study. *Lancet* 2001; **358**:527–533. https://doi.org/10.1016/s0140-6736(01)05701-4
264. Palmerini T, Della Riva D, Benedetto U, Bacchi Reggiani L, Feres F, Abizaid A, *et al.* Three, six, or twelve months of dual antiplatelet therapy after DES implantation in patients with or without acute coronary syndromes: an individual patient data pairwise and network meta-analysis of six randomized trials and 11 473 patients. *Eur Heart J* 2017;**38**:1034–1043. https://doi.org/10.1093/eurheartj/ehw627
265. Hahn J-Y, Song YB, Oh J-H, Cho D-K, Lee JB, Doh J-H, *et al.* 6-month versus 12-month or longer dual antiplatelet therapy after percutaneous coronary intervention in patients with acute coronary syndrome (SMART-DATE): a randomised, open-label, non-inferiority trial. *Lancet* 2018;**391**:1274–1284. https://doi.org/10.1016/s0140-6736(18)30493-8
266. Kedhi E, Fabris E, van der Ent M, Buszman P, von Birgelen C, Roolvink V, *et al.* Six months versus 12 months dual antiplatelet therapy after drug-eluting stent implantation in ST-elevation myocardial infarction (DAPT-STEMI): randomised, multicentre, non-inferiority trial. *BMJ* 2018;**363**:k3793. https://doi.org/10.1136/bmj.k3793
267. De Luca G, Damen SA, Camaro C, Benit E, Verdoia M, Rasoul S, *et al.* Final results of the randomised evaluation of short-term dual antiplatelet therapy in patients with acute coronary syndrome treated with a new-generation stent (REDUCE trial). *EuroIntervention* 2019;**15**:e990–e998. https://doi.org/10.4244/eij-d-19-00539
268. Hahn J-Y, Song YB, Oh J-H, Chun WJ, Park YH, Jang WJ, *et al.* Effect of P2Y12 inhibitor monotherapy vs dual antiplatelet therapy on cardiovascular events in patients undergoing percutaneous coronary intervention: the SMART-CHOICE randomized clinical trial. *JAMA* 2019;**321**:2428–2437. https://doi.org/10.1001/jama.2019.8146
269. Vranckx P, Valgimigli M, Jüni P, Hamm C, Steg PG, Heg D, *et al.* Ticagrelor plus aspirin for 1 month, followed by ticagrelor monotherapy for 23 months vs aspirin plus clopidogrel or ticagrelor for 12 months, followed by aspirin monotherapy for 12 months after implantation of a drug-eluting stent: a multicentre, open-label, randomised superiority trial. *Lancet* 2018;**392**:940–949. https://doi.org/10.1016/s0140-6736(18)31858-0
270. Watanabe H, Domei T, Morimoto T, Natsuaki M, Shiomi H, Toyota T, *et al.* Effect of 1-month dual antiplatelet therapy followed by clopidogrel vs 12-month dual antiplatelet therapy on cardiovascular and bleeding events in patients receiving PCI: the STOPDAPT-2 randomized clinical trial. *JAMA* 2019;**321**:2414–2427. https://doi.org/10.1001/jama.2019.8145
271. Mehran R, Baber U, Sharma SK, Cohen DJ, Angiolillo DJ, Briguori C, *et al.* Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;**381**: 2032–2042. https://doi.org/10.1056/NEJMoa1908419
272. Baber U, Dangas G, Angiolillo DJ, Cohen DJ, Sharma SK, Nicolas J, *et al.* Ticagrelor alone vs. ticagrelor plus aspirin following percutaneous coronary intervention in patients with non-ST-segment elevation acute coronary syndromes: TWILIGHT-ACS. *Eur Heart J* 2020;**41**:3533–3545. https://doi.org/10.1093/eurheartj/ehaa670
273. Kim BK, Hong SJ, Cho YH, Yun KHo, Kim YH, Suh Y, *et al.* Effect of ticagrelor monotherapy vs ticagrelor with aspirin on major bleeding and cardiovascular events in patients with acute coronary syndrome: the TICO randomized clinical trial. *JAMA* 2020;**323**:2407–2416. https://doi.org/10.1001/jama.2020.7580
274. Giacoppo D, Matsuda Y, Fovino LN, D'Amico G, Gargiulo G, Byrne RA, *et al.* Short dual antiplatelet therapy followed by P2Y12 inhibitor monotherapy vs. prolonged dual antiplatelet therapy after percutaneous coronary intervention with second-generation drug-eluting stents: a systematic review and meta-analysis of randomized clinical trials. *Eur Heart J* 2021;**42**:308–319. https://doi.org/10.1093/eurheartj/ehaa739
275. Watanabe H, Morimoto T, Natsuaki M, Yamamoto K, Obayashi Y, Ogita M, *et al.* Comparison of clopidogrel monotherapy after 1 to 2 months of dual antiplatelet therapy with 12 months of dual antiplatelet therapy in patients with acute coronary syndrome: the STOPDAPT-2 ACS randomized clinical trial. *JAMA Cardiol* 2022;**7**: 407–417. https://doi.org/10.1001/jamacardio.2021.5244
276. Valgimigli M, Frigoli E, Heg D, Tijssen J, Jüni P, Vranckx P, *et al.* Dual antiplatelet therapy after PCI in patients at high bleeding risk. *N Engl J Med* 2021;**385**:1643–1655. https://doi.org/10.1056/NEJMoa2108749
277. Zettler ME, Peterson ED, McCoy LA, Effron MB, Anstrom KJ, Henry TD, *et al.* Switching of adenosine diphosphate receptor inhibitor after hospital discharge among myocardial infarction patients: insights from the treatment with adenosine diphosphate receptor inhibitors: longitudinal assessment of treatment patterns and events after acute coronary syndrome (TRANSLATE-ACS) observational study. *Am Heart J* 2017;**183**:62–68. https://doi.org/10.1016/j.ahj.2016.10.006
278. Angiolillo DJ, Rollini F, Storey RF, Bhatt DL, James S, Schneider DJ, *et al.* International expert consensus on switching platelet P2Y(12) receptor-inhibiting therapies. *Circulation* 2017;**136**:1955–1975. https://doi.org/10.1161/circulationaha.117.031164
279. Sibbing D, Aradi D, Jacobshagen C, Gross L, Trenk D, Geisler T, *et al.* Guided de-escalation of antiplatelet treatment in patients with acute coronary syndrome undergoing percutaneous coronary intervention (TROPICAL-ACS): a randomised, open-label, multicentre trial. *Lancet* 2017;**390**:1747–1757. https://doi.org/10.1016/s0140-6736(17)32155-4
280. Claassens DMF, Vos GJA, Bergmeijer TO, Hermanides RS, van't Hof AWJ, van der Harst P, *et al.* A genotype-guided strategy for oral P2Y(12) inhibitors in primary PCI. *N Engl J Med* 2019;**381**:1621–1631. https://doi.org/10.1056/NEJMoa1907096
281. Cuisset T, Deharo P, Quilici J, Johnson TW, Deffarges S, Bassez C, *et al.* Benefit of switching dual antiplatelet therapy after acute coronary syndrome: the TOPIC (timing of platelet inhibition after acute coronary syndrome) randomized study. *Eur Heart J* 2017;**38**:3070–3078. https://doi.org/10.1093/eurheartj/ehx175
282. Kim CJ, Park MW, Kim MC, Choo EH, Hwang B-H, Lee KY, *et al.* Unguided de-escalation from ticagrelor to clopidogrel in stabilised patients with acute myocardial infarction undergoing percutaneous coronary intervention (TALOS-AMI): an investigator-initiated, open-label, multicentre, non-inferiority, randomised trial. *Lancet* 2021;**398**:1305–1316. https://doi.org/10.1016/s0140-6736(21)01445-8

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

283. Kim HS, Kang J, Hwang D, Han J-K, Yang H-M, Kang H-J, *et al.* Prasugrel-based de-escalation of dual antiplatelet therapy after percutaneous coronary intervention in patients with acute coronary syndrome (HOST-REDUCE-POLYTECH-ACS): an open-label, multicentre, non-inferiority randomised trial. *Lancet* 2020;**396**:1079–1089. https://doi.org/10.1016/s0140-6736(20)31791-8
284. Antithrombotic Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ* 2002;**324**:71–86. https://doi.org/10.1136/bmj.324.7329.71
285. Antithrombotic Trialists' (ATT) Collaboration. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* 2009;**373**:1849–1860. https://doi.org/10.1016/s0140-6736(09)60503-1
286. Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK, *et al.* Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 2001;**345**:494–502. https://doi.org/10.1056/NEJMoa010746
287. Bhatt DL, Cryer BL, Contant CF, Cohen M, Lanas A, Schnitzer TJ, *et al.* Clopidogrel with or without omeprazole in coronary artery disease. *N Engl J Med* 2010;**363**:1909–1917. https://doi.org/10.1056/NEJMoa1007964
288. Gargiulo G, Costa F, Ariotti S, Biscaglia S, Campo G, Esposito G, *et al.* Impact of proton pump inhibitors on clinical outcomes in patients treated with a 6- or 24-month dual-antiplatelet therapy duration: insights from the PROlonging Dual-antiplatelet treatment after Grading stent-induced Intimal hyperplasia studY trial. *Am Heart J* 2016;**174**:95–102. https://doi.org/10.1016/j.ahj.2016.01.015
289. Mehta SR, Tanguay JF, Eikelboom JW, Jolly SS, Joyner CD, Granger CB, *et al.* Double-dose versus standard-dose clopidogrel and high-dose versus low-dose aspirin in individuals undergoing percutaneous coronary intervention for acute coronary syndromes (CURRENT-OASIS 7): a randomised factorial trial. *Lancet* 2010;**376**:1233–1243. https://doi.org/10.1016/s0140-6736(10)61088-4
290. Coughlan JJ, Aytekin A, Lahu S, Ndrepepa G, Menichelli M, Mayer K, *et al.* Ticagrelor or prasugrel for patients with acute coronary syndrome treated with percutaneous coronary intervention: a prespecified subgroup analysis of a randomized clinical trial. *JAMA Cardiol* 2021;**6**:1121–1129. https://doi.org/10.1001/jamacardio.2021.2228
291. Szummer K, Montez-Rath ME, Alfredsson J, Erlinge D, Lindahl B, Hofmann R, *et al.* Comparison between ticagrelor and clopidogrel in elderly patients with an acute coronary syndrome: insights from the SWEDEHEART registry. *Circulation* 2020;**142**:1700–1708. https://doi.org/10.1161/circulationaha.120.050645
292. Giugliano RP, White JA, Bode C, Armstrong PW, Montalescot G, Lewis BS, *et al.* Early versus delayed, provisional eptifibatide in acute coronary syndromes. *N Engl J Med* 2009;**360**:2176–2190. https://doi.org/10.1056/NEJMoa0901316
293. Dworeck C, Redfors B, Angerås O, Haraldsson I, Odenstedt J, Ioanes D, *et al.* Association of pretreatment with P2Y12 receptor antagonists preceding percutaneous coronary intervention in non-ST-segment elevation acute coronary syndromes with outcomes. *JAMA Netw Open* 2020;**3**:e2018735. https://doi.org/10.1001/jamanetworkopen.2020.18735
294. Steinhubl SR, Berger PB, Mann JT III, Fry ETA, DeLago A, Wilmer C, *et al.* Early and sustained dual oral antiplatelet therapy following percutaneous coronary intervention: a randomized controlled trial. *JAMA* 2002;**288**:2411–2420. https://doi.org/10.1001/jama.288.19.2411
295. Dawson LP, Chen D, Dagan M, Bloom J, Taylor A, Duffy SJ, *et al.* Assessment of pretreatment with oral P2Y12 inhibitors and cardiovascular and bleeding outcomes in patients with non-ST elevation acute coronary syndromes: a systematic review and meta-analysis. *JAMA Netw Open* 2021;**4**:e2134322. https://doi.org/10.1001/jamanetworkopen.2021.34322
296. Oler A, Whooley MA, Oler J, Grady D. Adding heparin to aspirin reduces the incidence of myocardial infarction and death in patients with unstable angina. A meta-analysis. *JAMA* 1996;**276**:811–815. doi: doi:10.1001/jama.1996.03540100055028
297. Antman EM, Cohen M, Radley D, McCabe C, Rush J, Premmereur J, *et al.* Assessment of the treatment effect of enoxaparin for unstable angina/non-Q-wave myocardial infarction: TIMI 11B-ESSENCE meta-analysis. *Circulation* 1999;**100**:1602–1608. https://doi.org/10.1161/01.cir.100.15.1602
298. Collet JP, Huber K, Cohen M, Zeymer U, Goldstein P, Pollack C, *et al.* A direct comparison of intravenous enoxaparin with unfractionated heparin in primary percutaneous coronary intervention (from the ATOLL trial). *Am J Cardiol* 2013;**112**:1367–1372. https://doi.org/10.1016/j.amjcard.2013.07.003
299. Stone GW, McLaurin BT, Cox DA, Bertrand ME, Lincoff AM, Moses JW, *et al.* Bivalirudin for patients with acute coronary syndromes. *N Engl J Med* 2006;**355**:2203–2216. https://doi.org/10.1056/NEJMoa062437
300. Valgimigli M, Frigoli E, Leonardi S, Rothenbühler M, Gagnor A, Calabrò P, *et al.* Bivalirudin or unfractionated heparin in acute coronary syndromes. *N Engl J Med* 2015;**373**:997–1009. https://doi.org/10.1056/NEJMoa1507854
301. Kastrati A, Neumann FJ, Schulz S, Massberg S, Byrne RA, Ferenc M, *et al.* Abciximab and heparin versus bivalirudin for non-ST-elevation myocardial infarction. *N Engl J Med* 2011;**365**:1980–1989. https://doi.org/10.1056/NEJMoa1109596
302. Erlinge D, Omerovic E, Fröbert O, Linder R, Danielewicz M, Hamid M, *et al.* Bivalirudin versus heparin monotherapy in myocardial infarction. *N Engl J Med* 2017;**377**:1132–1142. https://doi.org/10.1056/NEJMoa1706443
303. Han Y, Guo J, Zheng Y, Zang H, Su X, Wang Y, *et al.* Bivalirudin vs heparin with or without tirofiban during primary percutaneous coronary intervention in acute myocardial infarction: the BRIGHT randomized clinical trial. *JAMA* 2015;**313**:1336–1346. https://doi.org/10.1001/jama.2015.2323
304. Steg PG, Jolly SS, Mehta SR, Afzal R, Xavier D, Rupprecht H-J, *et al.* Low-dose vs standard-dose unfractionated heparin for percutaneous coronary intervention in acute coronary syndromes treated with fondaparinux: the FUTURA/OASIS-8 randomized trial. *JAMA* 2010;**304**:1339–1349. https://doi.org/10.1001/jama.2010.1320
305. Cannon CP, Bhatt DL, Oldgren J, Lip GYH, Ellis SG, Kimura T, *et al.* Dual antithrombotic therapy with dabigatran after PCI in atrial fibrillation. *N Engl J Med* 2017;**377**:1513–1524. https://doi.org/10.1056/NEJMoa1708454
306. Dewilde WJ, Oirbans T, Verheugt FW, Kelder JC, De Smet BJGL, Herrman J-P, *et al.* Use of clopidogrel with or without aspirin in patients taking oral anticoagulant therapy and undergoing percutaneous coronary intervention: an open-label, randomised, controlled trial. *Lancet* 2013;**381**:1107–1115. https://doi.org/10.1016/s0140-6736(12)62177-1
307. Gibson CM, Mehran R, Bode C, Halperin J, Verheugt FW, Wildgoose P, *et al.* Prevention of bleeding in patients with atrial fibrillation undergoing PCI. *N Engl J Med* 2016;**375**:2423–2434. https://doi.org/10.1056/NEJMoa1611594
308. Lopes RD, Heizer G, Aronson R, Vora AN, Massaro T, Mehran R, *et al.* Antithrombotic therapy after acute coronary syndrome or PCI in atrial fibrillation. *N Engl J Med* 2019;**380**:1509–1524. https://doi.org/10.1056/NEJMoa1817083
309. Vranckx P, Valgimigli M, Eckardt L, Tijssen J, Lewalter T, Gargiulo G, *et al.* Edoxaban-based versus vitamin K antagonist-based antithrombotic regimen after successful coronary stenting in patients with atrial fibrillation (ENTRUST-AF PCI): a randomised, open-label, phase 3b trial. *Lancet* 2019;**394**:1335–1343. https://doi.org/10.1016/s0140-6736(19)31872-0
310. Gargiulo G, Goette A, Tijssen J, Eckardt L, Lewalter T, Vranckx P, *et al.* Safety and efficacy outcomes of double vs. triple antithrombotic therapy in patients with atrial fibrillation following percutaneous coronary intervention: a systematic review and meta-analysis of non-vitamin K antagonist oral anticoagulant-based randomized clinical trials. *Eur Heart J* 2019;**40**:3757–3767. https://doi.org/10.1093/eurheartj/ehz732
311. Dewilde WJ, Janssen PW, Kelder JC, Verheugt FWA, De Smet BJGL, Adriaenssens T, *et al.* Uninterrupted oral anticoagulation versus bridging in patients with long-term oral anticoagulation during percutaneous coronary intervention: subgroup analysis from the WOEST trial. *EuroIntervention* 2015;**11**:381–390. https://doi.org/10.4244/eijy14m06_07
312. Windecker S, Lopes RD, Massaro T, Jones-Burton C, Granger CB, Aronson R, *et al.* Antithrombotic therapy in patients with atrial fibrillation and acute coronary syndrome treated medically or with percutaneous coronary intervention or undergoing elective percutaneous coronary intervention: insights from the AUGUSTUS trial. *Circulation* 2019;**140**:1921–1932. https://doi.org/10.1161/circulationaha.119.043308
313. Smits PC, Frigoli E, Tijssen J, Jüni P, Vranckx P, Ozaki Y, *et al.* Abbreviated antiplatelet therapy in patients at high bleeding risk with or without oral anticoagulant therapy after coronary stenting: an open-label, randomized, controlled trial. *Circulation* 2021;**144**:1196–1211. https://doi.org/10.1161/circulationaha.121.056680
314. Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, Shestakovska O, *et al.* Rivaroxaban with or without aspirin in stable cardiovascular disease. *N Engl J Med* 2017;**377**:1319–1330. https://doi.org/10.1056/NEJMoa1709118
315. Connolly SJ, Eikelboom JW, Bosch J, Dagenais G, Dyal L, Lanas F, *et al.* Rivaroxaban with or without aspirin in patients with stable coronary artery disease: an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2018;**391**:205–218. https://doi.org/10.1016/s0140-6736(17)32458-3
316. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE, Steg PG, *et al.* Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med* 2014;**371**:2155–2166. https://doi.org/10.1056/NEJMoa1409312
317. Bonaca MP, Bhatt DL, Steg PG, Storey RF, Cohen M, Im K, *et al.* Ischaemic risk and efficacy of ticagrelor in relation to time from P2Y12 inhibitor withdrawal in patients with prior myocardial infarction: insights from PEGASUS-TIMI 54. *Eur Heart J* 2016;**37**:1133–1142. https://doi.org/10.1093/eurheartj/ehv531
318. Bonaca MP, Bhatt DL, Cohen M, Steg PG, Storey RF, Jensen EC, *et al.* Long-term use of ticagrelor in patients with prior myocardial infarction. *N Engl J Med* 2015;**372**:1791–1800. https://doi.org/10.1056/NEJMoa1500857
319. Steg PG, Bhatt DL, Simon T, Fox K, Mehta SR, Harrington RA, *et al.* Ticagrelor in patients with stable coronary disease and diabetes. *N Engl J Med* 2019;**381**:1309–1320. https://doi.org/10.1056/NEJMoa1908077
320. Valgimigli M, Gragnano F, Branca M, Franzone A, Baber U, Jang Y, *et al.* P2Y12 inhibitor monotherapy or dual antiplatelet therapy after coronary revascularisation: individual patient level meta-analysis of randomised controlled trials. *BMJ* 2021;**373**:n1332. https://doi.org/10.1136/bmj.n1332
321. Shoji S, Kuno T, Fujisaki T, Takagi H, Briasoulis A, Deharo P, *et al.* De-escalation of dual antiplatelet therapy in patients with acute coronary syndromes. *J Am Coll Cardiol* 2021;**78**:763–777. https://doi.org/10.1016/j.jacc.2021.06.012

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

322. Laudani C, Greco A, Occhipinti G, Ingala S, Calderone D, Scalia L, *et al.* Short duration of DAPT versus de-escalation after percutaneous coronary intervention for acute coronary syndromes. *JACC Cardiovasc Interv* 2022;**15**:268–277. https://doi.org/10.1016/j.jcin.2021.11.028
323. Giustino G, Mehran R, Dangas GD, Kirtane AJ, Redfors B, Généreux P, *et al.* Characterization of the average daily ischemic and bleeding risk after primary PCI for STEMI. *J Am Coll Cardiol* 2017;**70**:1846–1857. https://doi.org/10.1016/j.jacc.2017.08.018
324. Yasuda S, Kaikita K, Akao M, Ako J, Matoba T, Nakamura M, *et al.* Antithrombotic therapy for atrial fibrillation with stable coronary disease. *N Engl J Med* 2019;**381**:1103–1113. https://doi.org/10.1056/NEJMoa1904143
325. Matsumura-Nakano Y, Shizuta S, Komasa A, Morimoto T, Masuda H, Shiomi H, *et al.* Open-label randomized trial comparing oral anticoagulation with and without single antiplatelet therapy in patients with atrial fibrillation and stable coronary artery disease beyond 1 year after coronary stent implantation. *Circulation* 2019;**139**:604–616. https://doi.org/10.1161/circulationaha.118.036768
326. Chiarito M, Sanz-Sánchez J, Cannata F, Cao D, Sturla M, Panico C, *et al.* Monotherapy with a P2Y(12) inhibitor or aspirin for secondary prevention in patients with established atherosclerosis: a systematic review and meta-analysis. *Lancet* 2020;**395**:1487–1495. https://doi.org/10.1016/s0140-6736(20)30315-9
327. Koo BK, Kang J, Park KW, Rhee T-M, Yang H-M, Won KB, *et al.* Aspirin versus clopidogrel for chronic maintenance monotherapy after percutaneous coronary intervention (HOST-EXAM): an investigator-initiated, prospective, randomised, open-label, multicentre trial. *Lancet* 2021;**397**:2487–2496. https://doi.org/10.1016/s0140-6736(21)01063-1
328. Gilard M, Blanchard D, Helft G, Carrier D, Eltchaninoff H, Belle L, *et al.* Antiplatelet therapy in patients with anticoagulants undergoing percutaneous coronary stenting (from STENTIng and oral antiCOagulants [STENTICO]). *Am J Cardiol* 2009;**104**:338–342. https://doi.org/10.1016/j.amjcard.2009.03.053
329. Lip GY, Windecker S, Huber K, Kirchhof P, Marin F, Ten Berg JM, *et al.* Management of antithrombotic therapy in atrial fibrillation patients presenting with acute coronary syndrome and/or undergoing percutaneous coronary or valve interventions: a joint consensus document of the European Society of Cardiology Working Group on Thrombosis, European Heart Rhythm Association (EHRA), European Association of Percutaneous Cardiovascular Interventions (EAPCI) and European Association of Acute Cardiac Care (ACCA) endorsed by the Heart Rhythm Society (HRS) and Asia-Pacific Heart Rhythm Society (APHRS). *Eur Heart J* 2014;**35**:3155–3179. https://doi.org/10.1093/eurheartj/ehu298
330. Ruiz-Nodar JM, Marín F, Hurtado JA, Valencia J, Pinar E, Pineda J, *et al.* Anticoagulant and antiplatelet therapy use in 426 patients with atrial fibrillation undergoing percutaneous coronary intervention and stent implantation implications for bleeding risk and prognosis. *J Am Coll Cardiol* 2008;**51**:818–825. https://doi.org/10.1016/j.jacc.2007.11.035
331. Beyer-Westendorf J, Gelbricht V, Förster K, Ebertz F, Kohler C, Werth S, *et al.* Peri-interventional management of novel oral anticoagulants in daily care: results from the prospective Dresden NOAC registry. *Eur Heart J* 2014;**35**:1888–1896. https://doi.org/10.1093/eurheartj/eht557
332. Kiviniemi T, Karjalainen P, Pietilä M, Ylitalo A, Niemelä M, Vikman S, *et al.* Comparison of additional versus no additional heparin during therapeutic oral anticoagulation in patients undergoing percutaneous coronary intervention. *Am J Cardiol* 2012;**110**:30–35. https://doi.org/10.1016/j.amjcard.2012.02.045
333. Fiedler KA, Maeng M, Mehilli J, Schulz-Schüpke S, Byrne RA, Sibbing D, *et al.* Duration of triple therapy in patients requiring oral anticoagulation after drug-eluting stent implantation: the ISAR-TRIPLE trial. *J Am Coll Cardiol* 2015;**65**:1619–1629. https://doi.org/10.1016/j.jacc.2015.02.050
334. Lopes RD, Leonardi S, Wojdyla DM, Vora AN, Thomas L, Storey RF, *et al.* Stent thrombosis in patients with atrial fibrillation undergoing coronary stenting in the AUGUSTUS trial. *Circulation* 2020;**141**:781–783. https://doi.org/10.1161/circulationaha.119.044584
335. Alexander JH, Wojdyla D, Vora AN, Thomas L, Granger CB, Goodman SG, *et al.* Risk/benefit tradeoff of antithrombotic therapy in patients with atrial fibrillation early and late after an acute coronary syndrome or percutaneous coronary intervention: insights from AUGUSTUS. *Circulation* 2020;**141**:1618–1627. https://doi.org/10.1161/circulationaha.120.046534
336. Lopes RD, Hong H, Harskamp RE, Bhatt DL, Mehran R, Cannon CP, *et al.* Safety and efficacy of antithrombotic strategies in patients with atrial fibrillation undergoing percutaneous coronary intervention: a network meta-analysis of randomized controlled trials. *JAMA Cardiol* 2019;**4**:747–755. https://doi.org/10.1001/jamacardio.2019.1880
337. Lopes RD, Hong H, Harskamp RE, Bhatt DL, Mehran R, Cannon CP, *et al.* Optimal antithrombotic regimens for patients with atrial fibrillation undergoing percutaneous coronary intervention: an updated network meta-analysis. *JAMA Cardiol* 2020;**5**:582–589. https://doi.org/10.1001/jamacardio.2019.6175
338. Capodanno D, Di Maio M, Greco A, Bhatt DL, Gibson CM, Goette A, *et al.* Safety and efficacy of double antithrombotic therapy with non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation undergoing percutaneous coronary intervention: a systematic review and meta-analysis. *J Am Heart Assoc* 2020;**9**:e017212. https://doi.org/10.1161/jaha.120.017212
339. Lip GYH, Collet JP, Haude M, Byrne R, Chung EH, Fauchier L, *et al.* 2018 Joint European consensus document on the management of antithrombotic therapy in atrial fibrillation patients presenting with acute coronary syndrome and/or undergoing percutaneous cardiovascular interventions: a joint consensus document of the European Heart Rhythm Association (EHRA), European Society of Cardiology Working Group on Thrombosis, European Association of Percutaneous Cardiovascular Interventions (EAPCI), and European Association of Acute Cardiac Care (ACCA) endorsed by the Heart Rhythm Society (HRS), Asia-Pacific Heart Rhythm Society (APHRS), Latin America Heart Rhythm Society (LAHRS), and Cardiac Arrhythmia Society of Southern Africa (CASSA). *Europace* 2019;**21**:192–193. https://doi.org/10.1093/europace/euy174
340. ISIS-2 Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;**2**:349–360.
341. Chen ZM, Jiang LX, Chen YP, Xie JX, Pan HC, Peto R, *et al.* Addition of clopidogrel to aspirin in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. *Lancet* 2005;**366**:1607–1621. https://doi.org/10.1016/s0140-6736(05)67660-x
342. Sabatine MS, Cannon CP, Gibson CM, López-Sendón JL, Montalescot G, Theroux P, *et al.* Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Eng J Med* 2005;**352**:1179–1189. https://doi.org/10.1056/NEJMoa050522
343. Osman M, Kheiri B, Shigle AJ, Saleem M, Osman K, Sengupta PP, *et al.* Ticagrelor after pharmacological thrombolysis in patients with ST-segment elevation myocardial infarctions: insight from a trial sequential analysis. *J Thromb Thrombolysis* 2019;**48**:661–667. https://doi.org/10.1007/s11239-019-01953-3
344. Berwanger O, Lopes RD, Moia DDF, Fonseca FA, Jiang L, Goodman SG, *et al.* Ticagrelor versus clopidogrel in patients with STEMI treated with fibrinolysis: TREAT trial. *J Am Coll Cardiol* 2019;**73**:2819–2828. https://doi.org/10.1016/j.jacc.2019.03.011
345. Kheiri B, Osman M, Abdalla A, Haykal T, Barbarawi M, Zayed Y, *et al.* Ticagrelor versus clopidogrel after fibrinolytic therapy in patients with ST-elevation myocardial infarction: a systematic review and meta-analysis of randomized clinical trials. *J Thromb Thrombolysis* 2018;**46**:299–303. https://doi.org/10.1007/s11239-018-1706-2
346. Sánchez PL, Gimeno F, Ancillo P, Sanz JJ, Alonso-Briales JH, Bosa F, *et al.* Role of the paclitaxel-eluting stent and tirofiban in patients with ST-elevation myocardial infarction undergoing postfibrinolysis angioplasty: the GRACIA-3 randomized clinical trial. *Circ Cardiovasc Interv* 2010;**3**:297–307. https://doi.org/10.1161/circinterventions.109.920868
347. ASSENT-3 Investigators. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: the ASSENT-3 randomised trial in acute myocardial infarction. *Lancet* 2001;**358**:605–613. https://doi.org/10.1016/s0140-6736(01)05775-0
348. Giraldez RR, Nicolau JC, Corbalan R, Gurfinkel EP, Juarez U, Lopez-Sendon J, *et al.* Enoxaparin is superior to unfractionated heparin in patients with ST elevation myocardial infarction undergoing fibrinolysis regardless of the choice of lytic: an ExTRACT-TIMI 25 analysis. *Eur Heart J* 2007;**28**:1566–1573. https://doi.org/10.1093/eurheartj/ehm179
349. White HD, Braunwald E, Murphy SA, Jacob AJ, Gotcheva N, Polonetsky L, *et al.* Enoxaparin vs. unfractionated heparin with fibrinolysis for ST-elevation myocardial infarction in elderly and younger patients: results from ExTRACT-TIMI 25. *Eur Heart J* 2007;**28**:1066–1071. https://doi.org/10.1093/eurheartj/ehm081
350. Peters RJ, Joyner C, Bassand JP, Afzal R, Chrolavicius S, Mehta SR, *et al.* The role of fondaparinux as an adjunct to thrombolytic therapy in acute myocardial infarction: a subgroup analysis of the OASIS-6 trial. *Eur Heart J* 2008;**29**:324–331. https://doi.org/10.1093/eurheartj/ehm616
351. White H. Thrombin-specific anticoagulation with bivalirudin versus heparin in patients receiving fibrinolytic therapy for acute myocardial infarction: the HERO-2 randomised trial. *Lancet* 2001;**358**:1855–1863. https://doi.org/10.1016/s0140-6736(01)06887-8
352. Fernández-Avilés F, Alonso JJ, Peña G, Blanco J, Alonso-Briales J, Lopez-Mesa J, *et al.* Primary angioplasty vs. early routine post-fibrinolysis angioplasty for acute myocardial infarction with ST-segment elevation: the GRACIA-2 non-inferiority, randomized, controlled trial. *Eur Heart J* 2007;**28**:949–960. https://doi.org/10.1093/eurheartj/ehl461
353. Björklund E, Stenestrand U, Lindbäck J, Svensson L, Wallentin L, Lindahl B. Pre-hospital thrombolysis delivered by paramedics is associated with reduced time delay and mortality in ambulance-transported real-life patients with ST-elevation myocardial infarction. *Eur Heart J* 2006;**27**:1146–1152. https://doi.org/10.1093/eurheartj/ehi886
354. Bonnefoy E, Steg PG, Boutitie F, Dubien P-Y, Lapostolle F, Roncalli J, *et al.* Comparison of primary angioplasty and pre-hospital fibrinolysis in acute myocardial infarction (CAPTIM) trial: a 5-year follow-up. *Eur Heart J* 2009;**30**:1598–1606. https://doi.org/10.1093/eurheartj/ehp156
355. Bonnefoy E, Lapostolle F, Leizorovicz A, Steg G, McFadden EP, Dubien PY, *et al.* Primary angioplasty versus prehospital fibrinolysis in acute myocardial infarction: a

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

randomised study. *Lancet* 2002;**360**:825–829. https://doi.org/10.1016/s0140-6736(02)09963-4
356. Assessment of the Safety and Efficacy of a New Thrombolytic (ASSENT-2) Investigators. Single-bolus tenecteplase compared with front-loaded alteplase in acute myocardial infarction: the ASSENT-2 double-blind randomised trial. *Lancet* 1999;**354**:716–722. https://doi.org/10.1016/s0140-6736(99)07403-6
357. GUSTO investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med* 1993;**329**:673–682. https://doi.org/10.1056/nejm199309023291001
358. Wallentin L, Goldstein P, Armstrong PW, Granger CB, Adgey AAJ, Arntz HR, *et al.* Efficacy and safety of tenecteplase in combination with the low-molecular-weight heparin enoxaparin or unfractionated heparin in the prehospital setting: the Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 PLUS randomized trial in acute myocardial infarction. *Circulation* 2003;**108**:135–142. https://doi.org/10.1161/01.Cir.0000081659.72985.A8
359. Ross AM, Molhoek P, Lundergan C, Knudtson M, Draoui Y, Regalado L, *et al.* Randomized comparison of enoxaparin, a low-molecular-weight heparin, with unfractionated heparin adjunctive to recombinant tissue plasminogen activator thrombolysis and aspirin: second trial of Heparin and Aspirin Reperfusion Therapy (HART II). *Circulation* 2001;**104**:648–652. https://doi.org/10.1161/hc3101.093866
360. Antman EM, Louwerenburg HW, Baars HF, Wesdorp JCL, Hamer B, Bassand J-P, *et al.* Enoxaparin as adjunctive antithrombin therapy for ST-elevation myocardial infarction: results of the ENTIRE-Thrombolysis in Myocardial Infarction (TIMI) 23 Trial. *Circulation* 2002;**105**:1642–1649. https://doi.org/10.1161/01.cir.0000013402.34759.46
361. James SK, Roe MT, Cannon CP, Cornel JH, Horrow J, Husted S, *et al.* Ticagrelor versus clopidogrel in patients with acute coronary syndromes intended for non-invasive management: substudy from prospective randomised PLATelet inhibition and patient Outcomes (PLATO) trial. *BMJ* 2011;**342**:d3527. https://doi.org/10.1136/bmj.d3527
362. Wiviott SD, White HD, Ohman EM, Fox KAA, Armstrong PW, Prabhakaran D, *et al.* Prasugrel versus clopidogrel for patients with unstable angina or non-ST-segment elevation myocardial infarction with or without angiography: a secondary, prespecified analysis of the TRILOGY ACS trial. *Lancet* 2013;**382**:605–613. https://doi.org/10.1016/s0140-6736(13)61451-8
363. Savonitto S, Ferri LA, Piatti L, Grosseto D, Piovaccari G, Morici N, *et al.* Comparison of reduced-dose prasugrel and standard-dose clopidogrel in elderly patients with acute coronary syndromes undergoing early percutaneous revascularization. *Circulation* 2018;**137**:2435–2445. https://doi.org/10.1161/circulationaha.117.032180
364. Patterson T, Perkins GD, Hassan Y, Moschonas K, Gray H, Curzen N, *et al.* Temporal trends in identification, management, and clinical outcomes after out-of-hospital cardiac arrest: insights from the myocardial ischaemia national audit project database. *Circ Cardiovasc Interv* 2018;**11**:e005346. https://doi.org/10.1161/circinterventions.117.005346
365. Byrne R, Constant O, Smyth Y, Callagy G, Nash P, Daly K, *et al.* Multiple source surveillance incidence and aetiology of out-of-hospital sudden cardiac death in a rural population in the West of Ireland. *Eur Heart J* 2008;**29**:1418–1423. https://doi.org/10.1093/eurheartj/ehn155
366. Kroupa J, Knot J, Ulman J, Bednar F, Dohnalova A, Motovska Z. Characteristics and survival determinants in patients after out-of-hospital cardiac arrest in the era of 24/7 coronary intervention facilities. *Heart Lung Circ* 2017;**26**:799–807. https://doi.org/10.1016/j.hlc.2016.11.012
367. Perkins GD, Graesner JT, Semeraro F, Olasveengen T, Soar J, Lott C, *et al.* European Resuscitation Council Guidelines 2021: executive summary. *Resuscitation* 2021;**161**:1–60. https://doi.org/10.1016/j.resuscitation.2021.02.003
368. Dumas F, Cariou A, Manzo-Silberman S, Grimaldi D, Vivien B, Rosencher J, *et al.* Immediate percutaneous coronary intervention is associated with better survival after out-of-hospital cardiac arrest: insights from the PROCAT (Parisian Region Out of hospital Cardiac ArresT) registry. *Circ Cardiovasc Interv* 2010;**3**:200–207. https://doi.org/10.1161/circinterventions.109.913665
369. Nolan JP, Sandroni C, Böttiger BW, Cariou A, Cronberg T, Friberg H, *et al.* European Resuscitation Council and European Society of Intensive Care Medicine guidelines 2021: post-resuscitation care. *Intensive Care Med* 2021;**47**:369–421. https://doi.org/10.1007/s00134-021-06368-4
370. Rab T, Kern KB, Tamis-Holland JE, Henry TD, McDaniel M, Dickert NW, *et al.* Cardiac arrest: a treatment algorithm for emergent invasive cardiac procedures in the resuscitated comatose patient. *J Am Coll Cardiol* 2015;**66**:62–73. https://doi.org/10.1016/j.jacc.2015.05.009
371. Gorjup V, Radsel P, Kocjancic ST, Erzen D, Noc M. Acute ST-elevation myocardial infarction after successful cardiopulmonary resuscitation. *Resuscitation* 2007;**72**:379–385. https://doi.org/10.1016/j.resuscitation.2006.07.013
372. Lemkes JS, Janssens GN, van der Hoeven NW, Jewbali LSD, Dubois EA, Meuwissen MM, *et al.* Coronary angiography after cardiac arrest without ST segment elevation: one-year outcomes of the COACT randomized clinical trial. *JAMA Cardiol* 2020;**5**:1358–1365. https://doi.org/10.1001/jamacardio.2020.3670
373. Desch S, Freund A, Akin I, Behnes M, Preusch MR, Zelniker TA, *et al.* Angiography after out-of-hospital cardiac arrest without ST-segment elevation. *N Engl J Med* 2021;**385**:2544–2553. https://doi.org/10.1056/NEJMoa2101909
374. Lemkes JS, Janssens GN, van der Hoeven NW, Jewbali LSD, Dubois EA, Meuwissen M, *et al.* Coronary angiography after cardiac arrest without ST-segment elevation. *N Engl J Med* 2019;**380**:1397–1407. https://doi.org/10.1056/NEJMoa1816897
375. Kern KB, Radsel P, Jentzer JC, Seder DB, Lee KS, Lotun K, *et al.* Randomized pilot clinical trial of early coronary angiography versus no early coronary angiography after cardiac arrest without ST-segment elevation: the PEARL study. *Circulation* 2020;**142**:2002–2012. https://doi.org/10.1161/circulationaha.120.049569
376. Hauw-Berlemont C, Lamhaut L, Diehl JL, Andreotti C, Varenne O, Leroux P, *et al.* Emergency vs delayed coronary angiogram in survivors of out-of-hospital cardiac arrest: results of the randomized, multicentric EMERGE trial. *JAMA Cardiol* 2022;**7**:700–707. https://doi.org/10.1001/jamacardio.2022.1416
377. Viana-Tejedor A, Andrea-Riba R, Scardino C, Ariza-Solé A, Bañeras J, García-García C, *et al.* Coronary angiography in patients without ST-segment elevation following out-of-hospital cardiac arrest. *Rev Esp Cardiol (Engl Ed)* 2022;**76**:94–102. https://doi.org/10.1016/j.rec.2022.05.013
378. The Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;**346**:549–556. https://doi.org/10.1056/NEJMoa012689
379. Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, *et al.* Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;**346**:557–563. https://doi.org/10.1056/NEJMoa003289
380. Belliard G, Catez E, Charron C, Caille V, Aegerter P, Dubourg O, *et al.* Efficacy of therapeutic hypothermia after out-of-hospital cardiac arrest due to ventricular fibrillation. *Resuscitation* 2007;**75**:252–259. https://doi.org/10.1016/j.resuscitation.2007.04.014
381. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, *et al.* Targeted temperature management at 33°C versus 36°C after cardiac arrest. *N Engl J Med* 2013;**369**:2197–2206. https://doi.org/10.1056/NEJMoa1310519
382. Vaahersalo J, Hiltunen P, Tiainen M, Oksanen T, Kaukonen K-M, Kurola J, *et al.* Therapeutic hypothermia after out-of-hospital cardiac arrest in Finnish intensive care units: the FINNRESUSCI study. *Intensive Care Med* 2013;**39**:826–837. https://doi.org/10.1007/s00134-013-2868-1
383. Nolan JP, Sandroni C, Andersen LW, Böttiger BW, Cariou A, Cronberg T, *et al.* ERC-ESICM guidelines on temperature control after cardiac arrest in adults. *Resuscitation* 2022;**172**:229–236. https://doi.org/10.1016/j.resuscitation.2022.01.009
384. Hassager C, Schmidt H, Møller JE, Grand J, Mølstrøm S, Beske RP, *et al.* Duration of device-based fever prevention after cardiac arrest. *N Engl J Med* 2023;**388**:888–897. https://doi.org/10.1056/NEJMoa2212528
385. Wolfrum S, Roedl K, Hanebutte A, Pfeifer R, Kurowski V, Riessen R, *et al.* Temperature control after in-hospital cardiac arrest: a randomized clinical trial. *Circulation* 2022;**146**:1357–1366. https://doi.org/10.1161/circulationaha.122.060106
386. Sandroni C, Geocadin RG. Neurological prognostication after cardiac arrest. *Curr Opin Crit Care* 2015;**21**:209–214. https://doi.org/10.1097/mcc.0000000000000202
387. Garot P, Lefevre T, Eltchaninoff H, Morice M-C, Tamion F, Abry B, *et al.* Six-month outcome of emergency percutaneous coronary intervention in resuscitated patients after cardiac arrest complicating ST-elevation myocardial infarction. *Circulation* 2007;**115**:1354–1362. https://doi.org/10.1161/circulationaha.106.657619
388. Spaulding CM, Joly LM, Rosenberg A, Monchi M, Weber SN, Dhainaut J-FA, *et al.* Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med* 1997;**336**:1629–1633. https://doi.org/10.1056/nejm199706053362302
389. Dankiewicz J, Cronberg T, Lilja G, Jakobsen JC, Levin H, Ullén S, *et al.* Hypothermia versus normothermia after out-of-hospital cardiac arrest. *N Engl J Med* 2021;**384**:2283–2294. https://doi.org/10.1056/NEJMoa2100591
390. Callaway CW, Schmicker R, Kampmeyer M, Powell J, Rea TD, Daya MR, *et al.* Receiving hospital characteristics associated with survival after out-of-hospital cardiac arrest. *Resuscitation* 2010;**81**:524–529. https://doi.org/10.1016/j.resuscitation.2009.12.006
391. Wnent J, Seewald S, Heringlake M, Lemke H, Brauer K, Lefering R, *et al.* Choice of hospital after out-of-hospital cardiac arrest – a decision with far-reaching consequences: a study in a large German city. *Crit Care* 2012;**16**:R164. https://doi.org/10.1186/cc11516
392. Kragholm K, Malta Hansen C, Dupre ME, Xian Y, Strauss B, Tyson C, *et al.* Direct transport to a percutaneous cardiac intervention center and outcomes in patients with out-of-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2017;**10**:e003414. https://doi.org/10.1161/circoutcomes.116.003414
393. Yeo JW, Ng ZHC, Goh AXC, Gao JF, Liu N, Lam SWS, *et al.* Impact of cardiac arrest centers on the survival of patients with nontraumatic out-of-hospital cardiac arrest: a systematic review and meta-analysis. *J Am Heart Assoc* 2022;**11**:e023806. https://doi.org/10.1161/jaha.121.023806
394. Hochman JS, Sleeper LA, Webb JG, Sanborn TA, White HD, Talley JD, *et al.* Early revascularization in acute myocardial infarction complicated by cardiogenic shock. *N Engl J Med* 1999;**341**:625–634. https://doi.org/10.1056/nejm199908263410901
395. White HD, Assmann SF, Sanborn TA, Jacobs AK, Webb JG, Sleeper LA, *et al.* Comparison of percutaneous coronary intervention and coronary artery bypass

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

grafting after acute myocardial infarction complicated by cardiogenic shock: results from the Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock (SHOCK) trial. *Circulation* 2005;**112**:1992–2001. https://doi.org/10.1161/circulationaha.105.540948
396. Hochman JS, Sleeper LA, Webb JG, Dzavik V, Buller CE, Aylward P, *et al.* Early revascularization and long-term survival in cardiogenic shock complicating acute myocardial infarction. *JAMA* 2006;**295**:2511–2515. https://doi.org/10.1001/jama.295.21.2511
397. Liakopoulos OJ, Schlachtenberger G, Wendt D, Choi Y-H, Slottosch I, Welp H, *et al.* Early clinical outcomes of surgical myocardial revascularization for acute coronary syndromes complicated by cardiogenic shock: a report from the north-Rhine-Westphalia surgical myocardial infarction registry. *J Am Heart Assoc* 2019;**8**:e012049. https://doi.org/10.1161/jaha.119.012049
398. Thielmann M, Wendt D, Slottosch I, Welp H, Schiller W, Tsagakis K, *et al.* Coronary artery bypass graft surgery in patients with acute coronary syndromes after primary percutaneous coronary intervention: a current report from the north-Rhine Westphalia surgical myocardial infarction registry. *J Am Heart Assoc* 2021;**10**:e021182. https://doi.org/10.1161/jaha.121.021182
399. Thiele H, Zeymer U, Neumann FJ, Ferenc M, Olbrich H-G, Hausleiter J, *et al.* Intraaortic balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med* 2012;**367**:1287–1296. https://doi.org/10.1056/NEJMoa1208410
400. Schrage B, Ibrahim K, Loehn T, Werner N, Sinning J-M, Pappalardo F, *et al.* Impella support for acute myocardial infarction complicated by cardiogenic shock. *Circulation* 2019;**139**:1249–1258. https://doi.org/10.1161/circulationaha.118.036614
401. Miller PE, Bromfield SG, Ma Q, Crawford G, Whitney J, DeVries A, *et al.* Clinical outcomes and cost associated with an intravascular microaxial left ventricular assist device vs intra-aortic balloon pump in patients presenting with acute myocardial infarction complicated by cardiogenic shock. *JAMA Intern Med* 2022;**182**:926–933. https://doi.org/10.1001/jamainternmed.2022.2735
402. Ostadal P, Rokyta R, Karasek J, Kruger A, Vondrakova D, Janotka M, *et al.* Extracorporeal membrane oxygenation in the therapy of cardiogenic shock: results of the ECMO-CS randomized clinical trial. *Circulation* 2022;**147**:454–464. https://doi.org/10.1161/circulationaha.122.062949
403. Kim Y, Shapero K, Ahn SS, Goldsweig AM, Desai N, Altin SE. Outcomes of mechanical circulatory support for acute myocardial infarction complicated by cardiogenic shock. *Catheter Cardiovasc Interv* 2022;**99**:658–663. https://doi.org/10.1002/ccd.29834
404. Thiele H, Akin I, Sandri M, Fuernau G, de Waha S, Meyer-Saraei R, *et al.* PCI strategies in patients with acute myocardial infarction and cardiogenic shock. *N Engl J Med* 2017;**377**:2419–2432. https://doi.org/10.1056/NEJMoa1710261
405. Thiele H, Zeymer U, Neumann FJ, Ferenc M, Olbrich H-G, Hausleiter J, *et al.* Intra-aortic balloon counterpulsation in acute myocardial infarction complicated by cardiogenic shock (IABP-SHOCK II): final 12 month results of a randomised, open-label trial. *Lancet* 2013;**382**:1638–1645. https://doi.org/10.1016/s0140-6736(13)61783-3
406. Unverzagt S, Buerke M, de Waha A, Haerting J, Pietzner D, Seyfarth M, *et al.* Intra-aortic balloon pump counterpulsation (IABP) for myocardial infarction complicated by cardiogenic shock. *Cochrane Database Syst Rev* 2015;**2015**:Cd007398. https://doi.org/10.1002/14651858.CD007398.pub3
407. Thiele H, Zeymer U, Thelemann N, Neumann F-J, Hausleiter J, Abdel-Wahab M, *et al.* Intraaortic balloon pump in cardiogenic shock complicating acute myocardial infarction: long-term 6-year outcome of the randomized IABP-SHOCK II trial. *Circulation* 2018;**139**:395–403. https://doi.org/10.1161/circulationaha.118.038201
408. Bonnefoy-Cudraz E, Bueno H, Casella G, De Maria E, Fitzsimons D, Halvorsen S, *et al.* Editor's Choice – acute cardiovascular care association position paper on intensive cardiovascular care units: an update on their definition, structure, organisation and function. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:80–95. https://doi.org/10.1177/2048872617724269
409. Winkler C, Funk M, Schindler DM, Hemsey JZ, Lampert R, Drew BJ. Arrhythmias in patients with acute coronary syndrome in the first 24 hours of hospitalization. *Heart Lung* 2013;**42**:422–427. https://doi.org/10.1016/j.hrtlng.2013.07.010
410. Wasfy JH, Kennedy KF, Masoudi FA, Ferris TG, Arnold SV, Kini V, *et al.* Predicting length of stay and the need for postacute care after acute myocardial infarction to improve healthcare efficiency. *Circ Cardiovasc Qual Outcomes* 2018;**11**:e004635. https://doi.org/10.1161/circoutcomes.118.004635
411. Melberg T, Jørgensen M, Ørn S, Solli T, Edland U, Dickstein K. Safety and health status following early discharge in patients with acute myocardial infarction treated with primary PCI: a randomized trial. *Eur J Prev Cardiol* 2015;**22**:1427–1434. https://doi.org/10.1177/2047487314559276
412. Berger AK, Duval S, Jacobs DR Jr, Barber C, Vazquez G, Lee S, *et al.* Relation of length of hospital stay in acute myocardial infarction to postdischarge mortality. *Am J Cardiol* 2008;**101**:428–434. https://doi.org/10.1016/j.amjcard.2007.09.090
413. Grines CL, Marsalese DL, Brodie B, Griffin J, Donohue B, Costantini CR, *et al.* Safety and cost-effectiveness of early discharge after primary angioplasty in low risk patients with acute myocardial infarction. *J Am Coll Cardiol* 1998;**31**:967–972. https://doi.org/10.1016/s0735-1097(98)00031-x
414. De Luca G, Suryapranata H, van't Hof AW, de Boer M-J, Hoorntje JCA, Dambrink J-HE, *et al.* Prognostic assessment of patients with acute myocardial infarction treated with primary angioplasty: implications for early discharge. *Circulation* 2004;**109**:2737–2743. https://doi.org/10.1161/01.Cir.0000131765.73959.87
415. Novobilsky K, Stipal R, Cerny P, Horak I, Kaucak V, Mrozek J, *et al.* Safety of early discharge in low risk patients after acute ST-segment elevation myocardial infarction, treated with primary percutaneous coronary intervention. Open label, randomized trial. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2019;**163**:61–66. https://doi.org/10.5507/bp.2018.041
416. Albanese M, Alpaslan K, Ouarrak T, Merguet P, Schneider S, Schöls W. In-hospital major arrhythmias, arrhythmic death and resuscitation after successful primary percutaneous intervention for acute transmural infarction: a retrospective single-centre cohort study. *BMC Cardiovasc Disord* 2018;**18**:116. https://doi.org/10.1186/s12872-018-0851-z
417. Yndigegn T, Gilje P, Dankiewicz J, Mokhtari A, Isma N, Holmqvist J, *et al.* Safety of early hospital discharge following admission with ST-elevation myocardial infarction treated with percutaneous coronary intervention: a nationwide cohort study. *EuroIntervention* 2022;**17**:1091–1099. https://doi.org/10.4244/eij-d-21-00501
418. Seto AH, Shroff A, Abu-Fadel M, Blankenship JC, Boudoulas KD, Cigarroa JE, *et al.* Length of stay following percutaneous coronary intervention: an expert consensus document update from the society for cardiovascular angiography and interventions. *Catheter Cardiovasc Interv* 2018;**92**:717–731. https://doi.org/10.1002/ccd.27637
419. Estévez-Loureiro R, Calviño-Santos R, Vázquez JM, Barge-Caballero E, Salgado-Fernández J, Piñeiro M, *et al.* Safety and feasibility of returning patients early to their originating centers after transfer for primary percutaneous coronary intervention. *Rev Esp Cardiol* 2009;**62**:1356–1364. https://doi.org/10.1016/s1885-5857(09)73529-7
420. Morrow DA, Antman EM, Charlesworth A, Cairns R, Murphy SA, de Lemos JA, *et al.* TIMI risk score for ST-elevation myocardial infarction: a convenient, bedside, clinical score for risk assessment at presentation: an intravenous nPA for treatment of infarcting myocardium early II trial substudy. *Circulation* 2000;**102**:2031–2037. https://doi.org/10.1161/01.cir.102.17.2031
421. Fox KA, Dabbous OH, Goldberg RJ, Pieper KS, Eagle KA, Van de Werf F, *et al.* Prediction of risk of death and myocardial infarction in the six months after presentation with acute coronary syndrome: prospective multinational observational study (GRACE). *BMJ* 2006;**333**:1091. https://doi.org/10.1136/bmj.38985.646481.55
422. Aragam KG, Tamhane UU, Kline-Rogers E, Li J, Fox KAA, Goodman SG, *et al.* Does simplicity compromise accuracy in ACS risk prediction? A retrospective analysis of the TIMI and GRACE risk scores. *PLoS One* 2009;**4**:e7947. https://doi.org/10.1371/journal.pone.0007947
423. D'Ascenzo F, Biondi-Zoccai G, Moretti C, Bollati M, Omedè P, Sciuto F, *et al.* TIMI, GRACE and alternative risk scores in Acute Coronary Syndromes: a meta-analysis of 40 derivation studies on 216,552 patients and of 42 validation studies on 31,625 patients. *Contemp Clin Trials* 2012;**33**:507–514. https://doi.org/10.1016/j.cct.2012.01.001
424. Gale CP, Manda SO, Weston CF, Birkhead JS, Batin PD, Hall AS. Evaluation of risk scores for risk stratification of acute coronary syndromes in the Myocardial Infarction National Audit Project (MINAP) database. *Heart* 2009;**95**:221–227. https://doi.org/10.1136/hrt.2008.144022
425. D'Ascenzo F, De Filippo O, Gallone G, Mittone G, Deriu MA, Iannaccone M, *et al.* Machine learning-based prediction of adverse events following an acute coronary syndrome (PRAISE): a modelling study of pooled datasets. *Lancet* 2021;**397**:199–207. https://doi.org/10.1016/s0140-6736(20)32519-8
426. Ng VG, Lansky AJ, Meller S, Witzenbichler B, Guagliumi G, Peruga JZ, *et al.* The prognostic importance of left ventricular function in patients with ST-segment elevation myocardial infarction: the HORIZONS-AMI trial. *Eur Heart J Acute Cardiovasc Care* 2014;**3**:67–77. https://doi.org/10.1177/2048872613507149
427. de Waha S, Eitel I, Desch S, Fuernau G, Lurz P, Stiermaier T, *et al.* Prognosis after ST-elevation myocardial infarction: a study on cardiac magnetic resonance imaging versus clinical routine. *Trials* 2014;**15**:249. https://doi.org/10.1186/1745-6215-15-249
428. Larose E, Côté J, Rodés-Cabau J, Noël B, Barbeau G, Bordeleau E, *et al.* Contrast-enhanced cardiovascular magnetic resonance in the hyperacute phase of ST-elevation myocardial infarction. *Int J Cardiovasc Imaging* 2009;**25**:519–527. https://doi.org/10.1007/s10554-009-9451-4
429. Stiermaier T, Jobs A, de Waha S, Fuernau G, Pöss J, Desch S, *et al.* Optimized prognosis assessment in ST-segment-elevation myocardial infarction using a cardiac magnetic resonance imaging risk score. *Circ Cardiovasc Imaging* 2017;**10**:e006774. https://doi.org/10.1161/circimaging.117.006774
430. de Waha S, Desch S, Eitel I, Fuernau G, Zachrau J, Leuschner A, *et al.* Impact of early vs. late microvascular obstruction assessed by magnetic resonance imaging on long-term outcome after ST-elevation myocardial infarction: a comparison with traditional prognostic markers. *Eur Heart J* 2010;**31**:2660–2668. https://doi.org/10.1093/eurheartj/ehq247
431. van Kranenburg M, Magro M, Thiele H, de Waha S, Eitel I, Cochet A, *et al.* Prognostic value of microvascular obstruction and infarct size, as measured by CMR in STEMI patients. *JACC Cardiovasc Imaging* 2014;**7**:930–939. https://doi.org/10.1016/j.jcmg.2014.05.010
432. van Loon RB, Veen G, Baur LHB, Kamp O, Bronzwaer JGF, Twisk JWR, *et al.* Improved clinical outcome after invasive management of patients with recent myocardial

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

infarction and proven myocardial viability: primary results of a randomized controlled trial (VIAMI-trial). *Trials* 2012;**13**:1. https://doi.org/10.1186/1745-6215-13-1
433. van Loon RB, Veen G, Baur LH, Twisk JW, van Rossum AC. Long-term follow-up of the viability guided angioplasty after acute myocardial infarction (VIAMI) trial. *Int J Cardiol* 2015;**186**:111–116. https://doi.org/10.1016/j.ijcard.2015.03.152
434. Priori SG, Blomström-Lundqvist C, Mazzanti A, Blom N, Borggrefe M, Camm J, *et al.* 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: the Task Force for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J* 2015;**36**:2793–2867. https://doi.org/10.1093/eurheartj/ehv316
435. Ibanez B, Aletras AH, Arai AE, Arheden H, Bax J, Berry C, *et al.* Cardiac MRI endpoints in myocardial infarction experimental and clinical trials: JACC Scientific Expert Panel. *J Am Coll Cardiol* 2019;**74**:238–256. https://doi.org/10.1016/j.jacc.2019.05.024
436. Stone GW, Selker HP, Thiele H, Patel MR, Udelson JE, Ohman EM, *et al.* Relationship between infarct size and outcomes following primary PCI: patient-level analysis from 10 randomized trials. *J Am Coll Cardiol* 2016;**67**:1674–1683. https://doi.org/10.1016/j.jacc.2016.01.069
437. Ibáñez B, Heusch G, Ovize M, Van de Werf F. Evolving therapies for myocardial ischemia/reperfusion injury. *J Am Coll Cardiol* 2015;**65**:1454–1471. https://doi.org/10.1016/j.jacc.2015.02.032
438. Eitel I, de Waha S, Wöhrle J, Fuernau G, Lurz P, Pauschinger M, *et al.* Comprehensive prognosis assessment by CMR imaging after ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2014;**64**:1217–1226. https://doi.org/10.1016/j.jacc.2014.06.1194
439. Haaf P, Reichlin T, Twerenbold R, Hoeller R, Rubini Gimenez M, Zellweger C, *et al.* Risk stratification in patients with acute chest pain using three high-sensitivity cardiac troponin assays. *Eur Heart J* 2014;**35**:365–375. https://doi.org/10.1093/eurheartj/eht218
440. Tveit SH, Myhre PL, Hoff NJS, Le TM, Seljeflot I, Røysland R, *et al.* Superiority of high sensitivity cardiac troponin T vs. I for long-term prognostic value in patients with chest pain; data from the Akershus Cardiac Examination (ACE) 3 study. *Clin Biochem* 2020;**78**:10–17. https://doi.org/10.1016/j.clinbiochem.2019.12.016
441. Welsh P, Preiss D, Hayward C, Shah ASV, McAllister D, Briggs A, *et al.* Cardiac troponin T and troponin I in the general population. *Circulation* 2019;**139**:2754–2764. https://doi.org/10.1161/circulationaha.118.038529
442. Thygesen K, Mair J, Giannitsis E, Mueller C, Lindahl B, Blankenberg S, *et al.* How to use high-sensitivity cardiac troponins in acute cardiac care. *Eur Heart J* 2012;**33**:2252–2257. https://doi.org/10.1093/eurheartj/ehs154
443. Anavekar NS, McMurray JJ, Velazquez EJ, Solomon SD, Kober L, Rouleau J-L, *et al.* Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med* 2004;**351**:1285–1295. https://doi.org/10.1056/NEJMoa041365
444. Thygesen K, Mair J, Mueller C, Huber K, Weber M, Plebani M, *et al.* Recommendations for the use of natriuretic peptides in acute cardiac care: a position statement from the Study Group on Biomarkers in Cardiology of the ESC Working Group on Acute Cardiac Care. *Eur Heart J* 2012;**33**:2001–2006. https://doi.org/10.1093/eurheartj/ehq509
445. Ducrocq G, Schulte PJ, Budaj A, Cornel JH, Held C, Himmelmann A, *et al.* Balancing the risk of spontaneous ischemic and major bleeding events in acute coronary syndromes. *Am Heart J* 2017;**186**:91–99. https://doi.org/10.1016/j.ahj.2017.01.010
446. Valgimigli M, Costa F, Lokhnygina Y, Clare RM, Wallentin L, Moliterno DJ, *et al.* Trade-off of myocardial infarction vs. bleeding types on mortality after acute coronary syndrome: lessons from the Thrombin Receptor Antagonist for Clinical Event Reduction in Acute Coronary Syndrome (TRACER) randomized trial. *Eur Heart J* 2017;**38**:804–810. https://doi.org/10.1093/eurheartj/ehw525
447. Newby LK, Hasselblad V, Armstrong PW, Van de Werf F, Mark DB, White HD, *et al.* Time-based risk assessment after myocardial infarction. Implications for timing of discharge and applications to medical decision-making. *Eur Heart J* 2003;**24**:182–189. https://doi.org/10.1016/s0195-668x(02)00301-9
448. Kwok CS, Khan MA, Rao SV, Kinnaird T, Sperrin M, Buchan I, *et al.* Access and non-access site bleeding after percutaneous coronary intervention and risk of subsequent mortality and major adverse cardiovascular events: systematic review and meta-analysis. *Circ Cardiovasc Interv* 2015;**8**:e001645. https://doi.org/10.1161/circinterventions.114.001645
449. Ndrepepa G, Neumann FJ, Richardt G, Schulz S, Tölg R, Stoyanov KM, *et al.* Prognostic value of access and non-access sites bleeding after percutaneous coronary intervention. *Circ Cardiovasc Interv* 2013;**6**:354–361. https://doi.org/10.1161/circinterventions.113.000433
450. Rao SV, Cohen MG, Kandzari DE, Bertrand OF, Gilchrist IC. The transradial approach to percutaneous coronary intervention: historical perspective, current concepts, and future directions. *J Am Coll Cardiol* 2010;**55**:2187–2195. https://doi.org/10.1016/j.jacc.2010.01.039
451. Jolly SS, Yusuf S, Cairns J, Niemelä K, Xavier D, Widimsky P, *et al.* Radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): a randomised, parallel group, multicentre trial. *Lancet* 2011;**377**:1409–1420. https://doi.org/10.1016/s0140-6736(11)60404-2
452. Valgimigli M, Gagnor A, Calabró P, Frigoli E, Leonardi S, Zaro T, *et al.* Radial versus femoral access in patients with acute coronary syndromes undergoing invasive management: a randomised multicentre trial. *Lancet* 2015;**385**:2465–2476. https://doi.org/10.1016/s0140-6736(15)60292-6
453. Vranckx P, Frigoli E, Rothenbühler M, Tomassini F, Garducci S, Andò G, *et al.* Radial versus femoral access in patients with acute coronary syndromes with or without ST-segment elevation. *Eur Heart J* 2017;**38**:1069–1080. https://doi.org/10.1093/eurheartj/ehx048
454. Lee P, Liew D, Brennan A, Stub D, Lefkovits J, Reid CM, *et al.* Cost-effectiveness of radial access percutaneous coronary intervention in acute coronary syndrome. *Am J Cardiol* 2021;**156**:44–51. https://doi.org/10.1016/j.amjcard.2021.06.034
455. Kerensky RA, Wade M, Deedwania P, Boden WE, Pepine CJ. Revisiting the culprit lesion in non-Q-wave myocardial infarction. Results from the VANQWISH trial angiographic core laboratory. *J Am Coll Cardiol* 2002;**39**:1456–1463. https://doi.org/10.1016/s0735-1097(02)01770-9
456. Johnson TW, Räber L, di Mario C, Bourantas C, Jia H, Mattesini A, *et al.* Clinical use of intracoronary imaging. Part 2: acute coronary syndromes, ambiguous coronary angiography findings, and guiding interventional decision-making: an expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J* 2019;**40**:2566–2584. https://doi.org/10.1093/eurheartj/ehz332
457. di Mario C, Koskinas KC, Räber L. Clinical benefit of IVUS guidance for coronary stenting: the ULTIMATE step toward definitive evidence? *J Am Coll Cardiol* 2018;**72**:3138–3141. https://doi.org/10.1016/j.jacc.2018.10.029
458. Gao XF, Wang ZM, Wang F, Gu Y, Ge Z, Kong X-Q, *et al.* Intravascular ultrasound guidance reduces cardiac death and coronary revascularization in patients undergoing drug-eluting stent implantation: results from a meta-analysis of 9 randomized trials and 4724 patients. *Int J Cardiovasc Imaging* 2019;**35**:239–247. https://doi.org/10.1007/s10554-019-01555-3
459. Darmoch F, Alraies MC, Al-Khadra Y, Moussa Pacha H, Pinto DS, Osborn EA. Intravascular ultrasound imaging-guided versus coronary angiography-guided percutaneous coronary intervention: a systematic review and meta-analysis. *J Am Heart Assoc* 2020;**9**:e013678. https://doi.org/10.1161/jaha.119.013678
460. Jia H, Dai J, He L, Xu Y, Shi Y, Zhao L, *et al.* EROSION III: a multicenter RCT of OCT-guided reperfusion in STEMI with early infarct artery patency. *JACC Cardiovasc Interv* 2022;**15**:846–856. https://doi.org/10.1016/j.jcin.2022.01.298
461. Cuculi F, De Maria GL, Meier P, Dall'Armellina E, de Caterina AR, Channon KM, *et al.* Impact of microvascular obstruction on the assessment of coronary flow reserve, index of microcirculatory resistance, and fractional flow reserve after ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2014;**64**:1894–1904. https://doi.org/10.1016/j.jacc.2014.07.987
462. De Bruyne B, Pijls NH, Bartunek J, Kulecki K, Bech J-W, De Winter H, *et al.* Fractional flow reserve in patients with prior myocardial infarction. *Circulation* 2001;**104**:157–162. https://doi.org/10.1161/01.cir.104.2.157
463. Bønaa KH, Mannsverk J, Wiseth R, Aaberge L, Myreng Y, Nygård O, *et al.* Drug-eluting or bare-metal stents for coronary artery disease. *N Engl J Med* 2016;**375**:1242–1252. https://doi.org/10.1056/NEJMoa1607991
464. Räber L, Kelbæk H, Ostojic M, Baumbach A, Heg D, Tüller D, *et al.* Effect of biolimus-eluting stents with biodegradable polymer vs bare-metal stents on cardiovascular events among patients with acute myocardial infarction: the COMFORTABLE AMI randomized trial. *JAMA* 2012;**308**:777–787. https://doi.org/10.1001/jama.2012.10065
465. Sabate M, Cequier A, Iñiguez A, Serra A, Hernandez-Antolin R, Mainar V, *et al.* Everolimus-eluting stent versus bare-metal stent in ST-segment elevation myocardial infarction (EXAMINATION): 1 year results of a randomised controlled trial. *Lancet* 2012;**380**:1482–1490. https://doi.org/10.1016/s0140-6736(12)61223-9
466. Sabaté M, Brugaletta S, Cequier A, Iñiguez A, Serra A, Jiménez-Quevedo P, *et al.* Clinical outcomes in patients with ST-segment elevation myocardial infarction treated with everolimus-eluting stents versus bare-metal stents (EXAMINATION): 5-year results of a randomised trial. *Lancet* 2016;**387**:357–366. https://doi.org/10.1016/s0140-6736(15)00548-6
467. Brugaletta S, Gomez-Lara J, Ortega-Paz L, Jimenez-Diaz V, Jimenez M, Jiménez-Quevedo P, *et al.* 10-Year follow-up of patients with everolimus-eluting versus bare-metal stents after ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2021;**77**:1165–1178. https://doi.org/10.1016/j.jacc.2020.12.059
468. Räber L, Yamaji K, Kelbæk H, Engstrøm T, Baumbach A, Roffi M, *et al.* Five-year clinical outcomes and intracoronary imaging findings of the COMFORTABLE AMI trial: randomized comparison of biodegradable polymer-based biolimus-eluting stents with bare-metal stents in patients with acute ST-segment elevation myocardial infarction. *Eur Heart J* 2019;**40**:1909–1919. https://doi.org/10.1093/eurheartj/ehz074
469. Vos NS, Fagel ND, Amoroso G, Herrman J-PR, Patterson MS, Piers LH, *et al.* Paclitaxel-coated balloon angioplasty versus drug-eluting stent in acute myocardial infarction: the REVELATION randomized trial. *JACC Cardiovasc Interv* 2019;**12**:1691–1699. https://doi.org/10.1016/j.jcin.2019.04.016
470. Scheller B, Ohlow MA, Ewen S, Kische S, Rudolph TK, Clever YP, *et al.* Bare metal or drug-eluting stent versus drug-coated balloon in non-ST-elevation myocardial

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

infarction: the randomised PEPCAD NSTEMI trial. *EuroIntervention* 2020;**15**: 1527–1533. https://doi.org/10.4244/eij-d-19-00723
471. Belkacemi A, Agostoni P, Nathoe HM, Voskuil M, Shao CL, Van Belle E, *et al.* First results of the DEB-AMI (drug eluting balloon in acute ST-segment elevation myocardial infarction) trial: a multicenter randomized comparison of drug-eluting balloon plus bare-metal stent versus bare-metal stent versus drug-eluting stent in primary percutaneous coronary intervention with 6-month angiographic, intravascular, functional, and clinical outcomes. *J Am Coll Cardiol* 2012;**59**:2327–2337. https://doi.org/10.1016/j.jacc.2012.02.027
472. Fröbert O, Lagerqvist B, Olivecrona GK, Omerovic E, Gudnason T, Maeng M, *et al.* Thrombus aspiration during ST-segment elevation myocardial infarction. *N Engl J Med* 2013;**369**:1587–1597. https://doi.org/10.1056/NEJMoa1308789
473. Lagerqvist B, Fröbert O, Olivecrona GK, Gudnason T, Maeng M, Alström P, *et al.* Outcomes 1 year after thrombus aspiration for myocardial infarction. *N Engl J Med* 2014;**371**:1111–1120. https://doi.org/10.1056/NEJMoa1405707
474. Jolly SS, Cairns JA, Yusuf S, Meeks B, Pogue J, Rokoss MJ, *et al.* Randomized trial of primary PCI with or without routine manual thrombectomy. *N Engl J Med* 2015;**372**: 1389–1398. https://doi.org/10.1056/NEJMoa1415098
475. Jolly SS, James S, Džavík V, Cairns JA, Mahmoud KD, Zijlstra F, *et al.* Thrombus aspiration in ST-segment-elevation myocardial infarction: an individual patient meta-analysis: thrombectomy trialists collaboration. *Circulation* 2017;**135**:143–152. https://doi.org/10.1161/circulationaha.116.025371
476. Jolly SS, Cairns JA, Lavi S, Cantor WJ, Bernat I, Cheema AN, *et al.* Thrombus aspiration in patients with high thrombus burden in the TOTAL trial. *J Am Coll Cardiol* 2018;**72**: 1589–1596. https://doi.org/10.1016/j.jacc.2018.07.047
477. Thiele H, de Waha S, Zeymer U, Desch S, Scheller B, Lauer B, *et al.* Effect of aspiration thrombectomy on microvascular obstruction in NSTEMI patients: the TATORT-NSTEMI trial. *J Am Coll Cardiol* 2014;**64**:1117–1124. https://doi.org/10.1016/j.jacc.2014.05.064
478. de Waha S, Patel MR, Granger CB, Ohman EM, Maehara A, Eitel I, *et al.* Relationship between microvascular obstruction and adverse events following primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: an individual patient data pooled analysis from seven randomized trials. *Eur Heart J* 2017;**38**: 3502–3510. https://doi.org/10.1093/eurheartj/ehx414
479. Corrigendum to: 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019;**40**:3096. https://doi.org/10.1093/eurheartj/ehz507
480. Sabatine MS, Bergmark BA, Murphy SA, O'Gara PT, Smith PK, Serruys PW, *et al.* Percutaneous coronary intervention with drug-eluting stents versus coronary artery bypass grafting in left main coronary artery disease: an individual patient data meta-analysis. *Lancet* 2021;**398**:2247–2257. https://doi.org/10.1016/s0140-6736(21)02334-5
481. Head SJ, Milojevic M, Daemen J, Ahn J-M, Boersma E, Christiansen EH, *et al.* Mortality after coronary artery bypass grafting versus percutaneous coronary intervention with stenting for coronary artery disease: a pooled analysis of individual patient data. *Lancet* 2018;**391**:939–948. https://doi.org/10.1016/s0140-6736(18)30423-9
482. Adlam D, Alfonso F, Maas A, Vrints C. European Society of Cardiology, Acute Cardiovascular Care Association, SCAD study group: a position paper on spontaneous coronary artery dissection. *Eur Heart J* 2018;**39**:3353–3368. https://doi.org/10.1093/eurheartj/ehy080
483. Alfonso F, de la Torre Hernández JM, Ibáñez B, Sabaté M, Pan M, Gulati R, *et al.* Rationale and design of the BA-SCAD (Beta-blockers and Antiplatelet agents in patients with Spontaneous Coronary Artery Dissection) randomized clinical trial. *Rev Esp Cardiol (Engl Ed)* 2022;**75**:515–522. https://doi.org/10.1016/j.rec.2021.08.003
484. Jackson R, Al-Hussaini A, Joseph S, van Soest G, Wood A, Macaya F, *et al.* Spontaneous coronary artery dissection: pathophysiological insights from optical coherence tomography. *JACC Cardiovasc Imaging* 2019;**12**:2475–2488. https://doi.org/10.1016/j.jcmg.2019.01.015
485. Hayes SN, Kim ESH, Saw J, Adlam D, Arslanian-Engoren C, Economy KE, *et al.* Spontaneous coronary artery dissection: current state of the science: a scientific statement from the American Heart Association. *Circulation* 2018;**137**:e523–e557. https://doi.org/10.1161/cir.0000000000000564
486. Tweet MS, Eleid MF, Best PJM, Lennon RJ, Lerman A, Rihal CS, *et al.* Spontaneous coronary artery dissection: revascularization versus conservative therapy. *Circ Cardiovasc Interv* 2014;**7**:777–786. https://doi.org/10.1161/circinterventions.114.001659
487. Saw J, Aymong E, Sedlak T, Buller CE, Starovoytov A, Ricci D, *et al.* Spontaneous coronary artery dissection: association with predisposing arteriopathies and precipitating stressors and cardiovascular outcomes. *Circ Cardiovasc Interv* 2014;**7**:645–655. https://doi.org/10.1161/circinterventions.114.001760
488. Lettieri C, Zavalloni D, Rossini R, Morici N, Ettori F, Leonzi O, *et al.* Management and long-term prognosis of spontaneous coronary artery dissection. *Am J Cardiol* 2015;**116**:66–73. https://doi.org/10.1016/j.amjcard.2015.03.039
489. Hayes SN, Tweet MS, Adlam D, Kim ESH, Gulati R, Price JE, *et al.* Spontaneous coronary artery dissection: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2020;**76**: 961–984. https://doi.org/10.1016/j.jacc.2020.05.084
490. Nordmann AJ, Hengstler P, Harr T, Young J, Bucher HC. Clinical outcomes of primary stenting versus balloon angioplasty in patients with myocardial infarction: a meta-analysis of randomized controlled trials. *Am J Med* 2004;**116**:253–262. https://doi.org/10.1016/j.amjmed.2003.08.035
491. Stone GW, Grines CL, Cox DA, Garcia E, Tcheng JE, Griffin JJ, *et al.* Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 2002;**346**:957–966. https://doi.org/10.1056/NEJMoa013404
492. Belle L, Motreff P, Mangin L, Rangé G, Marcaggi X, Marie A, *et al.* Comparison of immediate with delayed stenting using the minimalist immediate mechanical intervention approach in acute ST-segment-elevation myocardial infarction: the MIMI study. *Circ Cardiovasc Interv* 2016;**9**:e003388. https://doi.org/10.1161/circinterventions.115.003388
493. Kelbæk H, Høfsten DE, Køber L, Helqvist S, Kløvgaard L, Holmvang L, *et al.* Deferred versus conventional stent implantation in patients with ST-segment elevation myocardial infarction (DANAMI 3-DEFER): an open-label, randomised controlled trial. *Lancet* 2016;**387**:2199–2206. https://doi.org/10.1016/s0140-6736(16)30072-1
494. Carrick D, Oldroyd KG, McEntegart M, Haig C, Petrie MC, Eteiba H, *et al.* A randomized trial of deferred stenting versus immediate stenting to prevent no- or slow-reflow in acute ST-segment elevation myocardial infarction (DEFER-STEMI). *J Am Coll Cardiol* 2014;**63**:2088–2098. https://doi.org/10.1016/j.jacc.2014.02.530
495. Hong SJ, Kim BK, Shin DH, Nam CM, Kim JS, Ko Y-G, *et al.* Effect of intravascular ultrasound-guided vs angiography-guided everolimus-eluting stent implantation: the IVUS-XPL randomized clinical trial. *JAMA* 2015;**314**:2155–2163. https://doi.org/10.1001/jama.2015.15454
496. Zhang J, Gao X, Kan J, Ge Z, Han L, Lu S, *et al.* Intravascular ultrasound versus angiography-guided drug-eluting stent implantation: the ULTIMATE trial. *J Am Coll Cardiol* 2018;**72**:3126–3137. https://doi.org/10.1016/j.jacc.2018.09.013
497. Gao XF, Ge Z, Kong XQ, Kan J, Han L, Lu S, *et al.* 3-Year outcomes of the ULTIMATE trial comparing intravascular ultrasound versus angiography-guided drug-eluting stent implantation. *JACC Cardiovasc Interv* 2021;**14**:247–257. https://doi.org/10.1016/j.jcin.2020.10.001
498. Meneveau N, Souteyrand G, Motreff P, Caussin C, Amabile N, Ohlmann P, *et al.* Optical coherence tomography to optimize results of percutaneous coronary intervention in patients with non-ST-elevation acute coronary syndrome: results of the multicenter, randomized DOCTORS study (Does Optical Coherence Tomography Optimize Results of Stenting). *Circulation* 2016;**134**:906–917. https://doi.org/10.1161/circulationaha.116.024393
499. Kala P, Cervinka P, Jakl M, Kanovsky J, Kupec A, Spacek R, *et al.* OCT guidance during stent implantation in primary PCI: a randomized multicenter study with nine months of optical coherence tomography follow-up. *Int J Cardiol* 2018;**250**:98–103. https://doi.org/10.1016/j.ijcard.2017.10.059
500. Secemsky EA, Butala N, Raja A, Khera R, Wang Y, Curtis JP, *et al.* Temporal changes and institutional variation in use of percutaneous coronary intervention for ST-elevation myocardial infarction with multivessel coronary artery disease in the United States: an NCDR research to practice project. *JAMA Cardiol* 2021;**6**: 574–580. https://doi.org/10.1001/jamacardio.2020.5354
501. Holmes DR Jr, Berger PB, Hochman JS, Granger CB, Thompson TD, Califf RM, *et al.* Cardiogenic shock in patients with acute ischemic syndromes with and without ST-segment elevation. *Circulation* 1999;**100**:2067–2073. https://doi.org/10.1161/01.cir.100.20.2067
502. Thiele H, Ohman EM, de Waha-Thiele S, Zeymer U, Desch S. Management of cardiogenic shock complicating myocardial infarction: an update 2019. *Eur Heart J* 2019;**40**: 2671–2683. https://doi.org/10.1093/eurheartj/ehz363
503. Papolos AI, Kenigsberg BB, Berg DD, Alviar CL, Bohula E, Burke JA, *et al.* Management and outcomes of cardiogenic shock in cardiac ICUs with versus without shock teams. *J Am Coll Cardiol* 2021;**78**:1309–1317. https://doi.org/10.1016/j.jacc.2021.07.044
504. Rab T, Ratanapo S, Kern KB, Basir MB, McDaniel M, Meraj P, *et al.* Cardiac shock care centers: JACC review topic of the week. *J Am Coll Cardiol* 2018;**72**:1972–1980. https://doi.org/10.1016/j.jacc.2018.07.074
505. Thiele H, Akin I, Sandri M, de Waha-Thiele S, Meyer-Saraei R, Fuernau G, *et al.* One-year outcomes after PCI strategies in cardiogenic shock. *N Engl J Med* 2018;**379**:1699–1710. https://doi.org/10.1056/NEJMoa1808788
506. Sorajja P, Gersh BJ, Cox DA, McLaughlin MG, Zimetbaum P, Costantini C, *et al.* Impact of multivessel disease on reperfusion success and clinical outcomes in patients undergoing primary percutaneous coronary intervention for acute myocardial infarction. *Eur Heart J* 2007;**28**:1709–1716. https://doi.org/10.1093/eurheartj/ehm184
507. Dziewierz A, Siudak Z, Rakowski T, Zasada W, Dubiel JS, Dudek D. Impact of multivessel coronary artery disease and noninfarct-related artery revascularization on outcome of patients with ST-elevation myocardial infarction transferred for primary percutaneous coronary intervention (from the EUROTRANSFER Registry). *Am J Cardiol* 2010;**106**:342–347. https://doi.org/10.1016/j.amjcard.2010.03.029
508. Wald DS, Morris JK, Wald NJ, Chase AJ, Edwards RJ, Hughes LO, *et al.* Randomized trial of preventive angioplasty in myocardial infarction. *N Engl J Med* 2013;**369**: 1115–1123. https://doi.org/10.1056/NEJMoa1305520
509. Engstrøm T, Kelbæk H, Helqvist S, Høfsten DE, Kløvgaard L, Holmvang L, *et al.* Complete revascularisation versus treatment of the culprit lesion only in patients with ST-segment elevation myocardial infarction and multivessel disease

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

(DANAMI-3—PRIMULTI): an open-label, randomised controlled trial. *Lancet* 2015; **386**:665–671. https://doi.org/10.1016/s0140-6736(15)60648-1
510. Smits PC, Abdel-Wahab M, Neumann FJ, Boxma-de Klerk BM, Lunde K, Schotborgh CE, *et al.* Fractional flow reserve-guided multivessel angioplasty in myocardial infarction. *N Engl J Med* 2017;**376**:1234–1244. https://doi.org/10.1056/NEJMoa1701067
511. Mehta SR, Wood DA, Storey RF, Mehran R, Bainey KR, Nguyen H, *et al.* Complete revascularization with multivessel PCI for myocardial infarction. *N Engl J Med* 2019; **381**:1411–1421. https://doi.org/10.1056/NEJMoa1907775
512. Bainey KR, Engstrøm T, Smits PC, Gershlick AH, James SK, Storey RF, *et al.* Complete vs culprit-lesion-only revascularization for ST-segment elevation myocardial infarction: a systematic review and meta-analysis. *JAMA Cardiol* 2020;**5**:881–888. https://doi.org/10.1001/jamacardio.2020.1251
513. Sardella G, Lucisano L, Garbo R, Pennacchi M, Cavallo E, Stio RE, *et al.* Single-staged compared with multi-staged PCI in multivessel NSTEMI patients: the SMILE trial. *J Am Coll Cardiol* 2016;**67**:264–272. https://doi.org/10.1016/j.jacc.2015.10.082
514. Siebert VR, Borgaonkar S, Jia X, Nguyen HL, Birnbaum Y, Lakkis NM, *et al.* Meta-analysis comparing multivessel versus culprit coronary arterial revascularization for patients with non-ST-segment elevation acute coronary syndromes. *Am J Cardiol* 2019;**124**:1501–1511. https://doi.org/10.1016/j.amjcard.2019.07.071
515. Rathod KS, Koganti S, Jain AK, Astroulakis Z, Lim P, Rakhit R, *et al.* Complete versus culprit-only lesion intervention in patients with acute coronary syndromes. *J Am Coll Cardiol* 2018;**72**:1989–1999. https://doi.org/10.1016/j.jacc.2018.07.089
516. Hanratty CG, Koyama Y, Rasmussen HH, Nelson GIC, Hansen PS, Ward MR. Exaggeration of nonculprit stenosis severity during acute myocardial infarction: implications for immediate multivessel revascularization. *J Am Coll Cardiol* 2002;**40**:911–916. https://doi.org/10.1016/s0735-1097(02)02049-1
517. Gibson CM, Ryan KA, Murphy SA, Mesley R, Marble SJ, Giugliano RP, *et al.* Impaired coronary blood flow in nonculprit arteries in the setting of acute myocardial infarction. *J Am Coll Cardiol* 1999;**34**:974–982. https://doi.org/10.1016/s0735-1097(99)00335-6
518. Ntalianis A, Sels JW, Davidavicius G, Tanaka N, Muller O, Trana C, *et al.* Fractional flow reserve for the assessment of nonculprit coronary artery stenoses in patients with acute myocardial infarction. *JACC Cardiovasc Interv* 2010;**3**:1274–1281. https://doi.org/10.1016/j.jcin.2010.08.025
519. Musto C, De Felice F, Rigattieri S, Chin D, Marra A, Nazzaro MS, *et al.* Instantaneous wave-free ratio and fractional flow reserve for the assessment of nonculprit lesions during the index procedure in patients with ST-segment elevation myocardial infarction: the WAVE study. *Am Heart J* 2017;**193**:63–69. https://doi.org/10.1016/j.ahj.2017.07.017
520. Erbay A, Penzel L, Abdelwahed YS, Klotsche J, Schatz A-S, Steiner J, *et al.* Feasibility and diagnostic reliability of quantitative flow ratio in the assessment of non-culprit lesions in acute coronary syndrome. *Int J Cardiovasc Imaging* 2021;**37**:1815–1823. https://doi.org/10.1007/s10554-021-02195-2
521. Tonino PA, Fearon WF, De Bruyne B, Oldroyd KG, Leesar MA, Ver Lee PN, *et al.* Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol* 2010;**55**:2816–2821. https://doi.org/10.1016/j.jacc.2009.11.096
522. Van Belle E, Baptista SB, Raposo L, Henderson J, Rioufol G, Santos L, *et al.* Impact of routine fractional flow reserve on management decision and 1-year clinical outcome of patients with acute coronary syndromes: PRIME-FFR (Insights From the POST-IT [Portuguese Study on the Evaluation of FFR-Guided Treatment of Coronary Disease] and R3F [French FFR Registry] Integrated Multicenter Registries – Implementation of FFR [Fractional Flow Reserve] in Routine Practice). *Circ Cardiovasc Interv* 2017;**10**:e004296. https://doi.org/10.1161/circinterventions.116.004296
523. Sels JW, Tonino PA, Siebert U, Fearon WF, Van't Veer M, De Bruyne B, *et al.* Fractional flow reserve in unstable angina and non-ST-segment elevation myocardial infarction experience from the FAME (Fractional flow reserve versus Angiography for Multivessel Evaluation) study. *JACC Cardiovasc Interv* 2011;**4**:1183–1189. https://doi.org/10.1016/j.jcin.2011.08.008
524. Puymirat E, Cayla G, Simon T, Steg PG, Montalescot G, Durand-Zaleski I, *et al.* Multivessel PCI guided by FFR or angiography for myocardial infarction. *N Engl J Med* 2021;**385**:297–308. https://doi.org/10.1056/NEJMoa2104650
525. Wald DS, Hadyanto S, Bestwick JP. Should fractional flow reserve follow angiographic visual inspection to guide preventive percutaneous coronary intervention in ST-elevation myocardial infarction? *Eur Heart J Qual Care Clin Outcomes* 2020;**6**:186–192. https://doi.org/10.1093/ehjqcco/qcaa012
526. Gallone G, Angelini F, Fortuni F, Gnecchi M, De Filippo O, Baldetti L, *et al.* Angiography- vs. physiology-guided complete revascularization in patients with ST-elevation myocardial infarction and multivessel disease: who is the better gatekeeper in this setting? A meta-analysis of randomized controlled trials. *Eur Heart J Qual Care Clin Outcomes* 2020;**6**:199–200. https://doi.org/10.1093/ehjqcco/qcaa007
527. Kobayashi Y, Lønborg J, Jong A, Nishi T, De Bruyne B, Høfsten DE, *et al.* Prognostic value of the residual SYNTAX score after functionally complete revascularization in ACS. *J Am Coll Cardiol* 2018;**72**:1321–1329. https://doi.org/10.1016/j.jacc.2018.06.069
528. Lee JM, Kim HK, Park KH, Choo EH, Kim CJ, Lee SH, *et al.* Fractional flow reserve versus angiography-guided strategy in acute myocardial infarction with multivessel disease: a randomized trial. *Eur Heart J* 2023;**44**:473–484. https://doi.org/10.1093/eurheartj/ehac763
529. Harskamp RE, Bonatti JO, Zhao DX, Puskas JD, de Winter RJ, Alexander JH, *et al.* Standardizing definitions for hybrid coronary revascularization. *J Thorac Cardiovasc Surg* 2014;**147**:556–560. https://doi.org/10.1016/j.jtcvs.2013.10.019
530. Doenst T, Haverich A, Serruys P, Bonow RO, Kappetein P, Falk V, *et al.* PCI and CABG for treating stable coronary artery disease: JACC review topic of the week. *J Am Coll Cardiol* 2019;**73**:964–976. https://doi.org/10.1016/j.jacc.2018.11.053
531. Gershlick AH, Khan JN, Kelly DJ, Greenwood JP, Sasikaran T, Curzen N, *et al.* Randomized trial of complete versus lesion-only revascularization in patients undergoing primary percutaneous coronary intervention for STEMI and multivessel disease: the CvLPRIT trial. *J Am Coll Cardiol* 2015;**65**:963–972. https://doi.org/10.1016/j.jacc.2014.12.038
532. Layland J, Oldroyd KG, Curzen N, Sood A, Balachandran K, Das R, *et al.* Fractional flow reserve vs. angiography in guiding management to optimize outcomes in non-ST-segment elevation myocardial infarction: the British Heart Foundation FAMOUS-NSTEMI randomized trial. *Eur Heart J* 2015;**36**:100–111. https://doi.org/10.1093/eurheartj/ehu338
533. Pasupathy S, Air T, Dreyer RP, Tavella R, Beltrame JF. Systematic review of patients presenting with suspected myocardial infarction and nonobstructive coronary arteries. *Circulation* 2015;**131**:861–870. https://doi.org/10.1161/circulationaha.114.011201
534. Agewall S, Beltrame JF, Reynolds HR, Niessner A, Rosano G, Caforio ALP, *et al.* ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J* 2017;**38**:143–153. https://doi.org/10.1093/eurheartj/ehw149
535. Tamis-Holland JE, Jneid H, Reynolds HR, Agewall S, Brilakis ES, Brown TM, *et al.* Contemporary diagnosis and management of patients with myocardial infarction in the absence of obstructive coronary artery disease: a scientific statement from the American Heart Association. *Circulation* 2019;**139**:e891–e908. https://doi.org/10.1161/cir.0000000000000670
536. Matta A, Nader V, Canitrot R, Delmas C, Bouisset F, Lhermusier T, *et al.* Myocardial bridging is significantly associated to myocardial infarction with non-obstructive coronary arteries. *Eur Heart J Acute Cardiovasc Care* 2022;**11**:501–507. https://doi.org/10.1093/ehjacc/zuac047
537. Pargaonkar VS, Kimura T, Kameda R, Tanaka S, Yamada R, Schwartz JG, *et al.* Invasive assessment of myocardial bridging in patients with angina and no obstructive coronary artery disease. *EuroIntervention* 2021;**16**:1070–1078. https://doi.org/10.4244/eij-d-20-00779
538. Ford TJ, Ong P, Sechtem U, Beltrame J, Camici PG, Crea F, *et al.* Assessment of vascular dysfunction in patients without obstructive coronary artery disease: why, how, and when. *JACC Cardiovasc Interv* 2020;**13**:1847–1864. https://doi.org/10.1016/j.jcin.2020.05.052
539. Occhipinti G, Bucciarelli-Ducci C, Capodanno D. Diagnostic pathways in myocardial infarction with non-obstructive coronary artery disease (MINOCA). *Eur Heart J Acute Cardiovasc Care* 2021;**10**:813–822. https://doi.org/10.1093/ehjacc/zuab049
540. Eitel I, Behrendt F, Schindler K, Kivelitz D, Gutberlet M, Schuler G, *et al.* Differential diagnosis of suspected apical ballooning syndrome using contrast-enhanced magnetic resonance imaging. *Eur Heart J* 2008;**29**:2651–2659. https://doi.org/10.1093/eurheartj/ehn433
541. Eitel I, von Knobelsdorff-Brenkenhoff F, Bernhardt P, Carbone I, Muellerleile K, Aldrovandi A, *et al.* Clinical characteristics and cardiovascular magnetic resonance findings in stress (takotsubo) cardiomyopathy. *JAMA* 2011;**306**:277–286. https://doi.org/10.1001/jama.2011.992
542. Ferreira VM, Schulz-Menger J, Holmvang G, Kramer CM, Carbone I, Sechtem U, *et al.* Cardiovascular magnetic resonance in nonischemic myocardial inflammation: expert recommendations. *J Am Coll Cardiol* 2018;**72**:3158–3176. https://doi.org/10.1016/j.jacc.2018.09.072
543. Lurz P, Luecke C, Eitel I, Föhrenbach F, Frank C, Grothoff M, *et al.* Comprehensive cardiac magnetic resonance imaging in patients with suspected myocarditis: the MyoRacer-trial. *J Am Coll Cardiol* 2016;**67**:1800–1811. https://doi.org/10.1016/j.jacc.2016.02.013
544. Reynolds HR, Maehara A, Kwong RY, Sedlak T, Saw J, Smilowitz NR, *et al.* Coronary optical coherence tomography and cardiac magnetic resonance imaging to determine underlying causes of myocardial infarction with nonobstructive coronary arteries in women. *Circulation* 2021;**143**:624–640. https://doi.org/10.1161/circulationaha.120.052008
545. Pathik B, Raman B, Mohd Amin NH, Mahadavan D, Rajendran S, McGavigan AD, *et al.* Troponin-positive chest pain with unobstructed coronary arteries: incremental diagnostic value of cardiovascular magnetic resonance imaging. *Eur Heart J Cardiovasc Imaging* 2016;**17**:1146–1152. https://doi.org/10.1093/ehjci/jev289
546. Lyon AR, Bossone E, Schneider B, Sechtem U, Citro R, Underwood SR, *et al.* Current state of knowledge on Takotsubo syndrome: a Position Statement from the Taskforce on Takotsubo Syndrome of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2016;**18**:8–27. https://doi.org/10.1002/ejhf.424
547. Ghadri JR, Wittstein IS, Prasad A, Sharkey S, Dote K, Akashi YJ, *et al.* International expert consensus document on Takotsubo syndrome (Part II): diagnostic workup,

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

outcome, and management. *Eur Heart J* 2018;**39**:2047–2062. https://doi.org/10.1093/eurheartj/ehy077
548. Adler Y, Charron P, Imazio M, Badano L, Barón-Esquivias G, Bogaert J, *et al.* 2015 ESC Guidelines for the diagnosis and management of pericardial diseases: the Task Force for the diagnosis and management of pericardial diseases of the European Society of Cardiology (ESC) endorsed by: the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2015;**36**:2921–2964. https://doi.org/10.1093/eurheartj/ehv318
549. Caforio AL, Pankuweit S, Arbustini E, Basso C, Gimeno-Blanes J, Felix SB, *et al.* Current state of knowledge on aetiology, diagnosis, management, and therapy of myocarditis: a position statement of the European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2013;**34**:2636–2648,2648a-2648d. https://doi.org/10.1093/eurheartj/eht210
550. Ford TJ, Stanley B, Good R, Rocchiccioli P, McEntegart M, Watkins S, *et al.* Stratified medical therapy using invasive coronary function testing in angina: the CorMicA trial. *J Am Coll Cardiol* 2018;**72**:2841–2855. https://doi.org/10.1016/j.jacc.2018.09.006
551. Kunadian V, Chieffo A, Camici PG, Berry C, Escaned J, Maas AHEM, *et al.* An EAPCI expert consensus document on ischaemia with non-obstructive coronary arteries in collaboration with European Society of Cardiology Working Group on Coronary Pathophysiology & Microcirculation Endorsed by Coronary Vasomotor Disorders International Study Group. *Eur Heart J* 2020;**41**:3504–3520. https://doi.org/10.1093/eurheartj/ehaa503
552. Cerrato E, Giacobbe F, Quadri G, Macaya F, Bianco M, Mori R, *et al.* Antiplatelet therapy in patients with conservatively managed spontaneous coronary artery dissection from the multicentre DISCO registry. *Eur Heart J* 2021;**42**:3161–3171. https://doi.org/10.1093/eurheartj/ehab372
553. McCarthy CP, Raber I, Chapman AR, Sandoval Y, Apple FS, Mills NL, *et al.* Myocardial injury in the era of high-sensitivity cardiac troponin assays: a practical approach for clinicians. *JAMA Cardiol* 2019;**4**:1034–1042. https://doi.org/10.1001/jamacardio.2019.2724
554. Bahit MC, Lopes RD, Clare RM, Newby LK, Pieper KS, Van de Werf F, *et al.* Heart failure complicating non-ST-segment elevation acute coronary syndrome: timing, predictors, and clinical outcomes. *JACC Heart Fail* 2013;**1**:223–229. https://doi.org/10.1016/j.jchf.2013.02.007
555. Chioncel O, Mebazaa A, Harjola VP, Coats AJ, Piepoli MF, Crespo-Leiro MG, *et al.* Clinical phenotypes and outcome of patients hospitalized for acute heart failure: the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail* 2017;**19**:1242–1254. https://doi.org/10.1002/ejhf.890
556. Steg PG, Dabbous OH, Feldman LJ, Cohen-Solal A, Aumont MC, López-Sendón J, *et al.* Determinants and prognostic impact of heart failure complicating acute coronary syndromes: observations from the Global Registry of Acute Coronary Events (GRACE). *Circulation* 2004;**109**:494–499. https://doi.org/10.1161/01.Cir.0000109691.16944.Da
557. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, *et al.* 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;**42**:3599–3726. https://doi.org/10.1093/eurheartj/ehab368
558. Chioncel O, Parissis J, Mebazaa A, Thiele H, Desch S, Bauersachs J, *et al.* Epidemiology, pathophysiology and contemporary management of cardiogenic shock—a position statement from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2020;**22**:1315–1341. https://doi.org/10.1002/ejhf.1922
559. Mullens W, Damman K, Harjola VP, Mebazaa A, Brunner-La Rocca HP, Martens P, *et al.* The use of diuretics in heart failure with congestion—a position statement from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2019;**21**:137–155. https://doi.org/10.1002/ejhf.1369
560. Lancellotti P, Price S, Edvardsen T, Cosyns B, Neskovic AN, Dulgheru R, *et al.* The use of echocardiography in acute cardiovascular care: recommendations of the European Association of Cardiovascular Imaging and the Acute Cardiovascular Care Association. *Eur Heart J Cardiovasc Imaging* 2015;**16**:119–146. https://doi.org/10.1093/ehjci/jeu210
561. Thiele H, Jobs A, Ouweneel DM, Henriques JPS, Seyfarth M, Desch S, *et al.* Percutaneous short-term active mechanical support devices in cardiogenic shock: a systematic review and collaborative meta-analysis of randomized trials. *Eur Heart J* 2017;**38**:3523–3531. https://doi.org/10.1093/eurheartj/ehx363
562. Amin AP, Spertus JA, Curtis JP, Desai N, Masoudi FA, Bach RG, *et al.* The evolving landscape of impella use in the United States among patients undergoing percutaneous coronary intervention with mechanical circulatory support. *Circulation* 2020;**141**:273–284. https://doi.org/10.1161/circulationaha.119.044007
563. Dhruva SS, Ross JS, Mortazavi BJ, Hurley NC, Krumholz HM, Curtis JP, *et al.* Association of use of an intravascular microaxial left ventricular assist device vs intra-aortic balloon pump with in-hospital mortality and major bleeding among patients with acute myocardial infarction complicated by cardiogenic shock. *JAMA* 2020;**323**:734–745. https://doi.org/10.1001/jama.2020.0254
564. Elbadawi A, Elgendy IY, Mahmoud K, Barakat AF, Mentias A, Mohamed AH, *et al.* Temporal trends and outcomes of mechanical complications in patients with acute myocardial infarction. *JACC Cardiovasc Interv* 2019;**12**:1825–1836. https://doi.org/10.1016/j.jcin.2019.04.039
565. Schrage B, Becher PM, Goßling A, Savarese G, Dabboura S, Yan I, *et al.* Temporal trends in incidence, causes, use of mechanical circulatory support and mortality in cardiogenic shock. *ESC Heart Fail* 2021;**8**:1295–1303. https://doi.org/10.1002/ehf2.13202
566. Watkins AC, Maassel NL, Ghoreishi M, Dawood MY, Pham SM, Kon ZN, *et al.* Preoperative venoarterial extracorporeal membrane oxygenation slashes risk score in advanced structural heart disease. *Ann Thorac Surg* 2018;**106**:1709–1715. https://doi.org/10.1016/j.athoracsur.2018.07.038
567. Ronco D, Matteucci M, Ravaux JM, Marra S, Torchio F, Corazzari C, *et al.* Mechanical circulatory support as a bridge to definitive treatment in post-infarction ventricular septal rupture. *JACC Cardiovasc Interv* 2021;**14**:1053–1066. https://doi.org/10.1016/j.jcin.2021.02.046
568. Matteucci M, Fina D, Jiritano F, Meani P, Raffa GM, Kowalewski M, *et al.* The use of extracorporeal membrane oxygenation in the setting of postinfarction mechanical complications: outcome analysis of the Extracorporeal Life Support Organization Registry. *Interact Cardiovasc Thorac Surg* 2020;**31**:369–374. https://doi.org/10.1093/icvts/ivaa108
569. Assenza GE, McElhinney DB, Valente AM, Pearson DD, Volpe M, Martucci G, *et al.* Transcatheter closure of post-myocardial infarction ventricular septal rupture. *Circ Cardiovasc Interv* 2013;**6**:59–67. https://doi.org/10.1161/circinterventions.112.972711
570. Kilic A, Sultan I, Chu D, Wang Y, Gleason TG. Mitral valve surgery for papillary muscle rupture: outcomes in 1342 patients from the society of thoracic surgeons database. *Ann Thorac Surg* 2020;**110**:1975–1981. https://doi.org/10.1016/j.athoracsur.2020.03.097
571. Valle JA, Miyasaka RL, Carroll JD. Acute mitral regurgitation secondary to papillary muscle tear: is transcatheter edge-to-edge mitral valve repair a new paradigm? *Circ Cardiovasc Interv* 2017;**10**:e005050. https://doi.org/10.1161/circinterventions.117.005050
572. Terashima M, Fujiwara S, Yaginuma GY, Takizawa K, Kaneko U, Meguro T. Outcome of percutaneous intrapericardial fibrin-glue injection therapy for left ventricular free wall rupture secondary to acute myocardial infarction. *Am J Cardiol* 2008;**101**:419–421. https://doi.org/10.1016/j.amjcard.2007.09.086
573. Damluji AA, van Diepen S, Katz JN, Menon V, Tamis-Holland JE, Bakitas M, *et al.* Mechanical complications of acute myocardial infarction: a scientific statement from the American Heart Association. *Circulation* 2021;**144**:e16–e35. https://doi.org/10.1161/cir.0000000000000985
574. Gong FF, Vaitenas I, Malaisrie SC, Maganti K. Mechanical complications of acute myocardial infarction: a review. *JAMA Cardiol* 2021;**6**:341–349. https://doi.org/10.1001/jamacardio.2020.3690
575. Robinson AA, Jain A, Gentry M, McNamara RL. Left ventricular thrombi after STEMI in the primary PCI era: a systematic review and meta-analysis. *Int J Cardiol* 2016;**221**:554–559. https://doi.org/10.1016/j.ijcard.2016.07.069
576. Levine GN, McEvoy JW, Fang JC, Ibeh C, McCarthy CP, Misra A, *et al.* Management of patients at risk for and with left ventricular thrombus: a scientific statement from the American Heart Association. *Circulation* 2022;**146**:e205–e223. https://doi.org/10.1161/cir.0000000000001092
577. Bulluck H, Chan MHH, Paradies V, Yellon RL, Ho HH, Chan MY, *et al.* Incidence and predictors of left ventricular thrombus by cardiovascular magnetic resonance in acute ST-segment elevation myocardial infarction treated by primary percutaneous coronary intervention: a meta-analysis. *J Cardiovasc Magn Reson* 2018;**20**:72. https://doi.org/10.1186/s12968-018-0494-3
578. Velangi PS, Choo C, Chen KA, Kazmirczak F, Nijjar PS, Farzaneh-Far A, *et al.* Long-term embolic outcomes after detection of left ventricular thrombus by late gadolinium enhancement cardiovascular magnetic resonance imaging: a matched cohort study. *Circ Cardiovasc Imaging* 2019;**12**:e009723. https://doi.org/10.1161/circimaging.119.009723
579. Funke Küpper AJ, Verheugt FW, Peels CH, Galema TW, Roos JP. Left ventricular thrombus incidence and behavior studied by serial two-dimensional echocardiography in acute anterior myocardial infarction: left ventricular wall motion, systemic embolism and oral anticoagulation. *J Am Coll Cardiol* 1989;**13**:1514–1520. https://doi.org/10.1016/0735-1097(89)90341-0
580. Zhang Z, Si D, Zhang Q, Jin L, Zheng H, Qu M, *et al.* Prophylactic rivaroxaban therapy for left ventricular thrombus after anterior ST-segment elevation myocardial infarction. *JACC Cardiovasc Interv* 2022;**15**:861–872. https://doi.org/10.1016/j.jcin.2022.01.285
581. Dalia T, Lahan S, Ranka S, Goyal A, Zoubek S, Gupta K, *et al.* Warfarin versus direct oral anticoagulants for treating left ventricular thrombus: a systematic review and meta-analysis. *Thromb J* 2021;**19**:7. https://doi.org/10.1186/s12959-021-00259-w
582. Verma BR, Montane B, Chetrit M, Khayata M, Furqan MM, Ayoub C, *et al.* Pericarditis and post-cardiac injury syndrome as a sequelae of acute myocardial infarction. *Curr Cardiol Rep* 2020;**22**:127. https://doi.org/10.1007/s11886-020-01371-5
583. Schmitt J, Duray G, Gersh BJ, Hohnloser SH. Atrial fibrillation in acute myocardial infarction: a systematic review of the incidence, clinical features and prognostic implications. *Eur Heart J* 2009;**30**:1038–1045. https://doi.org/10.1093/eurheartj/ehn579
584. Batra G, Svennblad B, Held C, Jernberg T, Johanson P, Wallentin L, *et al.* All types of atrial fibrillation in the setting of myocardial infarction are associated with impaired outcome. *Heart* 2016;**102**:926–933. https://doi.org/10.1136/heartjnl-2015-308678
585. Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C, *et al.* 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). The Task Force for the diagnosis and management of atrial fibrillation of the European

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

Society of Cardiology (ESC). Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J* 2021;**42**:373–498. https://doi.org/10.1093/eurheartj/ehaa612
586. Jabre P, Jouven X, Adnet F, Thabut G, Bielinski SJ, Weston SA, *et al.* Atrial fibrillation and death after myocardial infarction: a community study. *Circulation* 2011;**123**:2094–2100. https://doi.org/10.1161/circulationaha.110.990192
587. Siu CW, Jim MH, Ho HH, Miu R, Lee SWL, Lau CP, *et al.* Transient atrial fibrillation complicating acute inferior myocardial infarction: implications for future risk of ischemic stroke. *Chest* 2007;**132**:44–49. https://doi.org/10.1378/chest.06-2733
588. Piccini JP, Schulte PJ, Pieper KS, Mehta RH, White HD, Van de Werf F, *et al.* Antiarrhythmic drug therapy for sustained ventricular arrhythmias complicating acute myocardial infarction. *Crit Care Med* 2011;**39**:78–83. https://doi.org/10.1097/CCM.0b013e3181fd6ad7
589. Piccini JP, Hranitzky PM, Kilaru R, Rouleau JL, White HD, Aylward PE, *et al.* Relation of mortality to failure to prescribe beta blockers acutely in patients with sustained ventricular tachycardia and ventricular fibrillation following acute myocardial infarction (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry). *Am J Cardiol* 2008;**102**:1427–1432. https://doi.org/10.1016/j.amjcard.2008.07.033
590. Wolfe CL, Nibley C, Bhandari A, Chatterjee K, Scheinman M. Polymorphous ventricular tachycardia associated with acute myocardial infarction. *Circulation* 1991;**84**:1543–1551. https://doi.org/10.1161/01.cir.84.4.1543
591. Liang JJ, Fender EA, Cha YM, Lennon RJ, Prasad A, Barsness GW, *et al.* Long-term outcomes in survivors of early ventricular arrhythmias after acute ST-elevation and non-ST-elevation myocardial infarction treated with percutaneous coronary intervention. *Am J Cardiol* 2016;**117**:709–713. https://doi.org/10.1016/j.amjcard.2015.12.002
592. Mehta RH, Yu J, Piccini JP, Tcheng JE, Farkouh ME, Reiffel J, *et al.* Prognostic significance of postprocedural sustained ventricular tachycardia or fibrillation in patients undergoing primary percutaneous coronary intervention (from the HORIZONS-AMI Trial). *Am J Cardiol* 2012;**109**:805–812. https://doi.org/10.1016/j.amjcard.2011.10.043
593. Masuda M, Nakatani D, Hikoso S, Suna S, Usami M, Matsumoto S, *et al.* Clinical impact of ventricular tachycardia and/or fibrillation during the acute phase of acute myocardial infarction on in-hospital and 5-year mortality rates in the percutaneous coronary intervention era. *Circ J* 2016;**80**:1539–1547. https://doi.org/10.1253/circj.CJ-16-0183
594. Podolecki T, Lenarczyk R, Kowalczyk J, Jedrzejczyk-Patej E, Chodor P, Mazurek M, *et al.* Prognostic significance of complex ventricular arrhythmias complicating ST-segment elevation myocardial infarction. *Am J Cardiol* 2018;**121**:805–809. https://doi.org/10.1016/j.amjcard.2017.12.036
595. Komatsu Y, Hocini M, Nogami A, Maury P, Peichl P, Iwasaki YK, *et al.* Catheter ablation of refractory ventricular fibrillation storm after myocardial infarction. *Circulation* 2019;**139**:2315–2325. https://doi.org/10.1161/circulationaha.118.037997
596. Terkelsen CJ, Sørensen JT, Kaltoft AK, Nielsen SS, Thuesen L, Bøtker HE, *et al.* Prevalence and significance of accelerated idioventricular rhythm in patients with ST-elevation myocardial infarction treated with primary percutaneous coronary intervention. *Am J Cardiol* 2009;**104**:1641–1646. https://doi.org/10.1016/j.amjcard.2009.07.037
597. Mehran R, Pocock SJ, Nikolsky E, Clayton T, Dangas GD, Kirtane AJ, *et al.* A risk score to predict bleeding in patients with acute coronary syndromes. *J Am Coll Cardiol* 2010;**55**:2556–2566. https://doi.org/10.1016/j.jacc.2009.09.076
598. Mehran R, Pocock SJ, Stone GW, Clayton TC, Dangas GD, Feit F, *et al.* Associations of major bleeding and myocardial infarction with the incidence and timing of mortality in patients presenting with non-ST-elevation acute coronary syndromes: a risk model from the ACUITY trial. *Eur Heart J* 2009;**30**:1457–1466. https://doi.org/10.1093/eurheartj/ehp110
599. Rao SV. The conundrum of reducing ischemic and bleeding events after PCI. *J Am Coll Cardiol* 2015;**65**:1421–1423. https://doi.org/10.1016/j.jacc.2015.02.012
600. Kwok CS, Sherwood MW, Watson SM, Nasir SB, Sperrin M, Nolan J, *et al.* Blood transfusion after percutaneous coronary intervention and risk of subsequent adverse outcomes: a systematic review and meta-analysis. *JACC Cardiovasc Interv* 2015;**8**:436–446. https://doi.org/10.1016/j.jcin.2014.09.026
601. Mehran R, Baber U, Steg PG, Ariti C, Weisz G, Witzenbichler B, *et al.* Cessation of dual antiplatelet treatment and cardiac events after percutaneous coronary intervention (PARIS): 2 year results from a prospective observational study. *Lancet* 2013;**382**:1714–1722. https://doi.org/10.1016/s0140-6736(13)61720-1
602. Reidenberg MM. Drug discontinuation effects are part of the pharmacology of a drug. *J Pharmacol Exp Ther* 2011;**339**:324–328. https://doi.org/10.1124/jpet.111.183285
603. Abdelnabi M, Saleh Y, Fareed A, Nossikof A, Wang L, Morsi M, *et al.* Comparative study of oral anticoagulation in left ventricular thrombi (no-LVT trial). *J Am Coll Cardiol* 2021;**77**:1590–1592. https://doi.org/10.1016/j.jacc.2021.01.049
604. Weinsaft JW, Kim J, Medicherla CB, Ma CL, Codella NCF, Kukar N, *et al.* Echocardiographic algorithm for post-myocardial infarction LV thrombus: a gatekeeper for thrombus evaluation by delayed enhancement CMR. *JACC Cardiovasc Imaging* 2016;**9**:505–515. https://doi.org/10.1016/j.jcmg.2015.06.017
605. Segal JB, McNamara RL, Miller MR, Kim N, Goodman SN, Powe NR, *et al.* The evidence regarding the drugs used for ventricular rate control. *J Fam Pract* 2000;**49**:47–59
606. Hou ZY, Chang MS, Chen CY, Tu MS, Lin SL, Chiang HT, *et al.* Acute treatment of recent-onset atrial fibrillation and flutter with a tailored dosing regimen of intravenous amiodarone: a randomized, digoxin-controlled study. *Eur Heart J* 1995;**16**:521–528. https://doi.org/10.1093/oxfordjournals.eurheartj.a060945
607. Deedwania PC, Singh BN, Ellenbogen K, Fisher S, Fletcher R, Singh SN. Spontaneous conversion and maintenance of sinus rhythm by amiodarone in patients with heart failure and atrial fibrillation: observations from the veterans affairs Congestive Heart failure Survival Trial of Antiarrhythmic Therapy (CHF-STAT). *Circulation* 1998;**98**:2574–2579. https://doi.org/10.1161/01.cir.98.23.2574
608. Hofmann R, Steinwender C, Kammler J, Kypta A, Wimmer G, Leisch F, *et al.* Intravenous amiodarone bolus for treatment of atrial fibrillation in patients with advanced congestive heart failure or cardiogenic shock. *Wien Klin Wochenschr* 2004;**116**:744–749. https://doi.org/10.1007/s00508-004-0264-0
609. Moss AJ, Zareba W, Hall WJ, Klein H, Wilber DJ, Cannom DS, *et al.* Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med* 2002;**346**:877–883. https://doi.org/10.1056/NEJMoa013474
610. Bardy GH, Lee KL, Mark DB, Poole JE, Packer DL, Boineau R, *et al.* Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med* 2005;**352**:225–237. https://doi.org/10.1056/NEJMoa043399
611. Nademanee K, Taylor R, Bailey WE, Rieders DE, Kosar EM. Treating electrical storm: sympathetic blockade versus advanced cardiac life support-guided therapy. *Circulation* 2000;**102**:742–747. https://doi.org/10.1161/01.cir.102.7.742
612. Miwa Y, Ikeda T, Mera H, Miyakoshi M, Hoshida K, Yanagisawa R, *et al.* Effects of landiolol, an ultra-short-acting beta1-selective blocker, on electrical storm refractory to class III antiarrhythmic drugs. *Circ J* 2010;**74**:856–863. https://doi.org/10.1253/circj.cj-09-0772
613. Kudenchuk PJ, Cobb LA, Copass MK, Cummins RO, Doherty AM, Fahrenbruch CE, *et al.* Amiodarone for resuscitation after out-of-hospital cardiac arrest due to ventricular fibrillation. *N Engl J Med* 1999;**341**:871–878. https://doi.org/10.1056/nejm199909163411203
614. Levine JH, Massumi A, Scheinman MM, Winkle RA, Platia EV, Chilson DA, *et al.* Intravenous amiodarone for recurrent sustained hypotensive ventricular tachyarrhythmias. *J Am Coll Cardiol* 1996;**27**:67–75. https://doi.org/10.1016/0735-1097(95)00427-0
615. Gorenek B, Blomström Lundqvist C, Brugada Terradellas J, Camm AJ, Hindricks G, Huber K, *et al.* Cardiac arrhythmias in acute coronary syndromes: position paper from the joint EHRA, ACCA, and EAPCI task force. *Europace* 2014;**16**:1655–1673. https://doi.org/10.1093/europace/euu208
616. Kalarus Z, Svendsen JH, Capodanno D, Dan GA, De Maria E, Gorenek B, *et al.* Cardiac arrhythmias in the emergency settings of acute coronary syndrome and revascularization: an European Heart Rhythm Association (EHRA) consensus document, endorsed by the European Association of Percutaneous Cardiovascular Interventions (EAPCI), and European Acute Cardiovascular Care Association (ACCA). *Europace* 2019;**21**:1603–1604. https://doi.org/10.1093/europace/euz163
617. Feigl D, Ashkenazy J, Kishon Y. Early and late atrioventricular block in acute inferior myocardial infarction. *J Am Coll Cardiol* 1984;**4**:35–38. https://doi.org/10.1016/s0735-1097(84)80315-0
618. Brady WJ, Swart G, DeBehnke DJ, Ma OJ, Aufderheide TP. The efficacy of atropine in the treatment of hemodynamically unstable bradycardia and atrioventricular block: prehospital and emergency department considerations. *Resuscitation* 1999;**41**:47–55. https://doi.org/10.1016/s0300-9572(99)00032-5
619. Kusumoto FM, Calkins H, Boehmer J, Buxton AE, Chung MK, Gold MR, *et al.* HRS/ACC/AHA expert consensus statement on the use of implantable cardioverter-defibrillator therapy in patients who are not included or not well represented in clinical trials. *Circulation* 2014;**130**:94–125. https://doi.org/10.1161/cir.0000000000000056
620. Glikson M, Nielsen JC, Kronborg MB, Michowitz Y, Auricchio A, Barbash IM, *et al.* 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Eur Heart J* 2021;**42**:3427–3520. https://doi.org/10.1093/eurheartj/ehab364
621. Jim MH, Chan AO, Tse HF, Barold SS, Lau CP. Clinical and angiographic findings of complete atrioventricular block in acute inferior myocardial infarction. *Ann Acad Med Singap* 2010;**39**:185–190. https://doi.org/10.47102/annals-acadmedsg.V39N3p185
622. Gang UJ, Hvelplund A, Pedersen S, Iversen A, Jons C, Abildstrom SZ, *et al.* High-degree atrioventricular block complicating ST-segment elevation myocardial infarction in the era of primary percutaneous coronary intervention. *Europace* 2012;**14**:1639–1645. https://doi.org/10.1093/europace/eus161
623. Vicente-Ibarra N, Marín F, Pernías-Escrig V, Sandín-Rollán M, Núñez-Martínez L, Lozano T, *et al.* Impact of anemia as risk factor for major bleeding and mortality in patients with acute coronary syndrome. *Eur J Intern Med* 2019;**61**:48–53. https://doi.org/10.1016/j.ejim.2018.12.004
624. Younge JO, Nauta ST, Akkerhuis KM, Deckers JW, van Domburg RT. Effect of anemia on short- and long-term outcome in patients hospitalized for acute coronary syndromes. *Am J Cardiol* 2012;**109**:506–510. https://doi.org/10.1016/j.amjcard.2011.09.046
625. Bassand JP, Afzal R, Eikelboom J, Wallentin L, Peters R, Budaj A, *et al.* Relationship between baseline haemoglobin and major bleeding complications in acute coronary syndromes. *Eur Heart J* 2010;**31**:50–58. https://doi.org/10.1093/eurheartj/ehp401

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

626. Carson JL, Guyatt G, Heddle NM, Grossman BJ, Cohn CS, Fung MK, *et al.* Clinical practice guidelines from the AABB: red blood cell transfusion thresholds and storage. *JAMA* 2016;**316**:2025–2035. https://doi.org/10.1001/jama.2016.9185
627. Chatterjee S, Wetterslev J, Sharma A, Lichstein E, Mukherjee D. Association of blood transfusion with increased mortality in myocardial infarction: a meta-analysis and diversity-adjusted study sequential analysis. *JAMA Intern Med* 2013;**173**:132–139. https://doi.org/10.1001/2013.jamainternmed.1001
628. Carson JL, Stanworth SJ, Dennis JA, Trivella M, Roubinian N, Fergusson DA, *et al.* Transfusion thresholds for guiding red blood cell transfusion. *Cochrane Database Syst Rev* 2021;**12**:CD002042. https://doi.org/10.1002/14651858.CD002042.pub5
629. Cooper HA, Rao SV, Greenberg MD, Rumsey MP, McKenzie M, Alcorn KW, *et al.* Conservative versus liberal red cell transfusion in acute myocardial infarction (the CRIT Randomized Pilot Study). *Am J Cardiol* 2011;**108**:1108–1111. https://doi.org/10.1016/j.amjcard.2011.06.014
630. Alexander KP, Chen AY, Wang TY, Rao SV, Newby LK, LaPointe NMA, *et al.* Transfusion practice and outcomes in non-ST-segment elevation acute coronary syndromes. *Am Heart J* 2008;**155**:1047–1053. https://doi.org/10.1016/j.ahj.2008.01.009
631. Ducrocq G, Gonzalez-Juanatey JR, Puymirat E, Lemesle G, Cachanado M, Durand-Zaleski I, *et al.* Effect of a restrictive vs liberal blood transfusion strategy on major cardiovascular events among patients with acute myocardial infarction and anemia: the REALITY randomized clinical trial. *JAMA* 2021;**325**:552–560. https://doi.org/10.1001/jama.2021.0135
632. Gonzalez-Juanatey JR, Lemesle G, Puymirat E, Ducrocq G, Cachanado M, Arnaiz JA, *et al.* One-year major cardiovascular events after restrictive versus liberal blood transfusion strategy in patients with acute myocardial infarction and anemia: the REALITY randomized trial. *Circulation* 2022;**145**:486–488. https://doi.org/10.1161/circulationaha.121.057909
633. Gore JM, Spencer FA, Gurfinkel EP, López-Sendón J, Steg PG, Granger CB, *et al.* Thrombocytopenia in patients with an acute coronary syndrome (from the Global Registry of Acute Coronary Events [GRACE]). *Am J Cardiol* 2009;**103**:175–180. https://doi.org/10.1016/j.amjcard.2008.08.055
634. Vora AN, Chenier M, Schulte PJ, Goodman S, Peterson ED, Pieper K, *et al.* Long-term outcomes associated with hospital acquired thrombocytopenia among patients with non-ST-segment elevation acute coronary syndrome. *Am Heart J* 2014;**168**:189–-196.e1. https://doi.org/10.1016/j.ahj.2014.04.010
635. Szummer K, Lundman P, Jacobson SH, Schön S, Lindbäck J, Stenestrand U, *et al.* Relation between renal function, presentation, use of therapies and in-hospital complications in acute coronary syndrome: data from the SWEDEHEART register. *J Intern Med* 2010;**268**:40–49. https://doi.org/10.1111/j.1365-2796.2009.02204.x
636. Berger AK, Duval S, Krumholz HM. Aspirin, beta-blocker, and angiotensin-converting enzyme inhibitor therapy in patients with end-stage renal disease and an acute myocardial infarction. *J Am Coll Cardiol* 2003;**42**:201–208. https://doi.org/10.1016/s0735-1097(03)00572-2
637. Panchal HB, Zheng S, Devani K, White CJ, Leinaar EF, Mukherjee D, *et al.* Impact of chronic kidney disease on revascularization and outcomes in patients with ST-elevation myocardial infarction. *Am J Cardiol* 2021;**150**:15–23. https://doi.org/10.1016/j.amjcard.2021.03.057
638. Matsushita K, van der Velde M, Astor BC, Woodward M, Levey AS, de Jong PE, *et al.* Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet* 2010;**375**:2073–2081. https://doi.org/10.1016/s0140-6736(10)60674-5
639. Santopinto JJ, Fox KA, Goldberg RJ, Budaj A, Piñero G, Avezum A, *et al.* Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE). *Heart* 2003;**89**:1003–1008. https://doi.org/10.1136/heart.89.9.1003
640. Szummer K, Lundman P, Jacobson SH, Schön S, Lindbäck J, Stenestrand U, *et al.* Influence of renal function on the effects of early revascularization in non-ST-elevation myocardial infarction: data from the Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). *Circulation* 2009;**120**:851–858. https://doi.org/10.1161/circulationaha.108.838169
641. Huang HD, Alam M, Hamzeh I, Virani S, Deswal A, Aguilar D, *et al.* Patients with severe chronic kidney disease benefit from early revascularization after acute coronary syndrome. *Int J Cardiol* 2013;**168**:3741–3746. https://doi.org/10.1016/j.ijcard.2013.06.013
642. Kume K, Yasuoka Y, Adachi H, Noda Y, Hattori S, Araki R, *et al.* Impact of contrast-induced acute kidney injury on outcomes in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Cardiovasc Revasc Med* 2013;**14**:253–257. https://doi.org/10.1016/j.carrev.2013.07.009
643. Bangalore S, Briguori C. Preventive strategies for contrast-induced acute kidney injury: and the winner is…. *Circ Cardiovasc Interv* 2017;**10**:e005262. https://doi.org/10.1161/circinterventions.117.005262
644. Davenport MS, Perazella MA, Yee J, Dillman JR, Fine D, McDonald RJ, *et al.* Use of intravenous iodinated contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Kidney Med* 2020;**2**:85–93. https://doi.org/10.1016/j.xkme.2020.01.001
645. Schweiger MJ, Chambers CE, Davidson CJ, Zhang S, Blankenship J, Bhalla NP, *et al.* Prevention of contrast induced nephropathy: recommendations for the high risk patient undergoing cardiovascular procedures. *Catheter Cardiovasc Interv* 2007;**69**:135–140. https://doi.org/10.1002/ccd.20964
646. Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Bäck M, *et al.* 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021;**42**:3227–3337. https://doi.org/10.1093/eurheartj/ehab484
647. Ängerud KH, Brulin C, Näslund U, Eliasson M. Longer pre-hospital delay in first myocardial infarction among patients with diabetes: an analysis of 4266 patients in the northern Sweden MONICA Study. *BMC Cardiovasc Disord* 2013;**13**:6. https://doi.org/10.1186/1471-2261-13-6
648. Fu R, Li S-D, Song C-X, Yang J-A, Xu H-Y, Gao X-J, *et al.* Clinical significance of diabetes on symptom and patient delay among patients with acute myocardial infarction–an analysis from China Acute Myocardial Infarction (CAMI) registry. *J Geriatr Cardiol* 2019;**16**:395–400. https://doi.org/10.11909/j.issn.1671-5411.2019.05.002
649. Rossello X, Ferreira JP, McMurray JJ, Aguilar D, Pfeffer MA, Pitt B, *et al.* Editor's choice–impact of insulin-treated diabetes on cardiovascular outcomes following high-risk myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2019;**8**:231–241. https://doi.org/10.1177/2048872618803701
650. Wallert J, Mitchell A, Held C, Hagström E, Leosdottir M, Olsson EMG. Cardiac rehabilitation goal attainment after myocardial infarction with versus without diabetes: a nationwide registry study. *Int J Cardiol* 2019;**292**:19–24. https://doi.org/10.1016/j.ijcard.2019.04.049
651. Ritsinger V, Jensen J, Ohm D, Omerovic E, Koul S, Fröbert O, *et al.* Elevated admission glucose is common and associated with high short-term complication burden after acute myocardial infarction: insights from the VALIDATE-SWEDEHEART study. *Diab Vasc Dis Res* 2019;**16**:582–584. https://doi.org/10.1177/1479164119871540
652. Weston C, Walker L, Birkhead J. Early impact of insulin treatment on mortality for hyperglycaemic patients without known diabetes who present with an acute coronary syndrome. *Heart* 2007;**93**:1542–1546. https://doi.org/10.1136/hrt.2006.108696
653. Ritsinger V, Malmberg K, Mårtensson A, Rydén L, Wedel H, Norhammar A. Intensified insulin-based glycaemic control after myocardial infarction: mortality during 20 year follow-up of the randomised Diabetes Mellitus Insulin Glucose Infusion in Acute Myocardial Infarction (DIGAMI 1) trial. *Lancet Diabetes Endocrinol* 2014;**2**:627–633. https://doi.org/10.1016/s2213-8587(14)70088-9
654. Finfer S, Chittock DR, Su SY, Blair D, Foster D, Dhingra V, *et al.* Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;**360**:1283–1297. https://doi.org/10.1056/NEJMoa0810625
655. Rossello X, Yellon DM. A new era in the management of type 2 diabetes: is cardioprotection at long last a reality? *Int J Cardiol* 2017;**228**:198–200. https://doi.org/10.1016/j.ijcard.2016.11.246
656. Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, *et al.* Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015;**373**:2117–2128. https://doi.org/10.1056/NEJMoa1504720
657. Marso SP, Daniels GH, Brown-Frandsen K, Kristensen P, Mann JFE, Nauck MA, *et al.* Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016;**375**:311–322. https://doi.org/10.1056/NEJMoa1603827
658. Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, *et al.* 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2020;**41**:255–323. https://doi.org/10.1093/eurheartj/ehz486
659. Sinclair H, Batty JA, Qiu W, Kunadian V. Engaging older patients in cardiovascular research: observational analysis of the ICON-1 study. *Open Heart* 2016;**3**:e000436. https://doi.org/10.1136/openhrt-2016-000436
660. Rosengren A, Wallentin L, Simoons M, Gitt AK, Behar S, Battler A, *et al.* Age, clinical presentation, and outcome of acute coronary syndromes in the Euroheart acute coronary syndrome survey. *Eur Heart J* 2006;**27**:789–795. https://doi.org/10.1093/eurheartj/ehi774
661. Lopes RD, White JA, Tricoci P, White HD, Armstrong PW, Braunwald E, *et al.* Age, treatment, and outcomes in high-risk non-ST-segment elevation acute coronary syndrome patients: insights from the EARLY ACS trial. *Int J Cardiol* 2013;**167**:2580–2587. https://doi.org/10.1016/j.ijcard.2012.06.053
662. Reiter M, Twerenbold R, Reichlin T, Haaf P, Peter F, Meissner J, *et al.* Early diagnosis of acute myocardial infarction in the elderly using more sensitive cardiac troponin assays. *Eur Heart J* 2011;**32**:1379–1389. https://doi.org/10.1093/eurheartj/ehr033
663. Mills GB, Ratcovich H, Adams-Hall J, Beska B, Kirkup E, Raharjo DE, *et al.* Is the contemporary care of the older persons with acute coronary syndrome evidence-based? *European Heart Journal Open* 2021;**2**:oeab044. https://doi.org/10.1093/ehjopen/oeab044
664. Tegn N, Abdelnoor M, Aaberge L, Endresen K, Smith P, Aakhus S, *et al.* Invasive versus conservative strategy in patients aged 80 years or older with non-ST-elevation myocardial infarction or unstable angina pectoris (After Eighty study): an open-label randomised controlled trial. *Lancet* 2016;**387**:1057–1065. https://doi.org/10.1016/s0140-6736(15)01166-6
665. Bueno H, Betriu A, Heras M, Alonso JJ, Cequier A, Garcia EJ, *et al.* Primary angioplasty vs. fibrinolysis in very old patients with acute myocardial infarction: TRIANA

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

(TRatamiento del Infarto Agudo de miocardio eN Ancianos) randomized trial and pooled analysis with previous studies. *Eur Heart J* 2011;**32**:51–60. https://doi.org/10.1093/eurheartj/ehq375

666. Kunadian V, Qiu W, Ludman P, Redwood S, Curzen N, Stables R, *et al.* Outcomes in patients with cardiogenic shock following percutaneous coronary intervention in the contemporary era: an analysis from the BCIS database (British Cardiovascular Intervention Society). *JACC Cardiovasc Interv* 2014;**7**:1374–1385. https://doi.org/10.1016/j.jcin.2014.06.017
667. Kunadian V, Bawamia B, Maznyczka A, Zaman A, Qiu W. Outcomes following primary percutaneous coronary intervention in the setting of cardiac arrest: a registry database study. *Eur Heart J Acute Cardiovasc Care* 2015;**4**:6–15. https://doi.org/10.1177/2048872614534079
668. Richter D, Guasti L, Walker D, Lambrinou E, Lionis C, Abreu A, *et al.* Frailty in cardiology: definition, assessment and clinical implications for general cardiology. A consensus document of the Council for Cardiology Practice (CCP), Association for Acute Cardio Vascular Care (ACVC), Association of Cardiovascular Nursing and Allied Professions (ACNAP), European Association of Preventive Cardiology (EAPC), European Heart Rhythm Association (EHRA), Council on Valvular Heart Diseases (VHD), Council on Hypertension (CHT), Council of Cardio-Oncology (CCO), Working Group (WG) Aorta and Peripheral Vascular Diseases, WG e-Cardiology, WG Thrombosis, of the European Society of Cardiology, European Primary Care Cardiology Society (EPCCS). *Eur J Prev Cardiol* 2022;**29**:216–227. https://doi.org/10.1093/eurjpc/zwaa167
669. Walker DM, Gale CP, Lip G, Martin-Sanchez FJ, McIntyre HF, Mueller C, *et al.* Editor's choice–frailty and the management of patients with acute cardiovascular disease: a position paper from the Acute Cardiovascular Care Association. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:176–193. https://doi.org/10.1177/2048872618758931
670. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet* 2013;**381**:752–762. https://doi.org/10.1016/s0140-6736(12)62167-9
671. Chung KJNC, Wilkinson C, Veerasamy M, Kunadian V. Frailty scores and their utility in older patients with cardiovascular disease. *Interv Cardiol* 2021;**16**:e05. https://doi.org/10.15420/icr.2020.18
672. Gu SZ, Qiu W, Batty JA, Sinclair H, Veerasamy M, Brugaletta S, *et al.* Coronary artery lesion phenotype in frail older patients with non-ST-elevation acute coronary syndrome undergoing invasive care. *EuroIntervention* 2019;**15**:e261–e268. https://doi.org/10.4244/eij-d-18-00848
673. Batty J, Qiu W, Gu S, Sinclair H, Veerasamy M, Beska B, *et al.* One-year clinical outcomes in older patients with non-ST elevation acute coronary syndrome undergoing coronary angiography: an analysis of the ICON1 study. *Int J Cardiol* 2019;**274**:45–51. https://doi.org/10.1016/j.ijcard.2018.09.086
674. Beska B, Coakley D, MacGowan G, Adams-Hall J, Wilkinson C, Kunadian V, *et al.* Frailty and quality of life after invasive management for non-ST elevation acute coronary syndrome. *Heart* 2022;**108**:203–211. https://doi.org/10.1136/heartjnl-2021-319064
675. Beska B, Mills GB, Ratcovich H, Wilkinson C, Damluji AA, Kunadian V, *et al.* Impact of multimorbidity on long-term outcomes in older adults with non-ST elevation acute coronary syndrome in the North East of England: a multi-centre cohort study of patients undergoing invasive care. *BMJ Open* 2022;**12**:e061830. https://doi.org/10.1136/bmjopen-2022-061830
676. Gu SZ, Beska B, Chan D, Neely D, Batty JA, Adams-Hall J, *et al.* Cognitive decline in older patients with non-ST elevation acute coronary syndrome. *J Am Heart Assoc* 2019;**8**:e011218. https://doi.org/10.1161/jaha.118.011218
677. Ismail S, Wong C, Rajan P, Vidovich MI. ST-elevation acute myocardial infarction in pregnancy: 2016 update. *Clin Cardiol* 2017;**40**:399–406. https://doi.org/10.1002/clc.22655
678. Regitz-Zagrosek V, Roos-Hesselink JW, Bauersachs J, Blomström-Lundqvist C, Cífková R, De Bonis M, *et al.* 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. *Eur Heart J* 2018;**39**:3165–3241. https://doi.org/10.1093/eurheartj/ehy340
679. Roth A, Elkayam U. Acute myocardial infarction associated with pregnancy. *J Am Coll Cardiol* 2008;**52**:171–180. https://doi.org/10.1016/j.jacc.2008.03.049
680. Elkayam U, Jalnapurkar S, Barakkat MN, Khatri N, Kealey AJ, Mehra A, *et al.* Pregnancy-associated acute myocardial infarction: a review of contemporary experience in 150 cases between 2006 and 2011. *Circulation* 2014;**129**:1695–1702. https://doi.org/10.1161/circulationaha.113.002054
681. Elkayam U, Goland S, Pieper PG, Silverside CK. High-risk cardiac disease in pregnancy: Part I. *J Am Coll Cardiol* 2016;**68**:396–410. https://doi.org/10.1016/j.jacc.2016.05.048
682. Bharadwaj A, Potts J, Mohamed MO, Parwani P, Swamy P, Lopez-Mattei JC, *et al.* Acute myocardial infarction treatments and outcomes in 6.5 million patients with a current or historical diagnosis of cancer in the USA. *Eur Heart J* 2020;**41**:2183–2193. https://doi.org/10.1093/eurheartj/ehz851
683. Velders MA, Boden H, Hofma SH, Osanto S, van der Hoeven BL, Heestermans AACM, *et al.* Outcome after ST elevation myocardial infarction in patients with cancer treated with primary percutaneous coronary intervention. *Am J Cardiol* 2013;**112**:1867–1872. https://doi.org/10.1016/j.amjcard.2013.08.019
684. Lyon AR, López-Fernández T, Couch LS, Asteggiano R, Aznar MC, Bergler-Klein J, *et al.* 2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS): developed by the Task Force on cardio-oncology of the European Society of Cardiology (ESC). *Eur Heart J* 2022;**43**:4229–4361. https://doi.org/10.1093/eurheartj/ehac244
685. Potts JE, Iliescu CA, Lopez Mattei JC, Martinez SC, Holmvang L, Ludman P, *et al.* Percutaneous coronary intervention in cancer patients: a report of the prevalence and outcomes in the United States. *Eur Heart J* 2019;**40**:1790–1800. https://doi.org/10.1093/eurheartj/ehy769
686. Pothineni NV, Shah NN, Rochlani Y, Saad M, Kovelamudi S, Marmagkiolis K, *et al.* Temporal trends and outcomes of acute myocardial infarction in patients with cancer. *Ann Transl Med* 2017;**5**:482. https://doi.org/10.21037/atm.2017.11.29
687. Gevaert SA, Halvorsen S, Sinnaeve PR, Sambola A, Gulati G, Lancellotti P, *et al.* Evaluation and management of cancer patients presenting with acute cardiovascular disease: a Consensus Document of the Acute CardioVascular Care (ACVC) association and the ESC Council of Cardio-Oncology-Part 1: acute coronary syndromes and acute pericardial diseases. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:947–959. https://doi.org/10.1093/ehjacc/zuab056
688. Lancellotti P, Suter TM, López-Fernández T, Galderisi M, Lyon AR, Van der Meer P, *et al.* Cardio-Oncology Services: rationale, organization, and implementation. *Eur Heart J* 2019;**40**:1756–1763. https://doi.org/10.1093/eurheartj/ehy453
689. Mohamed MO, Van Spall HGC, Kontopantelis E, Alkhouli M, Barac A, Elgendy IY, *et al.* Effect of primary percutaneous coronary intervention on in-hospital outcomes among active cancer patients presenting with ST-elevation myocardial infarction: a propensity score matching analysis. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:829–839. https://doi.org/10.1093/ehjacc/zuaa032
690. Guddati AK, Joy PS, Kumar G. Analysis of outcomes of percutaneous coronary intervention in metastatic cancer patients with acute coronary syndrome over a 10-year period. *J Cancer Res Clin Oncol* 2016;**142**:471–479. https://doi.org/10.1007/s00432-015-2056-5
691. Aspelin P, Aubry P, Fransson SG, Strasser R, Willenbrock R, Berg KJ. Nephrotoxic effects in high-risk patients undergoing angiography. *N Engl J Med* 2003;**348**:491–499. https://doi.org/10.1056/NEJMoa021833
692. Jo SH, Youn TJ, Koo BK, Park JS, Kang HJ, Cho YS, *et al.* Renal toxicity evaluation and comparison between visipaque (Iodixanol) and hexabrix (Ioxaglate) in patients with renal insufficiency undergoing coronary angiography: the RECOVER study: a randomized controlled trial. *J Am Coll Cardiol* 2006;**48**:924–930. https://doi.org/10.1016/j.jacc.2006.06.047
693. Solomon RJ, Natarajan MK, Doucet S, Sharma SK, Staniloae CS, Katholi RE, *et al.* Cardiac Angiography in Renally Impaired Patients (CARE) study: a randomized double-blind trial of contrast-induced nephropathy in patients with chronic kidney disease. *Circulation* 2007;**115**:3189–3196. https://doi.org/10.1161/circulationaha.106.671644
694. Brar SS, Shen AY, Jorgensen MB, Kotlewski A, Aharonian VJ, Desai N, *et al.* Sodium bicarbonate vs sodium chloride for the prevention of contrast medium-induced nephropathy in patients undergoing coronary angiography: a randomized trial. *JAMA* 2008;**300**:1038–1046. https://doi.org/10.1001/jama.300.9.1038
695. Brar SS, Aharonian V, Mansukhani P, Moore N, Shen AYJ, Jorgensen M, *et al.* Haemodynamic-guided fluid administration for the prevention of contrast-induced acute kidney injury: the POSEIDON randomised controlled trial. *Lancet* 2014;**383**:1814–1823. https://doi.org/10.1016/s0140-6736(14)60689-9
696. Giacoppo D, Gargiulo G, Buccheri S, Aruta P, Byrne RA, Cassese S, *et al.* Preventive strategies for contrast-induced acute kidney injury in patients undergoing percutaneous coronary procedures: evidence from a hierarchical Bayesian network meta-analysis of 124 trials and 28240 patients. *Circ Cardiovasc Interv* 2017;**10**:e004383. https://doi.org/10.1161/circinterventions.116.004383
697. Nijssen EC, Rennenberg RJ, Nelemans PJ, Essers BA, Janssen MM, Vermeeren MA, *et al.* Prophylactic hydration to protect renal function from intravascular iodinated contrast material in patients at high risk of contrast-induced nephropathy (AMACING): a prospective, randomised, phase 3, controlled, open-label, non-inferiority trial. *Lancet* 2017;**389**:1312–1322. https://doi.org/10.1016/s0140-6736(17)30057-0
698. Zelniker TA, Wiviott SD, Raz I, Im K, Goodrich EL, Bonaca MP, *et al.* SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet* 2019;**393**:31–39. https://doi.org/10.1016/s0140-6736(18)32590-x
699. Kristensen SL, Rørth R, Jhund PS, Docherty KF, Sattar N, Preiss D, *et al.* Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet Diabetes Endocrinol* 2019;**7**:776–785. https://doi.org/10.1016/s2213-8587(19)30249-9
700. Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, *et al.* Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019;**380**:2295–2306. https://doi.org/10.1056/NEJMoa1811744
701. Heerspink HJL, Stefánsson BV, Correa-Rotter R, Chertow GM, Greene T, Hou FF, *et al.* Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;**383**:1436–1446. https://doi.org/10.1056/NEJMoa2024816

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

702. McMurray JJV, Solomon SD, Inzucchi SE, Køber L, Kosiborod MN, Martinez FA, *et al.* Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 2019;**381**:1995–2008. https://doi.org/10.1056/NEJMoa1911303
703. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, *et al.* Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* 2020;**383**:1413–1424. https://doi.org/10.1056/NEJMoa2022190
704. Packer M, Butler J, Zannad F, Filippatos G, Ferreira JP, Pocock SJ, *et al.* Effect of empagliflozin on worsening heart failure events in patients with heart failure and a preserved ejection fraction: the EMPEROR-preserved trial. *Circulation* 2021;**144**:1284–1294. https://doi.org/10.1161/circulationaha.121.056824
705. Ferrannini G, De Bacquer D, De Backer G, Kotseva K, Mellbin L, Wood D, *et al.* Screening for glucose perturbations and risk factor management in dysglycemic patients with coronary artery disease–a persistent challenge in need of substantial improvement: a report from ESC EORP EUROASPIRE V. *Diabetes Care* 2020;**43**:726–733. https://doi.org/10.2337/dc19-2165
706. Shahim B, De Bacquer D, De Backer G, Gyberg V, Kotseva K, Mellbin L, *et al.* The prognostic value of fasting plasma glucose, two-hour postload glucose, and HbA(1c) in patients with coronary artery disease: a report from EUROASPIRE IV: a survey from the European Society of Cardiology. *Diabetes Care* 2017;**40**:1233–1240. https://doi.org/10.2337/dc17-0245
707. Ritsinger V, Tanoglidi E, Malmberg K, Näsman P, Rydén L, Tenerz Å, *et al.* Sustained prognostic implications of newly detected glucose abnormalities in patients with acute myocardial infarction: long-term follow-up of the Glucose Tolerance in Patients with Acute Myocardial Infarction cohort. *Diab Vasc Dis Res* 2015;**12**:23–32. https://doi.org/10.1177/1479164114551746
708. Svensson AM, McGuire DK, Abrahamsson P, Dellborg M. Association between hyper- and hypoglycaemia and 2 year all-cause mortality risk in diabetic patients with acute coronary events. *Eur Heart J* 2005;**26**:1255–1261. https://doi.org/10.1093/eurheartj/ehi230
709. Pinto DS, Skolnick AH, Kirtane AJ, Murphy SA, Barron HV, Giugliano RP, *et al.* U-shaped relationship of blood glucose with adverse outcomes among patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2005;**46**:178–180. https://doi.org/10.1016/j.jacc.2005.03.052
710. Cholesterol Treatment Trialists' Collaboration. Efficacy and safety of statin therapy in older people: a meta-analysis of individual participant data from 28 randomised controlled trials. *Lancet* 2019;**393**:407–415. https://doi.org/10.1016/s0140-6736(18)31942-1
711. Bach RG, Cannon CP, Giugliano RP, White JA, Lokhnygina Y, Bohula EA, *et al.* Effect of simvastatin-ezetimibe compared with simvastatin monotherapy after acute coronary syndrome among patients 75 years or older: a secondary analysis of a randomized clinical trial. *JAMA Cardiol* 2019;**4**:846–854. https://doi.org/10.1001/jamacardio.2019.2306
712. Roe MT, Armstrong PW, Fox KA, White HD, Prabhakaran D, Goodman SG, *et al.* Prasugrel versus clopidogrel for acute coronary syndromes without revascularization. *N Engl J Med* 2012;**367**:1297–1309. https://doi.org/10.1056/NEJMoa1205512
713. Chang HM, Okwuosa TM, Scarabelli T, Moudgil R, Yeh ETH. Cardiovascular complications of cancer therapy: best practices in diagnosis, prevention, and management: Part 2. *J Am Coll Cardiol* 2017;**70**:2552–2565. https://doi.org/10.1016/j.jacc.2017.09.1095
714. Herrmann J. Vascular toxic effects of cancer therapies. *Nat Rev Cardiol* 2020;**17**:503–522. https://doi.org/10.1038/s41569-020-0347-2
715. Long M, Ye Z, Zheng J, Chen W, Li L. Dual anti-platelet therapy following percutaneous coronary intervention in a population of patients with thrombocytopenia at baseline: a meta-analysis. *BMC Pharmacol Toxicol* 2020;**21**:31. https://doi.org/10.1186/s40360-020-00409-2
716. Ambrosetti M, Abreu A, Corrà U, Davos CH, Hansen D, Frederix I, *et al.* Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020;**28**:460–495. https://doi.org/10.1177/2047487320913379
717. Abreu A, Frederix I, Dendale P, Janssen A, Doherty P, Piepoli MF, *et al.* Standardization and quality improvement of secondary prevention through cardiovascular rehabilitation programmes in Europe: the avenue towards EAPC accreditation programme: a position statement of the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2020;**28**:496–509. https://doi.org/10.1177/2047487320924912
718. Rossello X, Pocock SJ, Julian DG. Long-term use of cardiovascular drugs: challenges for research and for patient care. *J Am Coll Cardiol* 2015;**66**:1273–1285. https://doi.org/10.1016/j.jacc.2015.07.018
719. Frederix I, Dendale P, Schmid JP. Who needs secondary prevention? *Eur J Prev Cardiol* 2017;**24**:8–13. https://doi.org/10.1177/2047487317706112
720. Rea F, Ronco R, Pedretti RFE, Merlino L, Corrao G. Better adherence with out-of-hospital healthcare improved long-term prognosis of acute coronary syndromes: evidence from an Italian real-world investigation. *Int J Cardiol* 2020;**318**:14–20. https://doi.org/10.1016/j.ijcard.2020.06.017
721. Salzwedel A, Jensen K, Rauch B, Doherty P, Metzendorf MI, Hackbusch M, *et al.* Effectiveness of comprehensive cardiac rehabilitation in coronary artery disease patients treated according to contemporary evidence based medicine: update of the Cardiac Rehabilitation Outcome Study (CROS-II). *Eur J Prev Cardiol* 2020;**27**:1756–1774. https://doi.org/10.1177/2047487320905719
722. Santiago de Araújo Pio C, Marzolini S, Pakosh M, Grace SL. Effect of cardiac rehabilitation dose on mortality and morbidity: a systematic review and meta-regression analysis. *Mayo Clin Proc* 2017;**92**:1644–1659. https://doi.org/10.1016/j.mayocp.2017.07.019
723. van Halewijn G, Deckers J, Tay HY, van Domburg R, Kotseva K, Wood D. Lessons from contemporary trials of cardiovascular prevention and rehabilitation: a systematic review and meta-analysis. *Int J Cardiol* 2017;**232**:294–303. https://doi.org/10.1016/j.ijcard.2016.12.125
724. Dibben G, Faulkner J, Oldridge N, Rees K, Thompson DR, Zwisler AD, *et al.* Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2021;**11**:Cd001800. https://doi.org/10.1002/14651858.CD001800.pub4
725. Anderson L, Thompson DR, Oldridge N, Zwisler AD, Rees K, Martin N, *et al.* Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2016;**2016**:CD001800. https://doi.org/10.1002/14651858.CD001800.pub3
726. Benzer W, Rauch B, Schmid JP, Zwisler AD, Dendale P, Davos CH, *et al.* Exercise-based cardiac rehabilitation in twelve European countries: results of the European cardiac rehabilitation registry. *Int J Cardiol* 2017;**228**:58–67. https://doi.org/10.1016/j.ijcard.2016.11.059
727. Clark RA, Conway A, Poulsen V, Keech W, Tirimacco R, Tideman P. Alternative models of cardiac rehabilitation: a systematic review. *Eur J Prev Cardiol* 2015;**22**:35–74. https://doi.org/10.1177/2047487313501093
728. Kotseva K, De Backer G, De Bacquer D, Rydén L, Hoes A, Grobbee D, *et al.* Lifestyle and impact on cardiovascular risk factor control in coronary patients across 27 countries: results from the European Society of Cardiology ESC-EORP EUROASPIRE V registry. *Eur J Prev Cardiol* 2019;**26**:824–835. https://doi.org/10.1177/2047487318825350
729. Frederix I, Vanhees L, Dendale P, Goetschalckx K. A review of telerehabilitation for cardiac patients. *J Telemed Telecare* 2015;**21**:45–53. https://doi.org/10.1177/1357633x14562732
730. Conraads VM, Deaton C, Piotrowicz E, Santaularia N, Tierney S, Piepoli MF, *et al.* Adherence of heart failure patients to exercise: barriers and possible solutions: a position statement of the Study Group on Exercise Training in Heart Failure of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2012;**14**:451–458. https://doi.org/10.1093/eurjhf/hfs048
731. De Bacquer D, Astin F, Kotseva K, Pogosova N, De Smedt D, De Backer G, *et al.* Poor adherence to lifestyle recommendations in patients with coronary heart disease: results from the EUROASPIRE surveys. *Eur J Prev Cardiol* 2021;**29**:383–395. https://doi.org/10.1093/eurjpc/zwab115
732. Dalal HM, Taylor RS. Telehealth technologies could improve suboptimal rates of participation in cardiac rehabilitation. *Heart* 2016;**102**:1155–1156. https://doi.org/10.1136/heartjnl-2016-309429
733. Lavie CJ, Arena R, Franklin BA. Cardiac rehabilitation and healthy life-style interventions: rectifying program deficiencies to improve patient outcomes. *J Am Coll Cardiol* 2016;**67**:13–15. https://doi.org/10.1016/j.jacc.2015.09.103
734. Fors A, Taft C, Ulin K, Ekman I. Person-centred care improves self-efficacy to control symptoms after acute coronary syndrome: a randomized controlled trial. *Eur J Cardiovasc Nurs* 2016;**15**:186–194. https://doi.org/10.1177/1474515115623437
735. Frederix I, Caiani EG, Dendale P, Anker S, Bax J, Böhm A, *et al.* ESC e-Cardiology Working Group Position Paper: overcoming challenges in digital health implementation in cardiovascular medicine. *Eur J Prev Cardiol* 2019;**26**:1166–1177. https://doi.org/10.1177/2047487319832394
736. Rosselló X, Stanbury M, Beeri R, Kirchhof P, Casadei B, Kotecha D. Digital learning and the future cardiologist. *Eur Heart J* 2019;**40**:499–501. https://doi.org/10.1093/eurheartj/ehy884
737. Frederix I, Solmi F, Piepoli MF, Dendale P. Cardiac telerehabilitation: a novel cost-efficient care delivery strategy that can induce long-term health benefits. *Eur J Prev Cardiol* 2017;**24**:1708–1717. https://doi.org/10.1177/2047487317732274
738. Avila A, Claes J, Buys R, Azzawi M, Vanhees L, Cornelissen V. Home-based exercise with telemonitoring guidance in patients with coronary artery disease: does it improve long-term physical fitness? *Eur J Prev Cardiol* 2020;**27**:367–377. https://doi.org/10.1177/2047487319892201
739. Claes J, Cornelissen V, McDermott C, Moyna N, Pattyn N, Cornelis N, *et al.* Feasibility, acceptability, and clinical effectiveness of a technology-enabled cardiac rehabilitation platform (physical activity toward health-I): randomized controlled trial. *J Med Internet Res* 2020;**22**:e14221. https://doi.org/10.2196/14221
740. Kraal JJ, Peek N, Van den Akker-Van Marle ME, Kemps HM. Effects of home-based training with telemonitoring guidance in low to moderate risk patients entering cardiac rehabilitation: short-term results of the FIT@Home study. *Eur J Prev Cardiol* 2014;**21**:26–31. https://doi.org/10.1177/2047487314552606
741. Maddison R, Rawstorn JC, Stewart RAH, Benatar J, Whittaker R, Rolleston A, *et al.* Effects and costs of real-time cardiac telerehabilitation: randomised controlled non-inferiority trial. *Heart* 2019;**105**:122–129. https://doi.org/10.1136/heartjnl-2018-313189

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

742. Scherrenberg M, Wilhelm M, Hansen D, Völler H, Cornelissen V, Frederix I, *et al.* The future is now: a call for action for cardiac telerehabilitation in the COVID-19 pandemic from the secondary prevention and rehabilitation section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020;**28**:524–540. https://doi.org/10.1177/2047487320939671
743. Huang K, Liu W, He D, Huang B, Xiao D, Peng Y, *et al.* Telehealth interventions versus center-based cardiac rehabilitation of coronary artery disease: a systematic review and meta-analysis. *Eur J Prev Cardiol* 2015;**22**:959–971. https://doi.org/10.1177/2047487314561168
744. Wolf A, Vella R, Fors A. The impact of person-centred care on patients' care experiences in relation to educational level after acute coronary syndrome: secondary outcome analysis of a randomised controlled trial. *Eur J Cardiovasc Nurs* 2019;**18**:299–308. https://doi.org/10.1177/1474515118821242
745. Brown MT, Bussell J, Dutta S, Davis K, Strong S, Mathew S. Medication adherence: truth and consequences. *Am J Med Sci* 2016;**351**:387–399. https://doi.org/10.1016/j.amjms.2016.01.010
746. Arlt AD, Nestoriuc Y, Rief W. Why current drug adherence programs fail: addressing psychological risk factors of nonadherence. *Curr Opin Psychiatry* 2017;**30**:326–333. https://doi.org/10.1097/yco.0000000000000345
747. Easthall C, Taylor N, Bhattacharya D. Barriers to medication adherence in patients prescribed medicines for the prevention of cardiovascular disease: a conceptual framework. *Int J Pharm Pract* 2019;**27**:223–231. https://doi.org/10.1111/ijpp.12491
748. Seabury SA, Dougherty JS, Sullivan J. Medication adherence as a measure of the quality of care provided by physicians. *Am J Manag Care* 2019;**25**:78–83.
749. Pedretti RFE, Hansen D, Ambrosetti M, Back M, Berger T, Ferreira MC, *et al.* How to optimize the adherence to a guideline-directed medical therapy in the secondary prevention of cardiovascular diseases: a clinical consensus statement from the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2022;**30**:149–166. https://doi.org/10.1093/eurjpc/zwac204
750. Castellano JM, Sanz G, Peñalvo JL, Bansilal S, Fernández-Ortiz A, Alvarez L, *et al.* A polypill strategy to improve adherence: results from the FOCUS project. *J Am Coll Cardiol* 2014;**64**:2071–2082. https://doi.org/10.1016/j.jacc.2014.08.021
751. Selak V, Webster R, Stepien S, Bullen C, Patel A, Thom S, *et al.* Reaching cardiovascular prevention guideline targets with a polypill-based approach: a meta-analysis of randomised clinical trials. *Heart* 2019;**105**:42–48. https://doi.org/10.1136/heartjnl-2018-313108
752. Castellano JM, Fuster V, Jennings C, Prescott E, Bueno H. Role of the polypill for secondary prevention in ischaemic heart disease. *Eur J Prev Cardiol* 2017;**24**:44–51. https://doi.org/10.1177/2047487317707324
753. Castellano JM, Pocock SJ, Bhatt DL, Quesada AJ, Owen R, Fernandez-Ortiz A, *et al.* Polypill strategy in secondary cardiovascular prevention. *N Engl J Med* 2022;**387**:967–977. https://doi.org/10.1056/NEJMoa2208275
754. Palmer MJ, Barnard S, Perel P, Free C. Mobile phone-based interventions for improving adherence to medication prescribed for the primary prevention of cardiovascular disease in adults. *Cochrane Database Syst Rev* 2018;**6**:CD012675. https://doi.org/10.1002/14651858.CD012675.pub2
755. Guerriero C, Cairns J, Roberts I, Rodgers A, Whittaker R, Free C. The cost-effectiveness of smoking cessation support delivered by mobile phone text messaging: Txt2stop. *Eur J Health Econ* 2013;**14**:789–797. https://doi.org/10.1007/s10198-012-0424-5
756. Gandapur Y, Kianoush S, Kelli HM, Misra S, Urrea B, Blaha MJ, *et al.* The role of mHealth for improving medication adherence in patients with cardiovascular disease: a systematic review. *Eur Heart J Qual Care Clin Outcomes* 2016;**2**:237–244. https://doi.org/10.1093/ehjqcco/qcw018
757. Fortuna RJ, Nagel AK, Rocco TA, Legette-Sobers S, Quigley DD. Patient experience with care and its association with adherence to hypertension medications. *Am J Hypertens* 2018;**31**:340–345. https://doi.org/10.1093/ajh/hpx200
758. Keenan J. Improving adherence to medication for secondary cardiovascular disease prevention. *Eur J Prev Cardiol* 2017;**24**:29–35. https://doi.org/10.1177/2047487317708145
759. Geidl W, Schlesinger S, Mino E, Miranda L, Pfeifer K. Dose-response relationship between physical activity and mortality in adults with noncommunicable diseases: a systematic review and meta-analysis of prospective observational studies. *Int J Behav Nutr Phys Act* 2020;**17**:109. https://doi.org/10.1186/s12966-020-01007-5
760. Ekblom O, Ek A, Cider Å, Hambraeus K, Börjesson M. Increased physical activity post-myocardial infarction is related to reduced mortality: results from the SWEDEHEART registry. *J Am Heart Assoc* 2018;**7**:e010108. https://doi.org/10.1161/jaha.118.010108
761. Delgado-Lista J, Alcala-Diaz JF, Torres-Peña JD, Quintana-Navarro GM, Fuentes F, Garcia-Rios A, *et al.* Long-term secondary prevention of cardiovascular disease with a Mediterranean diet and a low-fat diet (CORDIOPREV): a randomised controlled trial. *Lancet* 2022;**399**:1876–1885. https://doi.org/10.1016/s0140-6736(22)00122-2
762. de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999;**99**:779–785. https://doi.org/10.1161/01.cir.99.6.779
763. Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: a systematic review. *JAMA* 2003;**290**:86–97. https://doi.org/10.1001/jama.290.1.86
764. Chow CK, Jolly S, Rao-Melacini P, Fox KAA, Anand SS, Yusuf S. Association of diet, exercise, and smoking modification with risk of early cardiovascular events after acute coronary syndromes. *Circulation* 2010;**121**:750–758. https://doi.org/10.1161/circulationaha.109.891523
765. United States Public Health Service Office of the Surgeon General. Smoking Cessation: A Report of the Surgeon General. In: US Department of Health and Human Services; 2020.
766. Levine GN, Bates ER, Bittl JA, Brindis RG, Fihn SD, Fleisher LA, *et al.* 2016 ACC/AHA Guideline Focused Update on Duration of Dual Antiplatelet Therapy in Patients With Coronary Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines: an Update of the 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention, 2011 ACCF/AHA Guideline for Coronary Artery Bypass Graft Surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease, 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction, 2014 AHA/ACC Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes, and 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. *Circulation* 2016;**134**:e123–e155. https://doi.org/10.1161/cir.0000000000000404
767. Cahill K, Stevens S, Perera R, Lancaster T. Pharmacological interventions for smoking cessation: an overview and network meta-analysis. *Cochrane Database Syst Rev* 2013;**2013**:CD009329. https://doi.org/10.1002/14651858.CD009329.pub2
768. Eisenberg MJ, Windle SB, Roy N, Old W, Grondin FR, Bata I, *et al.* Varenicline for smoking cessation in hospitalized patients with acute coronary syndrome. *Circulation* 2016;**133**:21–30. https://doi.org/10.1161/circulationaha.115.019634
769. Sterling LH, Windle SB, Filion KB, Touma L, Eisenberg MJ. Varenicline and adverse cardiovascular events: a systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc* 2016;**5**:e002849. https://doi.org/10.1161/jaha.115.002849
770. Windle SB, Dehghani P, Roy N, Old W, Grondin FR, Bata I, *et al.* Smoking abstinence 1 year after acute coronary syndrome: follow-up from a randomized controlled trial of varenicline in patients admitted to hospital. *CMAJ* 2018;**190**:E347–E354. https://doi.org/10.1503/cmaj.170377
771. Kavousi M, Pisinger C, Barthelemy JC, De Smedt D, Koskinas K, Marques-Vidal P, *et al.* Electronic cigarettes and health with special focus on cardiovascular effects: position paper of the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2020;**28**:1552–1566. https://doi.org/10.1177/2047487320941993
772. Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, *et al.* Primary prevention of cardiovascular disease with a mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med* 2018;**378**:e34. https://doi.org/10.1056/NEJMoa1800389
773. Wood AM, Kaptoge S, Butterworth AS, Willeit P, Warnakula S, Bolton T, *et al.* Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599 912 current drinkers in 83 prospective studies. *Lancet* 2018;**391**:1513–1523. https://doi.org/10.1016/s0140-6736(18)30134-x
774. Holmes MV, Dale CE, Zuccolo L, Silverwood RJ, Guo Y, Ye Z, *et al.* Association between alcohol and cardiovascular disease: Mendelian randomisation analysis based on individual participant data. *BMJ* 2014;**349**:g4164. https://doi.org/10.1136/bmj.g4164
775. Millwood IY, Walters RG, Mei XW, Guo Y, Yang L, Bian Z, *et al.* Conventional and genetic evidence on alcohol and vascular disease aetiology: a prospective study of 500 000 men and women in China. *Lancet* 2019;**393**:1831–1842. https://doi.org/10.1016/s0140-6736(18)31772-0
776. Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, *et al.* Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* 2019;**366**:l4570. https://doi.org/10.1136/bmj.l4570
777. Patterson R, McNamara E, Tainio M, de Sá TH, Smith AD, Sharp SJ, *et al.* Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. *Eur J Epidemiol* 2018;**33**:811–829. https://doi.org/10.1007/s10654-018-0380-1
778. WHO Guidelines Review Committee. *WHO Guidelines on Physical Activity and Sedentary Behaviour*. World Health Organization, 2020.
779. Hansen D, Abreu A, Ambrosetti M, Cornelissen V, Gevaert A, Kemps H, *et al.* Exercise intensity assessment and prescription in cardiovascular rehabilitation and beyond: why and how: a position statement from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2022;**29**:230–245. https://doi.org/10.1093/eurjpc/zwab007
780. De Schutter A, Kachur S, Lavie CJ, Menezes A, Shum KK, Bangalore S, *et al.* Cardiac rehabilitation fitness changes and subsequent survival. *Eur Heart J Qual Care Clin Outcomes* 2018;**4**:173–179. https://doi.org/10.1093/ehjqcco/qcy018
781. Baumeister H, Hutter N, Bengel J. Psychological and pharmacological interventions for depression in patients with coronary artery disease. *Cochrane Database Syst Rev* 2011;**2011**:Cd008012. https://doi.org/10.1002/14651858.CD008012.pub3

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

782. Richards SH, Anderson L, Jenkinson CE, Whalley B, Rees K, Davies P, *et al.* Psychological interventions for coronary heart disease: cochrane systematic review and meta-analysis. *Eur J Prev Cardiol* 2018;**25**:247–259. https://doi.org/10.1177/2047487317739978
783. Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, *et al.* 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020;**41**:111–188. https://doi.org/10.1093/eurheartj/ehz455
784. Ference BA, Ginsberg HN, Graham I, Ray KK, Packard CJ, Bruckert E, *et al.* Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur Heart J* 2017;**38**:2459–2472. https://doi.org/10.1093/eurheartj/ehx144
785. Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, *et al.* Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017;**376**:1713–1722. https://doi.org/10.1056/NEJMoa1615664
786. Schwartz GG, Steg PG, Szarek M, Bhatt DL, Bittner VA, Diaz R, *et al.* Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med* 2018;**379**:2097–2107. https://doi.org/10.1056/NEJMoa1801174
787. Navarese EP, Kowalewski M, Andreotti F, van Wely M, Camaro C, Kolodziejczak M, *et al.* Meta-analysis of time-related benefits of statin therapy in patients with acute coronary syndrome undergoing percutaneous coronary intervention. *Am J Cardiol* 2014;**113**:1753–1764. https://doi.org/10.1016/j.amjcard.2014.02.034
788. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, *et al.* Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;**372**:2387–2397. https://doi.org/10.1056/NEJMoa1410489
789. Koskinas KC, Windecker S, Pedrazzini G, Mueller C, Cook S, Matter CM, *et al.* Evolocumab for early reduction of LDL cholesterol levels in patients with acute coronary syndromes (EVOPACS). *J Am Coll Cardiol* 2019;**74**:2452–2462. https://doi.org/10.1016/j.jacc.2019.08.010
790. Trankle CR, Wohlford G, Buckley LF, Kadariya D, Ravindra K, Markley R, *et al.* Alirocumab in acute myocardial infarction: results from the Virginia Commonwealth University Alirocumab Response Trial (VCU-AlirocRT). *J Cardiovasc Pharmacol* 2019;**74**:266–269. https://doi.org/10.1097/fjc.0000000000000706
791. Iannuzzo G, Gentile M, Bresciani A, Mallardo V, Di Lorenzo A, Merone P, *et al.* Inhibitors of protein convertase subtilisin/kexin 9 (PCSK9) and acute coronary syndrome (ACS): the state-of-the-art. *J Clin Med* 2021;**10**:1510. https://doi.org/10.3390/jcm10071510
792. Räber L, Ueki Y, Otsuka T, Losdat S, Häner JD, Lonborg J, *et al.* Effect of alirocumab added to high-intensity statin therapy on coronary atherosclerosis in patients with acute myocardial infarction: the PACMAN-AMI randomized clinical trial. *JAMA* 2022;**327**:1771–1781. https://doi.org/10.1001/jama.2022.5218
793. Nicholls SJ, Kataoka Y, Nissen SE, Prati F, Windecker S, Puri R, *et al.* Effect of evolocumab on coronary plaque phenotype and burden in statin-treated patients following myocardial infarction. *JACC Cardiovasc Imaging* 2022;**15**:1308–1321. https://doi.org/10.1016/j.jcmg.2022.03.002
794. Schubert J, Lindahl B, Melhus H, Renlund H, Leosdottir M, Yari A, *et al.* Low-density lipoprotein cholesterol reduction and statin intensity in myocardial infarction patients and major adverse outcomes: a Swedish nationwide cohort study. *Eur Heart J* 2021;**42**:243–252. https://doi.org/10.1093/eurheartj/ehaa1011
795. Gencer B, Mach F, Murphy SA, De Ferrari GM, Huber K, Lewis BS, *et al.* Efficacy of evolocumab on cardiovascular outcomes in patients with recent myocardial infarction: a prespecified secondary analysis from the FOURIER trial. *JAMA Cardiol* 2020;**5**:952–957. https://doi.org/10.1001/jamacardio.2020.0882
796. O'Donoghue ML, Giugliano RP, Wiviott SD, Atar D, Keech A, Kuder JF, *et al.* Long-term evolocumab in patients with established atherosclerotic cardiovascular disease. *Circulation* 2022;**146**:1109–1119. https://doi.org/10.1161/circulationaha.122.061620
797. Bhatt DL, Steg PG, Miller M, Brinton EA, Jacobson TA, Ketchum SB, *et al.* Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med* 2019;**380**:11–22. https://doi.org/10.1056/NEJMoa1812792
798. Freemantle N, Cleland J, Young P, Mason J, Harrison J. β blockade after myocardial infarction: systematic review and meta regression analysis. *BMJ* 1999;**318**:1730–1737. https://doi.org/10.1136/bmj.318.7200.1730
799. Martínez-Milla J, Raposeiras-Roubín S, Pascual-Figal DA, Ibáñez B. Role of beta-blockers in cardiovascular disease in 2019. *Rev Esp Cardiol (Engl Ed)* 2019;**72**:844–852. https://doi.org/10.1016/j.rec.2019.04.014
800. Dahl Aarvik M, Sandven I, Dondo TB, Gale CP, Ruddox V, Munkhaugen J, *et al.* Effect of oral β-blocker treatment on mortality in contemporary post-myocardial infarction patients: a systematic review and meta-analysis. *Eur Heart J Cardiovasc Pharmacother* 2019;**5**:12–20. https://doi.org/10.1093/ehjcvp/pvy034
801. Dargie HJ. Effect of carvedilol on outcome after myocardial infarction in patients with left-ventricular dysfunction: the CAPRICORN randomised trial. *Lancet* 2001;**357**:1385–1390. https://doi.org/10.1016/s0140-6736(00)04560-8
802. Kim J, Kang D, Park H, Kang M, Park TK, Lee JM, *et al.* Long-term β-blocker therapy and clinical outcomes after acute myocardial infarction in patients without heart failure: nationwide cohort study. *Eur Heart J* 2020;**41**:3521–3529. https://doi.org/10.1093/eurheartj/ehaa376
803. Raposeiras-Roubín S, Abu-Assi E, Redondo-Diéguez A, González-Ferreiro R, López-López A, Bouzas-Cruz N, *et al.* Prognostic benefit of beta-blockers after acute coronary syndrome with preserved systolic function. Still relevant today? *Rev Esp Cardiol (Engl Ed)* 2015;**68**:585–591. https://doi.org/10.1016/j.rec.2014.07.028
804. Dondo TB, Hall M, West RM, Jernberg T, Lindahl B, Bueno H, *et al.* β-blockers and mortality after acute myocardial infarction in patients without heart failure or ventricular dysfunction. *J Am Coll Cardiol* 2017;**69**:2710–2720. https://doi.org/10.1016/j.jacc.2017.03.578
805. Watanabe H, Ozasa N, Morimoto T, Shiomi H, Bingyuan B, Suwa S, *et al.* Long-term use of carvedilol in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. *PLoS One* 2018;**13**:e0199347. https://doi.org/10.1371/journal.pone.0199347
806. Rossello X, Raposeiras-Roubin S, Latini R, Dominguez-Rodriguez A, Barrabés JA, Sánchez PL, *et al.* Rationale and design of the pragmatic clinical trial tREatment with Beta-blockers after myOcardial infarction withOut reduced ejection fracTion (REBOOT). *Eur Heart J Cardiovasc Pharmacother* 2021;**8**:291–301. https://doi.org/10.1093/ehjcvp/pvab060
807. Munkhaugen J, Ruddox V, Halvorsen S, Dammen T, Fagerland MW, Hernæs KH, *et al.* BEtablocker Treatment After acute Myocardial Infarction in revascularized patients without reduced left ventricular ejection fraction (BETAMI): rationale and design of a prospective, randomized, open, blinded end point study. *Am Heart J* 2019;**208**:37–46. https://doi.org/10.1016/j.ahj.2018.10.005
808. Kristensen AMD, Bovin A, Zwisler AD, Cerquira C, Torp-Pedersen C, Bøtker HE, *et al.* Design and rationale of the Danish trial of beta-blocker treatment after myocardial infarction without reduced ejection fraction: study protocol for a randomized controlled trial. *Trials* 2020;**21**:415. https://doi.org/10.1186/s13063-020-4214-6
809. Puymirat E, Riant E, Aissaoui N, Soria A, Ducrocq G, Coste P, *et al.* β blockers and mortality after myocardial infarction in patients without heart failure: multicentre prospective cohort study. *BMJ* 2016;**354**:i4801. https://doi.org/10.1136/bmj.i4801
810. Zeitouni M, Kerneis M, Lattuca B, Guedeney P, Cayla G, Collet JP, *et al.* Do patients need lifelong β-blockers after an uncomplicated myocardial infarction? *Am J Cardiovasc Drugs* 2019;**19**:431–438. https://doi.org/10.1007/s40256-019-00338-4
811. ISIS-4 Collaborative Group. ISIS-4: a randomised factorial trial assessing early oral captopril, oral mononitrate, and intravenous magnesium sulphate in 58,050 patients with suspected acute myocardial infarction. *Lancet* 1995;**345**:669–685. https://doi.org/10.1016/S0140-6736(95)90865-X
812. Held PH, Yusuf S, Furberg CD. Calcium channel blockers in acute myocardial infarction and unstable angina: an overview. *BMJ* 1989;**299**:1187–1192. https://doi.org/10.1136/bmj.299.6709.1187
813. Køber L, Torp-Pedersen C, Carlsen JE, Bagger H, Eliasen P, Lyngborg K, *et al.* A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 1995;**333**:1670–1676. https://doi.org/10.1056/nejm199512213332503
814. Pfeffer MA, Braunwald E, Moyé LA, Basta L, Brown EJ, Cuddy TE, *et al.* Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction–results of the survival and ventricular enlargement trial. *N Engl J Med* 1992;**327**:669–677. https://doi.org/10.1056/nejm199209033271001
815. The AIRE Study Investigators. Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet* 1993;**342**:821–828. https://doi.org/10.1016/0140-6736(93)92693-N
816. Fox KM. Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomised, double-blind, placebo-controlled, multicentre trial (the EUROPA study). *Lancet* 2003;**362**:782–788. https://doi.org/10.1016/s0140-6736(03)14286-9
817. The Heart Outcomes Prevention Evaluation Study Investigators; Yusuf S, Sleight P, Pogue Bosch J, Davies R, *et al.* Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;**342**:145–153. https://doi.org/10.1056/nejm200001203420301
818. ACE Inhibitor Myocardial Infarction Collaborative Group. Indications for ACE inhibitors in the early treatment of acute myocardial infarction: systematic overview of individual data from 100,000 patients in randomized trials. *Circulation* 1998;**97**:2202–2212. https://doi.org/10.1161/01.cir.97.22.2202
819. Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Køber L, Maggioni AP, *et al.* Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;**349**:1893–1906. https://doi.org/10.1056/NEJMoa032292
820. Swedberg K, Eneroth P, Kjekshus J, Snapinn S. Effects of enalapril and neuroendocrine activation on prognosis in severe congestive heart failure (follow-up of the CONSENSUS trial). *Am J Cardiol* 1990;**66**:D40–D45. https://doi.org/10.1016/0002-9149(90)90475-g
821. Garg R, Yusuf S. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. *JAMA* 1995;**273**:1450–1456. https://doi.org/10.1001/jama.1995.03520420066040

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

822. Packer M, Poole-Wilson PA, Armstrong PW, Cleland JGF, Horowitz JD, Massie BM, *et al.* Comparative effects of low and high doses of the angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. *Circulation* 1999; **100**:2312–2318. https://doi.org/10.1161/01.cir.100.23.2312
823. Yusuf S, Pitt B, Davis CE, Hood WB, Cohn JN. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991;**325**:293–302. https://doi.org/10.1056/nejm199108013250501
824. McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, *et al.* Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014; **371**:993–1004. https://doi.org/10.1056/NEJMoa1409077
825. Pfeffer MA, Claggett B, Lewis EF, Granger CB, Køber L, Maggioni AP, *et al.* Angiotensin receptor-neprilysin inhibition in acute myocardial infarction. *N Engl J Med* 2021;**385**: 1845–1855. https://doi.org/10.1056/NEJMoa2104508
826. Pitt B, Remme W, Zannad F, Neaton J, Martinez F, Roniker B, *et al.* Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003;**348**:1309–1321. https://doi.org/10.1056/NEJMoa030207
827. Montalescot G, Pitt B, Lopez de Sa E, Hamm CW, Flather M, Verheugt F, *et al.* Early eplerenone treatment in patients with acute ST-elevation myocardial infarction without heart failure: the Randomized Double-Blind Reminder Study. *Eur Heart J* 2014;**35**: 2295–2302. https://doi.org/10.1093/eurheartj/ehu164
828. Cowie MR, Fisher M. SGLT2 inhibitors: mechanisms of cardiovascular benefit beyond glycaemic control. *Nat Rev Cardiol* 2020;**17**:761–772. https://doi.org/10.1038/s41569-020-0406-8
829. Neal B, Perkovic V, Mahaffey KW, de Zeeuw D, Fulcher G, Erondu N, *et al.* Canagliflozin and cardiovascular and renal events in type 2 diabetes. *N Engl J Med* 2017;**377**:644–657. https://doi.org/10.1056/NEJMoa1611925
830. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, *et al.* Dapagliflozin and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2019;**380**:347–357. https://doi.org/10.1056/NEJMoa1812389
831. Nikolaus M, Massimo F, Katharina S, Dirk M-W, Ramzi AA, Manuel JA, *et al.* 2023 ESC Guidelines for the management of cardiovascular disease in patients with diabetes. *Eur Heart J* 2023; https://doi.org/10.1093/eurheartj/ehad192
832. Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, *et al.* Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;**385**:1451–1461. https://doi.org/10.1056/NEJMoa2107038
833. Solomon SD, McMurray JJV, Claggett B, de Boer RA, DeMets D, Hernandez AF, *et al.* Dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2022;**387**:1089–1098. https://doi.org/10.1056/NEJMoa2206286
834. von Lewinski D, Kolesnik E, Tripolt NJ, Pferschy PN, Benedikt M, Wallner M, *et al.* Empagliflozin in acute myocardial infarction: the EMMY trial. *Eur Heart J* 2022;**43**: 4421–4432. https://doi.org/10.1093/eurheartj/ehac494
835. Harrington J, Udell JA, Jones WS, Anker SD, Bhatt DL, Petrie MC, *et al.* Empagliflozin in patients post myocardial infarction rationale and design of the EMPACT-MI trial. *Am Heart J* 2022;**253**:86–98. https://doi.org/10.1016/j.ahj.2022.05.010
836. Lai KC, Lam SK, Chu KM, Wong BCY, Hui WM, Hu WHC, *et al.* Lansoprazole for the prevention of recurrences of ulcer complications from long-term low-dose aspirin use. *N Engl J Med* 2002;**346**:2033–2038. https://doi.org/10.1056/NEJMoa012877
837. Casado Arroyo R, Polo-Tomas M, Roncalés MP, Scheiman J, Lanas A. Lower GI bleeding is more common than upper among patients on dual antiplatelet therapy: long-term follow-up of a cohort of patients commonly using PPI co-therapy. *Heart* 2012; **98**:718–723. https://doi.org/10.1136/heartjnl-2012-301632
838. Small DS, Farid NA, Payne CD, Weerakkody GJ, Li YG, Brandt JT, *et al.* Effects of the proton pump inhibitor lansoprazole on the pharmacokinetics and pharmacodynamics of prasugrel and clopidogrel. *J Clin Pharmacol* 2008;**48**:475–484. https://doi.org/10.1177/0091270008315310
839. Sibbing D, Morath T, Stegherr J, Braun S, Vogt W, Hadamitzky M, *et al.* Impact of proton pump inhibitors on the antiplatelet effects of clopidogrel. *Thromb Haemost* 2009; **101**:714–719. https://doi.org/10.1160/TH08-12-0808
840. Gilard M, Arnaud B, Cornily JC, Le Gal G, Lacut K, Le Calvez G, *et al.* Influence of omeprazole on the antiplatelet action of clopidogrel associated with aspirin: the randomized, double-blind OCLA (Omeprazole CLopidogrel Aspirin) study. *J Am Coll Cardiol* 2008;**51**:256–260. https://doi.org/10.1016/j.jacc.2007.06.064
841. O'Donoghue ML, Braunwald E, Antman EM, Murphy SA, Bates ER, Rozenman Y, *et al.* Pharmacodynamic effect and clinical efficacy of clopidogrel and prasugrel with or without a proton-pump inhibitor: an analysis of two randomised trials. *Lancet* 2009;**374**: 989–997. https://doi.org/10.1016/s0140-6736(09)61525-7
842. Goodman SG, Clare R, Pieper KS, Nicolau JC, Storey RF, Cantor WJ, *et al.* Association of proton pump inhibitor use on cardiovascular outcomes with clopidogrel and ticagrelor: insights from the platelet inhibition and patient outcomes trial. *Circulation* 2012;**125**:978–986. https://doi.org/10.1161/circulationaha.111.032912
843. Yedlapati SH, Khan SU, Talluri S, Lone AN, Khan MZ, Khan MS, *et al.* Effects of influenza vaccine on mortality and cardiovascular outcomes in patients with cardiovascular disease: a systematic review and meta-analysis. *J Am Heart Assoc* 2021;**10**:e019636. https://doi.org/10.1161/jaha.120.019636
844. Liprandi ÁS, Liprandi MIS, Zaidel EJ, Aisenberg GM, Baranchuk A, Barbosa ECD, *et al.* Influenza vaccination for the prevention of cardiovascular disease in the Americas: consensus document of the Inter-American Society of Cardiology and the World Heart Federation. *Glob Heart* 2021;**16**:55. https://doi.org/10.5334/gh.1069
845. Gurfinkel EP, Leon de la Fuente R, Mendiz O, Mautner B. Flu vaccination in acute coronary syndromes and planned percutaneous coronary interventions (FLUVACS) study. *Eur Heart J* 2004;**25**:25–31. https://doi.org/10.1016/j.ehj.2003.10.018
846. Phrommintikul A, Kuanprasert S, Wongcharoen W, Kanjanavanit R, Chaiwarith R, Sukonthasarn A, *et al.* Influenza vaccination reduces cardiovascular events in patients with acute coronary syndrome. *Eur Heart J* 2011;**32**:1730–1735. https://doi.org/10.1093/eurheartj/ehr004
847. Fröbert O, Götberg M, Erlinge D, Akhtar Z, Christiansen EH, MacIntyre CR, *et al.* Influenza vaccination after myocardial infarction: a randomized, double-blind, placebo-controlled, multicenter trial. *Circulation* 2021;**144**:1476–1484. https://doi.org/10.1161/circulationaha.121.057042
848. Chen Y, Zhang H, Chen Y, Li M, Luo W, Liu Y, *et al.* Colchicine may become a new cornerstone therapy for coronary artery disease: a meta-analysis of randomized controlled trials. *Clin Rheumatol* 2022;**41**:1873–1887. https://doi.org/10.1007/s10067-022-06050-0
849. Razavi E, Ramezani A, Kazemi A, Attar A. Effect of treatment with colchicine after acute coronary syndrome on major cardiovascular events: a systematic review and meta-analysis of clinical trials. *Cardiovasc Ther* 2022;**2022**:8317011. https://doi.org/10.1155/2022/8317011
850. Tardif JC, Kouz S, Waters DD, Bertrand OF, Diaz R, Maggioni AP, *et al.* Efficacy and safety of low-dose colchicine after myocardial infarction. *N Engl J Med* 2019;**381**: 2497–2505. https://doi.org/10.1056/NEJMoa1912388
851. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, *et al.* Colchicine in patients with chronic coronary disease. *N Engl J Med* 2020;**383**:1838–1847. https://doi.org/10.1056/NEJMoa2021372
852. Opstal TSJ, Fiolet ATL, van Broekhoven A, Mosterd A, Eikelboom JW, Nidorf SM, *et al.* Colchicine in patients with chronic coronary disease in relation to prior acute coronary syndrome. *J Am Coll Cardiol* 2021;**78**:859–866. https://doi.org/10.1016/j.jacc.2021.06.037
853. Ji H, Fang L, Yuan L, Zhang Q. Effects of exercise-based cardiac rehabilitation in patients with acute coronary syndrome: a meta-analysis. *Med Sci Monit* 2019;**25**: 5015–5027. https://doi.org/10.12659/msm.917362
854. Candelaria D, Randall S, Ladak L, Gallagher R. Health-related quality of life and exercise-based cardiac rehabilitation in contemporary acute coronary syndrome patients: a systematic review and meta-analysis. *Qual Life Res* 2020;**29**:579–592. https://doi.org/10.1007/s11136-019-02338-y
855. Anthonisen NR, Skeans MA, Wise RA, Manfreda J, Kanner RE, Connett JE, *et al.* The effects of a smoking cessation intervention on 14.5-year mortality: a randomized clinical trial. *Ann Intern Med* 2005;**142**:233–239. https://doi.org/10.7326/0003-4819-142-4-200502150-00005
856. Becerra-Tomás N, Blanco Mejía S, Viguiliouk E, Khan T, Kendall CWC, Kahleova H, *et al.* Mediterranean diet, cardiovascular disease and mortality in diabetes: a systematic review and meta-analysis of prospective cohort studies and randomized clinical trials. *Crit Rev Food Sci Nutr* 2020;**60**:1207–1227. https://doi.org/10.1080/10408398.2019.1565281
857. Liu Y, Lee DC, Li Y, Zhu W, Zhang R, Sui X, *et al.* Associations of resistance exercise with cardiovascular disease morbidity and mortality. *Med Sci Sports Exerc* 2019;**51**: 499–508. https://doi.org/10.1249/mss.0000000000001822
858. Saeidifard F, Medina-Inojosa JR, West CP, Olson TP, Somers VK, Bonikowske AR, *et al.* The association of resistance training with mortality: a systematic review and meta-analysis. *Eur J Prev Cardiol* 2019;**26**:1647–1665. https://doi.org/10.1177/2047487319850718
859. Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T. Nicotine replacement therapy versus control for smoking cessation. *Cochrane Database Syst Rev* 2018;**5**: CD000146. https://doi.org/10.1002/14651858.CD000146.pub5
860. Howes S, Hartmann-Boyce J, Livingstone-Banks J, Hong B, Lindson N. Antidepressants for smoking cessation. *Cochrane Database Syst Rev* 2020;**4**:CD000031. https://doi.org/10.1002/14651858.CD000031.pub5
861. Cahill K, Lindson-Hawley N, Thomas KH, Fanshawe TR, Lancaster T. Nicotine receptor partial agonists for smoking cessation. *Cochrane Database Syst Rev* 2016;**2016**: CD006103. https://doi.org/10.1002/14651858.CD006103.pub7
862. Lindson N, Chepkin SC, Ye W, Fanshawe TR, Bullen C, Hartmann-Boyce J. Different doses, durations and modes of delivery of nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev* 2019;**4**:CD013308. https://doi.org/10.1002/14651858.Cd013308
863. Woolf KJ, Zabad MN, Post JM, McNitt S, Williams GC, Bisognano JD. Effect of nicotine replacement therapy on cardiovascular outcomes after acute coronary syndromes. *Am J Cardiol* 2012;**110**:968–970. https://doi.org/10.1016/j.amjcard.2012.05.028
864. Suissa K, Larivière J, Eisenberg MJ, Eberg M, Gore GC, Grad R, *et al.* Efficacy and safety of smoking cessation interventions in patients with cardiovascular disease: a network meta-analysis of randomized controlled trials. *Circ Cardiovasc Qual Outcomes* 2017;**10**: e002458. https://doi.org/10.1161/circoutcomes.115.002458

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

865. de Lemos JA, Blazing MA, Wiviott SD, Lewis EF, Fox KAA, White HD, *et al.* Early intensive vs a delayed conservative simvastatin strategy in patients with acute coronary syndromes: phase Z of the A to Z trial. *JAMA* 2004;**292**:1307–1316. https://doi.org/10.1001/jama.292.11.1307
866. Ray KK, Cannon CP, McCabe CH, Cairns R, Tonkin AM, Sacks FM, *et al.* Early and late benefits of high-dose atorvastatin in patients with acute coronary syndromes: results from the PROVE IT-TIMI 22 trial. *J Am Coll Cardiol* 2005;**46**:1405–1410. https://doi.org/10.1016/j.jacc.2005.03.077
867. Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF, Waters D, *et al.* Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *JAMA* 2001;**285**:1711–1718. https://doi.org/10.1001/jama.285.13.1711
868. Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhala N, *et al.* Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010;**376**:1670–1681. https://doi.org/10.1016/s0140-6736(10)61350-5
869. Fulcher J, O'Connell R, Voysey M, Emberson J, Blackwell L, Mihaylova B. Efficacy and safety of LDL-lowering therapy among men and women: meta-analysis of individual data from 174,000 participants in 27 randomised trials. *Lancet* 2015;**385**:1397–1405. https://doi.org/10.1016/s0140-6736(14)61368-4
870. CIBIS-II Investigators. The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. *Lancet* 1999;**353**:9–13. https://doi.org/10.1016/S0140-6736(98)11181-9
871. Packer M, Coats AJ, Fowler MB, Katus HA, Krum H, Mohacsi P, *et al.* Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med* 2001;**344**:1651–1658. https://doi.org/10.1056/nejm200105313442201
872. MERIT-HF Study Group. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* 1999;**353**:2001–2007. https://doi.org/10.1016/S0140-6736(99)04440-2
873. Chen ZM, Pan HC, Chen YP, Peto R, Collins R, Jiang LX, *et al.* Early intravenous then oral metoprolol in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. *Lancet* 2005;**366**:1622–1632. https://doi.org/10.1016/s0140-6736(05)67661-1
874. Bangalore S, Makani H, Radford M, Thakur K, Toklu B, Katz SD, *et al.* Clinical outcomes with β-blockers for myocardial infarction: a meta-analysis of randomized trials. *Am J Med* 2014;**127**:939–953. https://doi.org/10.1016/j.amjmed.2014.05.032
875. Huang BT, Huang FY, Zuo ZL, Liao YB, Heng Y, Wang PJ, *et al.* Meta-analysis of relation between oral β-blocker therapy and outcomes in patients with acute myocardial infarction who underwent percutaneous coronary intervention. *Am J Cardiol* 2015;**115**:1529–1538. https://doi.org/10.1016/j.amjcard.2015.02.057
876. Goldberger JJ, Bonow RO, Cuffe M, Liu L, Rosenberg Y, Shah PK, *et al.* Effect of beta-blocker dose on survival after acute myocardial infarction. *J Am Coll Cardiol* 2015;**66**:1431–1441. https://doi.org/10.1016/j.jacc.2015.07.047
877. Andersson C, Shilane D, Go AS, Chang TI, Kazi D, Solomon MD, *et al.* β-blocker therapy and cardiac events among patients with newly diagnosed coronary heart disease. *J Am Coll Cardiol* 2014;**64**:247–252. https://doi.org/10.1016/j.jacc.2014.04.042
878. Bangalore S, Steg G, Deedwania P, Crowley K, Eagle KA, Goto S, *et al.* β-blocker use and clinical outcomes in stable outpatients with and without coronary artery disease. *JAMA* 2012;**308**:1340–1349. https://doi.org/10.1001/jama.2012.12559
879. Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;**355**:253–259. https://doi.org/10.1016/S0140-6736(99)12323-7
880. Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, *et al.* The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999;**341**:709–717. https://doi.org/10.1056/nejm199909023411001
881. Hirpa M, Woreta T, Addis H, Kebede S. What matters to patients? A timely question for value-based care. *PLoS One* 2020;**15**:e0227845. https://doi.org/10.1371/journal.pone.0227845
882. Ebrahimi Z, Patel H, Wijk H, Ekman I, Olaya-Contreras P. A systematic review on implementation of person-centered care interventions for older people in out-of-hospital settings. *Geriatr Nurs* 2021;**42**:213–224. https://doi.org/10.1016/j.gerinurse.2020.08.004
883. Gluyas H. Patient-centred care: improving healthcare outcomes. *Nurs Stand* 2015;**30**:50–57, quiz 59. https://doi.org/10.7748/ns.30.4.50.e10186
884. Kok MM, von Birgelen C. Involving the patient's perspective and preferences concerning coronary angiography and percutaneous coronary intervention. *EuroIntervention* 2020;**15**:1228–1231. https://doi.org/10.4244/eijv15i14a221
885. Astin F, Stephenson J, Probyn J, Holt J, Marshall K, Conway D. Cardiologists' and patients' views about the informed consent process and their understanding of the anticipated treatment benefits of coronary angioplasty: a survey study. *Eur J Cardiovasc Nurs* 2020;**19**:260–268. https://doi.org/10.1177/1474515119879050
886. Flynn D, Knoedler MA, Hess EP, Murad MH, Erwin PJ, Montori VM, *et al.* Engaging patients in health care decisions in the emergency department through shared decision-making: a systematic review. *Acad Emerg Med* 2012;**19**:959–967. https://doi.org/10.1111/j.1553-2712.2012.01414.x
887. Grant EV, Summapund J, Matlock DD, Vaughan Dickson V, Iqbal S, Patel S, *et al.* Patient and cardiologist perspectives on shared decision making in the treatment of older adults hospitalized for acute myocardial infarction. *Med Decis Making* 2020;**40**:279–288. https://doi.org/10.1177/0272989x20912293
888. Shah P, Thornton I, Turrin D, Hipskind JE. Informed consent. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing, Copyright © 2022, StatPearls Publishing LLC, 2022.
889. Prochnow JA, Meiers SJ, Scheckel MM. Improving patient and caregiver new medication education using an innovative teach-back toolkit. *J Nurs Care Qual* 2019;**34**:101–106. https://doi.org/10.1097/ncq.0000000000000342
890. Klingbeil C, Gibson C. The Teach Back Project: a system-wide evidence based practice implementation. *J Pediatr Nurs* 2018;**42**:81–85. https://doi.org/10.1016/j.pedn.2018.06.002
891. Ha Dinh TT, Bonner A, Clark R, Ramsbotham J, Hines S. The effectiveness of the teach-back method on adherence and self-management in health education for people with chronic disease: a systematic review. *JBI Database System Rev Implement Rep* 2016;**14**:210–247. https://doi.org/10.11124/jbisrir-2016-2296
892. Dickert NW, Scicluna VM, Adeoye O, Angiolillo DJ, Blankenship JC, Devireddy CM, *et al.* Emergency consent: patients' and surrogates' perspectives on consent for clinical trials in acute stroke and myocardial infarction. *J Am Heart Assoc* 2019;**8**:e010905. https://doi.org/10.1161/jaha.118.010905
893. Dickert NW, Miller FG. Involving patients in enrolment decisions for acute myocardial infarction trials. *BMJ* 2015;**351**:h3791. https://doi.org/10.1136/bmj.h3791
894. Olsson A, Ring C, Josefsson J, Eriksson A, Rylance R, Fröbert O, *et al.* Patient experience of the informed consent process during acute myocardial infarction: a sub-study of the VALIDATE-SWEDEHEART trial. *Trials* 2020;**21**:246. https://doi.org/10.1186/s13063-020-4147-0
895. El-Haddad C, Hegazi I, Hu W. Understanding patient expectations of health care: a qualitative study. *J Patient Exp* 2020;**7**:1724–1731. https://doi.org/10.1177/2374373520921692
896. Scott JT, Thompson DR. Assessing the information needs of post-myocardial infarction patients: a systematic review. *Patient Educ Couns* 2003;**50**:167–177. https://doi.org/10.1016/s0738-3991(02)00126-x
897. Saczynski JS, McManus DD, Waring ME, Lessard D, Anatchkova MD, Gurwitz JH, *et al.* Change in cognitive function in the month after hospitalization for acute coronary syndromes: findings from TRACE-CORE (Transition, Risks, and Actions in Coronary Events-Center for Outcomes Research and Education). *Circ Cardiovasc Qual Outcomes* 2017;**10**:e001669. https://doi.org/10.1161/circoutcomes.115.001669
898. Goldman JD, Harte FM. Transition of care to prevent recurrence after acute coronary syndrome: the critical role of the primary care provider and pharmacist. *Postgrad Med* 2020;**132**:426–432. https://doi.org/10.1080/00325481.2020.1740512
899. Huriani E. Myocardial infarction patients' learning needs: perceptions of patients, family members and nurses. *Int J Nurs Sci* 2019;**6**:294–299. https://doi.org/10.1016/j.ijnss.2019.05.001
900. Messerli AW, Deutsch C. Implementation of institutional discharge protocols and transition of care following acute coronary syndrome. *Cardiovasc Revasc Med* 2020;**21**:1180–1188. https://doi.org/10.1016/j.carrev.2020.02.013
901. Schiele F, Lemesle G, Angoulvant D, Krempf M, Kownator S, Cheggour S, *et al.* Proposal for a standardized discharge letter after hospital stay for acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:788–801. https://doi.org/10.1177/2048872619844444
902. Murphy B, Le Grande M, Alvarenga M, Worcester M, Jackson A. Anxiety and depression after a cardiac event: prevalence and predictors. *Front Psychol* 2019;**10**:3010. https://doi.org/10.3389/fpsyg.2019.03010
903. Ceccarini M, Manzoni GM, Castelnuovo G. Assessing depression in cardiac patients: what measures should be considered? *Depress Res Treat* 2014;**2014**:148256. https://doi.org/10.1155/2014/148256
904. Moser DK. "The rust of life": impact of anxiety on cardiac patients. *Am J Crit Care* 2007;**16**:361–369. https://doi.org/10.4037/ajcc2007.16.4.361
905. Turgeon RD, Koshman SL, Dong Y, Graham MM. P2Y12 inhibitor adherence trajectories in patients with acute coronary syndrome undergoing percutaneous coronary intervention: prognostic implications. *Eur Heart J* 2022;**43**:2303–2313. https://doi.org/10.1093/eurheartj/ehac116
906. Poitras ME, Maltais ME, Bestard-Denommé L, Stewart M, Fortin M. What are the effective elements in patient-centered and multimorbidity care? A scoping review. *BMC Health Serv Res* 2018;**18**:446. https://doi.org/10.1186/s12913-018-3213-8
907. Hochhalter AK, Song J, Rush J, Sklar L, Stevens A. Making the Most of Your Healthcare intervention for older adults with multiple chronic illnesses. *Patient Educ Couns* 2010;**81**:207–213. https://doi.org/10.1016/j.pec.2010.01.018
908. Hess EP, Knoedler MA, Shah ND, Kline JA, Breslin M, Branda ME, *et al.* The chest pain choice decision aid: a randomized trial. *Circ Cardiovasc Qual Outcomes* 2012;**5**:251–259. https://doi.org/10.1161/circoutcomes.111.964791
909. Hess EP, Hollander JE, Schaffer JT, Kline JA, Torres CA, Diercks DB, *et al.* Shared decision making in patients with low risk chest pain: prospective randomized pragmatic trial. *BMJ* 2016;**355**:i6165. https://doi.org/10.1136/bmj.i6165
910. van Oosterhout REM, de Boer AR, Maas A, Rutten FH, Bots ML, Peters SAE. Sex differences in symptom presentation in acute coronary syndromes: a systematic review

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026

and meta-analysis. *J Am Heart Assoc* 2020;**9**:e014733. https://doi.org/10.1161/jaha.119.014733
911. Hedegaard U, Kjeldsen LJ, Pottegård A, Henriksen JE, Lambrechtsen J, Hangaard J, *et al.* Improving medication adherence in patients with hypertension: a randomized trial. *Am J Med* 2015;**128**:1351–1361. https://doi.org/10.1016/j.amjmed.2015.08.011
912. Bauer LK, Caro MA, Beach SR, Mastromauro CA, Lenihan E, Januzzi JL, *et al.* Effects of depression and anxiety improvement on adherence to medication and health behaviors in recently hospitalized cardiac patients. *Am J Cardiol* 2012;**109**:1266–1271. https://doi.org/10.1016/j.amjcard.2011.12.017
913. Van der Kooy K, van Hout H, Marwijk H, Marten H, Stehouwer C, Beekman A. Depression and the risk for cardiovascular diseases: systematic review and meta analysis. *Int J Geriatr Psychiatry* 2007;**22**:613–626. https://doi.org/10.1002/gps.1723
914. Redfors B, Angerås O, Råmunddal T, Petursson P, Haraldsson I, Dworeck C, *et al.* Trends in gender differences in cardiac care and outcome after acute myocardial infarction in Western Sweden: a report from the Swedish Web System for Enhancement of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). *J Am Heart Assoc* 2015;**4**:e001995. https://doi.org/10.1161/jaha.115.001995
915. Anand SS, Xie CC, Mehta S, Franzosi MG, Joyner C, Chrolavicius S, *et al.* Differences in the management and prognosis of women and men who suffer from acute coronary syndromes. *J Am Coll Cardiol* 2005;**46**:1845–1851. https://doi.org/10.1016/j.jacc.2005.05.091
916. Blomkalns AL, Chen AY, Hochman JS, Peterson ED, Trynosky K, Diercks DB, *et al.* Gender disparities in the diagnosis and treatment of non-ST-segment elevation acute coronary syndromes: large-scale observations from the CRUSADE (Can Rapid Risk Stratification of Unstable Angina Patients Suppress Adverse Outcomes With Early Implementation of the American College of Cardiology/American Heart Association Guidelines) National Quality Improvement Initiative. *J Am Coll Cardiol* 2005;**45**:832–837. https://doi.org/10.1016/j.jacc.2004.11.055
917. Bugiardini R, Yan AT, Yan RT, Fitchett D, Langer A, Manfrini O, *et al.* Factors influencing underutilization of evidence-based therapies in women. *Eur Heart J* 2011;**32**:1337–1344. https://doi.org/10.1093/eurheartj/ehr027
918. Samayoa L, Grace SL, Gravely S, Scott LB, Marzolini S, Colella TJF. Sex differences in cardiac rehabilitation enrollment: a meta-analysis. *Can J Cardiol* 2014;**30**:793–800. https://doi.org/10.1016/j.cjca.2013.11.007
919. Rossello X, Mas-Lladó C, Pocock S, Vicent L, van de Werf F, Chin CT, *et al.* Sex differences in mortality after an acute coronary syndrome increase with lower country wealth and higher income inequality. *Rev Esp Cardiol (Engl Ed)* 2022;**75**:392–400. https://doi.org/10.1016/j.rec.2021.05.006
920. Sardar MR, Badri M, Prince CT, Seltzer J, Kowey PR. Underrepresentation of women, elderly patients, and racial minorities in the randomized trials used for cardiovascular guidelines. *JAMA Internal Medicine* 2014;**174**:1868–1870. https://doi.org/10.1001/jamainternmed.2014.4758
921. Cho L, Vest AR, O'Donoghue ML, Ogunniyi MO, Sarma AA, Denby KJ, *et al.* Increasing participation of women in cardiovascular trials. *J Am College Cardiol* 2021;**78**:737–751. https://doi.org/10.1016/j.jacc.2021.06.022
922. Gong IY, Tan NS, Ali SH, Lebovic G, Mamdani M, Goodman SG, *et al.* Temporal trends of women enrollment in major cardiovascular randomized clinical trials. *Can J Cardiol* 2019;**35**:653–660. https://doi.org/10.1016/j.cjca.2019.01.010
923. Rossello X, Ferreira JP, Caimari F, Lamiral Z, Sharma A, Mehta C, *et al.* Influence of sex, age and race on coronary and heart failure events in patients with diabetes and post-acute coronary syndrome. *Clin Res Cardiol* 2021;**110**:1612–1624. https://doi.org/10.1007/s00392-021-01859-2
924. Mas-Llado C, González-Del-Hoyo M, Siquier-Padilla J, Blaya-Peña L, Coughlan JJ, García de la Villa B, *et al.* Representativeness in randomised clinical trials supporting acute coronary syndrome guidelines. *Eur Heart J Qual Care Clin Outcomes* 2023. https://doi.org/10.1093/ehjqcco/qcad007
925. Aktaa S, Batra G, Wallentin L, Baigent C, Erlinge D, James S, *et al.* European Society of Cardiology methodology for the development of quality indicators for the quantification of cardiovascular care and outcomes. *Eur Heart J Qual Care Clin Outcomes* 2022;**8**:4–13. https://doi.org/10.1093/ehjqcco/qcaa069
926. Minchin M, Roland M, Richardson J, Rowark S, Guthrie B. Quality of care in the United Kingdom after removal of financial incentives. *N Engl J Med* 2018;**379**:948–957. https://doi.org/10.1056/NEJMsa1801495
927. Song Z, Ji Y, Safran DG, Chernew ME. Health care spending, utilization, and quality 8 years into global payment. *N Engl J Med* 2019;**381**:252–263. https://doi.org/10.1056/NEJMsa1813621
928. Arbelo E, Aktaa S, Bollmann A, D'Avila A, Drossart I, Dwight J, *et al.* Quality indicators for the care and outcomes of adults with atrial fibrillation. *Europace* 2021;**23**:494–495. https://doi.org/10.1093/europace/euaa253
929. Schiele F, Aktaa S, Rossello X, Ahrens I, Claeys MJ, Collet JP, *et al.* 2020 Update of the quality indicators for acute myocardial infarction: a position paper of the Association for Acute Cardiovascular Care: the study group for quality indicators from the ACVC and the NSTE-ACS guideline group. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:224–233. https://doi.org/10.1093/ehjacc/zuaa037
930. Aktaa S, Abdin A, Arbelo E, Burri H, Vernooy K, Blomström-Lundqvist C, *et al.* European Society of Cardiology Quality Indicators for the care and outcomes of cardiac pacing: developed by the Working Group for Cardiac Pacing Quality Indicators in collaboration with the European Heart Rhythm Association of the European Society of Cardiology. *Europace* 2022;**24**:165–172. https://doi.org/10.1093/europace/euab193
931. Aktaa S, Gencer B, Arbelo E, Davos CH, Désormais I, Hollander M, *et al.* European Society of Cardiology Quality Indicators for Cardiovascular Disease Prevention: developed by the Working Group for Cardiovascular Disease Prevention Quality Indicators in collaboration with the European Association for Preventive Cardiology of the European Society of Cardiology. *Eur J Prev Cardiol* 2022;**29**:1060–1071. https://doi.org/10.1093/eurjpc/zwab160
932. Schiele F, Gale CP, Simon T, Fox KAA, Bueno H, Lettino M, *et al.* The 2020 ESC-ACVC quality indicators for the management of acute myocardial infarction applied to the FAST-MI registries. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:207–215. https://doi.org/10.1093/ehjacc/zuab010
933. Rossello X, Medina J, Pocock S, Van de Werf F, Chin CT, Danchin N, *et al.* Assessment of quality indicators for acute myocardial infarction management in 28 countries and use of composite quality indicators for benchmarking. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:911–922. https://doi.org/10.1177/2048872620911853
934. Rossello X, Massó-van Roessel A, Perelló-Bordoy A, Mas-Lladó C, Ramis-Barceló MF, Vives-Borrás M, *et al.* Assessment of the ESC quality indicators in patients with acute myocardial infarction: a systematic review. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:878–889. https://doi.org/10.1093/ehjacc/zuab042
935. Batra G, Aktaa S, Wallentin L, Maggioni AP, Wilkinson C, Casadei B, *et al.* Methodology for the development of international clinical data standards for common cardiovascular conditions: European Unified Registries for Heart Care Evaluation and Randomised Trials (EuroHeart). *Eur Heart J Qual Care Clin Outcomes* 2021;**9**:161–168. https://doi.org/10.1093/ehjqcco/qcab052
936. Rossello X, Massó-van Roessel A, Chioncel O, Tavazzi L, Ferrari R, Vahanian A, *et al.* EURObservational Research Programme: a bibliometric assessment of its scientific output. *Eur Heart J Qual Care Clin Outcomes* 2022;**8**:804–811. https://doi.org/10.1093/ehjqcco/qcac041

Downloaded from academic.oup.com/eurheartj/article/44/38/3720/7243210 by guest on 26 October 2026