Congratulations! You passed!

Grade received 90% Latest Submission Grade 90% To pass 80% or higher

Go to next item

1. On your next mission, while collecting rock samples, you observe a new crystal structure containing carbon, which could be key to life! You utilize the third spacecraft, Ingenuity, and meticulously collect enough rock samples to distribute within the weight limits of each spacecraft.

1/1 point

You place 2 basalt samples, 1 meteorite, and 5 crystal rock samples into the Perseverance rover, which all weigh 20 grams.

You then distribute 1 basalt, 2 meteorites, and 1 crystal into the Curiosity rover, with a weight of 10 grams in total.

Lastly, you place 2 basalt samples, 1 meteorite, and 3 crystals to Ingenuity, which together weigh 15 grams. Each rock sample is represented with variables b for basalt, m for meteorite, and c for crystal structures.

Which of the following systems of equations represents the correct information in the above system of sentences?

$$\begin{cases} m + 2b + 5 = 20 \\ 2b + m + c = 10 \\ b + 2m + 3c = 15 \end{cases}$$

$$\begin{cases} 2b+m+5=20\\ b+2m+c=10 \end{cases}$$

$$\begin{cases} 2b + m + 5c = 100 \\ b + 2m + c = 23 \\ 2b + m + 3c = 35 \end{cases}$$

$$\begin{cases} 2b+m+5c=20\\ b+2m+c=10\\ 2b+m+3c=15 \end{cases}$$

⊘ Correct

Correct! This system of equations represents the weights of each rock sample noted with variables b for basalt, m for meteorite, and c for the crystal structure. The first equation represents the weight of the samples on the Perseverance rover, the second on the Curiosity rover, and the third equation for Ingenuity.

2. Which of the following matrices represents the system of equations?

1/1 point

$$\begin{bmatrix} 2 & 1 & 5 \\ 1 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & 5 & 20 \\ 1 & 2 & 1 & 10 \\ 2 & 1 & 3 & 15 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 2 & 1 \end{bmatrix}$$

✓ Correct

 ${\tt Correct!\ This\ is\ the\ representation\ of\ the\ system\ of\ equations\ in\ matrix\ form.}$

the determinant, apply the information in a mount of the matrix found in Question in the matrix found in Question in the matrix found in Question in the matrix has linearly other rows.	culated the dete	erminant an	d identified the n	non singularity of the i	matrix.	1/1point
ngular ar ir ine! You have correctly cald the matrix found in Ques lependent. independent. be determined. ine! The matrix has linearly other rows.	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1 point
the matrix found in Questiependent. Independent. Independ	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1 point
the matrix found in Questiependent. Independent. Independ	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1 point
the matrix found in Questlependent. Independent. Independ	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1point
the matrix found in Questlependent. Independent. Independent. Independent in the determined. Independent in the matrix has linearly in the rows.	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1point
the matrix found in Questlependent. Independent. Independ	stion 2 has line	early depend	dent or independ	dent rows.	matrix.	1/1point
lependent. ndependent. be determined. ne! The matrix has linearly ther rows.						1/1 point
ndependent. be determined. ne! The matrix has linearly ther rows. bes each rock sample we	y independent r	rows. You ca	nnot obtain one r			
be determined. ne! The matrix has linearly other rows. pes each rock sample we	y independent r	rows. You ca	nnot obtain one r			
ne! The matrix has linearly other rows. Des each rock sample we	y independent r	rows. You ca	nnot obtain one r			
ne! The matrix has linearly other rows. Des each rock sample we	y independent r	rows. You ca	nnot obtain one i			
				row by using row ope	rations	
em of equations for each	igh?					1/1 point
	of the spacecra	ft.				
.5g, meteorite = 2.5g, crys						
	_					
g, meteorite = 1g, crystal :	= 3g					
sample = 2.5g						
.5g, meteorite = 3.5g, crys	stal = 2.5g					
2.5. In other words, the ba	asalt rock samp	le weighs 2.5				
ting of the first and secon	d row respective	ely.		erseverance and Curio	osity	0 / 1 point
ues in Ingenuity does th	e matrix have	linearly dep	endent rows?			
	$\begin{bmatrix} 2 \\ 1 \\ \mathbf{x} \end{bmatrix}$	$\begin{bmatrix} 1 & 5 \\ 2 & 1 \\ \mathbf{y} & \mathbf{z} \end{bmatrix}$				
weight samples: x - basalt	, y - meteorite, z	z- crystal.)				
3, z = 6						
3, z = 3						
t te. Linearly dependent rov					other	
	I The system of equations 2.5. In other words, the bans, and the crystal sample the following matrix with ting of the first and secondues in Ingenuity does the weight samples: x - basalt 3, z = 6 3, z = 3 2, z = 3 4 4 4 4 4 4 5 4 5 5 6 5 6 6 6 6 6 6 6 6	.5g, meteorite = 3.5g, crystal = 2.5g If the system of equations has a unique so 2.5. In other words, the basalt rock sampnes, and the crystal sample weighs 2.5 grather following matrix with the values for reting of the first and second row respective ting of the first and second row	If the system of equations has a unique solution at the 2.5. In other words, the basalt rock sample weighs 2.5 ms, and the crystal sample weighs 2.5 grams. The following matrix with the values for rock samplesting of the first and second row respectively. Solution in Ingenuity does the matrix have linearly dependent samples: x - basalt, y - meteorite, z- crystal.) Solution is a significant of the first and second row respectively. The following matrix with the values for rock samples ting of the first and second row respectively. Lives in Ingenuity does the matrix have linearly dependent samples: x - basalt, y - meteorite, z- crystal.) Solution is a significant form of the first and second row respectively.	If the system of equations has a unique solution at the point (2.5, 2.5, 2.5, 1). In other words, the basalt rock sample weighs 2.5 grams, the meters, and the crystal sample weighs 2.5 grams. The following matrix with the values for rock samples weighs in the Peting of the first and second row respectively. Liues in Ingenuity does the matrix have linearly dependent rows? \[\begin{array} 2 & 1 & 5 \\ 1 & 2 & 1 \\ x & y & z \end{array} \] Weight samples: x - basalt, y - meteorite, z- crystal.) 3, z = 6 3, z = 3 2, z = 3 televalues into the matrix and see if it makes the rows deputibility the samples into the matrix and see if it makes the rows deputibility the samples into the matrix and see if it makes the rows deputibility the samples into the matrix and see if it makes the rows deputibility the samples into the matrix and see if it makes the rows deputibility the samples into the matrix and see if it makes the rows deputible in the point (2.5, 2.5, 2.5).	If the system of equations has a unique solution at the point (2.5, 2.5, 2.5) where b = 2.5, m = 2.5. In other words, the basalt rock sample weighs 2.5 grams, the meteorite rock sample weighs, and the crystal sample weighs 2.5 grams. The following matrix with the values for rock samples weighs in the Perseverance and Curic ting of the first and second row respectively. Liues in Ingenuity does the matrix have linearly dependent rows? \[\begin{array}{cccccccccccccccccccccccccccccccccccc	If the system of equations has a unique solution at the point (2.5, 2.5, 2.5) where b = 2.5, m = 2.5, 2.5. In other words, the basalt rock sample weighs 2.5 grams, the meteorite rock sample weighs ns, and the crystal sample weighs 2.5 grams. The following matrix with the values for rock samples weighs in the Perseverance and Curiosity ting of the first and second row respectively. Liues in Ingenuity does the matrix have linearly dependent rows? \[\begin{array}{cccccccccccccccccccccccccccccccccccc

8. Select which of the following are true for non-singular matrices.

1/1 point

- ☐ In a non-singular matrix, rows are linearly dependent.
- In a non-singular matrix there is only a unique solution for the represented system of equations.
- ✓ Correct

Correct! Since the rows are linearly independent in the non-singular matrix, you can find a unique solution for the represented system of equations.


- ✓ In a non-singular matrix, rows are linearly independent.
- **⊘** Correct


Correct! Non-singular matrices have linearly independent rows.


- In a non-singular matrix a row can be a multiple of the other one.
- 9. To train your AI assistant to classify systems of equations, you're now asked to select the correct sequence of graphs that represents a system of linear equations with:

1/1 point

1. zero solutions, 2. just one solution, 3. infinitely many solutions.







✓ Correct

Correct! You can visually determine how many solutions a system of linear equations has following these rules: If the lines are parallel, the system has no solutions; if the lines intersect at just one point, the system has just one solution and the solution is the point where they intersect. If the lines totally overlap, the system has infinitely many solutions.

10. Select the correct sequence of graphs that shows:

1/1 point

 ${\bf 1.}\ {\bf zero}\ {\bf solutions}, {\bf 2.}\ {\bf just}\ {\bf one}\ {\bf solution}, {\bf 3.}\ {\bf infinitely}\ {\bf many}\ {\bf solutions}.$













○ Correct Correct! The first graph has no solution because there is no place where the three planes intersect. The second graph has one solution, the origin (0, 0, 0). And finally, you can see that in the third graph, there are infinitely many solutions (points where the lines intersect).