

✔ Congratulations! You passed!

Grade
received **85.71%**

Latest Submission
Grade 85.71%

To pass 75% or higher

[Go to next item](#)



Your computer's timezone does not seem to match your Coursera account's timezone setting of Asia/Jakarta.
[Change your Coursera timezone setting](#)

1. What is the difference between sample mean and population mean?

1 / 1 point

- ☒ Sample mean is the average of a subset of the population, while population mean is the average of the entire population.
- ☐ Sample mean is always smaller than population mean.
- ☐ There is no difference between sample mean and population mean.

✔ **Correct**

Correct! Sample mean is the average of a subset of the population, while population mean is the average of the entire population.

2. Which of the following method can be used to estimate the variance, mean, and proportion of a population?

1 / 1 point

- ☐ Sample mean
- ☐ Sample variance
- ☒ Point estimation
- ☐ Regression analysis

✔ **Correct**

Correct! Point estimation is a statistical method that estimates population parameters such as mean, variance, and proportion based on a sample.

[Change your Coursera timezone setting](#)

3. Which of the following statements best describes the law of large numbers?

1 / 1 point

- ☐ The law of large numbers states that as the sample size increases, the sample mean becomes more variable.
- ☒ The law of large numbers states that as the sample size increases, the sample mean approaches the population mean with increasing accuracy.
- ☐ The law of large numbers states that as the sample size increases, the sample variance approaches the population variance.
- ☐ The law of large numbers states that as the sample size increases, the sample becomes more biased.

✔ **Correct**

Correct! The law of large numbers states that as the sample size increases, the sample mean approaches the population mean with increasing accuracy.

4. Suppose you flip a coin 10 times and obtain 6 heads and 4 tails. What function needs to be maximized to find the maximum likelihood estimate of the probability of getting heads on a single coin toss? Let p be the probability of the coin being heads.

1 /

- ☐ $L(p) = p^{1/6}(1-p)^{1/4}$
- ☒ $L(p) = p^6(1-p)^4$
- ☐ $L(p) = p^4(1-p)^6$
- ☐ $L(p) = p^{10}(1-p)^0$

✔ **Correct**

Correct! The likelihood function for this problem is $L(p) = p^6(1-p)^4$.

[Change your Coursera timezone setting](#)

5. What is the purpose of regularization in machine learning?

1 / 1 point

- ☐ Regularization is used to make a model more complex and flexible, which can lead to better performance on the training data.
- ☒ Regularization is used to prevent overfitting and reduce the complexity of a model, by adding a penalty term to the loss function that encourages smaller parameter values.
- ☐ Regularization is used to increase the training error of a model, which can improve its generalization performance.
- ☐ Regularization is used to improve the interpretability of a model by reducing its complexity.

[Change your Coursera timezone setting](#)

✓ **Correct**

Correct! Regularization is used to prevent overfitting and reduce the complexity of a model, by adding a penalty term to the loss function that encourages smaller parameter values.

[Change your Coursera timezone setting](#)

6. Consider the following population: $[-2, -1, 0, 1, 2]$ and the following sample $[-1, 0, 2]$.

1 / 1 point

What is the **population mean**?

0

✓ **Correct**

Correct!

7. Consider the following population: $[-2, -1, 0, 1, 2]$ and the following sample $[-1, 0, 2]$.

0 / 1 point

What is the **sample variance**? (Use two decimal places in your answer)

3.83

✗ **Incorrect**

Incorrect! Please review your calculations! Remember that the **sample** variance is divided $n - 1$ instead of n , where n is the sample size.