

a
AQP4 antibody splits NMO from MS
NMO criteria (2006)
NMOSD criteria (2015)
treated as MS
NMO
NMOSD
eculizumab PNH pilot (~2002)
eculizumab approved for PNH (2007)
first NMOSD trial (2009)
approved for NMOSD (2019)
2000
2005
2010
2015
2020
≈2 years before first trial: NMO ranks 45 of 1,728 by DisAtlas for eculizumab (top 2.6%)
how this rank is computed
1 Known use
eculizumab
antibody that blocks C5 (complement)
PNH: pilot ~2002, phase 3 2004, approved 2007
2 Allow for popularity
1,728 candidate diseases, 2006
DisAtlas score
NMO
trials before 2007 (log scale)
NMO is compared with every disease not yet trialed with eculizumab, and popularity is adjusted for
3 Rank by DisAtlas score
NMO: rank 45 of 1,728
= top 2.6% of candidates
DisAtlas
popularity alone: rank 121 (top 7%)
top
chance
bottom
score = closeness to PNH in the map
no direct link to PNH in the map (2 hops apart)
b
Odds that a disease is the drug's next trial
DisAtlas, unadjusted
DisAtlas, adjusted for popularity
shuffled DisAtlas scores, adjusted
Odds ratio per 1 SD of score (log scale)
1.93
no effect
Decision year (evidence up to the year before)
pooled
c
Each method, same adjustment
DisAtlas
Shared genes
GO processes
HPO phenotypes
Weighted Adamic–Adar
Map connectivity
1.93
1.33
1.27
1.20
1.10
1.02
Odds ratio per 1 SD, adjusted for popularity (2000–2021 pooled)