

Chap 2: Linear regression with least square estimate

A. Rey, HES-SO MSE, 02.2022

close all
clear

Exercise 2.1 Ellipses

Description

We collect a dataset of measurements (for exemple positions) and they are plotted in a 2 axis graph shown bellow:




Figure 1

The goal is to find the best Ellipse approximation with the minimum of the L_2 norm. and then plot it like in the figure 2:




Figure 2

Theory

The ellipse equation is given by:

$$ax^2 + by^2 + cx + dy + exy = 1 \quad (1)$$

the constants values a, b, c, d, e are the parameters to be found, in this case they are the unknowns of the problem to solve. We need only 5 points $P(x_i, y_i)$ to determine the constants a, b, c, d, e , but here we want to find the best approximation that minimize the L_2 norm of the error between the points $P(x_i, y_i)$ and the best ellipse approximation.

For each point $P_i = P(x_i, y_i)$ we could write the equation above for N points we have the overdetermined system of equations with N equations and 5 unknowns.

Data Import and plot

First import the data from the file `EllipseData.mat`, the points are in the matrix `Pi`, the column number is the point number i the first row contain the x_i values and the second row contains the y_i values.

Next plot the points in a graph like this shown in figure 1.

```
load('Ellipse_data.mat');
xi = Pi(1,:)';
yi = Pi(2,:)';
N = length(xi);
```

```
figure
```

```

plot(xi, yi, 'r.')
xlabel('x')
ylabel('y')
grid;

```

Compute the parameters for the best ellipse

Build the matrix A and the vector b representing the system of N equations like the equation (1):

To find the best parameters we need to solve the over-determined system:

$$A\Theta = b \quad (2)$$

It is no solution, that satisfies all the conditions (all points) so we could only find an approximation, solving eq (2) by pseudo inverse will give:

$$\hat{\Theta} = (A^T A)^{-1} A^T b \quad (3)$$

```

A = [xi.^2, yi.^2, xi, yi, xi.*yi];
b = ones(N, 1);
%Thetahat = inv(A'*A)*A'*b % use below one to avoid warning
Thetahat = (A'*A)\A' * b

```

```

Thetahat = 5x1
    0.0162
    0.0395
   -0.0019
    0.0003
   -0.0296

```

Compute the best ellipse points

With those parameters it is possible to plot the approximation of the Ellipse that fit the measurement points P_i but it is not trivial because y coordinates depends of x coordinates as shown by the ellipse equation (1) so we need compute for each possible x (in the same range than in the dataset) the corresponding y on the best ellipse given by the approximation parameters. From the equation (1) we need to solve $\hat{y} = f(x, \hat{\Theta})$ and this is a second order equation with 2, 1 or 0 real possible values of y for each x .

Let's re-write the eq (1) like this:

$by^2 + (d + ex)y + (ax^2 + cx - 1) = 0$ we need to exprime y and to simplify the notation we call $V = d + ex$ and $W = (ax^2 + bx - 1)$ and $Z = b$

the equation would be $Zy^2 + Vy + W = 0$ so y is find by solving this second order equation:

$$y = \frac{-V \pm \sqrt{V^2 - 4ZW}}{2Z} \quad (4)$$

First create a x vectos with the enough points to plot an Ellipse trajectory and with the same span than the x_i values in the data set:

Create the variables Z , V and W with the data x_i and the parameters in the vector $\hat{\Theta}$:

Next compute the \hat{y}_i values by solving (4) for positive and negative case (be careful to not take the non-real values):

```
x = floor(min(xi)):0.001:max(xi);
% For helping parameters a,b,c,d,e are equal to:
% a = Thetahat(1)
% b = Thetahat(2)
% c = Thetahat(3)
% d = Thetahat(4)
% e = Thetahat(5)

V = Thetahat(4) + Thetahat(5).*x;
W = Thetahat(1).*x.^2 + Thetahat(3).*x - 1;
Z = Thetahat(2);

discriminant = V.^2 - 4.*Z.*W;

x_double = [x(discriminant >= 0), x(discriminant >= 0)];
y = [(-V(discriminant >= 0) + (sqrt(discriminant(discriminant >= 0)))) ./ (2*Z), ...
      (-V(discriminant >= 0) - (sqrt(discriminant(discriminant >= 0)))) ./ (2*Z)];
```

Plot the best ellipse

```
figure
plot(xi, yi, 'b.', ...
      x_double, y, 'r');
xlabel('x')
ylabel('y')
```

Eigen values and eigen vectors application

To find the principal axis of the Ellipse we could compute the covariance matrix of the coordinate points: The Covariance matrix C_p is give by, the size is 2×2

$$C_p = \frac{1}{N} (P_i^T - m_p)^T (P_i^T - m_p) \text{ where } m_p \text{ is the vector of mean values of the rows of } P_i$$

We obtain the same result with `cov ()` function

```
mp = mean(Pi, 2)'
```

```
mp = 1x2
    -0.0037    -0.0265
```

```
Cp = 1/N .* ((Pi'-mp)' * (Pi'-mp))
```

```
Cp = 2x2
    46.8203    18.5040
    18.5040    17.1023
```

```
% check
cov(Pi')
```

```
ans = 2x2
    46.8672    18.5225
    18.5225    17.1194
```

Now we compute the eigen vectors (columns of V) and the eigen values (elements of D):

```
[V,D] = eig(Cp')
```

```
V = 2x2
    0.4324    -0.9017
   -0.9017    -0.4324
D = 2x2
    8.2297     0
     0    55.6929
```

```
% same values as teacher but not sorted in the same way
```

Eigen vectors must be orthogonal:

```
if V(:,1)'*V(:,2) == 0
    disp('V(:,1) et V(:,2) sont orthogonaux.')
end
```

```
V(:,1) et V(:,2) sont orthogonaux.
```

Usually we place the eigenvalues and so the eigenvectors, from the bigger to the smaller

```
V2 = [-V(2,1) V(2,2); -V(2,2) -V(2,1)]
```

```
V2 = 2x2
    0.9017    -0.4324
    0.4324     0.9017
```

```
D2 = [D(2,2) 0; 0 D(1,1)]
```

```
D2 = 2x2
    55.6929     0
     0     8.2297
```

Remarque intéressante: Le vecteur propre qui correspond à la valeur propre la plus grande correspond à l'axe le plus large de l'ellipse.