

I. Multiple Choice Questions



▼ Medical Entrance Gallery

- 1** Viroids differ from viruses in having— [NEET '17]
- Ⓐ DNA molecules without protein coat
 - Ⓑ RNA molecules with protein coat

- Ⓒ RNA molecules without protein coat
- Ⓓ DNA molecules with protein coat

- 2** Which of the following are found in extreme saline conditions? [NEET T]
- Ⓐ Eubacteria
 - Ⓑ Cyanobacteria
 - Ⓒ Mycobacteria
 - Ⓓ Archaeobacteria

- 3** Which among the following are the smallest living cells, known without a definite cell wall, pathogenic to plants as well as animals and can survive without oxygen? [NEET '17]
 (A) *Pseudomonas* (B) *Mycoplasma*
 (C) *Nostoc* (D) *Bacillus*
- 4** The primary producers of the deep-sea hydrothermal vent ecosystem are— [NEET '16]
 (A) blue-green algae
 (B) coral reefs
 (C) green algae
 (D) chemosynthetic bacteria
- 5** The primary prokaryotes responsible for the production of biogas from the dung of ruminant animals, include the— [NEET '16]
 (A) thermoacidophiles (B) methanogens
 (C) eubacteria (D) halophiles
- 6** Which one of the following is wrong for fungi? [NEET '16]
 (A) They are heterotrophic
 (B) They are both unicellular and multicellular
 (C) They are eukaryotic
 (D) All fungi possess a purely cellulosic cell wall
- 7** Methanogens belong to— [NEET '16]
 (A) Dinoflagellates
 (B) Slime moulds
 (C) Eubacteria
 (D) Archaeobacteria
- 8** Select the wrong statement. [NEET '16]
 (A) Diatoms are chief producers in the oceans
 (B) Diatoms are microscopic and float passively in water
 (C) The walls of diatoms are easily destructible
 (D) 'Diatomaceous earth' is formed by the cell walls of diatoms.
- 9** Select the mismatch. [NEET '16]
 (A) Protists- Eukaryotes
 (B) Methanogens- Prokaryotes
 (C) Gas vacuoles- Green bacteria
 (D) Large central vacuoles- Animal cells
- 10** Select the wrong statement. [NEET '16]
 (A) Cyanobacteria lack flagellated cells
 (B) *Mycoplasma* is a wall-less microorganism
 (C) Bacterial cell wall is made up of peptidoglycan
 (D) Pill and fimbriae are mainly involved in motility of bacterial cells
- 11** Which of the following statements is wrong for viroids? [NEET '16]
 (A) They are smaller than viruses
 (B) They cause infections
 (C) Their RNA is of high molecular weight
 (D) They lack a protein coat
- 12** One of the major components of cell wall of most fungi is— [NEET '16]
 (A) Chitin (B) Peptidoglycan
 (C) Cellulose (D) Hemicellulose
- 13** Which of the following is not a feature of the plasmids? [NEET '16]
 (A) Circular structure (B) Transferable
 (C) Single-stranded
 (D) Independent replication
- 14** Which of the following would appear as the pioneer organisms on bare rocks? [NEET '16]
 (A) Lichens (B) Liverworts
 (C) Mosses (D) Green algae
- 15** Chrysophytes, Euglenoids, Dinoflagellates and Slime moulds are included in the kingdom — [NEET '16]
 (A) Monera (B) Protista (C) Fungi (D) Animalia
- 16** Choose the wrong statement— [RE-AIPMT '15]
 (A) Yeast is unicellular and useful in fermentation
 (B) *Penicillium* is multicellular and produces antibiotics
 (C) *Neurospora* is used in the study of biochemical genetics
 (D) Morels and truffles are poisonous mushrooms
- 17** Cell wall is absent in— [RE-AIPMT '15]
 (A) *Nostoc* (B) *Aspergillus*
 (C) *Funaria* (D) *Mycoplasma*
- 18** In which group of organisms the cells walls form two thin overlapping shells which fit together? [RE-AIPMT '15]
 (A) Slime moulds (B) Chrysophytes
 (C) Euglenoids (D) Dinoflagellates
- 19** Select the wrong statement— [RE-AIPMT '15]
 (A) Mosaic disease in tobacco and AIDS in human being are caused by viruses
 (B) The viroids were discovered by D.J. Ivanowski
 (C) W.M. Stanley showed that viruses could be crystallized
 (D) The term 'contagium vivum fluidum' was coined by M.W. Beijerinck



- 20** The structures that help some bacteria to attach to rocks and/or host tissues are— [RE-AIPMT '15]
 (A) Holdfast (B) Rhizoids
 (C) Fimbriae (D) Mesosomes
- 21** The imperfect fungi which are decomposers of litter and help in mineral cycling belong to— [RE-AIPMT '15]
 (A) Ascomycetes (B) Deuteromycetes
 (C) Basidiomycetes (D) Phycomycetes
- 22** Which the following are most suitable indicator of SO_2 pollution in the environment? [RE-AIPMT '15]
 (A) Fungi (B) Lichens (C) Conifers (D) Algae
- 23** Pick up the wrong statement— [RE-AIPMT '15]
 (A) Nuclear membrane is present in Monera
 (B) Cell wall is absent in Animalia
 (C) Protista have photosynthetic and heterotrophic modes of nutrition
 (D) Some fungi are edible
- 24** Five kingdom system of classification suggested by RH Whittaker is not based on [AIPMT '14]
 (A) presence or absence of a well defined nucleus
 (B) mode of reproduction
 (C) mode of nutrition
 (D) complexity of body organisation
- 25** Archaeobacteria differ from eubacteria in— [AIPMT '14]
 (A) cell membrane structure
 (B) mode of nutrition
 (C) cell shape
 (D) mode of reproduction
- 26** Which of the following shows coiled RNA strand and capsomeres? [AIPMT '14]
 (A) Polio virus (B) Tobacco mosaic virus
 (C) Measles virus (D) Retrovirus
- 27** Viruses have— [AIPMT '14]
 (A) DNA enclosed in a protein coat
 (B) prokaryotic nucleus
 (C) single chromosome (D) Both DNA and RNA
- 28** Identify the correct sequence of events in the viral replication process. [AFMC '14]
 (i) Eclipse (ii) Maturation
 (iii) Adsorption (iv) Assembly
 (v) Penetration (vi) Lysis
 (A) (i), (ii), (iii), (iv), (v), (vi)
 (B) (ii), (i), (iii), (iv), (v), (vi)
 (C) (iii), (v), (i), (ii), (iv), (vi)
 (D) (iii), (v), (vi), (iv), (ii), (i)
- 29** As per Whittaker's classification, an organism possessing eukaryotic cell structure, multicellular organisation, with a cell wall and nuclear membrane showing heterotrophic nutrition can be placed under the kingdom— [Kerala CEE '14]
 (A) Monera (B) Protista (C) Plantae (D) Fungi
 (E) Animalia
- 30** Which of the following does not apply to Ascomycetes? [Kerala CEE '14]
 (A) Mycelium coenocytic and aseptate
 (B) Commonly known as sac fungi
 (C) Asexual spores called conidia are produced exogenously
 (D) Sexual spores called ascospores are produced endogenously
 (E) They are saprophytic, decomposers, parasitic or coprophilous
- 31** Which of the following groups of organisms have a protein rich layer called pellicle? [Kerala CEE '14]
 (A) Chrysophytes (B) Euglenoids
 (C) Dinoflagellates (D) Slime moulds
 (E) Protozoans
- 32** Identify the correct pairs of combination. [EAMCET '14]
 (i) Viroid — Bovine spongiform encephalitis.
 (ii) Prion — Creutzfeldt-Jakob disease.
 (iii) Measles virus — Glycoprotein projections.
 (iv) Rabies virus Polyhedral symmetry.
 (A) (iii) and (iv) (B) (i) and (iii)
 (C) (i) and (ii) (D) (ii) and (iii)
- 33** The protozoan with heliopodia as locomotory structures— [EAMCET '14]
 (A) Actinophryx (B) Entamoeba
 (C) Elphidium (D) Euglypha
- 34** The type of syngamy seen in *Trichonympha* is— [EAMCET '14]
 (A) hologamy (B) anisogamy
 (C) isogamy (D) conjugation
- 35** Protists obtain their food as— [AIIMS '13]
 (A) photosynthesisers (B) chemosynthesisers
 (C) heterotrophs (D) Both (A) and (C)
- 36** Which of the following does not belong to the kingdom—Protista? [Kerala CEE '13]
 (A) Chrysophytes (B) Euglenoids
 (C) Ascomycetes (D) Dinoflagellates
 (E) Protozoans

37 Bacteria differ from plants in that they do not have— **[UP CPMT '13]**

(A) DNA

(B) RNA

(C) cell wall

(D) a well defined nucleus