

KZ WORKING GROUP 2026

ALEXEI LATYNTSEV, XINXANG TANG, AND WHOEVER ELSE WANTS TO JOIN

1. Plan

1.1. **Setting up definitions.** Write down

- (1) (a) the KZ equation (over any surface Σ) [ES; Fe; Iv],
(b) the boundary KZ equation,
(c) the dynamical KZ equation, [FMTV; LX]
- (2) (a) the q KZ equation (additive, multiplicative and heat versions [FV]),
(b) dynamical q KZ?
(c) the dynamical q KZ equation [LX],
- (3) the (q, t) KZ equation [AKMMMOZ; AKMMSZ].

Then write, without proof, a lot of Theorems of the form

Theorem A. *The X -limit of equation Y is equation Z .*

1.2. **Finite quantum groups and associators.** Prove

Theorem B. [ES, §15.2], [EK, §3] *For simple \mathfrak{g} , the KZ equations*

- (1) *induce a Drinfeld associator on $\text{Rep}_{k[[\hbar]]} U(\mathfrak{g})[[\hbar]] \rightarrow \text{Vect}$,*
- (2) *induce a braided monoidal structure on $\text{Rep}_{k[[\hbar]]} U(\mathfrak{g})[[\hbar]] \rightarrow \text{Vect}$,*
- (3) *hence a quasitriangular Hopf algebra structure on $U(\mathfrak{g})[[\hbar]]$.*

1.2.1. *Axiomatisation of the above Theorem.* Define constructible sheaves, then write a 1-categorical version of [Lu, 5.5.4.10]:

Theorem C. *There is a functor*

$\psi : \{\text{factorisable constructible sheaves of categories on } \text{Ran} \mathbf{R}^2\} \rightarrow \{\text{braided monoidal categories}\},$
there is a constructible sheaf of categories \mathcal{E} with

$$\psi(\mathcal{E}) = \text{Rep}_{k[[\hbar]]} U(\mathfrak{g})[[\hbar]]$$

and interpret the KZ associator and Drinfeld twist and braiding in terms of ψ .

1.3. Multiple zeta values.


Conjecture D. [ES] The KZ associator is (given in terms of? how?) multiple zeta values.

discuss relation to $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ iff it's relevant to KZ?

1.4. KZ from QFT.

1.4.1. *Mathematical formalisation.* Define a Atiyah-style QFT as a functor $\mathcal{Z} : \text{Cob}_n \rightarrow \text{Vect}_n$ with an enriched structure. Define spacetime symmetry of \mathcal{Z} . Give example of holomorphic-topological theories.


Theorem E. Given an n -dimensional $X \times \mathbf{R}$, the category $\mathcal{Z}(S_x^{n-2})$ of line operators naturally forms a (topological/holomorphic/...) factorisation category over X :



Conclude that for any sections $V_1, \dots, V_n \in \mathcal{Z}(S^{n-2})$ as x varies, we get a vector bundle (with connection of the appropriate type given by spacetime symmetries) over X^n with fibres $V_1 \otimes \dots \otimes V_n$.

1.4.2. *To prepare for Kazhdan–Lustztig.* Give a definition of topological QFT \mathcal{Z} with holomorphic boundary $\partial\mathcal{Z}$, show that there is a “Riemann–Hilbert” functor $\mathcal{Z}(S^{n-2}) \rightarrow \partial\mathcal{Z}(S^{n-3})$.

1.4.3. *Physics.* Discuss the Physics-Theorem due to [Co] that M-theory on a CY3 Y gives a 5d QFT on noncommutative spacetime. Sketch Kevin’s example where $Y = K_{\text{ADE}}$:



Discuss how 4d Chern–Simons can only be valued on an algebraic group [CWY].

Theorem F. Assuming X properties about 3, 4 and 5 Chern–Simons theory, we get the KZ, q KZ and (q, t) -KZ equations. The different variables q, t , etc. correspond to Y, Z , etc. in the above geometry.

1.5. Relation to Coulomb branches. Firstly, using quasimaps

$$\begin{array}{ccc} & \text{QMaps}_{p_1, p_2}^d(\Sigma, M) & \\ \text{ev}_2 \swarrow & & \searrow \text{ev}_1 \\ M & & M \end{array}$$

prove Okounkov's quasimaps solutions to KZ via *capping operators*,

Theorem 1.5.1. [Ok, Thm. 8.1.16, 8.2.20] When $\Sigma = \mathbf{P}^1$ and $M = G/B$, the capping operator

$$J(z) = \sum_{d \in H_2(M, \mathbf{Z})} q^d \cdot (\text{ev}_1 \times \text{ev}_2)_* \hat{\mathcal{O}}_{\text{vir}} \in K_{G \times G_{m_q}}(M)_{\text{loc}}^{\otimes 2} \otimes \mathbf{Q}[[q^d]]$$

1.5.2. KZ as a Gauss–Manin connection on quantum cohomology. Let \mathcal{M}_C denote the BFN Coulomb branch.

Theorem 1.5.3. [Da] For G simply laced the map

$$(V_{\lambda_1} \otimes \cdots \otimes V_{\lambda_n})_\mu \xrightarrow{\sim} H_{T,\bullet}^{\text{BM}}(\mathcal{M}_C^T) \xrightarrow{\text{Stab}_e} H_{T,\bullet}^{\text{BM}}(\mathcal{M}_C)_{T-\text{loc}}$$

of vector bundles over $\mathbf{C}^n \times \mathfrak{h}$ intertwines the KZ differential ∇_i^{KZ} and quantum multiplication $c_1(\mathcal{E}_i)^*$ by the first chern class, where V_λ is irreducible finite dimensional with highest weight λ .¹

1.5.4. *Question.* As the Coulomb branch is Spec of cohomology of the Hecke stack \mathcal{B}_z , we have taken cohomology twice,

$$H_{T,\bullet}^{\text{BM}}(\mathcal{M}_C) = H_{T,\bullet}^{\text{BM}}(\text{Spec} H_\bullet^{\text{BM}}(\mathcal{B}_z, m))$$

and therefore we can try to apply any pair of cohomology theories (labelled by one-dimensional formal groups):

	C	C[×]	E	(Σ)	
C					KZ
C[×]					qKZ
E					qKZB heat

What is the analogue of [Da]’s Theorem in each case? Is there a version for critical cohomology?

1.5.5. *Physics question.* Understand the physics relation between $4d$ Chern–Simons theory and $3d$ $\mathcal{N} = 4$ gauge theories whose Higgs/Coulomb branches are Nakajima quiver varieties. [Talk to Tianqing about this](#)

1.6. KZ via the Riemann–Hilbert correspondence: Kazhdan–Lusztig.

1.6.1. One can view the KZ equations as defining an element of either side of the Riemann–Hilbert correspondence

$$\text{RH} : \{\text{vector bundle with connection with regular singularity on } \mathbf{C}^n\} \xrightarrow{\sim} \{\text{perverse sheaf on } \mathbf{C}^n\}.$$

We show that this corresponds to two equivalent ways of viewing the KZ equations: via affine vertex algebras and via quantum groups.

1.6.2. Show vertex algebras equivalent to factorisable sheaves over $\text{Ran } \mathbf{C}$, and

Theorem G. FBZ Let V be a vertex algebra.

(1) If M_1, \dots, M_n are vertex modules over V , then there is a vector bundle with connection of their conformal blocks

$$C^0(M_1, \dots, M_n) \rightarrow \text{Conf}_n \mathbf{C}.$$

¹Here T is the framing torus and H the gauge torus.

(2) *There is a natural functor*

$$\psi_{\text{hol}} : \{\text{factorisable sheaves of QCoh-module categories on } \text{Ran}\mathbf{C}\} \rightarrow \{\text{categories}\}$$

and $\text{Rep}_{\text{VA}} V$ lies in the image.

1.6.3. Define a Riemann–Hilbert functor

$$\text{RH} : \{\text{sheaves of QCoh-module categories on } X \text{ with connection}\} \rightarrow \{\text{constructible sheaves of categories on } X\}$$

sending $(\mathcal{C}, \nabla) \mapsto \mathcal{C}^\nabla$, see [CF].

Theorem H. [CF, §1] *There is a factorisable sheaf of QCoh-module categories with flat connection (\mathcal{C}, ∇) with $\text{RH}(\mathcal{C}, \nabla) = \mathcal{E}$.*

1.6.4. *Kazhdan–Lusztig.* Define vertex algebras, and state without proof:

Theorem I. (Kazhdan–Lusztig; [CF]) *There is an equivalence of categories*

$$\widehat{\mathfrak{g}}\text{-Mod}_\kappa^{G(\mathcal{O})} \simeq U_q(\mathfrak{g})\text{-Mod}^{\text{f.d.}}$$

between integrable representations of the affine Lie algebra at level κ and finite-dimensional representations of the quantum group at $q = \exp(\pi i/(\kappa - h^\vee))$, for all nice enough κ .

References

- [AKMMMOZ] Hidetoshi Awata, Hiroaki Kanno, Andrei Mironov, Alexei Morozov, Andrey Morozov, Yusuke Ohkubo, and Yegor Zenkevich. “Generalized Knizhnik-Zamolodchikov equation for Ding-Iohara-Miki algebra”. In: *Physical Review D* 96.2 (2017), p. 026021.
- [AKMMSZ] Hidetoshi Awata, Hiroaki Kanno, Andrei Mironov, Alexei Morozov, Kazuma Suetake, and Yegor Zenkevich. “ (q, t) -KZ equations for quantum toroidal algebra and Nekrasov partition functions on ALE spaces”. In: *Journal of High Energy Physics* 2018.3 (2018), pp. 1–70.
- [CF] L. Chen and Y. Fu. “An extension of the Kazhdan-Lusztig equivalence”. In: *arXiv preprint arXiv:2111.14606* (2021).
- [Co] Kevin Costello. “M-theory in the Omega-background and 5-dimensional non-commutative gauge theory”. In: *arXiv preprint arXiv:1610.04144* (2016).
- [CWY] K. Costello, E. Witten, and M. Yamazaki. “Gauge theory and integrability, I”. In: *arXiv preprint arXiv:1709.09993* (2017).
- [Da] Ivan Danilenko. *Quantum cohomology of slices of the affine Grassmannian*. Columbia University, 2020.
- [EK] Pavel Etingof and David Kazhdan. “Quantization of Lie bialgebras, I”. In: *Selecta Mathematica* 2.1 (1996), pp. 1–41.
- [ES] P. Etingof and O. Schiffmann. *Lectures on Quantum Groups*.
- [Fe] G. Felder. “Conformal field theory and integrable systems associated to elliptic curves”. In: *Proceedings of the International Congress of Mathematicians: August 3–11, 1994 Zürich, Switzerland*. Birkhäuser Basel, 1995, pp. 1247–1255.
- [FMTV] G Felder, Ya Markov, V Tarasov, and A Varchenko. “Differential equations compatible with KZ equations”. In: *Mathematical Physics, Analysis and Geometry* 3 (2000), pp. 139–177.
- [FV] Giovanni Felder and Alexander Varchenko. “q-deformed KZB heat equation: completeness, modular properties and $SL(3, \mathbf{Z})$ ”. In: *Advances in Mathematics* 171.2 (2002), pp. 228–275.
- [GRZ] D. Gaiotto, M. Rapčák, and Y. Zhou. “Deformed Double Current Algebras, Matrix Extended \mathcal{W}_∞ Algebras, Coproducts, and Intertwiners from the M2-M5 Intersection”. In: *arXiv preprint arXiv:2309.16929* (2023).
- [Iv] D Ivanov. “Knizhnik-Zamolodchikov-Bernard equations on Riemann surfaces”. In: *International Journal of Modern Physics A* 10.17 (1995), pp. 2507–2536.
- [Lu] J. Lurie. *Higher algebra*. Preprint, available at <http://www.math.harvard.edu/~lurie>. 2017.
- [LX] V.T. Laredo and X. Xu. “Stokes phenomena, Poisson–Lie groups and quantum groups”. In: *Advances in Mathematics* 429 (2023), p. 109189.

- [Ok] Andrei Okounkov. “Lectures on K-theoretic computations in enumerative geometry”. In: *arXiv preprint arXiv:1512.07363* (2015).