



Eurytemora sample count by year (chronological)


Zero-height bars = years with no data (1994–2004 gap)




Mean CPUE by year (chronological, labels = n)




Channel_Station — counts
Showing top 25 of 1935 levels by frequency




CPUE by Channel_Station (log1p scale)




Channel_Station coverage by year (blank = no data)




Genus — counts



CPUE by Genus (log1p scale)



Genus coverage by year (blank = no data)



A histogram showing the distribution of latitude for the 1000 closest stations to the origin. The x-axis is labeled 'Latitude' and ranges from 37.8 to 38.6. The y-axis represents frequency, with a maximum value of 100. The distribution is unimodal and slightly right-skewed, with a peak frequency of approximately 100 occurring at a latitude of about 38.07. The data is spread between approximately 37.82 and 38.55.


Latitude (approx.)	Frequency (approx.)
37.82	10
37.84	10
37.94	10
37.96	20
37.98	25
38.00	20
38.02	40
38.04	70
38.06	100
38.08	70
38.10	40
38.12	30
38.14	15
38.16	15
38.18	30
38.20	5
38.22	10
38.24	20
38.26	10
38.28	10
38.30	5
38.32	10
38.34	10
38.40	5
38.46	5
38.52	5
38.54	2

Bar chart showing Mean CPUE (Y-axis, 0 to 2000) versus Latitude (X-axis, 15 bins). The values for each bin are: 288, 237, 248, 1917, 7500, 12198, 3967, 2343, 1201, 740, 584, 47, 154, 139, 139, 70.


Latitude Bin	Mean CPUE
[37.8,37.85]	288
(37.85,37.9]	237
(37.9,37.95]	248
(37.95,38]	1917
(38,38.05]	7500
(38.05,38.1]	12198
(38.1,38.15]	3967
(38.15,38.2]	2343
(38.2,38.25]	1201
(38.25,38.3]	740
(38.3,38.35]	584
(38.35,38.4]	47
(38.4,38.45]	154
(38.45,38.5]	139
(38.5,38.55]	139
(38.55,38.6]	70

The heatmap displays the distribution of observations across years and latitudes. The x-axis represents the year from 1975 to 2021, and the y-axis represents latitude from 37.8 to 38.6. The color intensity indicates the number of observations (n) for each year-latitude combination. A color bar on the right side of the plot provides a scale for 'n', ranging from 0 (dark purple) to 500 (yellow). The data shows a high concentration of observations in the early years (1975-1993) at latitudes 38.05 and 38.1, with a notable peak in 2019 at latitude 38.05.


Longitude — distribution




Mean CPUE by Longitude bin (labels = n per bin)




Longitude coverage by year (blank = no data)




Source — counts



CPUE by Source (log1p scale)



Source coverage by year (blank = no data)



TotalGrabs — distribution

Count

TotalGrabs


Mean CPUE by TotalGrabs bin (labels = n per bin)

Mean CPUE


TotalGrabs

TotalGrabs coverage by year (blank = no data)


TotalGrabs




OUT — distribution




Mean CPUE by OUT bin (labels = n per bin)




OUT coverage by year (blank = no data)




X2 — distribution



Mean CPUE by X2 bin (labels = n per bin)



X2 coverage by year (blank = no data)






A histogram showing the distribution of Secchi Depth. The x-axis is labeled 'Secchi Depth' and ranges from 0 to 300. The y-axis is labeled 'Count' and ranges from 0 to 100. The distribution is right-skewed, with the highest frequency (count ~105) occurring between 25 and 50. The count decreases as Secchi Depth increases, with a few outliers at higher depths.

Secchi Depth Range	Count
0 - 10	2
10 - 20	42
20 - 30	105
30 - 40	80
40 - 50	90
50 - 60	70
60 - 70	60
70 - 80	38
80 - 90	8
90 - 100	10
100 - 110	2
110 - 120	2
120 - 130	4
130 - 140	3
140 - 150	2
150 - 160	9
160 - 170	3
170 - 180	2
180 - 190	2
190 - 200	2
200 - 210	4
210 - 220	0
220 - 230	1
230 - 240	0
240 - 250	2
250 - 260	0
260 - 270	0
270 - 280	0
280 - 290	0
290 - 300	0
300 - 310	1

A bar chart illustrating the relationship between Secchi Depth and Mean CPUE. The x-axis represents Secchi Depth in meters, with intervals of 20 meters from 0 to 300. The y-axis represents Mean CPUE, ranging from 0 to 3000. The chart shows a general trend of decreasing CPUE as Secchi depth increases, with a notable peak at 0-20m and a small increase at 300m.


Secchi Depth (m)	Mean CPUE
[0,20]	95
(20,40]	219
(40,60]	145
(60,80]	41
(80,100]	9
(100,120]	5
(120,140]	13
(140,160]	6
(160,180]	3
(180,200]	5
(200,220]	1
(220,240]	1
(240,260]	2
(300,320]	1

The heatmap displays the distribution of observations across years and Secchi depths. The x-axis represents the Year from 1975 to 2021, and the y-axis represents the Secchi Depth in 20m intervals from 0,20 to 300,320. The color scale indicates the number of observations (n) for each cell, ranging from 0 (dark purple) to 15 (yellow). The data shows a high concentration of observations in the shallowest depth bins (0,20 and 20,40) for the years 2005 through 2021, with some cells reaching a count of 15. The number of observations generally decreases as the Secchi depth increases, with very few observations recorded in the deeper bins (above 100,120).



A histogram showing the distribution of Dissolved Oxygen % Saturation. The x-axis is labeled 'Dissolved Oxygen % Saturation' and ranges from approximately 75 to 105. The y-axis is labeled 'Count' and ranges from 0.0 to 10.0. The distribution is unimodal and slightly right-skewed, with a peak count of 10.0 at approximately 95% saturation.

Dissolved Oxygen % Saturation (Bin Center)	Count
76	1
78	1
80	1
82	1
84	3
86	2
88	5
90	2
92	4
94	7
96	10
98	2
100	2
102	1
104	1




A histogram showing the distribution of Final Temperature. The x-axis is labeled 'Final Temperature' and ranges from approximately 8 to 28, with major ticks at 10, 15, 20, 25, and 30. The y-axis is labeled 'Count' and ranges from 0 to 2500, with major ticks at 0, 500, 1000, 1500, 2000, and 2500. The distribution is unimodal and slightly right-skewed, with the highest frequency (count of approximately 2500) occurring at a final temperature of 20. The counts decrease as the temperature moves away from this peak in both directions.


Final Temperature (approx.)	Count (approx.)
8	10
9	20
10	100
11	150
12	200
13	350
14	400
15	500
16	550
17	900
18	1100
19	1250
20	1300
21	1550
22	1750
23	1800
24	2300
25	2400
26	2300
27	1700
28	1450
29	950
30	650
31	400
32	300
33	180
34	100
35	50
36	20
37	10

Final Temperature	Mean CPU
[6,7]	15
(7,8]	78
(8,9]	224
(9,10]	327
(10,11]	578
(11,12]	881
(12,13]	934
(13,14]	1317
(14,15]	1885
(15,16]	1965
(16,17]	2178
(17,18]	2413
(18,19]	2648
(19,20]	3416
(20,21]	3980
(21,22]	3819
(22,23]	2487
(23,24]	1268
(24,25]	704
(25,26]	300
(26,27]	138
(27,28]	44
(28,29]	7


Final_Ch1 — distribution



Mean CPUE by Final_Ch1 bin (labels = n per bin)



Final_Ch1 coverage by year (blank = no data)




A histogram showing the frequency distribution of 'Final DO' values. The x-axis is labeled 'Final DO' and has major tick marks at 0, 10, and 20. The y-axis is labeled 'Count' and has major tick marks at 0, 200, 400, and 600. The histogram consists of 15 bars, each with a width of 1 unit. The distribution is unimodal and slightly right-skewed, with the highest frequency (count of approximately 750) occurring at a Final DO value of 8.5. The counts decrease as the Final DO value moves away from the peak, with a count of approximately 100 at 10.5 and near-zero counts at 11.5 and 12.5.


Final DO (Bin Center)	Count (Approximate)
5.5	10
6.0	30
6.5	150
7.0	480
7.5	750
8.0	480
8.5	300
9.0	70
9.5	20
10.0	10
10.5	5
11.0	2
11.5	1
12.0	1

Final DO	Mean CPUE
[0,1]	1
(2,3]	3
(3,4]	2
(4,5]	7
(5,6]	19
(6,7]	133
(7,8]	732
(8,9]	932
(9,10]	439
(10,11]	67
(11,12]	20
(12,13]	8
(13,14]	5800
(26,27]	1


Final_pH — distribution



Mean CPUE by Final_pH bin (labels = n per bin)



Final_pH coverage by year (blank = no data)



A histogram showing the distribution of 'Final_Turbidity'. The x-axis is labeled 'Final_Turbidity' and ranges from 0 to 600. The y-axis is labeled 'Count' and ranges from 0 to 4000. The distribution is highly right-skewed, with a peak count of approximately 4500 for the first bin (0-20). The count drops sharply for subsequent bins, with values around 2600, 1700, 700, 300, 150, 100, and 50 for the next seven bins respectively. The distribution tapers off to near zero for values above 100.

Final_Turbidity Bin	Count
0 - 20	2600
20 - 40	4500
40 - 60	1700
60 - 80	700
80 - 100	300
100 - 120	150
120 - 140	100
140 - 160	50
160 - 180	20
180 - 200	10

