


The gas phase reaction $\text{CO} + \text{H}_2\text{O} \leftrightarrow \text{CO}_2 + \text{H}_2$ takes place in an equilibrium reactor. The feed to the reactor enters at 500°C with a 1 mol/s of CO and 1 mol/s of H_2O and no CO_2 or H_2 . The equilibrium relation for the reaction is:

$$64 = \frac{y_{\text{CO}_2} y_{\text{H}_2}}{y_{\text{CO}} y_{\text{H}_2\text{O}}}$$

- (a) Calculate the species molar flow rates of the exit stream in mol/s.
 (b) Calculate the heat exchanged with the surroundings in kJ/s if the reactor is isothermal.
 (c) Estimate the product gas temperature in $^\circ\text{C}$ if the reactor is adiabatic. Give the temperature in the table that is closest to the answer. You do not need to interpolate.

Data is on the back of this sheet.



$$\text{CO}_{\text{out}} = \text{CO}_{\text{in}} - \dot{\xi}$$

$$\text{H}_2\text{O}_{\text{out}} = \text{H}_2\text{O}_{\text{in}} - \dot{\xi}$$

$$\text{CO}_{2,\text{out}} = \dot{\xi}$$

$$\text{H}_{2,\text{out}} = \dot{\xi}$$

4 unk 3 ind
eqns

Equilibrium rxn

$$64 = \frac{\dot{\xi} \cdot \dot{\xi}}{(\text{CO}_{\text{in}} - \dot{\xi})(\text{H}_2\text{O}_{\text{in}} - \dot{\xi})}$$

$$64 = \frac{(\dot{\xi})^2}{(1 - 2\dot{\xi} + (\dot{\xi})^2)} \rightarrow 64 - 128\dot{\xi} + 64\dot{\xi}^2 = \dot{\xi}^2 \rightarrow 64 - 128\dot{\xi} + 63\dot{\xi}^2 = 0$$

$$\frac{128 \pm \sqrt{(128)^2 - 4 \cdot 63 \cdot 64}}{126}$$

+ root too large ($\text{CO}/\text{H}_2\text{O}$ out)

$$- \text{root} = 0.888 \text{ mol/s} = \dot{\xi} \text{ or } \frac{8}{9} \frac{\text{mol}}{\text{s}}$$

$$\frac{8}{9} \frac{\text{mol}}{\text{s}}$$

Species	in ($\frac{\text{mol}}{\text{s}}$)	out ($\frac{\text{mol}}{\text{s}}$)
CO	1	$\frac{1}{9}$
H_2O	1	$\frac{1}{9}$
CO_2	—	$\frac{8}{9}$
H_2	—	$\frac{8}{9}$
Tot	2	2

Table B.8 Specific Enthalpies of Selected Gases: SI Units

$\hat{h}$ (kJ/mol)							
Reference state: Gas, $P_{ref} = 1$ atm, $T_{ref} = 25^\circ\text{C}$							
T	Air	O ₂	N ₂	H ₂	CO	CO ₂	H ₂ O
0	-0.72	-0.73	-0.73	-0.72	-0.73	-0.92	-0.84
25	0.00	0.00	0.00	0.00	0.00	0.00	0.00
100	2.19	2.24	2.19	2.16	2.19	2.90	2.54
200	5.15	5.31	5.13	5.06	5.16	7.08	6.01
300	8.17	8.47	8.12	7.96	8.17	11.58	9.57
400	11.24	11.72	11.15	10.89	11.25	16.35	13.23
500	14.37	15.03	14.24	13.83	14.38	21.34	17.01
600	17.55	18.41	17.39	16.81	17.57	26.53	20.91
700	20.80	21.86	20.59	19.81	20.82	31.88	24.92
800	24.10	25.35	23.86	22.85	24.13	37.36	29.05
900	27.46	28.89	27.19	25.93	27.49	42.94	33.32
1000	30.86	32.47	30.56	29.04	30.91	48.60	37.69
1100	34.31	36.07	33.99	32.19	34.37	54.33	42.18
1200	37.81	39.70	37.46	35.39	37.87	60.14	46.78
1300	41.34	43.38	40.97	38.62	41.40	65.98	51.47
1400	44.89	47.07	44.51	41.90	44.95	71.89	56.25
1500	48.45	50.77	48.06	45.22	48.51	77.84	61.09

b) Energy balance

species	$\dot{n}_{in} (\frac{\text{mol}}{\text{s}})$	$\hat{H}_i (\frac{\text{kJ}}{\text{mol}})$	$\dot{n}_{out} (\frac{\text{mol}}{\text{s}})$	$\hat{H}_e (\frac{\text{kJ}}{\text{mol}})$
CO	1	$\hat{H}_1 = -96.2$	$\frac{1}{9}$	$\hat{H}_1 = -96.2$
H ₂ O	1	$\hat{H}_2 = -224.8$	$\frac{1}{9}$	$\hat{H}_2 = -224.8$
CO ₂	—	—	$\frac{8}{9}$	$\hat{H}_3 = -372.2$
H ₂	—	—	$\frac{8}{9}$	$\hat{H}_4 = 13.92$

$$\hat{H}_1 = \int_{25+273}^{500+273} C_p(\text{CO})_g dT + \Delta \hat{H}_{F,i}^{\circ} = \left(\frac{1}{1000} \right) \int_{298}^{773} 30.87 - 0.02285T + 2.789 \cdot 10^{-5} T^2 - 1.272 \cdot 10^{-8} T^3 dT$$

Table B17 and B3

$$= 14.33 + (-110.53) = -96.2 \frac{\text{kJ}}{\text{mol}} \quad \div 1000$$

$$\hat{H}_2 = \int_{298}^{773} 32.24 + 1.924 \cdot 10^{-3} T + 1.055 \cdot 10^{-5} T^2 - 3.596 \cdot 10^{-9} T^3 dT - 241.83 = -224.8 \frac{\text{kJ}}{\text{mol}}$$

← $\div 1000$

$$\hat{H}_3 = \int_{298}^{773} 19.8 + 0.07344 T - 5.602 \cdot 10^{-5} T^2 + 1.7115 \cdot 10^{-8} T^3 dT - 393.5 = -372.2 \frac{\text{kJ}}{\text{mol}}$$

← $\div 1000$

$$\hat{H}_4 = \int_{298}^{773} 27.14 + 0.0093 T - 1.381 \cdot 10^{-5} T^2 + 7.645 \cdot 10^{-9} T^3 dT + 0 = 13.92 \frac{\text{kJ}}{\text{mol}}$$

← $\div 1000$

$$\dot{Q} = \Delta \dot{H} = \left(\left(1 - \frac{1}{9}\right)(-96.2) + \left(1 - \frac{1}{9}\right)(-224.8) + \left(-\frac{8}{9}\right)(-372.1) + \left(-\frac{8}{9}\right)(13.92) \right)$$

$$\dot{Q} = -33.05 \frac{\text{kJ}}{\text{s}}$$

C) on back

$$T_1 = 500^\circ\text{C} \quad \frac{1}{9} \cdot 14.38 + \frac{1}{9} \cdot 17.01 + \frac{8}{9} \cdot 21.34 + \frac{8}{9} \cdot 13.83$$

$$+ 33.25 \text{ (from B)} = 67.8$$

$T_2 =$ Estimate at 940°C (roughly $\times 2$ ~~at~~ T_1)

*I don't see your strategy
to solve for T adiabatic*