Elranatamab (Elrexfio®)

Elranatamab is a cancer drug. It is used to treat myeloma.

On this page

- What is elranatamab (Elrexfio®)?
- How elranatamab is given
- About side effects
- Common side effects
- Other important information
- How we can help

What is elranatamab (Elrexfio®)?

Elranatamab is also called Elrexfio®. It is a type of targeted therapy drug called a monoclonal antibody. It is also called a bi-specific antibody.

Elranatamab is used to treat a type of blood cancer called myeloma.

It is best to read this information with our general information about targeted therapy drugs and the type of cancer you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree (consent) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

How elranatamab is given

You have elranatamab as an injection under the skin (subcutaneous injection) of your tummy or thigh.

During a course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor
- specialist nurse
- specialist pharmacist.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your injection.

Your course of treatment

Elranatamab can cause a side effect called cytokine release syndrome or CRS. It can also affect the brain and nervous system, causing a side effect known as ICANS. There is more information about CRS and ICANS in our section about common side effects.

To reduce the risk of these side effects, you start on a low dose of elranatamab. You also have extra medicines 1 hour before your first 3 injections. These help prevent or reduce any reaction.

On day 1, you have your first injection with a low dose of elranatamab.

On day 4, you have a higher dose elranatamab injection.

On day 8, you have a full dose injection.

You then continue to have elranatamab once a week. After 24 weeks of treatment, you have it once every 2 weeks.

Your first 3 injections are usually given during a stay in hospital. This means your cancer team can monitor you closely for any side effects and treat them quickly. After your first treatments, you may be able to have elranatamab in a day unit or clinic as an outpatient. Your cancer team will explain what to expect.

You usually have elranatamab for as long as it is effective and side effects are manageable. Your cancer team will discuss your treatment plan with you.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Common side effects

Cytokine Release Syndrome (CRS)

This treatment makes the body release large amounts of cytokines. Cytokines are chemical messengers that help your immune system protect the body from infection and illness.

You may have flu-like symptoms. These can range from mild to severe. You may need treatment and close monitoring until the symptoms improve. Some people will be monitored and treated in an intensive care unit.

CRS usually happens within 2 days of a treatment. It is most likely to happen after your first, second or third injection. But it can happen later. If you have any of the following symptoms at any time during this treatment, contact the hospital straight away on the 24-hour number:

- a high temperature
- chills
- a fast heart rate
- dizziness or light-headedness
- muscle or joint pain
- diarrhoea
- feeling sick or being sick (vomiting)
- difficulty breathing or shortness of breath.

Effects on the brain and ICANS

This treatment can cause temporary effects on the brain. This is called neurotoxicity or immune effector cell-associated neurotoxicity syndrome (ICANS).

Side effects usually happen within 3 days of a treatment. They are most likely to happen after your first, second or third injection. But they can happen later. You may need treatment and close monitoring until the symptoms improve. Some people will be monitored and treated in an intensive care unit.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

You must not drive or use heavy machinery:

- for at least 48 hours after each of the first 3 injections
- or if you have any of the symptoms below.

If you have any of the following symptoms after you go home, contact the hospital straight away on the 24-hour number:

- difficulty speaking
- difficulty writing or doing fiddly tasks, such as fastening buttons or tying shoelaces
- drowsiness
- confusion or disorientation
- reduced consciousness (being less alert)
- memory loss
- tremors or shaking
- muscle weakness
- loss of balance
- headaches
- seizures (fits).

Risk of infection

This treatment can reduce the number of white blood cells and antibodies in your blood. White blood cells and antibodies fight infection. If they are low, you may be more likely to get an infection. A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. Your doctor may give you antibiotics and other drugs to take during this treatment. They may give you an injection or drip to help increase low antibody levels. These treatments may help reduce your risk of getting an infection.

It is still important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

- feeling shivery and shaking
- a sore throat
- a cough
- breathlessness
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

diarrhoea

- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

Your white blood cell count will usually return to normal before your next treatment. You will have a blood test before having more treatment. If your white blood cell count is low, your doctor may delay your treatment for a short time, until your cell count increases.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Feeling tired (fatigue)

Feeling tired is a common side effect of this treatment. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Injection site reaction

The skin in the area that was injected may become:

- painful

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

- itchy
- red, if you have white skin
- darker, if you have black or brown skin.

Having the injection in a different place each time can help with this.

Loss of appetite

This treatment can affect your appetite. Don’t worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Skin changes

This treatment can affect your skin. It might feel dry. You may develop a rash, which may be itchy. Always tell your doctor, nurse or pharmacist about any skin changes. They can give you advice or prescribe creams or medicines to help.

If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

Numb or tingling hands or feet (peripheral neuropathy)

This treatment may affect the nerves in your fingers and toes. This can cause numbness, tingling or pain in your hands or feet. This is called peripheral neuropathy. You might find it hard to do fiddly tasks such as fastening buttons or tying shoelaces.

If you have these symptoms, always tell your doctor, nurse or pharmacist. They sometimes need to change the drug or the dose of the drug. The symptoms usually improve slowly after treatment ends. But for some people they continue and are a long-term side effect of treatment.

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 October 2024 | Next review: 01 October 2026

This content is currently being reviewed. New information will be coming soon.

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026

use short sentences

- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as ‘you’ so that everyone feels included. Where clinically necessary we use the terms ‘men’ and ‘women’ or ‘male’ and ‘female’. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information here.

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Macmillan Support Line

The Macmillan Support Line is a free and confidential service for people living with and affected by cancer. If you need to talk, we'll listen.

Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.

Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don’t want to call.

Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.

Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/elranatamab · retrieved 26 October 2026