Nivolumab-relatlimab (Opdualag®)

Nivolumab-relatlimab (Opdualag®) is a cancer drug. It is used to treat melanoma.

On this page

- What is nivolumab-relatlimab (Opdualag®)?
- How nivolumab-relatlimab is given
- About side effects
- Side effects while treatment is being given
- Immune system side effects
- Other possible side effects
- Other important information
- How we can help

What is nivolumab-relatlimab (Opdualag®)?

Nivolumab-relatlimab is sometimes called Opdualag®. It is 1 treatment, but it contains 2 different drugs, nivolumab and relatlimab. These are both a type of immunotherapy drug called a checkpoint inhibitor.

Nivolumab-relatlimab is used to treat melanoma.

It is best to read this information with our general information about immunotherapy drugs and the type of cancer you have.

Your cancer team will talk to you about this treatment and the possible side effects before you agree (consent) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

How nivolumab-relatlimab is given

You usually have nivolumab-relatlimab at a day unit or clinic as an outpatient.

During a course of treatment, you will meet someone from your cancer team, such as a:

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

cancer doctor

- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your treatment.

You have nivolumab-relatlimab as a drip into a vein (intravenous infusion). You have it over 30 minutes.

Your nurse will give you nivolumab-relatlimab through 1 of the following:

- a cannula – a short, thin tube the nurse puts into a vein in the arm or hand
- a central line – a fine tube that goes under the skin of the chest and into a vein close by
- a PICC line – a fine tube that is put into a vein in the arm and goes up into a vein in your chest
- an implantable port (portacath) – a disc that is put under the skin on the chest or arm and goes into a vein in your chest.

Your course of treatment

You have nivolumab-relatlimab once every 4 weeks. Your doctor, nurse or pharmacist will discuss your treatment plan with you.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common. This treatment can make the immune system too active. This can cause side effects that may be serious if not treated quickly.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Do not try to treat side effects yourself. Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects.
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Delayed side effects

This treatment may also cause delayed side effects. These can start weeks, months, or sometimes more than a year after you finish treatment.

If you have symptoms or side effects that start during treatment or after treatment ends, contact the hospital straight away on the 24-hour number. Side effects can become serious if they are not treated.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Your patient alert card

They will also give you a card with information about your treatment. It is important to keep the card with you during treatment and after treatment ends. You should show it to any doctor or healthcare professional caring for you.

Side effects while treatment is being given

Some people may have side effects while they are having this treatment or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your treatment. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

Pain along the vein

This treatment can cause pain:

- at the place where the drip (infusion) is given
- along the vein.

If you feel pain, tell your nurse straight away. They can check the site. They may give the drug more slowly or flush it through with more fluid to reduce pain.

Immune system side effects

This treatment can make the immune system too active. This can cause the following side effects. Some of them are rare. But it is important to know about them.

Do not try to treat these side effects yourself. If you have these side effects during treatment or after it ends, always contact the hospital straight away on the 24-hour number.

Effects on the lungs

This treatment can cause inflammation of the lungs. This is called pneumonitis. Contact the hospital straight away on the 24-hour number if you notice any of these changes during treatment or after it ends:

- breathlessness
- a cough that does not go away
- wheezing
- a fever, with a temperature over 37.5°C.

You should also tell them if any existing breathing problems get worse. You may have:

- tests to check your lungs
- treatment, if needed.

Diarrhoea or tummy pain

You may have diarrhoea or tummy (abdominal) pain. Sometimes it can cause serious inflammation of the bowel called colitis.

Contact the hospital straight away on the 24-hour number if you have any of the following symptoms during treatment, or after treatment ends:

- passing more stools (poo) than is normal for you
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

passing stools at night

- if you have a stoma, it is more active or needs emptied more often than usual
- watery or loose stools
- uncomfortable tummy cramps or pain
- a temperature over 37.5°C
- blood or mucus (slime) in your stool.

Effects on the liver

This treatment may affect how your liver works. It can cause liver inflammation called hepatitis. You will have regular blood tests to check how your liver is working. Tell your doctor, nurse or pharmacist if you have loss of appetite.

Sometimes liver changes can be serious. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- yellow skin or eyes
- feeling very sleepy
- dark urine (pee)
- unexplained bleeding or bruising
- pain in the right side of your tummy (abdomen).

If you have black or brown skin, changes in your skin colour may be less noticeable. You might first notice the whites of your eyes look yellow, or that the colour of your urine has changed.

Effects on the kidneys

This treatment can affect how the kidneys work. It may cause kidney inflammation called nephritis. You will have regular blood tests to check how your kidneys are working.

Contact the hospital straight away on the 24-hour number during treatment, or after it ends, if you:

- have blood in your urine (pee)
- are passing less urine or peeing less often than usual.

Hormone changes

This treatment may permanently affect glands in the body that produce hormones. Hormones control different processes in the body. You will have regular blood tests to check some hormone levels, such as your thyroid. You may need to take tablets daily to correct your hormone levels.

Contact the hospital on the 24-hour number if you have any of these side effects during treatment or after it ends:

- increased sweating
- problems with constipation
- weight gain or loss
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

dizziness or fainting

- feeling or being sick
- feeling more hungry or thirsty than usual
- loss of sex drive
- passing urine (peeing) more often than usual
- headaches that do not go away
- feeling more tired than usual (fatigue)
- mood changes
- changes to your heartbeat (palpitations).

Raised blood sugar levels

This treatment may raise your blood sugar levels and increase the risk of diabetes.

You will have regular blood tests to check this.

Contact the hospital on the 24-hour number if you have these symptoms during treatment or after it ends:

- feeling thirsty all the time
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.
- losing weight without trying.

If you have diabetes, your blood sugar levels may be higher than usual. You may need advice and support from your GP or diabetes team about managing this.

Muscle and joint pain

This treatment can cause sore or weak muscles. Your joints may also be stiff, sore or swollen. This is called arthritis. If you already have problems with joint pain, this may become worse.

Contact the hospital on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- new or worsening aches or pains
- new or worsening stiffness in a joint
- swollen joints or muscle weakness.

Skin changes

This treatment may cause inflammation in the skin. This is usually mild, but if it is not treated it can become serious. You may need creams, steroids or other medicines to treat skin problems.

Contact the hospital on the 24-hour number as soon as possible, if you notice any skin changes during treatment, or after treatment ends. This includes:

- a rash or bumps on the skin

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

- dry, tight or itchy skin
- areas that are a different colour to your usual skin tone.

Here are some things you can do to take care of your skin during treatment:

- Wash with lukewarm water and mild, unperfumed, soap-free cleansers.
- Moisturise your skin regularly with unperfumed moisturisers for dry skin conditions.
- Use sun cream of at least SPF 30 to protect your skin from the sun. Cover up with clothing and a hat.

Contact the hospital straight away on the 24-hour number if you have any of these serious skin symptoms:

- a rash that is spreading
- swollen, blistering or peeling skin
- flu-like symptoms, such as a high temperature and joint pain.

Effects on the heart

This treatment can affect how the heart works. You may have tests to check how well your heart is working. These may be done before, during and after treatment.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.
- flu-like symptoms, such as body aches, pains and tiredness.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call **999** if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Eyesight changes

This treatment can affect your eyes and eyesight. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- dry, itchy, painful, or swollen eyes
- eyes that are sensitive to light
- changes to your eyesight.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Effects on the pancreas

This treatment can cause inflammation of the pancreas (pancreatitis). You will have regular blood tests to check your pancreas.

Contact the hospital straight away on the 24-hour telephone number you have been given if you have any of these symptoms:

- severe pain in the centre of your stomach area (upper abdomen)
- feeling or being sick
- indigestion or heartburn
- a high temperature of 37.5°C or more (fever)
- fast heartbeat (tachycardia) or rapid breathing
- yellow skin or eyes
- tenderness or swelling of the tummy.

Effects on the nervous system

Rarely, this treatment can affect the brain and nerves. These effects may be mild at first but can become serious if not treated. Symptoms may include:

- weakness, numbness or tingling in your arms, legs or face
- a headache that does not get better
- seeing double or having drooping eyelids
- difficulty turning or bending your nec
- drowsiness or confusion
- problems with moving or speaking
- fits (seizures).

If you have any of these symptoms during treatment or after it ends, it is important to either:

- contact the hospital straight away on the 24-hour number
- go to the hospital straight away
- call **999**.

Do not drive yourself to hospital.

Other possible side effects

Feeling tired (fatigue)

Feeling tired is a common side effect of this treatment. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Sometimes, this treatment can affect glands that make hormones. This can make you feel tired. Contact the hospital on the 24-hour number if either of these things happen during treatment or after it ends:

- your tiredness gets worse or stops you doing your usual daily activities
- you feel weak or shaky.

If you feel sleepy, do not drive or use machinery.

Low number of white blood cells

This treatment can reduce the number of white cells in your blood. These cells fight infection. If you have any signs of an infection, tell your doctor, nurse or pharmacist. These signs include:

- a cough
- a sore throat
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Constipation

This treatment can cause constipation. Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- Eat high-fibre foods, such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Feeling sick

You may feel sick during this treatment, but this is usually mild. Your doctor, nurse or pharmacist may prescribe you anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to. If you feel sick, take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids.

If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice.

Loss of appetite

This treatment can affect your appetite. Don’t worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Changes to your taste

Some foods may taste different or have no taste. Try different foods to find out what tastes best to you. You may also get a bitter or metallic taste in your mouth. Your doctor, nurse or pharmacist can give you advice.

Sore mouth and throat

This treatment may cause a sore mouth and throat. You may also get mouth ulcers. This can make you more likely to get a mouth or throat infection. Use a soft toothbrush to clean your teeth or dentures in the morning, at night and after meals.

Contact the hospital straight away on the 24-hour number, if:

-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

a sore mouth or throat affects how much you can drink or eat

- your mouth, tongue, throat or lips have any blisters, ulcers or white patches.

They can give you advice, and mouthwash or medicines to help with the pain or to treat any infection. Follow their advice and make sure you:

- drink plenty of fluids
- avoid alcohol and tobacco
- avoid food or drinks that irritate your mouth and throat.

Hair loss

You might notice that your hair slowly gets thinner. But you are very unlikely to lose all the hair from your head. Your hair can also become more brittle. Your nurse can give you information about coping with hair loss. Your hair will usually go back to normal after treatment ends.

Fluid build-up

This treatment can cause a build-up of fluid in the body. This will slowly get better after treatment ends. Contact the hospital on the 24-hour number if you:

- are gaining weight
- have swelling in your face, legs or ankles.

They can give you advice and treatment to help.

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Show them your patient alert card. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist or optician, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 October 2024 | Next review: 01 October 2026

This content is currently being reviewed. New information will be coming soon.

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026

use short sentences

- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as ‘you’ so that everyone feels included. Where clinically necessary we use the terms ‘men’ and ‘women’ or ‘male’ and ‘female’. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information here.

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Macmillan Support Line

The Macmillan Support Line is a free and confidential service for people living with and affected by cancer. If you need to talk, we'll listen.

Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.

Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don’t want to call.

Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.

Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/nivolumab-relatlimab · retrieved 26 October 2026