Daratumumab (Darzalex®)

Daratumumab is a cancer drug. It is used to treat myeloma.

On this page

- What is daratumumab (Darzalex®)?
- How daratumumab is given
- About side effects
- Side effects while treatment is being given
- Very common side effects
- Other side effects
- Other important information
- How we can help

What is daratumumab (Darzalex®)?

Daratumumab is also called Darzalex®. It is used to treat a type of blood cancer called myeloma. It is a type of targeted therapy drug called a monoclonal antibody.

It is best to read this information with our general information about targeted therapy drugs and the type of cancer you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree (consent) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

How daratumumab is given

You usually have daratumumab in a day unit or clinic as an outpatient. Sometimes you have it during a stay in hospital. You may have it on its own or with other cancer drugs.

During a course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your targeted therapy.

You will have medicines before and after the daratumumab to reduce the chance of an allergic reaction. After the first treatment, you may be asked to wait in the hospital for 4 to 6 hours. This is to check for any signs of a reaction to the treatment.

You usually have daratumumab as an injection under the skin (subcutaneous). Your nurse will give this into your tummy area (abdomen). The injection takes 3 to 5 minutes. You have it in different areas of the tummy each time. This can help stop your skin from becoming sore.

Sometimes daratumuab is given into a vein as a drip (intravenous infusion) through 1 of the following:

- a cannula – a short, thin tube the nurse puts into a vein in the arm or hand
- a central line – a fine tube that goes under the skin of the chest and into a vein close by
- a PICC line – a fine tube that is put into a vein in the arm and goes up into a vein in the chest
- an implantable port (portacath) – a disc that is put under the skin on the chest or arm and goes into a vein in the chest.

Your course of treatment

To begin with, you will have daratumumab once a week. Then you usually have it once every 2 weeks. Or sometimes it is given every 3 weeks instead. This may depend on if you are having it with other cancer drugs. Eventually you only have it once a month. Your cancer team will discuss your treatment plan with you.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Side effects while treatment is being given

Some people may have side effects while they are having this treatment or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. Before treatment, you will have medicines to help prevent or reduce any reaction.

Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your treatment. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

Problems at the injection site

If you are having daratumumab as an injection in the tummy, the skin in the area that was injected may become:

- painful
- swollen
- red, if you have white skin
- darker, if you have black or brown skin.

Tell your doctor, nurse or pharmacist if this happens. They will give the injection in a different areas each time.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Risk of infection

This treatment can reduce the number of white blood cells in your blood. These cells fight infection. If your white blood cell count is low, you may be more likely to get an infection. A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. It is important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

- feeling shivery and shaking
- a sore throat
- a cough
- breathlessness
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

Your white blood cell count will usually return to normal before your next treatment. You will have a blood test before having more treatment. If your white blood cell count is low, your doctor may delay your treatment for a short time, until your cell count increases.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Feeling sick

You may feel sick during this treatment, but this is usually mild. Your doctor, nurse or pharmacist may prescribe you anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to. If you feel sick, take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids.

If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice.

Diarrhoea

This treatment may cause diarrhoea. Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Constipation

This treatment can cause constipation. Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- Eat high-fibre foods, such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Tummy pain

You may get pain in your tummy (abdomen), or have indigestion. Your doctor, nurse or pharmacist can give you advice or treatment to help. Contact the hospital straight away on the 24-hour number if your symptoms do not go away, or get worse.

Feeling tired

Feeling tired is a common side effect of this treatment. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Muscle or joint pain

You may get pain in your muscles or joints for a few days after treatment. If this happens, tell your doctor, nurse or pharmacist. They can give you painkillers and advice. They can also tell you if any of the painkillers you usually take are suitable.

Tell them if the pain does not get better. Having warm baths and resting regularly may help.

Effects on the lungs

This treatment can cause changes to the lungs. Tell your doctor, nurse or pharmacist if you develop:

- a cough that does not go away
- wheezing
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

breathlessness.

You should also tell them if any existing breathing problems get worse. You may have tests to check your lungs.

Numb or tingling hands or feet (peripheral neuropathy)

This treatment may affect the nerves in your fingers and toes. This can cause numbness, tingling or pain in your hands or feet. This is called peripheral neuropathy. You might find it hard to do fiddly tasks such as fastening buttons or tying shoelaces.

If you have these symptoms, always tell your doctor, nurse or pharmacist. They sometimes need to change the drug or the dose of the drug. The symptoms usually improve slowly after treatment ends. But for some people they continue and are a long-term side effect of treatment.

Headaches

This treatment may cause headaches. If you have headaches, tell your doctor, nurse or pharmacist. They can give you advice about painkillers that may help. Tell them if the headache does not get better, or gets worse.

Fluid build-up

This treatment can cause a build-up of fluid in the body. This will slowly get better after treatment ends. Contact the hospital on the 24-hour number if you:

- are gaining weight
- have swelling in your face, legs or ankles.

They can give you advice and treatment to help.

Loss of appetite

This treatment can affect your appetite. Don’t worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Skin changes

This treatment can affect your skin. It may cause a rash, which may be itchy. Your doctor or nurse can tell you what to expect. If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

Always tell your doctor or nurse about any skin changes. They can give you advice and may prescribe creams or medicines to help. Changes to your skin usually improve when treatment ends.

Difficulty sleeping (insomnia)

Sometimes this treatment can affect your sleep. Tell your doctor, nurse or pharmacist if cancer treatment makes it difficult to sleep.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Changes in blood pressure

Your blood pressure may become higher than normal. Tell your nurse or doctor if you have headaches or feel dizzy. These can be signs of blood pressure changes.

Effects on the heart

This treatment can affect how the heart works. You may have tests to check how well your heart is working. These may be done before, during and after treatment.

If the treatment is causing heart problems, your doctor may change the type of treatment you are having.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call **999** if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Raised blood sugar levels

This treatment may raise your blood sugar levels and increase the risk of diabetes.

You will have regular blood tests to check this.

Contact the hospital on the 24-hour number if you have these symptoms during treatment or after it ends:

- feeling thirsty all the time
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.
- losing weight without trying.

If you have diabetes, your blood sugar levels may be higher than usual. You may need advice and support from your GP or diabetes team about managing this.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

Hepatitis B reactivation

If you have had a liver infection called hepatitis B, this treatment can make it active again. Your doctor, nurse or pharmacist will talk to you about this. You may have tests for hepatitis B before, during, and after treatment.

Shingles

You may be more likely to develop a viral infection called shingles during this treatment. Your doctor may give you medicine to help prevent shingles.

Blood transfusions

Daratumumab can affect the result of a blood test called the indirect Coombs test. This is one of the tests that matches you to a donor's blood before you have a blood transfusion.

Before you start daratumumab, you will have the blood tests you need for a blood transfusion. Your doctor may also give you a card to carry that explains the treatment you are having.

If you need a blood transfusion during treatment or for 6 months after, tell the person arranging it that you have had daratumumab. You can show them the card your doctor has given you.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 December 2025 | Next review: 01 December 2027

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as 'you' so that everyone feels included. Where clinically necessary we use the terms 'men' and 'women' or 'male' and 'female'. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026

You can read more about how we produce our information here.

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Macmillan Support Line

The Macmillan Support Line is a free and confidential service for people living with and affected by cancer. If you need to talk, we'll listen.

Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.

Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don’t want to call.

Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.

Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/daratumumab · retrieved 19 October 2026