Durvalumab (Imfinzi®) and tremelimumab (Imjudo®)

Durvalumab (Imfinzi®) and tremelimumab (Imjudo®) is a combination of cancer drugs used to treat liver cancer.

On this page

- What is durvalumab and tremelimumab?
- How durvalumab and tremelimumab works
- How durvalumab and tremelimumab is given
- About side effects
- Side effects while treatment is being given
- Immune system side effects
- Other side effects (very common)
- Other side effects (less common)
- Other important information
- Get in touch

What is durvalumab and tremelimumab?

Durvalumab is also called Imfinzi®. Tremelimumab is also called Imjudo®. They are both a type of immunotherapy drug called a checkpoint inhibitor.

Durvalumab and tremelimumab is a combination of cancer drugs used to treat liver cancer. It may sometimes be used to treat other cancers.

It is best to read this information with our general information about immunotherapy drugs and the type of cancer you have.

Your cancer team will talk to you about this treatment and the possible side effects before you agree (consent) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

How durvalumab and tremelimumab works

The immune system protects the body against illness and infection. Immunotherapy drugs use the immune system to find and attack cancer cells.

Checkpoint inhibitors affect a type of white blood cell called a lymphocyte. Lymphocytes are an important part of the immune system. When they are active, they can attack other cells, including cancer cells. But if they receive a certain signal from the other cell, they switch off (become inactive) and do not attack it.

Checkpoint inhibitors block the signals that switch off lymphocytes. They do this by attaching to either the cancer cell or the lymphocyte. This means the lymphocyte stays active and can attack the cancer cell.

How durvalumab and tremelimumab is given

You usually have durvalumab and tremelimumab at a day unit or clinic as an outpatient.

During a course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor
- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your treatment.

First you have tremelimumab as a drip into a vein (intravenous infusion). You have it over one hour. You then have durvalumab as an intravenous infusion over one hour.

Your nurse will give you the drugs through one of the following:

- a cannula – a short, thin tube the nurse puts into a vein in your arm or hand
- a central line – a fine tube that goes under the skin of your chest and into a vein close by
- a PICC line – a fine tube that is put into a vein in your arm and goes up into a vein in your chest
- an implantable port (portacath) – a disc that is put under the skin on your chest or arm and goes into a vein in your chest.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Your course of treatment

You have tremelimumab with durvalumab on day one. 4 weeks later, you have durvalumab on its own. You have durvalumab on its own every 4 weeks for as long as it is working for you, and you are able to cope with the side effects.

Your doctor, nurse or pharmacist will discuss your treatment plan with you.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common. This treatment can make the immune system too active. This can cause side effects that may be serious if not treated quickly.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Do not try to treat side effects yourself. Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Delayed side effects

This treatment may also cause delayed side effects. These can start weeks, months, or sometimes more than a year after you finish treatment.

If you have symptoms or side effects that start during treatment or after treatment ends, contact the hospital straight away on the 24-hour number. Side effects can become serious if they are not treated.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Your patient alert card

They will also give you a card with information about your treatment. It is important to keep the card with you during treatment and after treatment ends. You should show it to any doctor or healthcare professional caring for you.

Side effects while treatment is being given

Some people may have side effects while they are having this treatment or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your treatment. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

Immune system side effects

This treatment can make the immune system too active. This can cause the following side effects. Some of them are rare. But it is important to know about them.

Do not try to treat these side effects yourself. If you have these side effects during treatment or after it ends, always contact the hospital straight away on the 24-hour number.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Hormone changes

This treatment may permanently affect glands in the body that produce hormones. Hormones control different processes in the body. You will have regular blood tests to check some hormone levels, such as your thyroid. You may need to take tablets daily to correct your hormone levels.

Contact the hospital on the 24-hour number if you have any of these side effects during treatment or after it ends:

- increased sweating
- problems with constipation
- weight gain or loss
- dizziness or fainting
- feeling or being sick
- feeling more hungry or thirsty than usual
- loss of sex drive
- passing urine (peeing) more often than usual
- headaches that do not go away
- feeling more tired than usual (fatigue)
- mood changes
- changes to your heartbeat (palpitations).

Effects on the lungs

This treatment can cause inflammation of the lungs. This is called pneumonitis. Contact the hospital straight away on the 24-hour number if you notice any of these changes during treatment or after it ends:

- breathlessness
- a cough that does not go away
- wheezing
- a fever, with a temperature over 37.5°C.

You should also tell them if any existing breathing problems get worse. You may have:

- tests to check your lungs
- treatment, if needed.

Diarrhoea or tummy pain

You may have diarrhoea or tummy (abdominal) pain. Sometimes it can cause serious inflammation of the bowel called colitis.

Contact the hospital straight away on the 24-hour number if you have any of the following symptoms during treatment, or after treatment ends:

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

- passing more stools (poo) than is normal for you
- passing stools at night
- if you have a stoma, it is more active or needs emptied more often than usual
- watery or loose stools
- uncomfortable tummy cramps or pain
- a temperature over 37.5°C
- blood or mucus (slime) in your stool.

Skin changes

This treatment may cause inflammation in the skin. This is usually mild, but if it is not treated it can become serious. You may need creams, steroids or other medicines to treat skin problems.

Contact the hospital on the 24-hour number as soon as possible, if you notice any skin changes during treatment, or after treatment ends. This includes:

- a rash or bumps on the skin
- dry, tight or itchy skin
- areas that are a different colour to your usual skin tone.

Here are some things you can do to take care of your skin during treatment:

- Wash with lukewarm water and mild, unperfumed, soap-free cleansers.
- Moisturise your skin regularly with unperfumed moisturisers for dry skin conditions.
- Use sun cream of at least SPF 30 to protect your skin from the sun. Cover up with clothing and a hat.

Contact the hospital straight away on the 24-hour number if you have any of these serious skin symptoms:

- a rash that is spreading
- swollen, blistering or peeling skin
- flu-like symptoms, such as a high temperature and joint pain.

Effects on the liver

This treatment may affect how your liver works. It can cause liver inflammation called hepatitis. You will have regular blood tests to check how your liver is working. Tell your doctor, nurse or pharmacist if you have loss of appetite.

Sometimes liver changes can be serious. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- yellow skin or eyes
- feeling very sleepy
- dark urine (pee)
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

unexplained bleeding or bruising

- pain in the right side of your tummy (abdomen).

If you have black or brown skin, changes in your skin colour may be less noticeable. You might first notice the whites of your eyes look yellow, or that the colour of your urine has changed.

Effects on the kidneys

This treatment can affect how the kidneys work. It may cause kidney inflammation called nephritis. You will have regular blood tests to check how your kidneys are working.

Contact the hospital straight away on the 24-hour number during treatment, or after it ends, if you:

- have blood in your urine (pee)
- are passing less urine or peeing less often than usual.

Muscle and joint pain

This treatment can cause sore or weak muscles. Your joints may also be stiff, sore or swollen. This is called arthritis. If you already have problems with joint pain, this may become worse.

Contact the hospital on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- new or worsening aches or pains
- new or worsening stiffness in a joint
- swollen joints or muscle weakness.

Effects on the heart

This treatment can affect how the heart works. You may have tests to check how well your heart is working. These may be done before, during and after treatment.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.
- flu-like symptoms, such as body aches, pains and tiredness.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call **999** if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Effects on the nervous system

Rarely, this treatment can affect the brain and nerves. These effects may be mild at first but can become serious if not treated. Symptoms may include:

- weakness, numbness or tingling in your arms, legs or face
- a headache that does not get better
- seeing double or having drooping eyelids
- difficulty turning or bending your nec
- drowsiness or confusion
- problems with moving or speaking
- fits (seizures).

If you have any of these symptoms during treatment or after it ends, it is important to either:

- contact the hospital straight away on the 24-hour number
- go to the hospital straight away
- call **999**.

Do not drive yourself to hospital.

Effects on the pancreas

This treatment can cause inflammation of the pancreas. This is called pancreatitis. You will have regular blood tests to check your pancreas.

Contact the hospital straight away on the 24-hour telephone number you have been given if you have any of these symptoms:

- severe pain in the centre of your tummy (abdomen)
- feeling or being sick (vomiting)
- indigestion
- a temperature above 37.5°C
- fast heartbeat (tachycardia) or rapid breathing
- yellow skin or eyes – this may be harder to notice if you have black or brown skin
- tenderness or swelling of the tummy.

Eyesight changes

This treatment can affect your eyes and eyesight. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- dry, itchy, painful, or swollen eyes
- eyes that are sensitive to light
- changes to your eyesight.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Raised blood sugar levels

This treatment may raise your blood sugar levels and increase the risk of diabetes.

You will have regular blood tests to check this.

Contact the hospital on the 24-hour number if you have these symptoms during treatment or after it ends:

- feeling thirsty all the time
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.
- losing weight without trying.

If you have diabetes, your blood sugar levels may be higher than usual. You may need advice and support from your GP or diabetes team about managing this.

Other side effects (very common)

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Fluid build-up

This treatment can cause a build-up of fluid in the body. This will slowly get better after treatment ends. Contact the hospital on the 24-hour number if you:

- are gaining weight
- have swelling in your face, legs or ankles.

They can give you advice and treatment to help.

Other side effects (less common)

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Pain passing urine

This treatment can cause inflammation of the bladder. Contact the hospital straight away on the 24-hour number if you have discomfort or pain when you pass urine (pee). You may have a test to check for infection. You may need steroids or other treatments.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Sore throat, colds and cough

You are more likely to get infections such as colds, sore throats or sinus problems with this treatment. It may also cause a cough.

Contact the hospital straight away if you have:

- a cough
- breathlessness
- cold symptoms
- a sore throat, hoarseness or blocked sinuses.

Mouth problems

You may get mouth problems, such as a sore mouth or infection. Always let your doctor or nurse know if you have any problems with your mouth, including dental problems. They can give you mouthwashes, or medicines and gels to heal ulcers and clear or prevent any infection. Your nurse will explain how to look after your mouth to reduce the risk of problems.

The most common mouth infection is called thrush (or candidiasis). It shows as either:

- white spots on your mouth and tongue
- your tongue and mouth lining become red and swollen.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

throbbing pain or swelling in a leg or arm

- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Show them your patient alert card. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist or optician, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 March 2026 | Next review: 01 March 2028

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as ‘you’ so that everyone feels included. Where clinically necessary we use the terms ‘men’ and ‘women’ or ‘male’ and ‘female’. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information here.

Get in touch

0300 1000 200

Our Support Line is available Monday to Friday, 9:30am-4:30pm

Email us

Email the Macmillan Fundraising team if you have any questions.

Fundraising FAQs

Answers to common questions our supporters have about supporting Macmillan – how to pay in money, changing your details, making a donation.

Fundraising tips

Get top tips and advice from Macmillan on how to maximise your fundraising.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/durvalumab-and-tremelimumab · retrieved 19 October 2026