Rituximab

Rituximab is a cancer drug. It is used to treat lymphoma and chronic lymphocytic leukaemia (CLL). It may sometimes be used to treat acute lymphoblastic leukaemia (ALL).

On this page

- What is rituximab?
- How rituximab is given
- About side effects
- Side effects while treatment is being given
- Very common side effects
- Other side effects
- Other information
- How we can help

What is rituximab?

Rituximab is a type of targeted therapy drug called a monoclonal antibody.

Rituximab is used to treat lymphoma and chronic lymphocytic leukaemia (CLL). It may sometimes be used to treat acute lymphoblastic leukaemia (ALL) and other cancers.

It is best to read this information with our general information about targeted therapy and the type of cancer you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree (consent) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

How rituximab is given

You usually have rituximab in a day unit or clinic as an outpatient, or during a stay in hospital.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

During a course of treatment, you will meet someone from your cancer team, such as a:

- haematologist – a doctor who diagnoses and treats blood disorders and cancers.
- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your targeted therapy.

You have rituximab as a drip into a vein (intravenous infusion). Some people may have it as an injection under the skin (subcutaneously). You can only have rituximab as an injection if you have already had at least one treatment as an intravenous infusion.

You may have it with other cancer drugs.

As a drip

If you are having rituximab into a vein, your nurse will give it to you through 1 of the following:

- a cannula – a short, thin tube the nurse puts into a vein in your arm or hand
- a central line – a fine tube that goes under the skin of your chest and into a vein close by
- a PICC line – a fine tube that is put into a vein in your arm and goes up into a vein in your chest
- an implantable port (portacath) – a disc that is put under the skin on your chest or arm and goes into a vein in your chest.

You will be given your treatment through a pump over a set time.

The first infusion will be given slowly because it can cause a reaction. Before treatment, you will be given medicines to help prevent or reduce any reaction. You may have the first infusion in hospital.

After the first treatment, you can usually have rituximab in the day unit or outpatient department. You can also have it over shorter periods of time.

As an injection under the skin

If you are having rituximab as a subcutaneous injection, the nurse will inject your treatment into your abdomen (tummy). It will be given slowly, over about 5 minutes.

Your course of treatment

You usually have a course of several cycles of treatment over a few months. Your cancer team will discuss your treatment plan with you.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Hepatitis B reactivation

If you have had hepatitis B (a liver infection) in the past, rituximab can make it active again. Your doctor or nurse will talk to you about this and test you for hepatitis B.

If your test shows that you have had it in the past you may be given treatment to prevent it becoming active again.

If you have active hepatitis B, you will not have treatment with rituximab.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Side effects while treatment is being given

Some people may have side effects while they are having this treatment or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. The first infusion is the most likely to cause a reaction, so it is usually given more slowly than later treatments. Before treatment, you will have medicines to help prevent or reduce any reaction.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your infusion. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

Blood pressure

Your blood pressure may fall when you are having rituximab. If you usually take medicine to lower your blood pressure, your doctor, nurse or pharmacist may ask you not to take it for 12 hours before having rituximab. Sometimes, rituximab can make your blood pressure go up. Your nurse will check your blood pressure regularly.

Tumour pain

During the rituximab infusion, some people have mild pain in the parts of the body where they have cancer. Painkillers can help with this.

Problems at the injection site

If you have rituximab as an injection, you may have some redness and swelling where it is given (injection site). Your nurse can give you advice on coping with this.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Risk of infection

This treatment can reduce the number of white blood cells in your blood. These cells fight infection. If your white blood cell count is low, you may be more likely to get an infection. A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. It is important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

- feeling shivery and shaking
- a sore throat
- a cough
- breathlessness
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

Your white blood cell count will usually return to normal before your next treatment. You will have a blood test before having more treatment. If your white blood cell count is low, your doctor may delay your treatment for a short time, until your cell count increases.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Feeling sick

Your doctor, nurse or pharmacist will prescribe anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to, even if you do not feel sick. It is easier to prevent sickness than to treat it after it has started.

If you feel sick, take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids. If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice. They may change your anti-sickness treatment. Let them know if you still feel sick.

Skin changes

This treatment may cause inflammation in the skin. This is usually mild, but if it is not treated it can become serious. You may need creams, steroids or other medicines to treat skin problems. If you notice skin changes during treatment, or after treatment ends, contact the hospital on the 24-hour number as soon as possible. Skin changes can include:

- a rash or bumps on the skin
- dry or itchy skin
- patches of white or paler skin.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms:

- a rash that is spreading
- blistering or peeling skin
- flu-like symptoms, such as a high temperature and joint pain.

These symptoms can be a sign of a serious skin reaction that needs to be treated immediately in hospital.

Hair loss

Your hair will get thinner. Or you may lose all the hair from your head. You may also lose your eyelashes and eyebrows, as well as other body hair. Hair loss usually starts after your first or second treatment.

If you want to cover up hair loss, there are different ways you can do this. Your nurse will give you information about coping with hair loss.

Remember to protect your skin from the sun. Use suncream with a sun protection factor (SPF) of at least 30 on your scalp. Or cover up with a hat or scarf.

Hair loss is almost always temporary. Your hair will usually grow back after treatment ends.

Effects on the lungs

This treatment can cause changes to the lungs. Tell your doctor, nurse or pharmacist if you develop:

- a cough that does not go away
- wheezing
- breathlessness.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

You should also tell them if any existing breathing problems get worse. You may have tests to check your lungs.

Effects on the heart

This treatment can affect how the heart works. You may have tests to check how well your heart is working. These may be done before, during and after treatment.

If the treatment is causing heart problems, your doctor may change the type of treatment you are having.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call **999** if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Raised blood sugar levels

This treatment may raise your blood sugar levels and increase the risk of diabetes.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

You will have regular blood tests to check this.

Contact the hospital on the 24-hour number if you have these symptoms during treatment or after it ends:

- feeling thirsty all the time
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.
- losing weight without trying.

If you have diabetes, your blood sugar levels may be higher than usual. You may need advice and support from your GP or diabetes team about managing this.

Muscle or joint pain

You may get pain in your muscles or joints for a few days after treatment. If this happens, tell your doctor, nurse or pharmacist. They can give you painkillers and advice. They can also tell you if any of the painkillers you usually take are suitable.

Tell them if the pain does not get better. Having warm baths and resting regularly may help.

Fluid build-up

This treatment can cause a build-up of fluid in the body. This will slowly get better after treatment ends. Contact the hospital on the 24-hour number if you:

- are gaining weight
- have swelling in your face, legs or ankles.

They can give you advice and treatment to help.

Constipation

This treatment can cause constipation. Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- Eat high-fibre foods, such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Diarrhoea

This treatment may cause diarrhoea. Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Loss of appetite

This treatment can affect your appetite. Don’t worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Changes to your taste

Some foods may taste different or have no taste. Try different foods to find out what tastes best to you. You may also get a bitter or metallic taste in your mouth. Your doctor, nurse or pharmacist can give you advice. It might help to try:

- sucking sugar-free sour or boiled sweets
- eating cold foods
- eating sharp-tasting fresh fruit.

Taste changes usually get better after treatment ends. We have more information about coping with changes to taste.

Hearing changes

This treatment may cause hearing changes, including hearing loss. You may have ringing in the ears. This is called tinnitus. You may also become unable to hear some high-pitched sounds. Hearing changes may get better after this treatment ends. But this does not always happen. If you notice any changes in your hearing, tell your doctor, nurse or pharmacist.

Effects on the eyes

Your eyes may become watery and feel sore. Your doctor, nurse or pharmacist can give you advice and eye drops to help with this.

Contact the hospital on the 24-hour number if:

- your eyes get red and inflamed (conjunctivitis)
- you have pain in your eyes
- you notice any change in your vision.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Mood changes or difficulty sleeping

You may feel low or depressed, or have mood swings, during this treatment. Talking to family and friends about how you feel might help. If mood changes last for more than a few weeks, tell your doctor, nurse or pharmacist. They can talk to you about different ways to manage low mood or depression.

Sometimes this treatment can affect your sleep. Tell your doctor, nurse or pharmacist if cancer treatment makes it difficult to sleep.

Tumour lysis syndrome (TLS)

Some people are at risk of developing a condition called tumour lysis syndrome (TLS). The risk is highest at the beginning of treatment. TLS happens when treatment makes large numbers of cancer cells die and break down quickly. This releases lots of waste products into the blood and can affect the kidneys and heart.

TLS can be prevented or treated. You will have regular blood tests to check for TLS. You may have:

- extra fluids through a drip
- medicines such as allopurinol tablets or rasburicase through a drip.

Drinking at least 2 litres (3½ pints) of fluid a day will also help.

Cytokine Release Syndrome (CRS)

Cytokines are chemical messengers that increase the effects of the immune system. Rituximab causes the body to release large amounts of cytokines. Very rarely, this can cause flu-like symptoms, which can range from mild to severe. Some people may be monitored and treated in intensive care until this improves.

Symptoms include:

- a high temperature
- a fast heart rate
- low blood pressure and dizziness
- muscle and joint pain
- diarrhoea
- feeling sick and being sick
- difficulty breathing.

If you have any of these symptoms during treatment or after it ends, it is important to either:

- contact the hospital straight away on the 24-hour number
- go to the hospital straight away or call 999.

You should not drive yourself to hospital.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

Progressive multifocal leukoencephalopathy (PML)

Very rarely, this treatment may cause the following symptoms:

- blurred vision, loss of vision or double vision
- difficulty speaking
- weakness in an arm or a leg
- a change in the way you walk or problems with your balance
- persistent numbness
- decreased sensation or loss of sensation
- memory loss or confusion.

These may all be symptoms of a serious and potentially fatal brain condition called progressive multifocal leukoencephalopathy (PML).

If you had these symptoms before treatment with rituximab, tell your doctor or nurse immediately about any change in these symptoms.

Other information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
-

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

vitamins or supplements

- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 June 2025 | Next review: 01 June 2027

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as 'you' so that everyone feels included. Where clinically necessary we use the terms 'men' and 'women' or 'male' and 'female'. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026

You can read more about how we produce our information here.

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Macmillan Support Line

The Macmillan Support Line is a free and confidential service for people living with and affected by cancer. If you need to talk, we'll listen.

Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.

Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don’t want to call.

Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.

Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/rituximab · retrieved 19 October 2026