Trastuzumab

Trastuzumab is a cancer drug. It is used to treat breast cancer and stomach cancer.

On this page

- What is trastuzumab?
- How trastuzumab is given
- About side effects
- Side effects while treatment is being given
- Very common side effects
- Other side effects
- Other important information
- How we can help

What is trastuzumab?

Trastuzumab is a type of targeted therapy drug called a monoclonal antibody.

Trastuzumab is used to treat breast cancer and stomach cancer. It is used if tests show that the cancer cells have too much of a protein called HER2. This is called HER2 positive cancer.

It is best to read this information with our general information about targeted therapy drugs and the type of cancer you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree (consent) to have treatment.

This information is **not** about:

- trastuzumab deruxtecan (enhertu®)
- trastuzumab emtansine (Kadcyla®)
- trastuzumab and pertuzumab combination treatment.

We have separate information about these treatments.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the electronic Medicines Compendium (eMC) website, which has patient information leaflets (PIL) for individual drugs.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

How trastuzumab is given

You usually have trastuzumab in a day unit or clinic as an outpatient.

During a course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor
- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If everything is ok the pharmacy will prepare your treatment.

You have trastuzumab as a drip into a vein (intravenous infusion) or an injection under the skin (subcutaneously). Which one you have depends on your treatment plan.

You may have it with other cancer drugs.

As a drip

Your nurse will give it to you through 1 of the following:

- a cannula – a short, thin tube the nurse puts into a vein in the arm or hand
- a central line – a fine tube that goes under the skin of the chest and into a vein close by
- a PICC line – a fine tube that is put into a vein in the arm and goes up into a vein in the chest
- an implantable port (portacath) – a disc that is put under the skin on the chest or arm and goes into a vein in the chest.

The nurses give you the first dose (infusion) slowly. This is usually over 1½ hours (90 minutes). For the first infusion, you stay in hospital for a few hours so they can monitor you for any reaction.

If the first infusion does not cause a reaction, you can have the following ones over 30 to 90 minutes. Then you go home about 1 hour later.

As an injection under the skin

A nurse gives you the injection under the skin on your thigh. This takes a few minutes. After the first injection, the nurse monitors you for a few hours to make sure you do not have a reaction. If the first injection does not cause a reaction, you can go home sooner after the following ones.

Your course of treatment

You usually have trastuzumab once every 3 weeks.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

How long you have trastuzumab for depends on your situation. Your doctor, nurse or pharmacist will talk to you about your treatment plan.

If the cancer is advanced, trastuzumab might be given for as long as it is controlling the cancer.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Side effects while treatment is being given

Some people may have side effects while they are having this treatment or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. The first infusion is the most likely to cause a reaction, so it is usually given more slowly than later treatments. Before treatment, you will have medicines to help prevent or reduce any reaction.

Signs of a reaction can include:

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your infusion. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

Injection site

If you have trastuzumab as an injection under the skin, the skin in the area that was injected may become:

- painful
- swollen
- red, if you have white skin
- darker, if you have black or brown skin.

If this happens, let your doctor know. Painkillers may help. The nurse will swap between your left and right thigh.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Risk of infection

This treatment can reduce the number of white blood cells in your blood. These cells fight infection. If your white blood cell count is low, you may be more likely to get an infection. A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. It is important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

- feeling shivery and shaking
- a sore throat
- a cough
- breathlessness
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

You will have regular blood tests during treatment. If your white blood cell count is low, your doctor may stop your treatment for a short time, until your cell count increases. Sometimes they will reduce the dose of your treatment.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Effects on the heart

This treatment can affect how the heart works. You will have tests to check how well your heart is working. These may be done before, during and after treatment.

If the treatment is causing heart problems, your doctor may give you tablets to manage these. They may change the type of cancer treatment you are having.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call 999 if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Feeling sick

This is usually mild, but your doctor, nurse or pharmacist can prescribe drugs to reduce sickness or vomiting. Tell your doctor if it does not get better.

Sore mouth and throat

This treatment may cause a sore mouth and throat. You may also get mouth ulcers. This can make you more likely to get a mouth or throat infection. Use a soft toothbrush to clean your teeth or dentures in the morning, at night and after meals.

Contact the hospital straight away on the 24-hour number, if:

- a sore mouth or throat affects how much you can drink or eat
- your mouth, tongue, throat or lips have any blisters, ulcers or white patches.

They can give you advice, and mouthwash or medicines to help with the pain or to treat any infection. Follow their advice and make sure you:

- drink plenty of fluids
- avoid alcohol and tobacco
- avoid food or drinks that irritate your mouth and throat.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Loss of appetite

This treatment can affect your appetite. Don’t worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Tummy pain

You may get pain in your tummy (abdomen), or have indigestion. Your doctor, nurse or pharmacist can give you advice or treatment to help. Contact the hospital straight away on the 24-hour number if your symptoms do not go away, or get worse.

Diarrhoea

This treatment may cause diarrhoea. Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Constipation

This treatment can cause constipation. Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- Eat high-fibre foods, such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Headaches

This treatment may cause headaches. If you have headaches, tell your doctor, nurse or pharmacist. They can give you advice about painkillers that may help. Tell them if the headache does not get better, or gets worse.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Muscle or joint pain

You may get pain in your muscles or joints for a few days after treatment. If this happens, tell your doctor, nurse or pharmacist. They can give you painkillers and advice. They can also tell you if any of the painkillers you usually take are suitable.

Tell them if the pain does not get better. Having warm baths and resting regularly may help.

Hot flushes and sweats

Hot flushes are a common side effect of this treatment. During a flush, your neck and face may feel warm. If you have white skin, your skin may look red. If you have black or brown skin, your skin colour may deepen. Flushes can last up to 5 minutes. You might also have sweats or feel anxious or irritable during a flush.

Here are some things you can do to reduce the effects of hot flushes:

- Wear natural fabrics, such as cotton.
- Wear layers so you can remove clothes as needed.
- Use cotton sheets and layers of bedding you can remove.
- Try cooling pads or pillows to keep you cool.
- Keep rooms cool or use a fan.
- Have cold drinks and avoid caffeine and alcohol.

If they do not improve, talk to your doctor. Certain drugs can help to improve hot flushes.

Effects on the eyes

Your eyes may become watery and feel sore. Your doctor, nurse or pharmacist can give you advice and eye drops to help with this.

Contact the hospital on the 24-hour number if:

- your eyes get red and inflamed (conjunctivitis)
- you have pain in your eyes
- you notice any change in your vision.

Blocked or runny nose, or nosebleeds

These side effects are not usually serious. But tell your doctor, nurse or pharmacist if you have them. Your doctor or nurse may be able to help if you find the side effects difficult to cope with.

Skin and nail changes

This treatment can affect your skin. If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

It may cause a rash, which might be itchy. Your doctor or nurse can tell you what to expect.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Always tell your doctor, nurse or pharmacist about any skin changes. They can give you advice or prescribe creams or medicines to help. Changes to your skin are usually temporary and improve when treatment ends.

Your nails also might become brittle and break easily. Here are some tips to look after your nails:

- Keep your nails clipped short and clean.
- Avoid using very hot water when washing your hands or bathing.
- Moisturise your nails and cuticles regularly.
- It is okay to use water-based nail polish - but ask your cancer team if you want to use false nails, gels or other acrylics during treatment.
- Wear gloves to protect your nails when working in the house or garden.
- If your toenails are affected, wear well-fitting shoes, or shoes with open toes to cushion them.

Tell your doctor or nurse about any changes to your nails. They can give you advice or arrange for you to see a podiatrist. They are a foot care specialist.

Hair loss

Your hair may get thinner. But you are unlikely to lose all the hair from your head. Hair loss usually starts after your first or second treatment. It is almost always temporary, and your hair will usually grow back after treatment ends.

Your nurse can talk to you about ways to cope with hair loss.

Hand-foot (palmar-plantar) syndrome

This treatment can affect the palms of your hands and the soles of your feet. This is called palmar-plantar or hand-foot syndrome.

If you have white skin these areas may become red. If you have black or brown skin, these areas might get darker.

The skin on the palms of your hands and the soles of your feet may:

- be sore
- be painful, tingle, or swell
- peel, crack or blister.

If you have any of these symptoms, contact the hospital straight away on the 24-hour number. They can give you advice. This is especially important if you have any broken skin or if walking is difficult. They can prescribe creams and painkillers to help.

You can care for your hands and feet by:

- keeping your hands and feet cool by washing in cool water
- gently moisturising your hands and feet regularly
- wearing gloves to protect your hands and nails when working in the house or garden
- wearing loose cotton socks and avoiding tight-fitting shoes and gloves.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Difficulty sleeping (insomnia)

Sometimes this treatment can affect your sleep. Tell your doctor, nurse or pharmacist if cancer treatment makes it difficult to sleep.

Feeling tired (fatigue)

Feeling tired is a common side effect. Try to pace yourself and plan your day so you have time to rest between activities.

Being physically active can help to manage tiredness and give you more energy. If you feel sleepy, do not drive or operate machinery.

Dizziness

If you feel dizzy or drowsy, tell your doctor, nurse or pharmacist. It is important not to drive or operate machinery if you feel this way.

Effects on the lungs

This treatment can cause inflammation of the lungs. This is called pneumonitis. Contact the hospital straight away on the 24-hour number if you notice any of these changes during treatment or after it ends:

- breathlessness
- a cough that does not go away
- wheezing
- a fever, with a temperature over 37.5°C.

You should also tell them if any existing breathing problems get worse. You may have:

- tests to check your lungs
- treatment, if needed.

Changes to blood pressure

This treatment can increase or decrease your blood pressure. A nurse will check your blood pressure regularly during your treatment. If you have headaches, nosebleeds, changes to your eyesight, shortness of breath or feel dizzy, let your doctor, nurse or pharmacist know.

Changes to your taste

Some foods may taste different or have no taste. Try different foods to find out what tastes best to you. You may also get a bitter or metallic taste in your mouth. Your doctor, nurse or pharmacist can give you advice. It might help to try:

- sucking sugar-free sour or boiled sweets
- eating cold foods
- eating sharp-tasting fresh fruit.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Taste changes usually get better after treatment ends. We have more information about coping with changes to taste.

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Numb or tingling hands or feet (peripheral neuropathy)

This treatment may affect the nerves in your fingers and toes. This can cause numbness, tingling or pain in your hands or feet. This is called peripheral neuropathy. You might find it hard to do fiddly tasks such as fastening buttons or tying shoelaces.

If you have these symptoms, always tell your doctor, nurse or pharmacist. The symptoms usually improve slowly after treatment ends. But for some people they continue and are a long-term side effect of treatment.

Mood changes

You may feel low or depressed, or have mood swings, during this treatment. Talking to family and friends about how you feel might help. If mood changes last for more than a few weeks, tell your doctor, nurse or pharmacist. They can talk to you about different ways to manage low mood or depression.

Effects on the liver and kidneys

This treatment can affect how your kidneys or liver work. This is usually mild and goes back to normal after treatment ends. You will have blood tests to check how well your kidneys and liver are working. Contact the hospital on the 24-hour number if you:

- have blood in your urine (pee)
- are passing less urine or peeing less often than usual.

Drinking fluids also helps protect your kidneys. The advice is usually to try to drink at least 2 litres (3½ pints) of fluid each day. But follow any advice from your doctor, nurse or pharmacist about how much is right for you.

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a blood clot. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- herbal drugs and complementary or homeopathic therapies
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and coronavirus (covid). They may also recommend other vaccines, such as Shingrix® for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to get pregnant or make someone pregnant while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

Breastfeeding

You are advised not to breastfeed while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can get pregnant or make someone pregnant. If you are worried about this, it is important to talk with your doctor before you start treatment.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the electronic Medicines Compendium (eMC) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

Cancer Information Development team

Author

Date reviewed

Reviewed: 01 April 2025 | Next review: 01 April 2027

Our cancer information meets the PIF TICK quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about how we produce our information.

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as ‘you’ so that everyone feels included. Where clinically necessary we use the terms ‘men’ and ‘women’ or ‘male’ and ‘female’. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information here.

How we can help

Clinical Information Nurse Specialists

Our Cancer Information Nurse Specialists are dedicated cancer nurses available to talk to on our Macmillan Cancer Support Line.

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.

Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don’t want to call.

Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.

Find cancer support services in your area

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help

© Macmillan Cancer Support. Reproduced with permission. Source: https://www.macmillan.org.uk/cancer-information-and-support/treatments-and-drugs/trastuzumab · retrieved 19 October 2026