

FOLFOX

FOLFOX is a cancer drug treatment. It is used to treat bowel cancer and some other types of cancer.

On this page

- [What is FOLFOX?](#)
- [DPD testing before treatment](#)
- [How FOLFOX is given](#)
- [About side effects](#)
- [Side effects while treatment is being given](#)
- [Very common side effects](#)
- [Other side effects](#)
- [Other important information](#)
- [Listen to an audio version of this information](#)
- [How we can help](#)

What is FOLFOX?

FOLFOX is also known as oxaliplatin de Gramont or modified oxaliplatin de Gramont (OxMdG).

FOLFOX is a combination of cancer drugs. It is used to treat [bowel cancer](#) and some other types of cancer.

It is best to read this information with our general information about [cancer drug treatments](#) and the [type of cancer](#) you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree ([consent](#)) to have treatment.

FOLFOX is named after the initials of the drugs used for treatment. The drugs are:

- FOL – [folinic acid](#) (sometimes called calcium folinate or leucovorin)
- F – [fluorouracil](#) (5FU)
- OX – [oxaliplatin](#).

In this information, we call fluorouracil by its more common name, 5FU.

Folinic acid is not a chemotherapy drug. It is often given with 5FU to make 5FU work better.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the [electronic Medicines Compendium \(eMC\)](#) website, which has patient information leaflets (PIL) for individual drugs.

DPD testing before treatment

Before starting this treatment, you should have a blood test. This test checks for gene changes that affect how your body produces DPD. DPD is an enzyme that helps the body break down 5FU.

If you have low DPD levels or no DPD (DPD deficiency), you have more risk of serious or life-threatening side effects from 5FU. To reduce your risk, your doctor may give you:

- a lower dose of 5FU to start with
- a different type of chemotherapy.

Having low DPD levels does not usually cause any symptoms. Without a test, you will not know whether your DPD levels are low. Testing finds most people who are affected, but not all. Your doctor, nurse or pharmacist can give you more information about your risk of DPD deficiency, before you start treatment.

How FOLFOX is given

You usually have FOLFOX in a chemotherapy day unit or clinic as an outpatient or during a stay in hospital.

During your course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor
- chemotherapy nurse or specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

Before or on the day of each treatment, you will have a blood test. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your cancer drugs.

Your nurse will usually give you anti-sickness drugs before the cancer drugs. You will have your cancer treatment through 1 of the following:

-

- a [cannula](#) – a short, thin tube the nurse puts into a vein in the arm or hand
- a [central line](#) – a fine tube that goes under the skin of the chest and into a vein close by
- a [PICC line](#) – a fine tube that is put into a vein in the arm and goes up into a vein in the chest
- an [implantable port](#) (portacath) – a disc that is put under the skin on the chest or arm and goes into a vein in the chest.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

Your course of treatment

You usually have a course of several cycles of treatment over a few months. Your cancer team will discuss your treatment plan with you. They may give you a copy of the plan to take home.

Each cycle is usually 2 weeks. On day 1 of each cycle, your nurse gives you:

- oxaliplatin as a drip over 2 hours
- folinic acid as a drip (infusion) over 30 minutes or longer
- 5FU as an injection into your line or cannula.

You then have more 5FU through a small pump that you carry on a belt or in a holder.

Usually, the pump is set to run for about 46 hours. Your chemotherapy nurse will explain what will happen. They will also tell you how to look after the pump and what to do if there is a problem. If you have a PICC or central line, you can go home once the pump has started.

When you have had all the 5FU, a nurse will disconnect the pump for you. This is usually done at the hospital, but sometimes a district nurse will do it.

If you are given this treatment through a cannula, you will need to stay in hospital to have the 5FU.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

If your body does not produce enough of an enzyme called DPD, you may have a higher risk of severe or life-threatening side effects.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Side effects while treatment is being given

Some people have side effects while they are having a cancer drug or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your treatment. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

The drug leaks outside the vein

Sometimes cancer drugs that are given into a vein may leak outside the vein. If this happens, some drugs can damage the tissue near the vein. This is called extravasation. Extravasation is not common, but it is important that it is dealt with quickly. If you have any of the following symptoms around the vein during or after your treatment, tell your nurse straight away:

-

stinging

- pain
- swelling
- if you have white skin, the area may become red
- if you have black or brown skin, the area may become darker.

Throat spasm

Rarely, oxaliplatin can cause a spasm in the throat area around the voicebox (larynx). This can cause difficulties with swallowing and breathing. This might happen during treatment, or in the first few days after treatment.

This side effect can be frightening, but it should only be temporary. If you have breathing difficulties, take long, deep breaths through your nose. This will have a calming effect and help your breathing return to normal.

This symptom may be worse in cold temperatures. During treatment, and for a few days afterwards, it may help to:

- avoid cold drinks and ice cubes
- wrap up warm and cover your nose and mouth in cold weather.

It is important to tell your doctor, nurse or pharmacist if you have this side effect. They may give the next doses of oxaliplatin over 4 to 6 hours to reduce the risk of throat spasms.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Risk of infection

This treatment can reduce the number of white blood cells in your blood. These cells fight infection. If your white blood cell count is low, you may be more likely to get an [infection](#). A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. It is important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

-

feeling shivery and shaking

- a sore throat
- a cough
- breathlessness
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

Your white blood cell count will usually return to normal before your next treatment. You will have a blood test before having more treatment. If your white blood cell count is low, your doctor may delay your treatment for a short time, until your cell count increases.

Bruising and bleeding

This treatment can reduce the number of platelets in your blood. Platelets are cells that help the blood to clot.

If the number of platelets is low, you may bruise or bleed easily. You may have:

- nosebleeds
- bleeding gums
- heavy periods
- blood in your urine (pee) or stools (poo)
- tiny red, brown or purple spots that may look like a rash – these spots can be harder to see if you have black or brown skin.

If you have any unexplained bruising or bleeding, contact the hospital straight away on the 24-hour number. You may need a drip to give you extra platelets. This is called a platelet transfusion.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Diarrhoea

This treatment may cause severe [diarrhoea](#). Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. If you have a stoma, it will be more active than usual.

Your doctor, nurse or pharmacist may give you anti-diarrhoea drugs to take at home.

If you have diarrhoea or a mild increase in stoma activity, follow their advice about:

- taking anti-diarrhoea drugs
- how much and what type of fluids to drink
- any changes to your diet that might help.

Contact the hospital straight away on the 24-hour number if any of the following things happen:

- you have diarrhoea at night
- you have uncomfortable stomach cramps
- you have diarrhoea 4 or more times in a day
- you have a moderate increase in stoma activity
- the anti-diarrhoea drugs do not work within 24 hours.

You may need to give a stool sample, or go to hospital to have fluids through a drip or antibiotics.

Effects on the heart

5FU can affect how the heart works. You may have [tests](#) to check how well your heart is working. These may be done before, during and after treatment.

You may develop [heart problems](#) even if these test results are normal. Very rarely, 5FU causes heart failure or a heart attack. The risk of this happening is very low. It happens to less than 1 in 100 (1%) of people who have 5FU. But it is important that you know about it.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.

Always call 999 if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Difficulty breathing

5FU can cause sudden tightness (spasm) in the muscles in your chest for a short time. Contact the hospital straight away on the 24-hour number if you have any of these symptoms during treatment:

- breathlessness
- a cough that does not go away
- wheezing.

Always call 999 if you have difficulty breathing.

Feeling sick

Your doctor, nurse or pharmacist will prescribe anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to, even if you do not feel sick. It is easier to prevent sickness than to treat it after it has started.

[If you feel sick](#), take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids. If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice. They may change your anti-sickness treatment. Let them know if you still feel sick.

Numb or tingling hands or feet (peripheral neuropathy)

This treatment may affect the nerves in your fingers and toes. This can cause numbness, tingling or pain in your hands or feet. This is called [peripheral neuropathy](#). You might find it hard to do fiddly tasks such as fastening buttons or tying shoelaces.

During treatment, these symptoms are often made worse by the cold. It can help to keep warm and avoid iced drinks. You may need to take care opening the fridge and use gloves to handle frozen food.

If you have these symptoms, always tell your doctor, nurse or pharmacist. They sometimes need to change the drug or the dose of the drug. The symptoms usually improve slowly after treatment ends. But for some people they continue and are a long-term side effect of treatment.

Sore mouth and throat

This treatment may cause a [sore mouth and throat](#). You may also get mouth ulcers. This can make you more likely to get a mouth or throat infection. Use a soft toothbrush to clean your teeth or dentures in the morning, at night and after meals.

Contact the hospital straight away on the 24-hour number, if:

- a sore mouth or throat affects how much you can drink or eat
- your mouth, tongue, throat or lips have any blisters, ulcers or white patches.

They can give you advice, and mouthwash or medicines to help with the pain or to treat any infection. Follow their advice and make sure you:

- drink plenty of fluids
- avoid alcohol and tobacco
- avoid food or drinks that irritate your mouth and throat.

Hand-foot (palmar-plantar) syndrome

This treatment can affect the palms of your hands and the soles of your feet. This is called palmar-plantar or hand-foot syndrome.

If you have white skin these areas may become red. If you have black or brown skin, these areas might get darker.

The skin on the palms of your hands and the soles of your feet may:

- be sore
- be painful, tingle, or swell
- peel, crack or blister.

If you have any of these symptoms, contact the hospital straight away on the 24-hour number. They can give you advice. This is especially important if you have any broken skin or if walking is difficult. They can prescribe creams and painkillers to help.

You can care for your hands and feet by:

- gently moisturising your hands and feet regularly
- wearing gloves to protect your hands and nails when working in the house or garden
- wearing loose cotton socks and avoiding tight-fitting shoes and gloves.

Tummy pain

You may get pain in your tummy (abdomen), or have indigestion. Your doctor, nurse or pharmacist can give you advice or treatment to help. Contact the hospital straight away on the 24-hour number if your symptoms do not go away, or get worse.

Loss of appetite

This treatment can affect your appetite. Don't worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Changes to your taste

Some foods may taste different or have no taste. Try different foods to find out what tastes best to you. You may also get a bitter or metallic taste in your mouth. Your doctor, nurse or pharmacist can give you advice. It might help to try:

- sucking sugar-free sour or boiled sweets
- eating cold foods
- eating sharp-tasting fresh fruit.

Taste changes usually get better after treatment ends. We have more information about [coping with changes to taste](#).

Feeling tired

[Feeling tired](#) is a common side effect of this treatment. It is often worse towards the end of treatment and for some weeks after it ends. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Hair loss

Your hair will get thinner. Or you may lose all the hair from your head. You may also lose your eyelashes and eyebrows, as well as other body hair. Hair loss usually starts after your first or second treatment.

If you want to cover up hair loss, there are different ways you can do this. Your nurse will give you information about [coping with hair loss](#).

Remember to [protect your skin from the sun](#). Use suncream with a sun protection factor (SPF) of at least 30 on your scalp. Or cover up with a hat or scarf.

Hair loss is almost always temporary. Your hair will usually grow back after treatment ends.

Muscle or joint pain

You may get pain in your muscles or joints for a few days after treatment. If this happens, tell your doctor, nurse or pharmacist. They can give you painkillers and advice. They can also tell you if any of the painkillers you usually take are suitable.

Tell them if the pain does not get better. Having warm baths and resting regularly may help.

Headaches

This treatment may cause headaches. If you have headaches, tell your doctor, nurse or pharmacist. They can give you advice about painkillers that may help.

Contact the hospital straight away on the 24-hour number if the headache does not get better, or gets worse.

Skin changes

Chemotherapy can affect your skin. If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

This treatment can also:

- cause a rash, which may be itchy
- make your skin darker in some areas
- make any area treated with radiotherapy become red or sore. If you have white skin the area will become red and if you have black or brown skin the area might become darker.
- make you more sensitive to the sun.

Your [skin may burn more easily](#) during treatment and for several months after. Use a sun cream of at least SPF 30. SPF stands for sun protection factor. Cover up with clothing and wear a hat.

Always tell your doctor, nurse or pharmacist about any [skin changes](#). They can give you advice or prescribe creams or medicines to help. Changes to your skin are usually temporary and improve when treatment ends.

Effects on the liver and kidneys

This treatment can affect how your kidneys or liver work. This is usually mild and goes back to normal after treatment ends. You will have blood tests to check how well your kidneys and liver are working. Contact the hospital on the 24-hour number if you:

- have blood in your urine (pee)
- are passing less urine or peeing less often than usual.

Drinking fluids also helps protect your kidneys. The advice is usually to try to drink at least 2 litres (3½ pints) of fluid each day. But follow any advice from your doctor, nurse or pharmacist about how much is right for you.

Raised blood sugar levels

This treatment can raise your blood sugar levels and increase the risk of diabetes.

Signs of raised blood sugar include:

- feeling thirsty
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.

Tell your doctor or nurse if you have these symptoms. They can do tests to check your blood sugar level and give you advice.

If you already have [diabetes](#), your blood sugars may be higher than usual. You may need to check them more often, or make changes to your diabetic treatment. Your diabetic doctor or nurse will talk to you about how to manage this.

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Nail changes

This treatment can affect [your nails](#). They may grow more slowly or break more easily. You might notice ridges or white or dark lines across your nails. These changes usually disappear as the nails grow out after treatment. Sometimes nails can become loose or fall off. Tell your doctor, nurse or pharmacist about any changes to your nails.

If the skin around your nails becomes sore and swollen, contact the hospital straight away on the 24-hour number. These might be signs of an infection.

Effects on the lungs

This treatment can cause changes to the lungs. Tell your doctor, nurse or pharmacist if you develop:

-

a cough that does not go away

- wheezing
- breathlessness.

You should also tell them if any existing breathing problems get worse. You may have tests to check your lungs.

Effects on the eyes

Your eyes may become watery and feel sore. Your doctor, nurse or pharmacist can give you advice and eye drops to help with this.

Contact the hospital on the 24-hour number if:

- your eyes get red and inflamed (conjunctivitis)
- you have pain in your eyes
- you notice any change in your vision.

Bladder effects

Oxaliplatin may cause you discomfort when you pass urine. You may also pass urine more often than usual. Make sure you drink plenty of fluids during the 24 hours following chemotherapy. Try to drink at least 2 litres (3½ pints).

It is also important to empty your bladder regularly and to try to pass urine as soon as you feel the need to go.

Contact the hospital straight away if you feel any discomfort or stinging when you pass urine, or if you notice any blood in it.

Constipation

This treatment can cause [constipation](#). Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- [Eat high-fibre foods](#), such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Hearing changes

This treatment may cause hearing changes, including hearing loss. You may have ringing in the ears. This is called tinnitus. You may also become unable to hear some high-pitched sounds. Hearing changes may get better after this treatment ends. But this does not always happen. If you notice any changes in your hearing, tell your doctor, nurse or pharmacist.

Effects on sleep or mood

Sometimes this treatment can affect your sleep. It can also affect your mood, such as feeling excited or agitated. Tell your doctor, nurse or pharmacist if you notice mood changes or if cancer treatment makes it difficult to sleep.

Effects on the brain

Rarely, this treatment causes a brain condition called encephalopathy that can be serious. It is important that it is diagnosed and treated quickly.

This condition can cause:

- a headache that does not get better
- unsteadiness such as loss of balance or difficulty walking
- drowsiness or confusion
- changes in eyesight
- fits (seizures).

If you have any of these symptoms, it is important to either:

- contact the hospital straight away on the 24-hour number
- go to the hospital straight away.

You should not drive yourself to hospital.

Other important information

Driving and using machines

If you feel sick, have blurred vision or notice any effects on your nervous system, do not drive or use machines. Your doctor can tell you more about this.

Blood clot risk

Cancer and some cancer treatments can increase the risk of a [blood clot](#). Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- [herbal drugs and complementary or homeopathic therapies](#)
- recreational drugs – for example, cannabis.

Lactose

Some preparations of oxaliplatin may contain lactose. If you have a lactose allergy or intolerance, ask your doctor, nurse or pharmacist for more information.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and [coronavirus \(covid\)](#). They may also recommend other vaccines, such as Shingrix[®] for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to [get pregnant](#) or [make someone pregnant](#) while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to [breastfeed](#) while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can [get pregnant](#) or [make someone pregnant](#). If you are worried about this, it is important to talk with your doctor before you start treatment.

Sex

It is possible that small amounts of chemotherapy may be passed on through vaginal fluids or semen. If you have sex in the first few days after treatment, your cancer team will usually advise using condoms or a dental dam to protect your partner.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

Listen to an audio version of this information

Audio version of FOLFOX

You can listen to an audio version of this information about FOLFOX.

Booklets and resources



01 Dec 2023

Information about FOLXFOX combined chemotherapy treatment

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References



Visit the [electronic Medicines Compendium \(eMC\)](#) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

[Cancer Information Development team](#)

Author

Date reviewed

Reviewed: 01 August 2025 | Next review: 01 August 2027

Our cancer information meets the [PIF TICK](#) quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about [how we produce our information](#).

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as 'you' so that everyone feels included. Where clinically necessary we use the terms 'men' and 'women' or 'male' and 'female'. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information [here](#).

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Clinical Information Nurse Specialists

Our Cancer Information Nurse Specialists are dedicated cancer nurses available to talk to on our Macmillan Cancer Support Line.



Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.



Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don't want to call.



Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.



Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help