

Gefitinib

Gefitinib is a cancer drug. It is used to treat non-small cell lung cancer (NSCLC) that has spread.

On this page

- [What is gefitinib?](#)
- [How gefitinib is given](#)
- [About side effects](#)
- [Very common side effects](#)
- [Other side effects](#)
- [Other important information](#)
- [How we can help](#)

What is gefitinib?

Gefitinib is a type of [targeted therapy](#) drug called a cancer growth inhibitor.

It is used to treat [non-small cell lung cancer \(NSCLC\)](#) that has started to spread to:

- surrounding tissues (locally advanced)
- other parts of the body (advanced or metastatic).

Gefitinib is used to treat cancers that have an abnormal form of a protein called epidermal growth factor receptor (EGFR). Tests are done on the cancer cells to check the level of EGFR. These cells might be taken from a biopsy, previous surgery or a blood sample. The test tells your doctor whether gefitinib is likely to work for you.

It is best to read this information with our general information about [targeted therapy drugs](#) and the [type of cancer](#) you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree ([consent](#)) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the [electronic Medicines Compendium \(eMC\)](#) website, which has patient information leaflets (PIL) for individual drugs.

How gefitinib is given

Gefitinib comes in tablets, so you can take it at home.

During a course of treatment, you will meet someone from your cancer team, such as a:

- cancer doctor
- specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

During your course of treatment, you will have regular blood tests. This is to check that it is safe for you to have treatment.

You will meet with a doctor, nurse or pharmacist before you have treatment. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your tablets.

Your course of treatment

You usually take gefitinib for as long as it is controlling the cancer. Your cancer team will discuss your treatment plan with you.

They will give you the tablets to take home. Always take them exactly as they tell you to. This is important to make sure they work as well as possible for you.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take all your capsules or tablets exactly as they tell you to.

Taking gefitinib tablets

You usually take gefitinib as a tablet once a day. Take it at about the same time each day. Swallow it whole with a glass of water. You can take it with or without food.

If you have trouble swallowing tablets, you can dissolve them in half a glass of still water. Do not crush gefitinib tablets and do not use any other type of liquid to dissolve them.

If you forget to take the tablets, take the missed dose as soon as you remember. But if there is less than 12 hours until your next dose, do not take the missed dose. Never take a double dose.

Antacids or medicines to reduce stomach acid can affect how gefitinib works. Talk to your doctor, nurse or pharmacist if you are taking other medicines (either over the counter or prescribed). They will tell you how to safely take these medicines with gefitinib.

Other things to remember about your tablets:

- Keep them in the original package and at room temperature, away from moisture, heat and direct sunlight.
- Keep them safe, where children cannot see or reach them.
- If you are sick (vomit) just after taking them, do not take an extra dose. Take your next dose at the usual time.

- Do not throw away unused tablets. Return them to your cancer team at the hospital.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Skin changes

This treatment can cause an acne-like rash on the face, neck, chest and back. The rash is most likely to start in the first 2 or 3 weeks of treatment. Your skin may also feel dry, itchy, scaly or tender.

Skin changes are often mild but can be more severe in some people. If you notice any skin changes, contact the hospital as soon as possible on the 24-hour number you have been given. You may need creams, steroids or antibiotics. Your doctor may stop your treatment until the skin changes improve.

Here are some things you can do to take care of your skin during treatment:

- Wash with lukewarm water and mild, unperfumed, soap-free cleansers.
- Do not use products containing alcohol on your skin, or anti-acne products (unless they have been prescribed).
- Moisturise your skin regularly with unperfumed moisturisers for dry skin conditions.
- Use sun cream of at least SPF 30 to protect your skin from the sun. Cover up with clothing and a hat.

Skin changes usually improve when treatment ends.

Rarely, this treatment can cause a serious skin reaction that needs to be treated immediately in hospital.

Do not take any more of this treatment and contact the hospital straight away on the 24-hour number if you have any of these symptoms:

- a skin rash that is spreading
- blistering or peeling skin
- flu-like symptoms, such as a high temperature and joint pain
- sores on your lips or in your mouth.

Diarrhoea

This treatment may cause [diarrhoea](#). Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Loss of appetite

This treatment can affect your appetite. Don't worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Sore or dry mouth and throat

This treatment may cause a [sore or dry mouth and throat](#). You may also get mouth ulcers. This can make you more likely to get a mouth or throat infection. Use a soft toothbrush to clean your teeth or dentures in the morning, at night and after meals.

Contact the hospital straight away on the 24-hour number, if:

-

- your sore mouth or throat affects how much you can drink or eat
- your mouth, tongue, throat or lips have any blisters, ulcers or white patches.

They can give you advice, and mouthwash or medicines to help with the pain or to treat any infection. Follow their advice and make sure you:

- drink plenty of fluids
- avoid alcohol and tobacco
- avoid food or drinks that irritate your mouth and throat.

Feeling sick

You may feel sick while having this treatment, but this is usually mild. Your doctor, nurse or pharmacist may prescribe you anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to. [If you feel sick](#), take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids.

If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice.

Feeling tired

[Feeling tired](#) is a common side effect of this treatment. It is often worse towards the end of treatment and for some weeks after it ends. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Effects on the liver

This treatment can affect how the liver works. This is usually mild. You will have regular blood tests to check this. Sometimes liver changes can be serious. Contact the hospital straight away on the 24-hour number if you have any of these symptoms:

- yellow skin or eyes
- feeling very sleepy
- dark urine (pee)
- unexplained bleeding or bruising
- pain in the right side of your tummy (abdomen).

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Bleeding problems

This treatment can cause unusual bleeding. Contact the hospital on the 24 hour number if you have any of the following symptoms:

- nosebleeds
- bleeding gums
- blood in your pee (urine).

Tell your doctor, nurse or pharmacist if you are taking medicine to thin your blood, such as warfarin. This can increase your risk of unusual bleeding while on gefitinib.

Eye problems

This treatment can cause eye problems. These are usually mild but can sometimes become serious and may need treatment. Contact the hospital on the 24-hour number if you have any of these symptoms during treatment or after it ends:

- dry, sore or watery eyes
- eyes that are red and inflamed (conjunctivitis)
- eyes that are sensitive to light
- changes to your eyesight.

Some people find that their eyelashes grow longer and more curly than usual, but this is not common.

Effects on the lungs

This treatment can cause changes to the lungs. Tell your doctor, nurse or pharmacist if you develop:

- a cough that does not go away
- wheezing
- breathlessness.

You should also tell them if any existing breathing problems get worse. You may have tests to check your lungs.

Allergic reaction

Some people have an allergic reaction while having this treatment. Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy

- swelling of your face or mouth
- pain in your back, tummy or chest.

If you develop any of these signs or feel unwell, contact the hospital straight away on the 24-hour number. Do not take any more of this treatment until you have spoken to the hospital.

Hair loss

You might notice that your hair slowly gets thinner. Your hair can also become more brittle. Your nurse can give you information about coping with hair loss . Your hair will usually go back to normal after treatment ends.

Nail changes

This treatment can affect [your nails](#). They may grow more slowly or break more easily. You might notice ridges or white or dark lines across your nails. These changes usually disappear as the nails grow out after treatment. Sometimes nails can become loose or fall off. Tell your doctor, nurse or pharmacist about any changes to your nails.

If the skin around your nails becomes sore and swollen, contact the hospital straight away on the 24-hour number. These might be signs of an infection.

Hand-foot (palmar-plantar) syndrome

This treatment can affect the palms of your hands and the soles of your feet. This is called palmar-plantar or hand-foot syndrome.

If you have white skin these areas may become red. If you have black or brown skin, these areas might get darker.

The skin on the palms of your hands and the soles of your feet may:

- be sore
- be painful, tingle, or swell
- peel, crack or blister.

If you have any of these symptoms, contact the hospital straight away on the 24-hour number. They can give you advice. This is especially important if you have any broken skin or if walking is difficult. They can prescribe creams and painkillers to help.

You can care for your hands and feet by:

- keeping your hands and feet cool by washing in cool water
- gently moisturising your hands and feet regularly
- wearing gloves to protect your hands and nails when working in the house or garden
- wearing loose cotton socks and avoiding tight-fitting shoes and gloves.

Bladder irritation

Your bladder may become irritated during your treatment. You may:

- have a burning feeling when you pass urine (pee)

- need to pee more often or more urgently
- have blood in your pee.

If you have these symptoms, contact the hospital on the 24-hour number. This could be a sign of a bladder infection (cystitis). You may need treatment for this.

Try to drink at least 2 litres (3½ pints) of fluids each day.

Severe tummy pain

Rarely, gefitinib can cause a hole (perforation) in the small bowel or an inflamed pancreas.

Contact the hospital on the 24-hour number straight away if you have severe pain in the tummy (abdomen).

It is also very important to let them know if you:

- are bleeding from the back passage (rectum)
- have black stools (poo)
- are vomiting up blood
- have vomit that looks dark and grainy, like coffee grounds.

Other important information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a [blood clot](#). Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- [herbal drugs and complementary or homeopathic therapies](#)
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and [coronavirus \(covid\)](#). They may also recommend other vaccines, such as Shingrix[®] for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Lactose

This treatment may contain lactose. If you have a lactose allergy or intolerance, ask your doctor, nurse or pharmacist for more information.

Fertility

Some cancer drugs can affect whether you can [get pregnant](#) or [make someone pregnant](#). If you are worried about this, it is important to talk with your doctor before you start treatment.

Contraception

Your doctor, nurse or pharmacist will advise you not to [get pregnant](#) or [make someone pregnant](#) while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to [breastfeed](#) while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References

Visit the [electronic Medicines Compendium \(eMC\)](#) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

[Cancer Information Development team](#)

Author

Date reviewed

Reviewed: 01 November 2024 | Next review: 01 November 2027

Our cancer information meets the [PIF TICK](#) quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about [how we produce our information](#).

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
-

use illustrations to explain text

- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as 'you' so that everyone feels included. Where clinically necessary we use the terms 'men' and 'women' or 'male' and 'female'. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information [here](#).

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Clinical Information Nurse Specialists

Our Cancer Information Nurse Specialists are dedicated cancer nurses available to talk to on our Macmillan Cancer Support Line.



Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.



Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don't want to call.



Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.



Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help