

Megestrol acetate (Megace®)

Megestrol acetate is a cancer drug. It is used to treat breast cancer and womb cancer.

On this page

- [What is megestrol acetate \(Megace®\)?](#)
- [How megestrol acetate works](#)
- [About side effects](#)
- [Very common side effects](#)
- [Other side effects](#)
- [Other information](#)
- [How we can help](#)

What is megestrol acetate (Megace®)?

Megestrol is a hormonal therapy drug. It is also called Megace[®]. It is used to treat [breast cancer](#) and [womb cancer](#). It may sometimes be used to treat other cancers.

It is best to read this information with our general information about [hormonal therapy](#) and the [type of cancer](#) you have.

Megestrol can also be used to improve your [appetite](#) and help you to put on weight.

Your cancer team will talk to you about this treatment and its possible side effects before you agree ([consent](#)) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the [electronic Medicines Compendium \(eMC\)](#) website, which has patient information leaflets (PIL) for individual drugs.

How megestrol acetate works

Hormones are chemicals that our bodies make. They act as messengers and help control how cells and organs work. Hormonal therapy drugs change how hormones are made or work in the body. Many cancers need hormones to grow.

Megestrol is similar to the sex hormone progesterone. It works by changing the hormone balance in the body, which may stop the cancer growing. It can also act directly on cancer cells so that they cannot grow.

Taking megestrol acetate tablets

You have megestrol acetate as tablets you can take at home. Your cancer team will discuss your treatment plan with you.

Always take the tablets exactly as they tell you to. This is important to make sure they work as well as possible for you. Make sure you:

- swallow them whole with a glass of water
- take them at the same time every day.

If you forget to take the tablets, take the missed dose as soon as possible within the same day. If a full day has passed, do not take a double dose.

Other things to remember about your tablets:

- Keep them in the original package and at room temperature, away from moisture, heat and direct sunlight.
- Keep them safe, where children cannot see or reach them.
- If you are sick (vomit) just after taking them, do not take an extra dose. Take your next dose at the usual time.
- Get a new prescription before you run out, and make sure you have plenty for holidays.
- Do not throw away unused tablets. Return them to a pharmacy.

Your cancer team may also give you other medicines to take home. Take all your medicines exactly as they tell you to. Do not stop taking any of your medicines unless your doctor tells you to.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

When you have this treatment, you will have regular appointments with a doctor, nurse or pharmacist. Always tell them about any side effects you have. They can give you:

- drugs to help control most side effects
- advice about managing side effects.

Most side effects can be managed. But sometimes side effects are harder to control. It is important not to stop taking hormonal therapy without telling your doctor. If side effects cannot be managed, your doctor may suggest a different type of hormonal therapy.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Increased appetite and weight gain

Megestrol can make you hungrier than usual. This may mean you gain weight. Your appetite should return to normal when you stop taking megestrol.

If you are worried about [gaining weight](#), your doctor, nurse or dietitian can give you more advice.

Because of this side effect, megestrol is sometimes used to treat people who have lost their appetite.

Hot flushes and sweats

Hot flushes are a common side effect of this treatment. During a flush, your neck and face may feel warm. If you have white skin, your skin may look red. If you have black or brown skin, your skin colour may deepen. Flushes can last up to 5 minutes. You might also have sweats or feel anxious or irritable during a flush.

Here are some things you can do to reduce the effects of hot flushes:

- Wear natural fabrics, such as cotton.
- Wear layers so you can remove clothes as needed.
- Use cotton sheets and layers of bedding you can remove.
- Try cooling pads or pillows to keep you cool.
- Keep rooms cool or use a fan.
- Have cold drinks and avoid caffeine and alcohol.

You might have fewer hot flushes as your body adjusts to this treatment. You could try therapies to help you [cope with hot flushes](#), such as:

- [talking therapies](#) like cognitive behaviour therapy (CBT)

- controlled breathing exercises
- yoga
- acupuncture.

If they do not improve, talk to your doctor. Certain drugs can help to improve hot flushes.

Flushes usually stop a few months after treatment ends. But some people continue to have them.

High blood pressure

While taking megestrol, your blood pressure will be checked regularly. If you have ever had any problems with your blood pressure, tell your doctor or nurse. You should also tell them if you have any headaches.

Feeling breathless

This treatment may make you feel breathless. If this happens, you should let your nurse or doctor know.

Constipation

This treatment can cause [constipation](#). Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- [Eat high-fibre foods](#), such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Rounded face

Your face may become rounded while taking megestrol. This is sometimes called a “moon face”.

Raised blood sugar levels

This treatment can raise your blood sugar levels and increase the risk of diabetes.

Signs of raised blood sugar include:

- feeling thirsty
- needing to pass urine (pee) more often than usual
- feeling more tired than usual.

Tell your doctor or nurse if you have these symptoms. They can do tests to check your blood sugar level and give you advice.

If you already have [diabetes](#), your blood sugars may be higher than usual. You may need to check them more often, or make changes to your diabetic treatment. Your diabetic doctor or nurse will talk to you about how to manage this.

Blood clot risk

Cancer and some cancer treatments can increase the risk of a [blood clot](#). Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Feeling sick and passing wind

If you feel sick when taking megestrol, tell your doctor or nurse. They can give you anti-sickness drugs to help stop you feeling sick. Take the drugs exactly as your nurse or pharmacist tells you. Megestrol can also cause you to pass wind more often than usual. Your nurse can give you advice on managing these side effects.

Diarrhoea

This treatment may cause [diarrhoea](#). Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Tumour flare

If you have symptoms caused by the cancer, they may get worse for a short time when you start taking it. This is called tumour flare. Tell your doctor, nurse or pharmacist if this happens, they can give you treatment to help with this.

Feeling tired

[Feeling tired](#) is a common side effect of this treatment. It is often worse towards the end of treatment and for some weeks after it ends. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Mood changes

You may feel low or depressed, or have mood swings, during this treatment. Talking to family and friends about how you feel might help. If mood changes last for more than a few weeks, tell your doctor, nurse or pharmacist. They can talk to you about different ways to manage low mood or [depression](#).

Memory and concentration

You may notice [changes in your memory](#). You may also find it harder to concentrate.

To help you remember things, try using:

- reminder alarms or notes on your phone
- a diary, calendar or phone app
- a notebook you carry with you.

Skin changes

This treatment can [affect your skin](#). It may cause a rash, which may be itchy. Your doctor or nurse can tell you what to expect. If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

Always tell your doctor or nurse about any skin changes. They can give you advice and may prescribe creams or medicines to help. Changes to your skin usually improve when treatment ends.

Hair thinning

While you are having this treatment, your hair may become thinner, drier and more brittle. This is usually mild. If you are worried about this, ask your nurse for advice on hair care.

Effects on the heart

This treatment might increase certain risk factors for [heart problems](#). Tell your doctor if you have:

- had any problems with your heart
-

high blood pressure

- high levels of cholesterol – a fatty substance in your blood.

They can monitor or treat these if needed. Talk to your doctor if you are worried about this.

Things you can do to take care of your heart include:

- not smoking
- maintaining a healthy weight
- being physically active.

Swollen hands, feet and ankles

Your hands, feet and ankles may swell because of fluid building up in them. This is not harmful, but can be uncomfortable. Tell your doctor or nurse if you notice any swelling. The swelling will get better after treatment ends.

Loss of sex drive

This treatment can affect your sex drive (libido). If you are worried about the [effects of treatment on your sex life](#), you can talk to your doctor, nurse or pharmacist. They can give you help and support.

Changes to periods

If you have periods, these may become irregular or stop while you are having this treatment. They might return after treatment, but this does not always happen. Your [menopause](#) may start sooner than it would have done. Your doctor, nurse or pharmacist can give you more information.

Other information

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- [herbal drugs and complementary or homeopathic therapies](#)
- recreational drugs – for example, cannabis.

Lactose

These tablets contain a type of sugar called lactose. If you have been told by a doctor that you cannot digest some sugars or are lactose intolerant, talk to your doctor before taking this drug.

Fertility

Some cancer drugs can affect whether you can [get pregnant](#) or [make someone pregnant](#). If you are worried about this, it is important to talk with your doctor before you start treatment.

Contraception

Your doctor, nurse or pharmacist will advise you not to [get pregnant](#) while having this treatment. The drug may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use, for example using barrier contraception instead of hormonal contraception like the oral pill
- how long after treatment you should continue to use contraception.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References



Visit the [electronic Medicines Compendium \(eMC\)](#) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

[Cancer Information Development team](#)

Author

Date reviewed

Reviewed: 01 December 2024 | Next review: 01 June 2027

Our cancer information meets the [PIF TICK](#) quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about [how we produce our information](#).

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Clinical Information Nurse Specialists

Our Cancer Information Nurse Specialists are dedicated cancer nurses available to talk to on our Macmillan Cancer Support Line.



Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.



Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don't want to call.



Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.



Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

Get help