

Nab-paclitaxel (Abraxane®)

Nab-paclitaxel (Abraxane®) is a cancer drug. It is used to treat different cancer types.

On this page

- [What is nab-paclitaxel \(Abraxane®\)?](#)
- [Having nab-paclitaxel](#)
- [About side effects](#)
- [Side effects while treatment is being given](#)
- [Very common side effects](#)
- [Other side effects](#)
- [Other information](#)
- [How we can help](#)

What is nab-paclitaxel (Abraxane®)?

Nab-paclitaxel is a type of chemotherapy. It is made up of the chemotherapy drug paclitaxel, which is combined with a protein called albumin. It is also called Abraxane®. It is used to treat [breast cancer](#), [non-small cell lung cancer](#) and [cancer of the pancreas](#). It may sometimes be used to treat other cancers.

It is best to read this information with our general information about [chemotherapy](#) and the [type of cancer](#) you have.

Your cancer team will talk to you about this treatment and its possible side effects before you agree ([consent](#)) to have treatment.

More information about this treatment

This information is correct at time of publishing. But sometimes the types of cancer this treatment is used for, or treatment side effects, may change between revision dates.

You can talk to your cancer team if you want more detailed information about this treatment. Or visit the [electronic Medicines Compendium \(eMC\)](#) website, which has patient information leaflets (PIL) for individual drugs.

Having nab-paclitaxel

You have nab-paclitaxel as a drip into a vein (intravenously). You usually have it in a chemotherapy day unit or clinic as an outpatient. You may have it with other cancer drugs.

During your course of treatment, you will meet with someone from your cancer team, such as a:

- cancer doctor
- chemotherapy nurse or specialist nurse
- specialist pharmacist.

This is who we mean when we mention doctor, nurse or pharmacist in this information.

Before or on the day of each treatment, you will have a blood test. This is to check that it is safe for you to have chemotherapy.

You will meet with a doctor, nurse or pharmacist before you have chemotherapy. They will talk to you about your blood results and ask how you have been feeling. If your blood results are okay, the pharmacy team will prepare your chemotherapy.

Your nurse will usually give you anti-sickness drugs before chemotherapy. You will have your chemotherapy through one of the following:

- a [cannula](#) – a short, thin tube the nurse puts into a vein in the arm or hand
- a [central line](#) – a fine tube that goes under the skin of the chest and into a vein close by
- a [PICC line](#) – a fine tube that goes into a vein in the arm and up into a vein in the chest
- an [implantable port](#) (portacath) – a disc that goes under the skin on the chest or arm and into a vein in the chest.

Your cancer team may also give you anti-sickness drugs and other medicines to take home. Take these exactly as they tell you to.

Your course of treatment

You usually have a course of several cycles of treatment over a few months. Your cancer team will discuss your treatment plan with you. They will explain how often you will have nab-paclitaxel based on your situation and type of cancer.

Nab-paclitaxel may be given as an infusion (drip):

- every 3 weeks
- once a week for 3 weeks, followed by 1 week of no chemotherapy.

It usually takes 30 minutes to give the infusion.

About side effects

We explain the most common side effects of this treatment here. We also include some that are less common.

You may get some of the side effects we mention, but you are unlikely to get all of them. And you may have some side effects, including rarer ones, that we have not listed here.

Other cancer treatments may cause different side effects. If you are also having other cancer treatment, you may have other side effects.

Always tell your doctor, nurse or pharmacist about any side effects you have. They can give you:

- drugs to help control some side effects
- advice about managing side effects.

It is important to take any drugs exactly as explained. This means they will be more likely to work for you.

Serious and life-threatening side effects

Some cancer treatments can cause serious side effects. Sometimes, these may be life-threatening. Your doctor, nurse or pharmacist can explain the risk of these side effects to you.

Contact the hospital

Your doctor, nurse or pharmacist will give you 24-hour contact numbers for the hospital. If you feel unwell or need advice, you can call at any time of the day or night. Save these numbers in your phone or keep them somewhere safe.

Side effects while treatment is being given

Some people have side effects while they are having chemotherapy or shortly after they have it.

Allergic reaction

Some people have an allergic reaction while having this treatment. Signs of a reaction can include:

- feeling hot or flushed
- shivering
- itching
- a skin rash
- feeling dizzy or sick
- a headache
- feeling breathless or wheezy
- swelling of your face or mouth
- pain in your back, tummy or chest.

Your nurse will check you for signs of a reaction during your treatment. If you feel unwell or have any of these signs, tell them straight away. If you do have a reaction, they can treat it quickly.

Sometimes a reaction happens a few hours after treatment. If you develop any of these signs or feel unwell after you get home, contact the hospital straight away on the 24-hour number.

Always call 999 if swelling happens suddenly or you are struggling to breathe.

The drug leaks outside the vein

Sometimes cancer drugs that are given into a vein may leak outside the vein. If this happens, some drugs can damage the tissue near the vein. This is called extravasation. Extravasation is not common, but it is important that it is dealt with quickly. If you have any of the following symptoms around the vein during or after your treatment, tell your nurse straight away:

- stinging
- pain
- swelling
- if you have white skin, the area may become red
- if you have black or brown skin, the area may become darker.

Pain along the vein

This treatment can cause pain:

- at the place where the drip (infusion) is given
- along the vein.

If you feel pain, tell your nurse straight away. They can check the site. They may give the drug more slowly or flush it through with more fluid to reduce pain.

Very common side effects

These side effects happen to 10 or more people in every 100 people (10% or more) who have this treatment.

Risk of infection

This treatment can reduce the number of white blood cells in your blood. These cells fight infection. If your white blood cell count is low, you may be more likely to get an [infection](#). A low white blood cell count is called neutropenia.

An infection can be very serious when the number of white blood cells is low. It is important to get any infection treated as soon as possible. If you have any of the following symptoms, contact the hospital straight away on the 24-hour number:

- a temperature above 37.5°C
- a temperature below 36°C
- you feel unwell, even with a normal temperature
- you have symptoms of an infection.

Symptoms of an infection include:

- feeling shivery and shaking

- a sore throat
- a cough
- breathlessness
- diarrhoea
- needing to pass urine (pee) often, or discomfort when you pass urine.

It is important to follow any specific advice your cancer treatment team gives you.

Your white blood cell count will usually return to normal before your next treatment. You will have a blood test before having more treatment. If your white blood cell count is low, your doctor may delay your treatment for a short time, until your cell count increases.

Anaemia (low number of red blood cells)

This treatment can reduce the number of red blood cells in your blood. Red blood cells carry oxygen around the body. If the number of red blood cells is low, this is called anaemia. You may feel:

- very low in energy
- breathless
- dizzy and light-headed.

If you have these symptoms, contact the hospital straight away on the 24-hour number. You may need treatment for anaemia. If you are very anaemic, you may need a drip to give you extra red blood cells. This is called a blood transfusion.

Feeling sick

Your doctor, nurse or pharmacist will prescribe anti-sickness drugs to help prevent or control sickness. Take the drugs exactly as they tell you to, even if you do not feel sick. It is easier to prevent sickness than to treat it after it has started.

[If you feel sick](#), take small sips of fluid often and eat small amounts regularly. It is important to drink enough fluids. If you continue to feel sick, or if you are sick (vomit) 1 to 2 times in 24 hours, contact the hospital on the 24-hour number as soon as possible. They will give you advice. They may change your anti-sickness treatment. Let them know if you still feel sick.

Feeling tired

[Feeling tired](#) is a common side effect of this treatment. It is often worse towards the end of treatment and for some weeks after it ends. Try to pace yourself and plan your day so you have time to rest. Gentle exercise, like short walks, can help you feel less tired.

If you feel sleepy, do not drive or use machinery.

Diarrhoea

This treatment may cause [diarrhoea](#). Diarrhoea means passing more stools (poo) than is normal for you, or having watery or loose stools. You may also have stomach cramps. If you have a stoma, it may be more active than usual.

If you are passing loose stools 3 or more times a day and this is not normal for you, contact the hospital as soon as possible on the 24-hour number. Follow the advice they give you about:

- taking anti-diarrhoea medicines
- drinking enough fluids to keep you hydrated and to replace lost salts and minerals
- any changes to your diet that might help.

They might also ask you for a specimen of your stool to check for infection.

Constipation

This treatment can cause [constipation](#). Constipation means that you are not able to pass stools (poo) as often as you normally do. It can become difficult or painful. Here are some tips that may help:

- Drink at least 2 litres (3½ pints) of fluids each day.
- [Eat high-fibre foods](#), such as fruit, vegetables and wholemeal bread.
- Do regular gentle exercise, like going for short walks.

If you have constipation, contact the hospital on the 24-hour number for advice. They can give you drugs called laxatives to help.

If you have not been able to pass stools for over 2 days and are being sick, contact the 24-hour number straight away.

Loss of appetite

This treatment can affect your appetite. Don't worry if you do not eat much for 1 or 2 days. But if your appetite does not come back after a few days, or if you are losing weight, tell your doctor, nurse or pharmacist. They can give you advice. They may give you food or drink supplements. Or they may suggest changes to your diet or eating habits to help.

Sore mouth and throat

This treatment may cause a [sore mouth and throat](#). You may also get mouth ulcers. This can make you more likely to get a mouth or throat infection. Use a soft toothbrush to clean your teeth or dentures in the morning, at night and after meals.

Contact the hospital straight away on the 24-hour number, if:

- a sore mouth or throat affects how much you can drink or eat
- your mouth, tongue, throat or lips have any blisters, ulcers or white patches.

They can give you advice, and mouthwash or medicines to help with the pain or to treat any infection. Follow their advice and make sure you:

- drink plenty of fluids
- avoid alcohol and tobacco
- avoid food or drinks that irritate your mouth and throat.

Numb or tingling hands or feet (peripheral neuropathy)

This treatment may affect the nerves in your fingers and toes. This can cause numbness, tingling or pain in your hands or feet. This is called [peripheral neuropathy](#). You might find it hard to do fiddly tasks such as fastening buttons or tying shoelaces.

If you have these symptoms, always tell your doctor, nurse or pharmacist. They sometimes need to change the drug or the dose of the drug. The symptoms usually improve slowly after treatment ends. But for some people they continue and are a long-term side effect of treatment.

Skin changes

Chemotherapy can affect your skin. It might feel dry. You may develop a rash, which may be itchy. Always tell your doctor, nurse or pharmacist about any [skin changes](#). They can give you advice or prescribe creams or medicines to help.

If your skin feels dry, try using soap-free cleansers and unperfumed moisturising cream every day.

Serious skin changes

Rarely, this treatment can cause a serious skin reaction that needs to be treated immediately in hospital. Contact the hospital straight away on the 24-hour number if you have any of these symptoms:

- a skin rash that is spreading
- blistering or peeling skin
- flu-like symptoms, such as a high temperature and joint pain
- sores on your lips or in your mouth.

Hair loss

You usually lose all the hair from your head. You may also lose your eyelashes and eyebrows, as well as other body hair. [Hair loss](#) usually starts after your first or second treatment.

[Scalp cooling](#) is a way of lowering the temperature of your scalp to help reduce hair loss. Your nurse can tell you whether this is an option for you.

If you want to cover up hair loss, there are different ways to do this. Your nurse can give you information about coping with hair loss. Remember to [protect your skin from the sun](#). Use suncream with a sun protection factor (SPF) of at least 30 on your scalp. Or cover up with a hat or scarf.

Hair loss is usually temporary. Your hair will usually grow back after treatment ends. Very rarely, hair may not grow back. Or it may grow back thinner than before. If you are worried about this, talk to your doctor, nurse or pharmacist.

Muscle or joint pain

You may get pain in your muscles or joints for a few days after treatment. If this happens, tell your doctor, nurse or pharmacist. They can give you painkillers and advice. They can also tell you if any of the painkillers you usually take are suitable.

If you have muscle or joint pain, try:

-

placing a heat pad or covered hot water bottle against the painful area

- taking warm baths
- planning your activities to include regular rests.

Other side effects

These side effects happen to less than 10 in 100 people (less than 10%) who have this treatment. Some of them are much rarer than this but they are still important to know about. Rare means a side effect that happens to less than 1 in 1,000 people (less than 0.1%).

Headaches

This treatment may cause headaches. If you have headaches, tell your doctor, nurse or pharmacist. They can give you advice about painkillers that may help. Tell them if the headache does not get better, or gets worse.

Effects on the eyes

Your eyes may become watery and feel sore. Your doctor, nurse or pharmacist can give you advice and eye drops to help with this.

Contact the hospital on the 24-hour number if:

- your eyes get red and inflamed (conjunctivitis)
- you have pain in your eyes
- you notice any change in your vision.

Effects on the liver

This treatment may affect how your liver works. This is usually mild and goes back to normal after treatment ends. You will have blood tests to check how well your liver is working.

Feeling dizzy

This treatment may cause dizziness. Tell your doctor or nurse if this is a problem. It is important you do not drive or operate machinery if you feel dizzy.

Changes to your taste

Some foods may taste different or have no taste. Try different foods to find out what tastes best to you. You may also get a bitter or metallic taste in your mouth. Your doctor, nurse or pharmacist can give you advice. It might help to try:

- sucking sugar-free sour or boiled sweets
- eating cold foods
- eating sharp-tasting fresh fruit.

Taste changes usually get better after treatment ends. We have more information about [coping with changes to taste](#).

Low levels of potassium

This treatment can sometimes affect the levels of potassium in your blood. You will have regular blood tests. Your doctor may advise you to eat foods that contain potassium or prescribe you supplements to take if the levels are too low.

Nail changes

This treatment can affect [your nails](#). They may grow more slowly or break more easily. You might notice ridges or white or dark lines across your nails. These changes usually disappear as the nails grow out after treatment. Sometimes nails can become loose or fall off. Tell your doctor, nurse or pharmacist about any changes to your nails.

If the skin around your nails becomes sore and swollen, contact the hospital straight away on the 24-hour number. These might be signs of an infection.

Effects on the lungs

This treatment can cause inflammation of the lungs. This is called pneumonitis. Contact the hospital straight away on the 24-hour number if you notice any of these changes during treatment or after it ends:

- breathlessness
- a cough that does not go away
- wheezing
- a fever, with a temperature over 37.5°C.

You should also tell them if any existing breathing problems get worse. You may have:

- tests to check your lungs
- treatment, if needed.

Effects on the heart

This treatment can affect how the heart works. You may have [tests](#) to check how well your heart is working. These may be done before, during and after treatment.

If the treatment is causing [heart problems](#), your doctor may change the type of treatment you are having.

Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- breathlessness
- dizziness
- changes to your heartbeat (palpitations)
- swollen feet and ankles.

Other conditions can cause these symptoms, but it is important to get them checked by a doctor.

Always call **999** if you have:

- chest pain, pressure, heaviness, tightness or squeezing across the chest
- difficulty breathing.

Fluid build-up

This treatment can cause a build-up of fluid in the body. This will slowly get better after treatment ends. Contact the hospital on the 24-hour number if you:

- are gaining weight
- have swelling in your face, legs or ankles.

They can give you advice and treatment to help.

Mood changes

You may feel low or depressed, or have mood swings, during this treatment. Talking to family and friends about how you feel might help. If mood changes last for more than a few weeks, tell your doctor, nurse or pharmacist. They can talk to you about different ways to manage low mood or [depression](#).

Tumour lysis syndrome (TLS)

Some people are at risk of developing a condition called tumour lysis syndrome (TLS). The risk is highest at the beginning of treatment. TLS happens when treatment makes large numbers of cancer cells die and break down quickly. This releases lots of waste products into the blood and can affect the kidneys and heart.

TLS can be prevented or treated. You will have regular blood tests to check for TLS. You may have:

- extra fluids through a drip
- medicines such as allopurinol tablets or [rasburicase](#) through a drip.

Drinking at least 2 litres (3½ pints) of fluid a day will also help.

Tummy pain

You may get pain in your tummy (abdomen), or have indigestion. Your doctor, nurse or pharmacist can give you advice or treatment to help. Contact the hospital straight away on the 24-hour number if your symptoms do not go away, or get worse.

Other information

Blood clot risk

Cancer and some cancer treatments can increase the risk of a [blood clot](#). Contact the hospital straight away on the 24-hour number if you have any of these symptoms during or after treatment:

- throbbing pain or swelling in a leg or arm
- reddening of the skin in the area – if you have black or brown skin, this can be harder to notice, but the skin might become darker
- suddenly feeling breathless or coughing.

Always call **999** if you have:

- chest pain
- difficulty breathing.

A blood clot is serious, but it can be treated with drugs called anticoagulants. These thin the blood. Your doctor, nurse or pharmacist can give you more information about preventing and treating blood clots.

Other medicines

Some medicines can affect how this treatment works or be harmful while you are having it. Always tell your cancer doctor, nurse or pharmacist about any drugs you are taking or planning to take, such as:

- medicines you have been prescribed
- medicines you buy in a shop, pharmacy or online
- vitamins or supplements
- [herbal drugs and complementary or homeopathic therapies](#)
- recreational drugs – for example, cannabis.

Vaccinations

Cancer doctors usually recommend that people with cancer have vaccinations for flu and [coronavirus \(covid\)](#). They may also recommend other vaccines, such as Shingrix[®] for shingles. These all help reduce your risk of serious illness from these infections. Most people can have these vaccines, including people with weak immune systems.

You should not have live vaccines if your immune system is weak. This includes if you are having or recently had chemotherapy, radiotherapy or other cancer treatments that affect your immune system. Live vaccines can make you unwell because they contain a very weak version of the illness they protect you against. There are several live vaccines, including the yellow fever vaccine.

It is important to ask your doctor, nurse or pharmacist for advice about having vaccinations. They can explain what vaccines are right for you and when it is best to have them.

Contraception

Your doctor, nurse or pharmacist will advise you not to [get pregnant](#) or [make someone pregnant](#) while having this treatment and for some time afterwards. The drugs may harm a developing baby. It is important to use contraception to prevent pregnancy. Follow their advice about:

- what types of contraception to use
- how long after treatment you should continue to use contraception.

Breastfeeding

You are advised not to [breastfeed](#) while having this treatment, or for some time after treatment ends. This is because the drugs could be passed to the baby through breast milk.

Your doctor, nurse or pharmacist can give you more information.

Fertility

Some cancer drugs can affect whether you can [get pregnant](#) or [make someone pregnant](#). If you are worried about this, it is important to talk with your doctor before you start treatment.

Sex

It is possible that small amounts of chemotherapy may be passed on through vaginal fluids or semen. If you have sex in the first few days after treatment, your cancer team will usually advise using condoms or a dental dam to protect your partner.

Medical and dental treatment

If you need medical treatment for any reason other than cancer, always tell the healthcare professional that you are having cancer treatment. Give them the contact details for your cancer doctor or cancer team so they can ask for advice.

If you have appointments with a dentist, always tell them you are having cancer treatment. Talk to your cancer team before you have any dental treatment.

About our information

This information has been written, revised and edited by Macmillan Cancer Support's Cancer Information Development team. It has been reviewed by expert medical and health professionals and people living with cancer.

References



Visit the [electronic Medicines Compendium \(eMC\)](#) to download a Patient Information Leaflet (PIL) for more detailed information. The leaflet lists all known side effects.

[Cancer Information Development team](#)

Author

Date reviewed

Reviewed: 01 May 2026 | Next review: 01 May 2028

Our cancer information meets the [PIF TICK](#) quality mark.

This means it is easy to use, up-to-date and based on the latest evidence. Learn more about [how we produce our information](#).

The language we use

We want everyone affected by cancer to feel our information is written for them.

We want our information to be as clear as possible. To do this, we try to:

- use plain English
- explain medical words
- use short sentences
- use illustrations to explain text
- structure the information clearly
- make sure important points are clear.

We use gender-inclusive language and talk to our readers as 'you' so that everyone feels included. Where clinically necessary we use the terms 'men' and 'women' or 'male' and 'female'. For example, we do so when talking about parts of the body or mentioning statistics or research about who is affected.

You can read more about how we produce our information [here](#).

How we can help

How to print a PDF from this website

Instructions on how you can make a PDF version of this page using the print function on your browser.

Clinical Information Nurse Specialists

Our Cancer Information Nurse Specialists are dedicated cancer nurses available to talk to on our Macmillan Cancer Support Line.



Call us free on 0808 808 0000

8am to 8pm, 7 days a week. You can talk to our cancer specialists about anything.



Email the Macmillan Support Line

Email the Macmillan Support Team if you have any questions.

Chat online

Available 8am to 7.45pm, 7 days a week. For confidential, real-time support if you don't want to call.



Online Community

Chat online anonymously to others who understand what you are going through. Our community is available 24/7 and has dedicated forums where you can get advice and ask our experts.



Find cancer support services in your area

What's going on near you? Find out about support groups, where to get information and how to get involved with Macmillan where you live.

[Get help](#)