

Time left 2:44:26

Hide

Which one of the following is an appropriate recommendation for women of childbearing potential/pregnant with epilepsy?

- ☐ a. Initiate folic acid 1–4 mg daily to all women of childbearing potential
- ☐ b. Use combination therapy whenever possible
- ☐ c. Replace other antiepileptic medications with low dose of valproate
- ☐ d. Use the maximum possible doses that control seizure

Next page

Time left 2:44:14

100 mg of drug "A" administered orally and 70 mg of this drug reach systemic circulation chemically unchanged. What is the bioavailability of drug A?

- ☐ a. 35%
- ☐ b. 17.5%
- ☐ c. 90%
- ☐ d. 70%

Next page

Time left 2:44:44

Hide

A 45-year-old female farmer is taken to the emergency room. She was found confused in the plantation and since then has lost consciousness. Her heart rate is 45, and her blood pressure is 78/45 mm Hg. she is sweating and salivating abundantly. Which drug is more appropriate to manage her condition?

- ☐ a. Methanechol
- ☐ b. Edrophonium
- ☐ c. Neostigmine
- ☐ d. Atropine

Next page

Select a disadvantage of coiled or helical shaped packed chromatographic column?

- ☐ a. It cannot be repacked easily
- ☐ b. It is not compact
- ☐ c. It is not easy to heat it evenly
- ☐ d. It cannot be packed uniformly

Identify the correct statement regarding the advantage of extended-release dosage forms

- ☐ a. They deliver medication in a slow
- ☐ b. Patients may take the medication frequently
- ☐ c. Increase chance of drug side effects
- ☐ d. Increase daily medication cost

Time left 2:45:35

Hide

Four diclofenac tablets were assayed for drug content, and the following results were obtained by high-pressure liquid chromatography analysis: 401 mg, 402 mg, 398 mg, and 399 mg. what is the mean diclofenac content?

- ☐ a. 398 mg
- ☐ b. 400 mg
- ☐ c. 402 mg
- ☐ d. 401 mg

Next page

Time left 2:45:22

Which of the following describe the principle that health professionals should act in the best interest of the patient

- ☐ a. Honesty
- ☐ b. Autonomy
- ☐ c. Beneficence
- ☐ d. Informed Consent

Next page

Which of the following activities can be described as a process that has a result of 100?

- ☐ a. Collecting
- ☐ b. Storing
- ☐ c. Transporting
- ☐ d. Distributing



Time left 2:45:43

on 24

ed  
question

Which of the following is a technique pharmacists can use to humanize themselves to their patients?

- ☐ a. Delegating as many tasks as possible to an assistant
- ☐ b. Providing business cards at the desk
- ☐ c. Reading the text of a medication bottle verbatim to a patient
- ☐ d. Using the patient's preferred name in conversation

as page

Next

An 8-year-child with acute otitis media brought to clinic by his mother. What is the preferred empirical antibiotic for this child?

- ☐ a. Benzathine penicillin
- ☐ b. Erythromycin
- ☐ c. Ampicillin
- ☐ d. Amoxicillin

[Next page](#)

Time left 2:46:

Which criteria are correctly matched to the percolation method?

- ☐ a. The solid ingredients are moistened with an appropriate amount of the specified menstruum
- ☐ b. In this process, the crude drug is boiled in unspecified volume of water
- ☐ c. The mixture is allowed to macerate in the closed V shaped percolator for 24 hr.
- ☐ d. Adding solvent through the bed only once

Time left 2:46:26

Hide

Mrs. GG came to the hospital to seek medication for her sick young child and the physician prescribed her child otic drops. What is your counseling to the mother on the use of the medication?

- ☐ a. Position child laying down in supine position
- ☐ b. If child is above 3 years of age, gently pull upward and back; apply drops
- ☐ c. If child above 3 years of age, gently pull downward and back; apply drops
- ☐ d. If child is less 3 years of age, gently pull upward and back; apply drops

Next Page

Identify the correct statement about the pathophysiology of schizophrenia?

- ☐ a. Hyperdopaminergic activity in the mesolimbic pathway contributes to positive symptoms of psychosis
- ☐ b. Hypodopaminergic activity of the mesocortical pathway in the prefrontal cortex may contribute to negative and positive symptoms
- ☐ c. Hyperdopaminergic activity in the mesolimbic pathway contributes to negative symptoms of psychosis
- ☐ d. Hypodopaminergic activity of the mesocortical pathway in the prefrontal cortex may contribute to positive symptoms

Next page

Which statement is true about Auxochrome

- ☐ a. It is only saturated compound
- ☐ b. It is Decreased intensity of radiation
- ☐ c. They are saturated and unsaturated compound
- ☐ d. It is only unsaturated compound

16

Which antihypertensive drug has direct vasodilator mechanism of action?

- ☐ a. Labetalol
- ☐ b. Fenoldopam
- ☐ c. Hydralazine
- ☐ d. Methyldopa

Question

Page

Time left 2:46:45

Hide

TA is a month old male neonate admitted with the diagnosis of neonatal sepsis. He is on ampicillin and gentamicin for the past 3 days and has a clinical improvement. Which one is the best laboratory parameter used to assess the safety of gentamicin?

- ☐ a. Prothrombin time
- ☐ b. Platelet count
- ☐ c. Serum creatinine
- ☐ d. Blood glucose level



Next page

In acute phase management of ischemic stroke, initiation of one of the following antiplatelet agent within 48 hours of onset results in a decrease in the rate of stroke recurrence risk?

- ☐ a. Prasugrel
- ☐ b. Aspirin
- ☐ c. Clopidogrel
- ☐ d. Warfarin

Next page

Time left 2:47:30

Hi

Question 13

Not yet  
answered

Flag question

Which one of the following is considered as an analytical member under the establishment of pharmacy law and act?

- ☐ a. Drug Consultative Committee
- ☐ b. Licensing Authorities
- ☐ c. Drug Control Laboratories
- ☐ d. Drug inspectors

Previous page

Next page

Which of the following is a reasonable duration of antimicrobial therapy for patients admitted to the hospital with the diagnosis of type-II diabetes mellitus and osteomyelitis?

- ☐ a. 1 week
- ☐ b. 6 months
- ☐ c. 2 weeks
- ☐ d. 6 weeks

Which dosage form is considered enteral?

- ☐ a. Intravenous injection
- ☐ b. Suppository
- ☐ c. Intramuscular injection
- ☐ d. Oral tablet

Time left 2:47:51

Question 12

yet  
covered

flag question

Which one of the following ordering approach is most commonly used to organize pharmaceuticals in hospital pharmacy?

- ☐ a. Consumption rate
- ☐ b. Pharmaceutical orders
- ☐ c. Pharmacological
- ☐ d. Alphabetical

previous page

Next page

Time left 2:25:44

Hide

A 45-year-old overweight man came to ambulatory care unit for a diabetes management. He has been treated with metformin with poor blood glucose control despite good medication adherence. The physician decided to add a one day dose of long-acting insulin. What is the most likely medication given to this patient?

- ☐ a. Insulin lispro
- ☐ b. Insulin aspart
- ☐ c. Insulin glargine
- ☐ d. NPH insulin

Next page

Time left 2:25:37

Which one of the following is true about medication allergies?

- ☐ a. Cephalosporins should be avoided in patients with a history of delayed type of penicillin allergy
- ☐ b. If the offending agent is considered medically necessary, induction of drug tolerance is an option.
- ☐ c. Cross-reactive allergy between sulfonamide antibiotics and nonantibiotics is high
- ☐ d. Anaphylaxis is a chronic, life-threatening allergic reaction

Next page

Time left 2:18:47

Which of the following characterize the outbound logistics?

- ☐ a. Focus on deployment of resources and raw materials, within the manufacturing plant Customer service
- ☐ b. Interaction between the supplier and the firm
- ☐ c. Relate to material management and procurement
- ☐ d. Interaction between the firm and customers

Next page

Select antidepressants which has more sexual dysfunction side effect than the others

- ☐ a. Citalopram
- ☒ b. Paroxetine
- ☐ c. Bupropion
- ☐ d. Mirtazapine

Clear my choice

Select antidepressants which has more sexual dysfunction side effect than the others

- ☐ a. Citalopram
- ☐ b. Paroxetine
- ☐ c. Bupropion
- ☐ d. Mirtazapine

Question 90

Not yet

answered

Flag question

Which one is an undesirable separation of the different components of the blend?

- ☐ a. Segregation
- ☐ b. Levigation
- ☐ c. Fluidization
- ☐ d. Trituration

Previous page

Time left 2:18:30

Question 89

Answer saved

Flag question

A 56-year-old black man with uncontrolled hypertension is brought to the emergency ward for evaluation. His blood pressure is 191/100 mm Hg. He currently takes no medications. Which of the following drug will have limited efficacy in the treatment of this patient?

- ☐ a. Methyldopa
- ☐ b. Pindolol
- ☒ c. Hydralazine
- ☐ d. Labetalol

Clear my choice

Previous page

Next page

Time left 2:44:03

Hide

You are intended to sterilize sealed containers that need a very high penetration power, which one of the following is a very effective sterilization technique that combines heat and moisture that fit for this purpose?

- ☐ a. Ethylene oxide sterilization
- ☐ b. Autoclave
- ☐ c. Lyophilization
- ☐ d. Dry heat

Next page

Differentiate the separation process used to separate immiscible liquids

- ☐ a. Distillation
- ☐ b. Centrifugation
- ☐ c. Precipitation
- ☐ d. Expression

Time left 2:44:26

Hide

Which one of the following is an appropriate recommendation for women of childbearing potential/pregnant with epilepsy?

- ☐ a. Initiate folic acid 1–4 mg daily to all women of childbearing potential
- ☐ b. Use combination therapy whenever possible
- ☐ c. Replace other antiepileptic medications with low dose of valproate
- ☐ d. Use the maximum possible doses that control seizure

Next page

Time left 2:43:53

Identify the correct criteria to define mild-persistent asthma presentation in Children 0–4 Years of Age

- ☐ a. SABA use for symptom control  $\leq 2$  days/week
- ☐ b. Minor Interference with normal activity
- ☐ c. No Interference with normal activity
- ☐ d. Symptom of asthma  $\leq 2$  days/week

Next page

Time left 2:44:03

Hide

You are intended to sterilize sealed containers that need a very high penetration power, which one of the following is a very effective sterilization technique that combines heat and moisture that fit for this purpose?

- ☐ a. Ethylene oxide sterilization
- ☐ b. Autoclave
- ☐ c. Lyophilization
- ☐ d. Dry heat

Next page

Differentiate the separation process used to separate immiscible liquids

- ☐ a. Distillation
- ☐ b. Centrifugation
- ☐ c. Precipitation
- ☐ d. Expression

In most syrup preparations in addition to active ingredients and purified water, sucrose is most commonly employed for the intention to:

- ☐ a. mask the taste of added medicinal agents
- ☐ b. decrease viscosity of the preparation
- ☐ c. enhance the growth bacteria
- ☐ d. make the preparation colorful and attractive

Which of the following is a role of Pharmacovigilance?

- ☐ a. Monitoring Drug safety
- ☐ b. Monitoring Drug price
- ☐ c. Monitoring pharmacy students
- ☐ d. Monitoring unethical practices

Question 54

Not yet

answered

Flag question

A 6-year-old boy is confirmed to have HIV and the physician is planning to start him on first-line combination ART. Which of the following combinations will be started for him?

- ☐ a. ABC + 3TC + DTG
- ☐ b. ABC + 3TC + LPV/r
- ☐ c. ABC + 3TC + EFV
- ☐ d. TDF + 3TC + DTG

Previous question

Time left

A 6-year-old boy is confirmed to have HIV and the physician is planning to start him on first-line ART. Which of the following combination ART will be started for him?

- ☐ a. ABC + 3TC + DTG
- ☐ b. ABC + 3TC + LPV/r
- ☐ c. ABC + 3TC + EFV
- ☐ d. TDF + 3TC + DTG

1. Choose the advantage of primary literature?

- ☐ a. Expensive to access
- ☐ b. Limited in scope
- ☐ c. Time consuming
- ☐ d. More detailed

Time left 1

Which of the following is not a result of activation of the intra-cardiac renin-angiotensin aldosterone system?

- ☐ a. Fibrosis
- ☐ b. Progressive hyperplasia
- ☐ c. Cardiac remodeling
- ☐ d. Progressive hypertrophy

Time left 1:36:35

GG is a 20-year-old man diagnosed with schizophrenia and put on 5 mg po olanzapine daily. When will be the time to assess the clinical response for GG?

- ☐ a. Within 4-6 months of treatment
- ☐ b. Within the 2-3 weeks of treatment
- ☐ c. Within 4-6 weeks of treatment
- ☐ d. Within 6 months of treatment

Next

Choose the rate-limiting step for bioavailability of most conventional solid drug products:

- ☐ a. The osmosis rate across the membrane
- ☐ b. The diffusion rate across the membrane
- ☐ c. The release of the drug from the delivery system
- ☐ d. Dissolution rate of the product

Which of the following is a diluent used in pharmaceutical industry?

- ☐ a. Magnesium stearate
- ☐ b. Honey
- ☐ c. Glucose
- ☐ d. Glycyrrhiza

A period of time where a drug stored under reasonable conditions is:

- ☐ a. Manufacturing date
- ☐ b. Half-life
- ☐ c. Shelf-life
- ☐ d. Expiration date

Time left 1:37

Which of the following compounds tend to accumulate at interface and reduce surface or interfacial tension

- ☐ a. Surfactants
- ☐ b. Antifoaming agents
- ☐ c. Wetting agents
- ☐ d. Detergents

Which form of drug requires metabolic transformation to change into pharmacologically active

- ☐ a. Teratogen drug
- ☐ b. Prodrug
- ☐ c. Sustain released drug
- ☐ d. Orphan drug

8 Identify the correct statement regarding the advantage of extended-release dosage forms

- ☐ a. They deliver medication in a slow
- ☐ b. Patients may take the medication frequently
- ☐ c. Increase chance of drug side effects
- ☐ d. Increase daily medication cost

Identify the role of indicator in a reaction

- ☐ a. To identify the end point of the titration
- ☐ b. To increase the rate of a reaction
- ☐ c. Assist reactants to react quickly
- ☐ d. To provide a surface for the reaction

Which of the following is an example of tertiary source of information?

- ☐ a. Pharmacotherapy textbook
- ☐ b. European journal of medicine
- ☐ c. Google scholar
- ☐ d. PubMed

DEEP

Time left 2:15:44

Choose the best combination of opioid withdrawal symptoms and its respective treatment regimen

- ☐ a. Diarrhea: clonidine 0.1–0.2 mg every 6–8 hours
- ☐ b. Insomnia: trazodone 75–200 mg at bedtime
- ☐ c. Abdominal cramps: acetaminophen 500–1000 mg every 6 hours
- ☐ d. Muscle aches: dicyclomine 10–20 mg every 6 hours

Next

n 56

d

question

Identify the first line medication for treatment of low back pains?

- ☐ a. Anticonvulsant
- ☐ b. Antidepressant
- ☐ c. Acetaminophen
- ☐ d. Morphine

ous page

Time left 2:15:51

A 6-year-old boy is confirmed to have HIV and the physician is planning to start him on first-line ART. Which of the following combination ART will be started for him?

- ☐ a. ABC + 3TC + DTG
- ☐ b. ABC + 3TC + LPV/r
- ☐ c. ABC + 3TC + EFV
- ☐ d. TDF + 3TC + DTG

Next page

Question 59

Not yet  
answered

Flag question

Which of the following department or service in a hospital is under the direction of qualified pharmacist which all medications are supplied to the nursing units and other services.

- ☐ a. Hospital pharmacy
- ☐ b. OPD pharmacy
- ☐ c. Community pharmacy
- ☐ d. Inpatient pharmacy

Previous page



Time left 2:16:02

Hide

GG is a 20-year-old man diagnosed with schizophrenia and put on 5 mg po olanzapine daily. When will be the right time to assess the clinical response for GG?

- ☐ a. Within 4-6 months of treatment
- ☒ b. Within the 2-3 weeks of treatment
- ☐ c. Within 4-6 weeks of treatment
- ☐ d. Within 6 months of treatment

Next page

Select antidepressants which has more sexual dysfunction side effect than the others

- ☐ a. Citalopram
- ☐ b. Paroxetine
- ☐ c. Bupropion
- ☐ d. Mirtazapine

Select antidepressants which has more sexual dysfunction side effect than the others

- ☐ a. Citalopram
- ☒ b. Paroxetine
- ☐ c. Bupropion
- ☐ d. Mirtazapine

Clear my choice

Time left 2:18:47

Which of the following characterize the outbound logistics?

- ☐ a. Focus on deployment of resources and raw materials, within the manufacturing plant Customer service
- ☐ b. Interaction between the supplier and the firm
- ☐ c. Relate to material management and procurement
- ☐ d. Interaction between the firm and customers

Next page

Question 90

Not yet  
answered

Flag question

Which one is an undesirable separation of the different components of the blend?

- ☐ a. Segregation
- ☐ b. Levigation
- ☐ c. Fluidization
- ☐ d. Trituration

Previous page

Time left 2:18:30

Question 89

Answer saved

Flag question

A 56-year-old black man with uncontrolled hypertension is brought to the emergency ward for evaluation. His blood pressure is 191/100 mm Hg. He currently takes no medications. Which of the following drug will have limited efficacy in the treatment of this patient?

- ☐ a. Methyldopa
- ☐ b. Pindolol
- ☒ c. Hydralazine
- ☐ d. Labetalol

Clear my choice

Previous page

Next page

Time left 2:16:10

Which of the following is not a result of activation of the intra-cardiac renin-angiotensin aldosterone system?

- ☐ a. Fibrosis
- ☐ b. Progressive hyperplasia
- ☐ c. Cardiac remodeling
- ☐ d. Progressive hypertrophy

Next page

Time left 2:25:44

Hide

A 45-year-old overweight man came to ambulatory care unit for a diabetes management. He has been treated with metformin with poor blood glucose control despite good medication adherence. The physician decided to add a one day dose of long-acting insulin. What is the most likely medication given to this patient?

- ☐ a. Insulin lispro
- ☐ b. Insulin aspart
- ☐ c. Insulin glargine
- ☐ d. NPH insulin

Next page

Time left 2:25:37

Which one of the following is true about medication allergies?

- ☐ a. Cephalosporins should be avoided in patients with a history of delayed type of penicillin allergy
- ☐ b. If the offending agent is considered medically necessary, induction of drug tolerance is an option.
- ☐ c. Cross-reactive allergy between sulfonamide antibiotics and nonantibiotics is high
- ☐ d. Anaphylaxis is a chronic, life-threatening allergic reaction

Next page

Time left 2:25:3

n 93

d

question

Mrs. GG is presented to ophthalmologic clinic for her complaint of open-angle glaucoma. Which drug is contraindicated to Mrs. GG?

- ☐ a. Tropicamide
- ☐ b. Physostigmine
- ☐ c. Atropine
- ☐ d. Pilocarpine

ous page

Next pa

Time left 2:26:11

Hide

In most syrup preparations in addition to active ingredients and purified water, sucrose is most commonly employed for the intention to:

- ☐ a. mask the taste of added medicinal agents
- ☐ b. decrease viscosity of the preparation
- ☐ c. enhance the growth bacteria
- ☐ d. make the preparation colorful and attractive

Next page

Which of the following is a role of Pharmacovigilance?

- ☐ a. Monitoring Drug safety
- ☐ b. Monitoring Drug price
- ☐ c. Monitoring pharmacy students
- ☐ d. Monitoring unethical practices

Time left 2:25

From the following non-pharmacologic treatment option for hypertensive patients, differentiate highly effective approach to reduce systolic pressure?

- ☐ a. Dietary sodium restriction
- ☐ b. Dietary approaches
- ☐ c. Moderation of alcohol consumption
- ☐ d. Weight reduction

Next

Which of the following characterize the outbound logistics?

- ☐ a. Focus on deployment of resources and raw materials, within the manufacturing plant
- ☐ b. Interaction between the supplier and the firm
- ☐ c. Relate to material management and procurement
- ☐ d. Interaction between the firm and customers

Time left 2:28:48

Child with suspected infection was admitted to pediatric care unit and treated with antibiotics with 3 consecutive days. However, her body temperature is persistently increased and currently she developed symptom of febrile seizure. What will your recommendation for this child?

- ☐ a. Waiting for antibiotic response
- ☐ b. Phenytoin
- ☐ c. Phenobarbital
- ☐ d. Suppository acetaminophen

Finish attempt

macy

Time

Outline the preferred chromatographic technique to separate macromolecules, ionic species, labile polymeric materials and high-molecular weight compounds from non-volatile and thermally stable

- ☐ a. Thin layer chromatography
- ☐ b. High Performance Liquid Chromatography
- ☐ c. Gas solid chromatography
- ☐ d. Paper chromatography

Which of the following is a required strategy to improve communication?

- ☐ a. Explain things in complex language
- ☐ b. Do not use "teach back" or "show me" technique to check understanding
- ☐ c. Focus on key messages and repeat
- ☐ d. Do not use patient-friendly educational materials to enhance interaction

Which of the following is prepared from any weak acid or base and are used to maintain a narrow range?

- ☐ a. Surfactant
- ☐ b. Wetting agent
- ☐ c. Detergent
- ☐ d. Buffer

Which the following is the disadvantage of ultrasonic extraction?

- ☐ a. Low cost
- ☐ b. Formation of free radical
- ☐ c. High speed
- ☐ d. Less time

Question 83

Not yet  
answered

Flag question

Separation of particles of one size in one area and another size in another area is called:

- ☐ a. Diffusive Mixing
- ☐ b. Convective Mixing
- ☐ c. Shear Mixing
- ☐ d. Segregation

Previous page

2  
A pharmacist acts with honesty and integrity in professional relationships. This is interpreted as:

- ☐ a. The pharmacist is responsible for fair balancing the needs of patients and society.
- ☐ b. The primary obligation of a pharmacist is to individual patients.
- ☐ c. The pharmacist has a duty to tell the truth and to act with conviction of conscience.
- ☐ d. The pharmacist considers needs stated by the patient as well as those defined by health science.

Which of the following are characteristics of O/W emulsions/creams formulations?

- ☐ a. Greasy in nature and hydrate the skin if applied topically.
- ☐ b. Oil is an internal phase and water is the dispersion medium.
- ☐ c. Water as dispersed phase and oil as dispersion medium.
- ☐ d. Creams are used for systemic effect of water-soluble drug applied on the skin.

A physician wants to switch a terminally-ill patient from slow release morphine sulphate tablets, 15 mg daily, to a liquid morphine sulphate dosage form because the patient has difficulty in swallowing tablets. A morphine sulphate solution containing 5 mg per mL is prescribed q4h, what volume should be dispensed for a 7-day supply to provide the same pain relief as the tablet regimen?

- ☐ a. 120 mL
- ☐ b. 20 mL
- ☐ c. 100 mL
- ☐ d. 60 mL

Back

Question 77

Not yet  
answered

Flag question

One of the drugs for the management of transient ischemic attacks is clopidogrel. Which of the threatening adverse effects?

- ☐ a. Stroke
- ☐ b. Hyperlipidemia
- ☐ c. Thrombotic thrombocytopenic purpura
- ☐ d. Hyperthermia

Previous page

Which one of the following term best explains the amount of moisture present in a material at ambient and it is dependent on the humidity, temperature, and nature of material.

- ☐ a. Equilibrium Moisture Content
- ☐ b. Percent Loss on Drying
- ☐ c. Total Moisture Content
- ☐ d. Bound Water

AG is 48-year-old patient admitted with the diagnosis of ST-elevation myocardial infarction, with contraindications for fibrinolysis. Within how many minutes of presentation is recommended for administration?

- ☐ a. 30 minutes
- ☐ b. 60 minutes
- ☐ c. 120 minutes
- ☐ d. 90 minutes

Which drug is a direct acting cholinomimetic?

- ☐ a. Bethanechol
- ☐ b. Echothiophate
- ☐ c. Endrophonium
- ☐ d. Physostigmine

Which of the following defines a list of drugs that have been approved for use in hospitals by the pharmacy and therapeutics committee and become standard stock carried by the pharmacy and other departments?

- ☐ a. Standard treatment guideline
- ☐ b. Formulary list
- ☐ c. Medicine consumption reporting
- ☐ d. Formulary manual

# pharmacy

Back

Time left 1:14:54

Hide

Question 67

Answer saved

Flag question

Which of the following drug is recommended as first-line for treatment of heart failure with reduced ejection fraction?

- ☒ a. Beta-blocker
- ☐ b. ACE inhibitors
- ☐ c. Loop diuretics
- ☐ d. Digoxin

Clear my choice

Exa



Moh

1

10

19

28

37

46



# Pharmacy

Back

Time left 1:15:05

Question 66

Not yet answered

Flag question

GR is a 60 year-old women, who is a known type-II diabetic patient and on metformin for the past 10 years. Currently, she is diagnosed with peripheral neuropathy. Which drug is recommended as first-line treatment for GR?

- ☐ a. Ibuprofen
- ☐ b. Acetaminophen
- ☐ c. Morphine
- ☐ d. Gabapentin

Previous page

Next page

Back

Time left 1:15:15

Question 65

Not yet answered

Flag question

A patient with an open-wound need an irrigation, which one of the following is an ideal important parameter for preparations used in this case?

- ☐ a. Must be solution and sterile
- ☐ b. Must be solution and Pyrogenic
- ☐ c. Must be semisolid and sterile
- ☐ d. Must be semisolid and pyrogenic

Previous page

Next page

Back

Question 64

Not yet  
answered

Flag question

Previous page

Time left 1:15:31

In critical appraisal of clinical literatures, which one of the following question is key to answer the generalizability of the results?

- ☐ a. Are the patients in the study representative of the patient(s) I am treating?
- ☐ b. Was the sample size adequate?
- ☐ c. Did the study results answer the research question?
- ☐ d. What is the research question?

Next page

1

10

19

28

37

46

55

64

73

# Pharmacy

Six

Time left 1:15:45

Question 63

Not yet  
answered

Flag question

Why low-molecular-weight heparins are preferred over unfractionated heparin to reduce the risk of deep vein thrombosis?

- ☐ a. Easier to manage for outpatients
- ☐ b. Better inhibitor of thrombin
- ☐ c. Does not cause HIT
- ☐ d. Carries no risk of bleeding

Next page

Time left 1:37

Which of the following compounds tend to accumulate at interface and reduce surface or interfacial tension

- ☐ a. Surfactants
- ☐ b. Antifoaming agents
- ☐ c. Wetting agents
- ☐ d. Detergents

Which of the following is a diluent used in pharmaceutical industry?

- ☐ a. Magnesium stearate
- ☐ b. Honey
- ☐ c. Glucose
- ☐ d. Glycyrrhiza

Choose the rate-limiting step for bioavailability of most conventional solid drug products:

- ☐ a. The osmosis rate across the membrane
- ☐ b. The diffusion rate across the membrane
- ☐ c. The release of the drug from the delivery system
- ☐ d. Dissolution rate of the product

1. Choose the advantage of primary literature?

- ☐ a. Expensive to access
- ☐ b. Limited in scope
- ☐ c. Time consuming
- ☐ d. More detailed

Time left 1

Which of the following is not a result of activation of the intra-cardiac renin-angiotensin aldosterone system?

- ☐ a. Fibrosis
- ☐ b. Progressive hyperplasia
- ☐ c. Cardiac remodeling
- ☐ d. Progressive hypertrophy

Time left 1:36:35

GG is a 20-year-old man diagnosed with schizophrenia and put on 5 mg po olanzapine daily. When will be the time to assess the clinical response for GG?

- ☐ a. Within 4-6 months of treatment
- ☐ b. Within the 2-3 weeks of treatment
- ☐ c. Within 4-6 weeks of treatment
- ☐ d. Within 6 months of treatment

Next

A 6-year-old boy is confirmed to have HIV and the physician is planning to start him on first-line ART. Which of the following combination ART will be started for him?

- ☐ a. ABC + 3TC + DTG
- ☐ b. ABC + 3TC + LPV/r
- ☐ c. ABC + 3TC + EFV
- ☐ d. TDF + 3TC + DTG

Question 54

Not yet  
answered

Flag question

A 6-year-old boy is confirmed to have HIV and the physician is planning to start him on first-line combination ART. Which of the following combinations will be started for him?

- ☐ a. ABC + 3TC + DDI
- ☐ b. ABC + 3TC + LPV/r
- ☒ c. ABC + 3TC + EFV
- ☐ d. ZDV + 3TC + DDI

Previous page

Choose the best combination of opioid withdrawal symptoms and its respective treatment regimen

- ☐ a. Diarrhea: clonidine 0.1–0.2 mg every 6–8 hours
- ☐ b. Insomnia: trazodone 75–200 mg at bedtime
- ☐ c. Abdominal cramps: acetaminophen 500–1000 mg every 6 hours
- ☐ d. Muscle aches: dicyclomine 10–20 mg every 6 hours

Which the following is the disadvantage of ultrasonic extraction?

- ☐ a. Low cost
- ☐ b. Formation of free radical
- ☐ c. High speed
- ☐ d. Less time

Which drug is a direct acting cholinomimetic?

- ☐ a. Bethanechol
- ☐ b. Echothiophate
- ☐ c. Endrophonium
- ☐ d. Physostigmine

Which drug is a direct acting cholinomimetic?

- ☐ a. Bethanechol
- ☐ b. Echothiophate
- ☐ c. Endrophonium
- ☐ d. Physostigmine

Which the following is the disadvantage of ultrasonic extraction?

- ☐ a. Low cost
- ☐ b. Formation of free radical
- ☐ c. High speed
- ☐ d. Less time

Time left 0:50:06

Hi

Which of the following defines a list of drugs that have been approved for use in hospitals by the pharmacy and therapeutics committee and become standard stock carried by the pharmacy and other departments?

- ☐ a. Standard treatment guideline
- ☐ b. Formulary list
- ☐ c. Medicine consumption reporting
- ☐ d. Formulary manual

Next page

Time left 0:50:06

Hi

Which of the following defines a list of drugs that have been approved for use in hospitals by the pharmacy and therapeutics committee and become standard stock carried by the pharmacy and other departments?

- ☐ a. Standard treatment guideline
- ☐ b. Formulary list
- ☐ c. Medicine consumption reporting
- ☐ d. Formulary manual

Next page

Which one of the following is correct about Integrated Pharmaceutical Logistic System inventory control system?

- ☐ a. There is no min/max inventory control system
- ☐ b. Facilities report not based the stock on hand
- ☐ c. The system is a forced order inventory control system
- ☐ d. Emergency order point and review period are the triggering points to report

59

Which of the following objective finding for diabetic mellitus is crucial to know the blood glucose level over past 3 month?

- ☐ a. Random plasma glucose level
- ☐ b. Hemoglobin A1c
- ☐ c. Plasma insulin level
- ☐ d. Oral glucose tolerance test

Time left 0:50:43

Hi

Question 66

Not yet  
answered

Flag question

GR is a 60-year-old woman, who is a known type-II diabetic patient and on metformin for the past 10 years. Currently, she is diagnosed with peripheral neuropathy. Which drug is recommended as first-line treatment for GR?

- ☐ a. Ibuprofen
- ☐ b. Acetaminophen
- ☐ c. Morphine
- ☐ d. Gabapentin

Previous page

Next page

Time left 0:50:58

H

A patient with an open-wound need an irrigation, which one of the following is an ideal important parameter for the preparations used in this case?

- ☐ a. Must be solution and sterile
- ☐ b. Must be solution and Pyrogenic
- ☐ c. Must be semisolid and sterile
- ☐ d. Must be semisolid and pyrogenic

Which one of the following term best explains the amount of moisture present in a material at ambient and it is dependent on the humidity, temperature, and nature of material.

- ☐ a. Equilibrium Moisture Content
- ☐ b. Percent Loss on Drying
- ☐ c. Total Moisture Content
- ☐ d. Bound Water

Back

Question 77

Not yet  
answered

Flag question

One of the drugs for the management of transient ischemic attacks is clopidogrel. Which of the threatening adverse effects?

- ☐ a. Stroke
- ☐ b. Hyperlipidemia
- ☐ c. Thrombotic thrombocytopenic purpura
- ☐ d. Hyperthermia

Previous page

A physician wants to switch a terminally-ill patient from slow release morphine sulphate tablets, 15 mg daily, to a liquid morphine sulphate dosage form because the patient has difficulty in swallowing tablets. If a morphine sulphate solution containing 5 mg per mL is prescribed q4h, what volume should be dispensed for a 7-day supply to provide the same pain relief as the tablet regimen?

- ☐ a. 120 mL
- ☐ b. 20 mL
- ☐ c. 100 mL
- ☐ d. 60 mL

Which of the following are characteristics of O/W emulsions/creams formulations?

- ☐ a. Greasy in nature and hydrate the skin if applied topically.
- ☐ b. Oil is an internal phase and water is the dispersion medium.
- ☐ c. Water as dispersed phase and oil as dispersion medium.
- ☐ d. Creams are used for systemic effect of water-soluble drug applied on the skin.

2  
A pharmacist acts with honesty and integrity in professional relationships. This is interpreted as:

- ☐ a. The pharmacist is responsible for fair balancing the needs of patients and society.
- ☐ b. The primary obligation of a pharmacist is to individual patients.
- ☐ c. The pharmacist has a duty to tell the truth and to act with conviction of conscience.
- ☐ d. The pharmacist considers needs stated by the patient as well as those defined by health science.

Question 83

Not yet  
answered

Flag question

Separation of particles of one size in one area and another size in another area is called:

- ☐ a. Diffusive Mixing
- ☐ b. Convective Mixing
- ☐ c. Shear Mixing
- ☐ d. Segregation

Previous page

Which the following is the disadvantage of ultrasonic extraction?

- ☐ a. Low cost
- ☐ b. Formation of free radical
- ☐ c. High speed
- ☐ d. Less time

Which of the following is prepared from any weak acid or base and are used to maintain a narrow range?

- ☐ a. Surfactant
- ☐ b. Wetting agent
- ☐ c. Detergent
- ☐ d. Buffer

Which of the following is a required strategy to improve communication?

- ☐ a. Explain things in complex language
- ☐ b. Do not use "teach back" or "show me" technique to check understanding
- ☐ c. Focus on key messages and repeat
- ☐ d. Do not use patient-friendly educational materials to enhance interaction

nacy

Time

Outline the preferred chromatographic technique to separate macromolecules, ionic species, labile polymeric materials and high-molecular weight compounds from non-volatile and thermally stable

- ☐ a. Thin layer chromatography
- ☐ b. High Performance Liquid Chromatography
- ☐ c. Gas solid chromatography
- ☐ d. Paper chromatography

01 Which of the following characterize the outbound logistics?

- ☐ a. Focus on deployment of resources and raw materials, within the manufacturing plant Custom
- ☐ b. Interaction between the supplier and the firm
- ☐ c. Relate to material management and procurement
- ☐ d. Interaction between the firm and customers

Time left 0:5

Question 64

yet  
answered

flag question

In critical appraisal of clinical literatures, which one of the following question is key to answer the generalization of the results?

- ☐ a. Are the patients in the study representative of the patient(s) I am treating?
- ☐ b. Was the sample size adequate?
- ☐ c. Did the study results answer the research question?
- ☐ d. What is the research question?

previous page

Next