

4. A 25-year-old woman came to a family planning clinic with complaints of irregular frequent menstrual bleeding after she took combined oral contraceptives. She has heavy bleeding that lasts more than seven days.

What is the most likely menstrual pattern for this woman?

- (A) Metorrhagia
- (B) Menorrhagia
- (C) Polymenorrhea
- (D) Menometrorrhagia

5. A married couple came to a hospital for infertility investigations. They were trying to achieve conception for the last two years. They do not have previous history of contraception usage. Accordingly the gynecologist ordered laboratory investigations for the couple.

What is the most likely first laboratory investigation to be done for this couple?

- (A) Urine HCG test
- (B) Semen analysis
- (C) Hormonal assay
- (D) Serum FSH level

6. A 30-year-old mother was admitted to gynecology ward with vesico-vaginal fistula secondary to obstructed labor. The mother gave birth at home before 2 months. She is scheduled for fistula repair after a week.

What is the most appropriate nursing intervention for this mother?

- (A) Emotional support
- (B) Insert catheter in the bladder
- (C) Maintain patient on clear liquids
- (D) Encourage frequent sitz baths

7. An 18-year-old lady came to a gynecology OPD with a complaint of abdominal distension for one month. She has abdominal swelling, and dull like abdominal pain. On examination, liver is enlarged, firm and nodular. The imaging result showed that the tumor is unilateral, solid, more than 10 cm in diameter which has originated from the primitive yolk sack.

What is the most likely diagnosis of this woman?

- (A) Dysgerminoma tumor
- (B) Choriocarcinoma tumor
- (C) Immature teratoma tumor
- (D) Endodermal sinus tumor

8. A 26-year-old non-pregnant woman came to a hospital with complaints of Inter-menstrual and post-coital vaginal bleeding for three months. CT scan investigation result showed a cervical tumor which involves lower third of the vagina, without extension to the pelvic wall. She was diagnosed for stage II.

What is the most likely manifestation that corresponds to this stage?

- (A) Dry and itchy vaginal orifice
- (B) Adenexial excitation tenderness
- (C) Offensive and watery vaginal discharge
- (D) Excessive fluid accumulation in culdosa-sac

30. A 45-year-old male patient came to an emergency department with complaints of abdominal pain, vomiting and constipation of 12 hours duration. The patient reports that he has previous history of hernia. On physical examination, the nurse identified distended abdomen, high pitched bowel sounds, but no detected masses.

What is the next course of the disorder if left untreated?

- (A) Passed flatus; small and formed stool
- (B) Intoxicated patient with disarranged vital sign
- (C) More intense pain and slowly dehydrated patient
- (D) Gradual decrease in vomiting and abdominal distention

31. During assessment of a patient with bilateral lower extremity edema, a nurse applies pressure on edematous area for 5 seconds using thumb and identified 3cm indentation.

What is the most likely edema grade of this patient?

- (A) One pitting edema
- (B) Two pitting edema
- (C) Three pitting edema
- (D) Four pitting edema

32. A 60-year-old female patient was admitted to a medical ward with a diagnosis of chronic renal failure. Laboratory result shows serum potassium level of 7mEq/L (normal value 3.5-5.0). On assessment she has ECG changes (tall-tented T waves and widened QRS). The health care provider diagnosed hyperkalemia.

What early signs and symptoms could support the diagnosis?

- (A) Muscle weakness and diarrhea
- (B) Blood pressure and fluid retention
- (C) Fever, chest pain and additive breath sound
- (D) RBC count, hemoglobin and hematocrit level

33. A 76-year-old woman was admitted to a medical ward with complaints of edema, rash, and shortness of breath. She urinate 120 ml of urine per 24 hours. The nurse diagnosed the patient with chronic renal failure empirically.

What is the possible additional manifestation which supported the nurse's decision?

- (A) Decrease BUN
- (B) Acute weight loss
- (C) Clear breath sound
- (D) Decrease hematocrit

25. A 23-year-old chronic liver disease patient was admitted to a medical ward with a diagnosis of hepatocellular carcinoma. The community where the patient came uses running water for drinking. He has no history of IV drug use and previous blood transfusion.

What is the most likely etiology for the patient's diagnosis?

- (A) Hepatitis A Virus
- (B) Hepatitis B Virus
- (C) Hepatitis C Virus
- (D) Hepatitis D Virus

26. A 36-year-old male patient is admitted in surgical ward with confirmed diagnosis of appendicitis. The patient is prepared for appendectomy and received preoperative care.

What is the most appropriate preoperative care for this patient?

- (A) Administer enema
- (B) Encourage ambulation
- (C) Administer antibiotic
- (D) Stimulates bowel activity

27. A 26-year-old male patient, who works in a construction firm, was brought to an emergency department. The patient was diagnosed with large bowel obstruction empirically.

What is the most likely clinical presentation that helps to diagnose this patient?

- (A) Vomiting
- (B) Dehydration
- (C) Gross abdominal distention
- (D) Bowel sound of 4/5 minute

28. A 64-year-old female patient is admitted to a medical ward with a complaint of unconsciousness for four days duration. The unconsciousness is due to head trauma that leads excessive blood loss. Naso-gastric intubation is option less to feed her. The available tube has no mark that the nurse should determine the length of NGT needed to reach the stomach.

What is the most likely technique to be used in this case?

- (A) From tip of the nose to the earlobe and from the earlobe to the umbilicus
- (B) From the earlobe to tip of the nose and from tip of nose to the xiphoid process
- (C) From tip of the nose to the earlobe and from the earlobe to the xiphoid process
- (D) From tip of the nose to the earlobe to half way between xiphoid process and umbilicus

29. A 62-year-old obese man came to a medical outpatient department seeking treatment. The nurse on duty diagnosed the patient with duodenal peptic ulcer disease based on the clinical manifestations and history of the patient.

What is the most likely clinical manifestation used by the nurse to diagnosis this patient?

- (A) Ingestion of food relieve pain
- (B) Pain occur one hour after meal
- (C) Being obese
- (D) Relieved by vomiting

9. A child is brought to an emergency department with dislodged fragment of toy part in the airway. He has violent paroxysms of coughing and choking accompanied by new onset wheezing, intercostal muscle retraction and cyanosis. His respiratory rate is 62/min and pulse rate is 100/min.

What is the most likely actual nursing diagnosis that needs priority for this child?

- (A) Impaired gas exchange
(B) Ineffective airway clearance
(C) Ineffective breathing pattern
(D) Impaired spontaneous ventilation
10. An unconscious seven-year-old child has sustained multiple body injury secondary to vehicle accident. The child was brought to hospital by bystanders and a nurse in charge suspected neck and back injury.

What is the initial step that the nurse should do for this child?

- (A) Perform jaw thrust
(B) Clear secretions from throat
(C) Inspect mouth and remove foreign body
(D) Check the airway by looking for chest movements
11. A preschool-age child with previous history of childhood asthma was brought to pediatric emergency unit. The child has a manifestation of peak expiratory flow 30%, oxygen saturation 78% and loud wheezing throughout inhalation and exhalation.

What is the actual nursing diagnosis that needs priority for this child?

- (A) Impaired gas exchange
(B) Ineffective airway clearance
(C) Ineffective breathing pattern
(D) Impaired spontaneous ventilation
12. A nine-year-old child was brought to a pediatric OPD with complaints of diarrhea, non-bilious and projectile vomiting. On examination, there was low grade fever.

What is the most likely diagnosis of this child?

- (A) Intussusceptions
(B) Pyloric stenosis
(C) Intestinal volvulus
(D) Hirschsprung disease
13. A nurse planned to give health education to a mother of a seven year-old-child who came with a disease called juvenile idiopathic arthritis.

What is the most appropriate advice the nurse could give to the mother?

- (A) Restrict from fatty foods during severe form of illness
(B) Restrict from exercise during moderate form of illness
(C) Restrict from protein foods during severe form of illness
(D) Avoid systemic steroids during moderate form of illness

34. A nurse on duty is planning nursing intervention for a patient admitted to medical ward with acute renal failure. The laboratory finding showed decreased serum phosphorus and calcium level.

What is the most important nursing intervention for this patient?

- (A) Administer blood components therapy
- (B) Assisting the patient with an exercise program
- (C) Teaching the patient to report sign of fluid overload
- (D) Encourage compliance with fluid restriction therapy

35. A nurse is assessing a male patient who is admitted in medical ward with acute renal failure. On physical examination, he has dry mucous membrane, poor skin turgor, flat neck vein and orthostatic hypotension. He is sick looking and irritable.

What is the primary nursing diagnosis for this patient?

- (A) Risk for altered skin integrity
- (B) Fluid volume deficit
- (C) Deficient knowledge
- (D) Activity intolerance

36. A 33-year-old patient came to a hospital with complaints of peripheral edema and cola-colored urine. The patient reported a history of frequent attack of tonsillitis and skin infections. On laboratory investigation, the patient's Glomerular Filtration Rate reads 100ml/minute/1.73m² (normal GFR is 125ml/min/1.73 m²).

What is the most likely fluid and electrolyte balance change expected from the patient?

- (A) Hypokalemia
- (B) Hypernatremia
- (C) Marked hypertension
- (D) Decreased urine output

37. A 63-year-old patient was brought to medical unit with complaints of sudden onset of visual disturbance, paralysis on one side of the body and severe headache. Assessment findings showed history of poorly controlled hypertension.

What is the most likely diagnosis of this patient?

- (A) Embolic stroke
- (B) Ischemic Stroke
- (C) Thrombotic stroke
- (D) Hemorrhagic Stroke

38. A 36-year-old patient was presented with complaints of severe headache, neck stiffness and fever. The nurse and the physician planned to do lumbar puncture to send sample for laboratory investigation.

What is the most likely contraindication for this procedure?

- (A) Decreased level of consciousness
- (B) Decreased intracranial pressure
- (C) Infection on proposed site
- (D) Increased blood pressure

39. A 72-year-old patient came to a medical unit with complaints of sudden onset of visual hemiparesis, confusion, trouble speaking, and impaired mobility. Assessment findings showed history of poorly controlled hypertension. The physician in charge of the patient stroke and put the patient on oxygen.

What is the initial treatment option for this patient?

- (A) Anti-platelet
- (B) Thrombolytic
- (C) Anti-hypertensive
- (D) Analgesics treatment

40. A 46-year-old female patient was admitted to surgical inpatient unit with diagnosis of subdural hematoma. Assessment findings showed decreased level of blood pressure and oxygen saturation. The nurse administered nasal oxygen and planned to evaluate the prognosis.

What is the most likely evidence for good prognosis of the given intervention?

- (A) SpO₂ of at least 90%
- (B) Respiratory rate of 28/min
- (C) PaCO₂ at least 52mmHg
- (D) PaO₂ is at least 70mmHg

41. A 72-year-old female patient was presented with complaints of numbness of the extremities, blurred vision and increased appetite. The patient reported that she has history of chronic alcohol consumption and visiting traditional healers for controlling blurred vision and other related health problems. Her random blood glucose level is 320mg/dL.

What is the priority nursing diagnosis for this woman?

- (A) Activity intolerance
- (B) Impaired skin integrity
- (C) Impaired verbal communication
- (D) Ineffective health maintenance

42. A 50-year-old male patient was transferred to surgical inpatient unit after thyroid surgery. The nurse starts to do post-operative assessment in order to identify early postoperative complication related to injury on the adjacent gland.

What is the most important action the nurse could do in this case?

- (A) Administering calcium gluconate
- (B) Monitoring symptoms of tetany
- (C) Monitoring hydration status
- (D) Monitoring input output

43. A 54-year-old female patient was presented to an outpatient department with complaints of shortness of breath, constipation, loss of appetite, weight gain and cold intolerance. The nurse suspected primary hypothyroidism and wants to verify with laboratory findings.

What is the most likely laboratory finding that helps to confirm the above diagnosis?

- (A) Normal thyroid hormone
- (B) Decreased thyroid hormone
- (C) Increased thyroid stimulating hormone
- (D) Decreased thyroid stimulating hormone

44. A 67-year-old known diabetic man presents with a complaint of infected foot ulcer with purulent discharge. He tried to treat it at home with over the-counter remedies. He was not aware of the long-term complications of diabetes mellitus and the prevention measures. The physician has made a diagnosis of gangrenous foot and recommended for an amputation.

What should be the priority patient expected outcome criteria for this patient?

- (A) Identifies signs and symptoms of hypoglycemia or hyperglycemia
- (B) Demonstrates proper technique for administering insulin injection
- (C) Demonstrates appropriate knowledge of diet selections
- (D) Maintains skin integrity and proper foot care

45. A 26-year-old male patient was presented to an ophthalmic clinic with complaint of burning pain on his eye for two days. On physical examination, the patient was unable to open the eye and has purulent discharge and lymphadenopathy.

What is the most likely type of conjunctivitis the patient developed?

- (A) Viral
- (B) Toxic
- (C) Allergic
- (D) Bacterial

46. A 24-year-old male patient came to a health center with complaints of itching, foreign body and burning sensation of both eyes. The patient reported exudates in the morning, purulent in nature that even makes difficult to open eyes. The patient is worried about his condition.

What is the priority nursing diagnosis for this patient?

- (A) Anxiety
- (B) Acute pain
- (C) Impaired mucosal lining
- (D) Disturbed sensory perception

47. A 47-year-old patient was presented to ophthalmic clinic with complaints of difficulty of reading and blurred vision. Ophthalmoscopic examination and tonometry revealed significant opacity and increased intra-ocular pressure.

What is the most appropriate management for this patient?

- (A) Surgery
- (B) Antibiotics
- (C) Eye irrigation
- (D) Corrective eyeglasses

What is the priority nursing intervention for a client with a sensory deficit?

(A) Pain management

(B) Disturbed sensory perception

60. A head nurse in a surgical unit is asked to use a piece of equipment immediately after use in order to abide with standard precautions. What type of standard precaution is applied in this context?

the head nurse will provide?

What is the most likely answer the head nurse will provide?

(A) Contact precaution
(B) Air born precaution
(C) Droplet precaution
(D) Isolation precaution

61. A male patient with closed arm fracture has stayed on traction for about two weeks. The healing is progressing well. The nurse prepares the patient for discharge from hospital with full arm cast. During discharge, the nurse wanted to give advice the patient about the problem he may face during his home stay.

52. A nurse working in a surgical inpatient unit has planned not to discuss assessment results and diagnostic test information of the patient with friends working in administration.

What is the most likely charting and documentation principle applied by the nurse?

- (A) Timing
- (B) Content
- (C) Accountability
- (D) Confidentiality

53. A 56-year-old patient has been newly enrolled to chronic illness clinic for follow-up. Immediately on arrival to the clinic, a nurse started to take vital signs where the environment is noisy due to re-innovation activity which makes the environment stressful.

What is the most likely abnormal finding expected of the patient?

- (A) Decreased pulse rate
- (B) Increased temperature
- (C) Increased blood pressure
- (D) Decreased respiratory rate

54. A 25-year-old patient was presented to an emergency medical unit with complaint of severe headache and blurring of vision. The nurse noticed blood pressure of 190/126mmHg and reported the result to the physician and they both agreed on the need for immediate action.

What is the most likely type of medication order in the above case?

- (A) Stat-Order
- (B) PRN-Order
- (C) Verbal Order
- (D) Standing protocol

55. A physician decided to discharge a 26-year-old mother who gave normal birth after six hours of follow-up. The head nurse requested the nurse in charge to facilitate the process as per the Ethiopian National Maternal Discharge Protocol.

What is the most likely initial nursing consideration to be taken?

- (A) Insuring clinician criteria for discharge
- (B) Communicating the planned discharge
- (C) Documenting Discharge checklist
- (D) Complete a discharge summary

56. A 34-year-old female patient was on blood transfusion as a management for severe anemia secondary to blood loss. The nurse started to monitor the patient during transfusion, observed signs of transfusion reaction.

What is the priority action the nurse should take in this case?

- (A) Stop transfusion
- (B) Check vital signs
- (C) Check the skin for rashes
- (D) Document transfusion reaction

48. A 60-year-old woman was admitted to an orthopedic ward due to fracture of tibia while walking. She has history of osteoporosis. Laboratory investigation of serum electrolyte showed calcium level of 5.6mg/dl (normal range 8.6-10.2 mg/dL) Phosphorus 5.78 mg/dl (normal range 2.5-4.5 mg/dl), sodium 130 mEq/l (normal range 135-145 mEq/L) Uric acid 9.65mg/dl(normal range 2.5-8 mg/dL)

What is the most likely explanation in this condition?

- (A) Increased serum uric acid (C) Decreased serum calcium
(B) Decreased serum sodium (D) Increased serum phosphorous

49. A 45-year-old client was presented to orthopedic unit after sustaining blunt trauma on the right upper limb. On physical examination, the patient has open fracture with large wound measuring 5cm and minor soft tissue damage.

What is the most likely grade of fracture in this patient?

- (A) Grade I (C) Grade III
(B) Grade II (D) Grade IV

50. A 66-year-old patient was admitted to inpatient unit with diagnosis of gout arthritis. The nurse initiated musculoskeletal examination to prepare patient teaching plan after admission. During patient teaching, the nurse advised the patient to increase water intake up to 3000ml per day.

What is the most likely anticipated complication by the nurse?

- (A) Urolithiasis (C) Dehydration
(B) Hypotension (D) Cholelithiasis

51. A 38-year-old woman was presented to a hospital with acute tenderness, swelling and paravertebral muscle spasm after being involved in deadly motor vehicle crash. She is conscious and has no open wound and bleeding. The pain is greater while moving and coughing.

What is the priority nursing intervention for this patient?

- (A) Immobilization and maintain spinal alignment
(B) Advise the patient to practice leg exercises
(C) Ambulate patient and maintain mobility
(D) Give pain medication as needed

BOOKLET CODE: 011

62. A 79-year-old man with some known cardiac and respiratory problems is admitted at a hospital. On assessment, he had crackles. Although he repeatedly attempts to cough and expectorate, he is too weak to remove the sputum. The nurse decides for postural drainage.

What typical condition can alert the nurse to stop doing the procedure for this client?

- (A) Greenish color for the sputum
(B) Increase distress

- (C) Increase sputum amount
(D) Increase temperature

63. An Emergency physician tried an endo-tracheal intubation for a 23-year-old female patient presented with apnea. After unsuccessful trial of intubation, he transferred her to emergency operating theatre for tracheotomy.

For what patient problem is this most likely indicative?

- (A) Upper airway obstruction
(B) Lower airway obstruction

- (C) Laryngeal cancer
(D) Vocal cord injury

64. A 35-year-old female patient was admitted to a gynecologic inpatient unit with the diagnosis of fistula. The gynecologist planned to do surgical intervention and requested the newly assigned nurses to prepare appropriate surgical equipments for this procedure.

What is the most likely category of equipment that should be prepared for this procedure?

- (A) Probing and dilating
(B) Grasping and clamping

- (C) Cutting and dissecting
(D) Exposing and retracting

65. A 55-year-old woman is scheduled to have a surgery on her lower abdomen for an inguinal hernia. The surgeon told that anesthetizing her lower body part is enough to perform the procedure safely. The Anesthetist decides to anesthetize the whole body part just below to the anesthesia injection site.

What kind of anesthesia is most appropriate in this case?

- (A) Spinal and epidural blocks
(B) General anesthesia

- (C) Local anesthesia
(D) Nerve blocks

66. A nurse performs the patient risk assessment and provides an intraoperative care for a patient undergoing laparotomy under general anesthesia.

What is the most likely complication that makes the patient most at risk?

- (A) Vomiting and nausea
(B) Peripheral edema

- (C) Atelectasis
(D) Anemia

BOOKLET CODE: 011

72. A 32-year-old man was presented with a complaint of watery diarrhea of two days duration. The diarrhea has a sudden onset of copious. He has severe muscle pain due to frequent diarrhea. The laboratory finding shows the presence of small, comma-shaped, motile, Gram-negative bacilli on stool.

What is the most effective prevention method for this case?

- (A) Sanitation
- (B) Antibiotics

- (C) Monitor output
- (D) Parenteral vaccine

73. A 45-year-old known HIV patient came to an outpatient department with complaint of nausea, vomiting and abdominal cramp. The nurse highly suspects Giardia lamblia infection.

What is the most likely additional assessment finding that supports the nurse's suspicion?

- (A) Foul-smelling greasy stools
- (B) Pale greasy bloody diarrhea

- (C) Pale greasy mucoid diarrhea
- (D) Prolonged watery diarrhea

74. A 16-year-old female patient visits her doctor with a complaint of cough and headache of three days duration. On history, she has sudden onset fever, runny nose, myalgia, chills and sore throat. The cough is severe and protracted.

What is the priority nursing diagnosis for this patient?

- (A) Acute pain
- (B) Hyperthermia

- (C) Impaired comfort
- (D) Impaired skin integrity

75. During a seminar presentation, a graduating student presents about the mode of transmission of leprosy. But students were confused about the main method of transmission and asked the presenter for clarification.

What is the best answer to be explained by the presenter?

- (A) Through sexual contact
- (B) Through respiratory secretion
- (C) Through broken skin and drainage
- (D) Through mother to child trans placental

BOOKLET CODE: 011

62. A 79-year-old man with some known cardiac and respiratory problems is admitted at a hospital. On assessment, he had crackles. Although he repeatedly attempts to cough and expectorate, he is too weak to remove the sputum. The nurse decides for postural drainage.

What typical condition can alert the nurse to stop doing the procedure for this client?

(A) Greenish color for the sputum
(B) Increase distress
(C) Increase sputum amount
(D) Increase temperature

63. An Emergency physician tried an endo-tracheal intubation for a 23-year-old female patient presented with apnea. After unsuccessful trial of intubation, he transferred her to emergency operating theatre for tracheotomy.

For what patient problem is this most likely indicative?

(A) Upper airway obstruction
(B) Lower airway obstruction
(C) Laryngeal cancer
(D) Vocal cord injury

64. A 35-year-old female patient was admitted to a gynecologic inpatient unit with the diagnosis of fistula. The gynecologist planned to do surgical intervention and requested the newly assigned nurses to prepare appropriate surgical equipments for this procedure.

What is the most likely category of equipment that should be prepared for this procedure?

(A) Probing and dilating
(B) Grasping and clamping
(C) Cutting and dissecting
(D) Exposing and retracting

65. A 55-year-old woman is scheduled to have a surgery on her lower abdomen for an inguinal hernia. The surgeon told that anesthetizing her lower body part is enough to perform the procedure safely. The Anesthetist decides to anesthetize the whole body part just below to the anesthesia injection site.

What kind of anesthesia is most appropriate in this case?

(A) Spinal and epidural blocks
(B) General anesthesia
(C) Local anesthesia
(D) Nerve blocks

66. A nurse performs the patient risk assessment and provides an intraoperative care for a patient undergoing laparotomy under general anesthesia.

What is the most likely complication that makes the patient most at risk?

(A) Vomiting and nausea
(B) Peripheral edema
(C) Atelectasis
(D) Anemia

48. A 60-year-old woman was admitted to an orthopedic ward due to fracture of tibia while walking. She has history of osteoporosis. Laboratory investigation of serum electrolyte showed calcium level of 5.6mg/dl (normal range 8.6-10.2 mg/dL) Phosphorus 5.78 mg/dl (normal range 2.5-4.5 mg/dl), sodium 130 mEq/l (normal range 135-145 mEq/L) Uric acid 9.65mg/dl(normal range 2.5-8 mg/dL)

What is the most likely explanation in this condition?

- (A) Increased serum uric acid (C) Decreased serum calcium
(B) Decreased serum sodium (D) Increased serum phosphorous

49. A 45-year-old client was presented to orthopedic unit after sustaining blunt trauma on the right upper limb. On physical examination, the patient has open fracture with large wound measuring 5cm and minor soft tissue damage.

What is the most likely grade of fracture in this patient?

- (A) Grade I (C) Grade III
(B) Grade II (D) Grade IV

50. A 66-year-old patient was admitted to inpatient unit with diagnosis of gout arthritis. The nurse initiated musculoskeletal examination to prepare patient teaching plan after admission. During patient teaching, the nurse advised the patient to increase water intake up to 3000ml per day.

What is the most likely anticipated complication by the nurse?

- (A) Urolithiasis (C) Dehydration
(B) Hypotension (D) Cholelithiasis

51. A 38-year-old woman was presented to a hospital with acute tenderness, swelling and paravertebral muscle spasm after being involved in deadly motor vehicle crash. She is conscious and has no open wound and bleeding. The pain is greater while moving and coughing.

What is the priority nursing intervention for this patient?

- (A) Immobilization and maintain spinal alignment
(B) Advise the patient to practice leg exercises
(C) Ambulate patient and maintain mobility
(D) Give pain medication as needed

52. A nurse working in a surgical inpatient unit has planned not to discuss assessment results and diagnostic test information of the patient with friends working in administration.

What is the most likely charting and documentation principle applied by the nurse?

- (A) Timing
- (B) Content
- (C) Accountability
- (D) Confidentiality

53. A 56-year-old patient has been newly enrolled to chronic illness clinic for follow-up. Immediately on arrival to the clinic, a nurse started to take vital signs where the environment is noisy due to re-innovation activity which makes the environment stressful.

What is the most likely abnormal finding expected of the patient?

- (A) Decreased pulse rate
- (B) Increased temperature
- (C) Increased blood pressure
- (D) Decreased respiratory rate

54. A 25-year-old patient was presented to an emergency medical unit with complaint of severe headache and blurring of vision. The nurse noticed blood pressure of 190/126mmHg and reported the result to the physician and they both agreed on the need for immediate action.

What is the most likely type of medication order in the above case?

- (A) Stat-Order
- (B) PRN-Order
- (C) Verbal Order
- (D) Standing protocol

55. A physician decided to discharge a 26-year-old mother who gave normal birth after six hours of follow-up. The head nurse requested the nurse in charge to facilitate the process as per the Ethiopian National Maternal Discharge Protocol.

What is the most likely initial nursing consideration to be taken?

- (A) Insuring clinician criteria for discharge
- (B) Communicating the planned discharge
- (C) Documenting Discharge checklist
- (D) Complete a discharge summary

56. A 34-year-old female patient was on blood transfusion as a management for severe anemia secondary to blood loss. The nurse started to monitor the patient during transfusion, observed signs of transfusion reaction.

What is the priority action the nurse should take in this case?

- (A) Stop transfusion
- (B) Check vital signs
- (C) Check the skin for rashes
- (D) Document transfusion reaction

SUBJECT CODE: 01

A 25-year-old patient was presented to a health center with complaint of constipation and laxative.

Assessment result revealed the management options after getting

difficult to pass stool. The nurse decided to change the management options after getting

but failed to pass stool. The nurse decided to change the management options after getting

evaluation of stool impaction

what is the most appropriate type of fluid to be recommended for this case?

(A) Normal saline

(B) Ringers' lactate Solution

(C) Normal saline

(D) Ringers' lactate Solution

58. A 25-year-old patient was admitted to surgical ward after major surgical procedure. The

nurse working in the unit planned to teach nursing students about the main events happening

in the wound healing process during defense phase and main personal precautions.

What is the most likely event that will happen at this stage?

(A) Tissue remodeling

(B) Collagen deposition

(C) Tissue remodeling

(D) Collagen deposition

59. A 39-year-old female patient came to ophthalmic clinic with complaint of vision problems

for the last two weeks. Examination result revealed that the patient used to avoid light; she

also stated using dark glass, rubbing her eyes repeatedly and squint eyes.

What is the priority nursing diagnosis for this patient?

(A) Disturbed body image

(B) Disturbed sensory perception

(C) Disturbed body image

(D) Situational low self-esteem

60. A head nurse in a surgical ward always warns nurses to decontaminate used medi-

equipment immediately after use in order to abide with standard precaution. A nurs-

student asked a head nurse "what type of standard precaution is applied in this context?"

What is the most likely answer the head nurse will provide?

(A) Droplet precaution

(B) Contact precaution

(C) Droplet precaution

(D) Isolation precaution

61. A male patient with closed arm fracture has stayed on traction for about two weeks. The

healing is progressing well. The nurse prepares the patient for discharge from hospital v-

full arm cast. During discharge, the nurse wanted to give advice the patient about the

problem he may face during his home stay.

What is the most likely problem that the nurse should focus while giving patient educa-

(A) Compartment syndrome

(B) Muscle atrophy

(C) Fat embolism

(D) Infection

44. A 67-year-old known diabetic man presents with a complaint of infected foot ulcer with purulent discharge. He tried to treat it at home with over the-counter remedies. He was not aware of the long-term complications of diabetes mellitus and the prevention measures. The physician has made a diagnosis of gangrenous foot and recommended for an amputation.

What should be the priority patient expected outcome criteria for this patient?

- (A) Identifies signs and symptoms of hypoglycemia or hyperglycemia
- (B) Demonstrates proper technique for administering insulin injection
- (C) Demonstrates appropriate knowledge of diet selections
- (D) Maintains skin integrity and proper foot care

45. A 26-year-old male patient was presented to an ophthalmic clinic with complaint of burning pain on his eye for two days. On physical examination, the patient was unable to open the eye and has purulent discharge and lymphadenopathy.

What is the most likely type of conjunctivitis the patient developed?

- (A) Viral
- (B) Toxic
- (C) Allergic
- (D) Bacterial

46. A 24-year-old male patient came to a health center with complaints of itching, foreign body and burning sensation of both eyes. The patient reported exudates in the morning, purulent in nature that even makes difficult to open eyes. The patient is worried about his condition.

What is the priority nursing diagnosis for this patient?

- (A) Anxiety
- (B) Acute pain
- (C) Impaired mucosal lining
- (D) Disturbed sensory perception

47. A 47-year-old patient was presented to ophthalmic clinic with complaints of difficulty of reading and blurred vision. Ophthalmoscopic examination and tonometry revealed significant opacity and increased intra-ocular pressure.

What is the most appropriate management for this patient?

- (A) Surgery
- (B) Antibiotics
- (C) Eye irrigation
- (D) Corrective eyeglasses

39. A 72-year-old patient came to a medical unit with complaints of sudden onset of visual hemiparesis, confusion, trouble speaking, and impaired mobility. Assessment findings showed history of poorly controlled hypertension. The physician in charge of the patient stroke and put the patient on oxygen.

What is the initial treatment option for this patient?

- (A) Anti-platelet
- (B) Thrombolytic
- (C) Anti-hypertensive
- (D) Analgesics treatment

40. A 46-year-old female patient was admitted to surgical inpatient unit with diagnosis of subdural hematoma. Assessment findings showed decreased level of blood pressure and oxygen saturation. The nurse administered nasal oxygen and planned to evaluate the prognosis.

What is the most likely evidence for good prognosis of the given intervention?

- (A) SpO₂ of at least 90%
- (B) Respiratory rate of 28/min
- (C) PaCO₂ at least 52mmHg
- (D) PaO₂ is at least 70mmHg

41. A 72-year-old female patient was presented with complaints of numbness of the extremities, blurred vision and increased appetite. The patient reported that she has history of chronic alcohol consumption and visiting traditional healers for controlling blurred vision and other related health problems. Her random blood glucose level is 320mg/dL.

What is the priority nursing diagnosis for this woman?

- (A) Activity intolerance
- (B) Impaired skin integrity
- (C) Impaired verbal communication
- (D) Ineffective health maintenance

42. A 50-year-old male patient was transferred to surgical inpatient unit after thyroid surgery. The nurse starts to do post-operative assessment in order to identify early postoperative complication related to injury on the adjacent gland.

What is the most important action the nurse could do in this case?

- (A) Administering calcium gluconate
- (B) Monitoring symptoms of tetany
- (C) Monitoring hydration status
- (D) Monitoring input output

43. A 54-year-old female patient was presented to an outpatient department with complaints of shortness of breath, constipation, loss of appetite, weight gain and cold intolerance. The nurse suspected primary hypothyroidism and wants to verify with laboratory findings.

What is the most likely laboratory finding that helps to confirm the above diagnosis?

- (A) Normal thyroid hormone
- (B) Decreased thyroid hormone
- (C) Increased thyroid stimulating hormone
- (D) Decreased thyroid stimulating hormone

39. A 72-year-old patient came to a medical unit with complaints of sudden onset of visual hemiparesis, confusion, trouble speaking, and impaired mobility. Assessment findings showed history of poorly controlled hypertension. The physician in charge of the patient stroke and put the patient on oxygen.

What is the initial treatment option for this patient?

- (A) Anti-platelet
- (B) Thrombolytic
- (C) Anti-hypertensive
- (D) Analgesics treatment

40. A 46-year-old female patient was admitted to surgical inpatient unit with diagnosis of subdural hematoma. Assessment findings showed decreased level of blood pressure and oxygen saturation. The nurse administered nasal oxygen and planned to evaluate the prognosis.

What is the most likely evidence for good prognosis of the given intervention?

- (A) SpO_2 of at least 90%
- (B) Respiratory rate of 28/min
- (C) PaCO_2 at least 52mmHg
- (D) PaO_2 is at least 70mmHg

41. A 72-year-old female patient was presented with complaints of numbness of the extremities, blurred vision and increased appetite. The patient reported that she has history of chronic alcohol consumption and visiting traditional healers for controlling blurred vision and other related health problems. Her random blood glucose level is 320mg/dL.

What is the priority nursing diagnosis for this woman?

- (A) Activity intolerance
- (B) Impaired skin integrity
- (C) Impaired verbal communication
- (D) Ineffective health maintenance

42. A 50-year-old male patient was transferred to surgical inpatient unit after thyroid surgery. The nurse starts to do post-operative assessment in order to identify early postoperative complication related to injury on the adjacent gland.

What is the most important action the nurse could do in this case?

- (A) Administering calcium gluconate
- (B) Monitoring symptoms of tetany
- (C) Monitoring hydration status
- (D) Monitoring input output

43. A 54-year-old female patient was presented to an outpatient department with complaints of shortness of breath, constipation, loss of appetite, weight gain and cold intolerance. The nurse suspected primary hypothyroidism and wants to verify with laboratory findings.

What is the most likely laboratory finding that helps to confirm the above diagnosis?

- (A) Normal thyroid hormone
- (B) Decreased thyroid hormone
- (C) Increased thyroid stimulating hormone
- (D) Decreased thyroid stimulating hormone

34. A nurse on duty is planning nursing intervention for a patient admitted to medical ward with acute renal failure. The laboratory finding showed decreased serum phosphorus and calcium level.

What is the most important nursing intervention for this patient?

- (A) Administer blood components therapy
- (B) Assisting the patient with an exercise program
- (C) Teaching the patient to report sign of fluid overload
- (D) Encourage compliance with fluid restriction therapy

35. A nurse is assessing a male patient who is admitted in medical ward with acute renal failure. On physical examination, he has dry mucous membrane, poor skin turgor, flat neck vein and orthostatic hypotension. He is sick looking and irritable.

What is the primary nursing diagnosis for this patient?

- (A) Risk for altered skin integrity
- (B) Fluid volume deficit
- (C) Deficient knowledge
- (D) Activity intolerance

36. A 33-year-old patient came to a hospital with complaints of peripheral edema and cola-colored urine. The patient reported a history of frequent attack of tonsillitis and skin infections. On laboratory investigation, the patient's Glomerular Filtration Rate reads 100ml/minute/1.73m² (normal GFR is 125ml/min/1.73 m²).

What is the most likely fluid and electrolyte balance change expected from the patient?

- (A) Hypokalemia
- (B) Hypernatremia
- (C) Marked hypertension
- (D) Decreased urine output

37. A 63-year-old patient was brought to medical unit with complaints of sudden onset of visual disturbance, paralysis on one side of the body and severe headache. Assessment findings showed history of poorly controlled hypertension.

What is the most likely diagnosis of this patient?

- (A) Embolic stroke
- (B) Ischemic Stroke
- (C) Thrombotic stroke
- (D) Hemorrhagic Stroke

38. A 36-year-old patient was presented with complaints of severe headache, neck stiffness and fever. The nurse and the physician planned to do lumbar puncture to send sample for laboratory investigation.

What is the most likely contraindication for this procedure?

- (A) Decreased level of consciousness
- (B) Decreased intracranial pressure
- (C) Infection on proposed site
- (D) Increased blood pressure

9. A child is brought to an emergency department with dislodged fragment of toy part in the airway. He has violent paroxysms of coughing and choking accompanied by new onset wheezing, intercostal muscle retraction and cyanosis. His respiratory rate is 62/min and pulse rate is 100/min.

What is the most likely actual nursing diagnosis that needs priority for this child?

- (A) Impaired gas exchange
(B) Ineffective airway clearance
(C) Ineffective breathing pattern
(D) Impaired spontaneous ventilation
10. An unconscious seven-year-old child has sustained multiple body injury secondary to vehicle accident. The child was brought to hospital by bystanders and a nurse in charge suspected neck and back injury.

What is the initial step that the nurse should do for this child?

- (A) Perform jaw thrust
(B) Clear secretions from throat
(C) Inspect mouth and remove foreign body
(D) Check the airway by looking for chest movements
11. A preschool-age child with previous history of childhood asthma was brought to pediatric emergency unit. The child has a manifestation of peak expiratory flow 30%, oxygen saturation 78% and loud wheezing throughout inhalation and exhalation.

What is the actual nursing diagnosis that needs priority for this child?

- (A) Impaired gas exchange
(B) Ineffective airway clearance
(C) Ineffective breathing pattern
(D) Impaired spontaneous ventilation
12. A nine-year-old child was brought to a pediatric OPD with complaints of diarrhea, non-bilious and projectile vomiting. On examination, there was low grade fever.

What is the most likely diagnosis of this child?

- (A) Intussusceptions
(B) Pyloric stenosis
(C) Intestinal volvulus
(D) Hirschsprung disease
13. A nurse planned to give health education to a mother of a seven year-old-child who came with a disease called juvenile idiopathic arthritis.

What is the most appropriate advice the nurse could give to the mother?

- (A) Restrict from fatty foods during severe form of illness
(B) Restrict from exercise during moderate form of illness
(C) Restrict from protein foods during severe form of illness
(D) Avoid systemic steroids during moderate form of illness

25. A 23-year-old chronic liver disease patient was admitted to a medical ward with a diagnosis of hepatocellular carcinoma. The community where the patient came uses running water for drinking. He has no history of IV drug use and previous blood transfusion.

What is the most likely etiology for the patient's diagnosis?

- (A) Hepatitis A Virus
- (B) Hepatitis B Virus
- (C) Hepatitis C Virus
- (D) Hepatitis D Virus

26. A 36-year-old male patient is admitted in surgical ward with confirmed diagnosis of appendicitis. The patient is prepared for appendectomy and received preoperative care.

What is the most appropriate preoperative care for this patient?

- (A) Administer enema
- (B) Encourage ambulation
- (C) Administer antibiotic
- (D) Stimulates bowel activity

27. A 26-year-old male patient, who works in a construction firm, was brought to an emergency department. The patient was diagnosed with large bowel obstruction empirically.

What is the most likely clinical presentation that helps to diagnose this patient?

- (A) Vomiting
- (B) Dehydration
- (C) Gross abdominal distention
- (D) Bowel sound of 4/5 minute

28. A 64-year-old female patient is admitted to a medical ward with a complaint of unconsciousness for four days duration. The unconsciousness is due to head trauma that leads excessive blood loss. Naso-gastric intubation is option less to feed her. The available tube has no mark that the nurse should determine the length of NGT needed to reach the stomach.

What is the most likely technique to be used in this case?

- (A) From tip of the nose to the earlobe and from the earlobe to the umbilicus
- (B) From the earlobe to tip of the nose and from tip of nose to the xiphoid process
- (C) From tip of the nose to the earlobe and from the earlobe to the xiphoid process
- (D) From tip of the nose to the earlobe to half way between xiphoid process and umbilicus

29. A 62-year-old obese man came to a medical outpatient department seeking treatment. The nurse on duty diagnosed the patient with duodenal peptic ulcer disease based on the clinical manifestations and history of the patient.

What is the most likely clinical manifestation used by the nurse to diagnosis this patient?

- (A) Ingestion of food relieve pain
- (B) Pain occur one hour after meal
- (C) Being obese
- (D) Relieved by vomiting

25. A 23-year-old chronic liver disease patient was admitted to a medical ward with a diagnosis of hepatocellular carcinoma. The community where the patient came uses running water for drinking. He has no history of IV drug use and previous blood transfusion.

What is the most likely etiology for the patient's diagnosis?

- (A) Hepatitis A Virus
- (B) Hepatitis B Virus
- (C) Hepatitis C Virus
- (D) Hepatitis D Virus

26. A 36-year-old male patient is admitted in surgical ward with confirmed diagnosis of appendicitis. The patient is prepared for appendectomy and received preoperative care.

What is the most appropriate preoperative care for this patient?

- (A) Administer enema
- (B) Encourage ambulation
- (C) Administer antibiotic
- (D) Stimulates bowel activity

27. A 26-year-old male patient, who works in a construction firm, was brought to an emergency department. The patient was diagnosed with large bowel obstruction empirically.

What is the most likely clinical presentation that helps to diagnose this patient?

- (A) Vomiting
- (B) Dehydration
- (C) Gross abdominal distention
- (D) Bowel sound of 4/5 minute

28. A 64-year-old female patient is admitted to a medical ward with a complaint of unconsciousness for four days duration. The unconsciousness is due to head trauma that leads excessive blood loss. Naso-gastric intubation is option less to feed her. The available tube has no mark that the nurse should determine the length of NGT needed to reach the stomach.

What is the most likely technique to be used in this case?

- (A) From tip of the nose to the earlobe and from the earlobe to the umbilicus
- (B) From the earlobe to tip of the nose and from tip of nose to the xiphoid process
- (C) From tip of the nose to the earlobe and from the earlobe to the xiphoid process
- (D) From tip of the nose to the earlobe to half way between xiphoid process and umbilicus

29. A 62-year-old obese man came to a medical outpatient department seeking treatment. The nurse on duty diagnosed the patient with duodenal peptic ulcer disease based on the clinical manifestations and history of the patient.

What is the most likely clinical manifestation used by the nurse to diagnosis this patient?

- (A) Ingestion of food relieve pain
- (B) Pain occur one hour after meal
- (C) Being obese
- (D) Relieved by vomiting

30. A 45-year-old male patient came to an emergency department with complaints of abdominal pain, vomiting and constipation of 12 hours duration. The patient reports that he has previous history of hernia. On physical examination, the nurse identified distended abdomen, high pitched bowel sounds, but no detected masses.

What is the next course of the disorder if left untreated?

- (A) Passed flatus; small and formed stool
- (B) Intoxicated patient with disarranged vital sign
- (C) More intense pain and slowly dehydrated patient
- (D) Gradual decrease in vomiting and abdominal distention

31. During assessment of a patient with bilateral lower extremity edema, a nurse applies pressure on edematous area for 5 seconds using thumb and identified 3cm indentation.

What is the most likely edema grade of this patient?

- (A) One pitting edema
- (B) Two pitting edema
- (C) Three pitting edema
- (D) Four pitting edema

32. A 60-year-old female patient was admitted to a medical ward with a diagnosis of chronic renal failure. Laboratory result shows serum potassium level of 7mEq/L (normal value 3.5-5.0). On assessment she has ECG changes (tall-tented T waves and widened QRS). The health care provider diagnosed hyperkalemia.

What early signs and symptoms could support the diagnosis?

- (A) Muscle weakness and diarrhea
- (B) Blood pressure and fluid retention
- (C) Fever, chest pain and additive breath sound
- (D) RBC count, hemoglobin and hematocrit level

33. A 76-year-old woman was admitted to a medical ward with complaints of edema, rash, and shortness of breath. She urinate 120 ml of urine per 24 hours. The nurse diagnosed the patient with chronic renal failure empirically.

What is the possible additional manifestation which supported the nurse's decision?

- (A) Decrease BUN
- (B) Acute weight loss
- (C) Clear breath sound
- (D) Decrease hematocrit

**Ethiopian Health Professionals' Licensure Examination
Nursing Exam
Tir 2015/January 2023
PART TWO**

BOOKLET CODE: 012
Number of Items: 100

SUBJECT CODE: 02
Time Allowed: 2:30 hours

DIRECTIONS: Each of the following questions is followed by four possible alternative answers. Read each question carefully and **blacken** the letter of your best choice on the separate answer sheet provided.

1. A 25-year-old lactating mother came to MCH unit for family planning services. On assessment, the mother has given birth before six weeks. She is on exclusive breast feeding, her menstruation is not returned, her BP: 120/78mmHg, pulse rate 85/min, weight: 60kg. She has no headache, vaginal discharge and fever. The nurse working in the unit counseled her about all contraception options and their effect.

What is the most appropriate family planning method for this mother?

- (A) Calendar method (C) Lactational amenorrhea method
(B) Progesterone only pills (D) Combined oral contraceptive pills

2. An HIV positive woman came to a family planning clinic to take contraceptive method. On assessment, the woman was on treatment for gonorrhea and she was on HAART for the last three years.

What is the most appropriate contraceptive method for this woman?

- (A) IUCD with condom (C) Diaphragm with condom
(B) Implant with condom (D) Spermicide with condom

3. A 25-year-old woman came to a health center for family planning service. A care provider working in the unit counseled her about all methods and screened her for eligibility. Finally, she preferred to take injectable depot medroxy-progesterone acetate (DMPA). Then, the care provider provided the methods as per the standard and appointed her for the next injection.

What is the most likely return visit for next injection?

- (A) Every month (C) Every nine month
(B) Every six month (D) Every three month

4. A 25-year-old woman came to a family planning clinic with complaints of irregular frequent menstrual bleeding after she took combined oral contraceptives. She has heavy bleeding that lasts more than seven days.

What is the most likely menstrual pattern for this woman?

- (A) Metorrhagia
- (B) Menorrhagia
- (C) Polymenorrhea
- (D) Menometrorrhagia

5. A married couple came to a hospital for infertility investigations. They were trying to achieve conception for the last two years. They do not have previous history of contraception usage. Accordingly the gynecologist ordered laboratory investigations for the couple.

What is the most likely first laboratory investigation to be done for this couple?

- (A) Urine HCG test
- (B) Semen analysis
- (C) Hormonal assay
- (D) Serum FSH level

6. A 30-year-old mother was admitted to gynecology ward with vesico-vaginal fistula secondary to obstructed labor. The mother gave birth at home before 2 months. She is scheduled for fistula repair after a week.

What is the most appropriate nursing intervention for this mother?

- (A) Emotional support
- (B) Insert catheter in the bladder
- (C) Maintain patient on clear liquids
- (D) Encourage frequent sitz baths

7. An 18-year-old lady came to a gynecology OPD with a complaint of abdominal distension for one month. She has abdominal swelling, and dull like abdominal pain. On examination, liver is enlarged, firm and nodular. The imaging result showed that the tumor is unilateral, solid, more than 10 cm in diameter which has originated from the primitive yolk sack.

What is the most likely diagnosis of this woman?

- (A) Dysgerminoma tumor
- (B) Choriocarcinoma tumor
- (C) Immature teratoma tumor
- (D) Endodermal sinus tumor

8. A 26-year-old non-pregnant woman came to a hospital with complaints of inter-menstrual and post-coital vaginal bleeding for three months. CT scan investigation result showed a cervical tumor which involves lower third of the vagina, without extension to the pelvic wall. She was diagnosed for stage II.

What is the most likely manifestation that corresponds to this stage?

- (A) Dry and itchy vaginal orifice
- (B) Adnexal excitation tenderness
- (C) Offensive and watery vaginal discharge
- (D) Excessive fluid accumulation in culdosa-sac

98. A postpartum mother was admitted to a hospital for complaints of pain and swelling on vulva for one day. On evaluation, there is hematoma formation on the vulva, the episiotomy wound appears to be moist, swollen and has pus containing discharge.

What is the most likely nursing diagnosis of this mother?

- (A) Self-care deficit
- (B) Impaired sexuality
- (C) Delayed wound healing
- (D) Impaired tissue integrity

99. A nurse was providing a care for a 2 days postnatal mother. The nursing care plan for the mother focused on an intervention that helps the nurse to promote health and prevent disease for both the mother and the new born.

What is the most appropriate intervention that should be included?

- (A) Episiotomy care for the mother
- (B) Treatment of postnatal infection
- (C) Inspecting the vagina for bleeding
- (D) Providing iron folic acid supplementation

100. A pregnant mother gave birth to a 3.8kg male neonate with vacuum. After one hour of delivery, the uterus was well contracted and there was a bright red vaginal bleeding. On examination, there is laceration of the vagina and cervix. The nurse shouted for help and started an IV fluid transfusion.

What is the most appropriate next management for this woman?

- (A) Repairing the tears
- (B) Give Oxytocin 10 units IM
- (C) Removal of the placenta manually
- (D) Bimanual compression of the uterus

THE END

BOOKLET CODE: 011

94. A 34-year-old multi-gravida mother is admitted in labor ward with a complaint of pushing down pain for eight hours. On evaluation, she had adequate pelvis with vertex presentation, the cervix was 6 cm dilated with 3 mild uterine contractions. After four hours, she has unchanged cervical dilation with two mild uterine contractions.

What is the most likely management for this mother?

- (A) Induction of labor
- (B) Start using partograph
- (C) Prepare the mother for CS
- (D) Start augmentation protocol

95. A 19-year-old woman came to a hospital with vaginal bleeding and lower abdominal pain for six hours. She has history of fall before 12 hours and she was amenorrhic for the past two months. Her blood pressure is 70/40mmHg and pulse rate 100/min. She is in emergency unit and her priority nursing diagnosis is fluid volume deficit.

What is the most appropriate outcome criterion for the identified nursing diagnosis?

- (A) Blood loss will decrease by 500ml/day
- (B) urine output will increase by 30ml/day
- (C) Patient will walk safely for short distance
- (D) Fluid volume will be maintained at fractional level

96. A mother visited a postnatal clinic on the fifth day of postpartum with a complaint of breast pain for one day. On evaluation, the left breast is tender, dry, swollen and red with wedge-shaped area. Her pulse rate is 88/min, respiratory rate is 22/min and temperature is 38°C.

What is the most likely diagnosis of this mother?

- (A) Superficial cellulites
- (B) Breast engorgement
- (C) Acute mastitis
- (D) Breast abscess

97. A 20-year-old primiparous mother gave birth to a 2.8kg female neonate with spontaneous vaginal delivery. The health care provider advises the mother about good attachment and positioning.

What is the most appropriate positioning the nurse should teach her?

- (A) Infant should face away from the mother
- (B) Infant head should be turn toward the mother
- (C) Infants body should touch mother's abdomen
- (D) Infant should be perpendicular with the mother lap

90. A 22-year-old primigravida mother came to a hospital with a complaint of labor pain for five hours. On evaluation, she had three regular uterine contractions and cervix is 8cm dilated. After an hour, the cervix became fully dilated, fetal heart beat is 90/min and there is three regular uterine contractions. The nurse decided to deliver the fetus using vacuum.

What is the most likely indication for the above procedure?

- (A) Tight perineum
- (B) Prolonged second stage of labor
- (C) Poor maternal expulsive effort
- (D) Non reassuring fetal heart rate

91. A nurse administered three doses of 2 IU of oxytocin in 1000 ml of normal saline solution for a pregnant mother who was indicated for labor induction. The dose was started with 20 drop per minute and increased every 30 minutes till adequate uterine contraction is achieved. After few hours, the uterus became tense, strong and highly contracted.

What is the most likely complication the mother could develop?

- (A) Uterine inertia
- (B) Uterine rupture
- (C) Retained placenta
- (D) Chorioamnionitis

92. A 31-year-old primigravida mother visited a health center for delivery. On examination, the contraction was adequate with vertex presentation, the head was visible on the vulva and there was a bulged thinned perineum. The nurse conducted delivery of the baby by performing episiotomy.

What is the most likely indication used to decide this procedure?

- (A) Vertex presentation
- (B) Uterine contraction
- (C) Thinned perineum
- (D) Head visible in vulva

93. A 20-year-old primigravida mother visited a health center for delivery. On arrival, she is on active first stage of labor with 8 cm cervical dilatation and three strong contractions in 10 minute. She shouts when she had pain secondary to uterine contraction.

What is the most appropriate pain management for this mother?

- (A) Provide tocolytics to rest labor
- (B) Teach labor breathing technique
- (C) Encourage movement with attendant
- (D) Establish a relationship with the mother

BOOKLET CODE: 011

86. A pregnant woman came to ANC clinic with complaints of headache and difficulty of breathing for one month. On examination, she has pale conjunctiva, pulse 120/min, and respiratory rate 30/min. Her hemoglobin level is 10g/dl. The health care provider gave treatment for anemia.

- What is most appropriate outcome criterion used to measure the woman's improvement?
- (A) Hemoglobin level of 13g/dl
 - (B) Pulse rate of 100/min
 - (C) Absence of headache
 - (D) Respiratory rate of 24/min

87. A 24-year-old pregnant woman came to ANC clinic for antenatal visit at her 36-weeks gestation. Her MUAC measurement is 16.5cm. The ultrasound finding highly suggests an estimated fetal weight less than the tenth percentile and inadequate fetal head and body growth.

- What is the most appropriate management for this woman?
- (A) Provide bed rest
 - (B) Terminate the pregnancy
 - (C) Decrease fluid intake
 - (D) Improve nutrition intake

88. A 23-year-old 34 weeks pregnant mother came to a hospital for her third ANC visit. On assessment, her blood pressure is 150/90mmHg measured two times six hours apart and her pulse rate 75/min. abdominal assessment shows cephalic presentation and positive fetal heart beat. Laboratory investigation revealed 2+ proteins in urine, negative for glucose and ketone in the urine. She was diagnosed with mild preeclampsia.

- What is the most appropriate management for this mother?
- (A) Admit the mother and administer diuretics
 - (B) Follow the mother as an outpatient twice a week
 - (C) Admit the mother and initiate magnesium sulfate
 - (D) Give anticonvulsants and antihypertensive before discharge

89. A 32 weeks gestational age pregnant woman is admitted in obstetric ward for pregnancy-induced hypertension. During admission, the woman had blurred vision, blood pressure 160/110mmHg, protein urea 0.3gm and oliguria. The woman was conscious and she had no any seizure. She also took the recommended intervention to manage her problem.

- What is the most appropriate outcome criterion to evaluate the woman's prognosis?
- (A) Proteinuria 1+
 - (B) Blood pressure 140/90mmHg
 - (C) Absence of blurred vision
 - (D) Urine output 600ml/24hrs

BOOKLET CODE: 012

40. A nurse who is providing vaccines for children, wanted to follow the safety precautions of vaccinations to avoid the possible transmission of infections due to unsafe injections.

What is the most important precaution that the nurse implement primarily to meet her goal?
(A) Avoid reuse of needles
(B) Avoid recapping of needles
(C) Avoid touching the needle
(D) Avoid injecting at the incorrect site

41. A nurse found that the cold chain temperature of the vaccine has fallen below 0°C for five hours. The nurse moves vaccines to the safe storage and checked some freeze sensitive vaccines using a shake test. The shake test was performed using the "TEST" vial and "FROZEN" vial.

What is the most likely condition to determine the vaccine is safe to use?

(A) If sedimentation is similar in both vials
(B) If the TEST vial sediments are faster than the FROZEN vial
(C) If the TEST vial sediments are slower than the FROZEN vial
(D) If there is no sedimentation in the TEST vial, but in the FROZEN vial

42. A nine-month-old child is brought to the immunization clinic for vaccination. The nurse checked that the child has taken all previous doses before giving the present dose. Then, the nurse gives measles vaccine and appointed the mother for the next dose.

What is the most likely vaccine to be provided in the next schedule?

(A) Measles
(B) Pertussis
(C) Tetanus toxoid
(D) H.inflenza type b

43. A mother of a school age child came to a hospital to consult the nurse about the development of her child. The nurse asked the mother about the behavioral indicators of development and suspected that the child may have attention-deficit disorder.

What could be the most likely indicator for the nurse's suspicion?

(A) Impulsivity in school
(B) Over engagement in tasks
(C) Sitting in class for long time
(D) Impairments in communication

44. A three-year-old child is brought to the well-baby clinic due to poor communication skill. On assessment, the child is found to have poor social interaction and the nurse suspected that the child is autistic.

What is the most appropriate management for this child?

(A) Psychosocial intervention
(B) Advise the mother for stimulants
(C) Advise the mother for antipsychotics
(D) Advise the mother for mood stabilizers

76. A 43-year-old male patient comes to a hospital with complaint of penetrated skin of two days duration. He also complains hematuria and dysuria. On laboratory, there are cercariae ova in urine. Most of the time the patient has swimming habit from their local river.

What is the most appropriate management for this case?

- (A) Administration of Diclofenac
- (B) Administration of Praziquantel
- (C) Avoid swimming in water bodies
- (D) Clearing of vegetation in water bodies

77. A 34-year-old man came to a hospital from malaria endemic area with the complaints of fatigue, headache and fever. The nurse performed further assessment and diagnosed the patient with malaria.

What clinical finding is most helpful to diagnose the patient is in complication?

- (A) Fever
- (B) Myalgia
- (C) Jaundice
- (D) Headache

78. A 36-year-old female patient comes to a health institution with complaint of skin rash on her genital area for two months. She has hard symmetrical chancre but do not itch and the serological test (VDRL) is positive. The chancre was painless and disappeared within one month. Her husband tells her that he had similar case for the last six months.

What is the best clinical feature to diagnose this case?

- (A) Non itchy chancre
- (B) Chancre on genital area
- (C) Positive serological test
- (D) Hard symmetrical chancre

79. A 27-year-old patient comes to a hospital with the complaints of acute onset profuse and purulent urethral discharge, urgency, itching sensation of the urethra and burning sensation during micturition. The patient have also painless and regular ulcer on genitalia.

What is the most likely diagnosis of this patient?

- (A) Syphilis
- (B) Chancroid
- (C) Gonorrhea
- (D) Herpes genitalia

80. An old man came to a hospital and was diagnosed with rabies after dog bite. The patient has open and deep wound, paraesthesia, and tingling sensations at the site of the bite.

What is the initial management for this patient?

- (A) Suture the wound
- (B) Administer antibiotics
- (C) Administer post exposure prophylaxis
- (D) Wash the wound with soap and water

81. A group of people came to a hospital with increasing fever, dyspnea, stridor, hypoxia, and hypotension after eating raw meat prepared for the cultural celebration. Blood culture is positive for *Bacillus anthracis* bacteria.

What is the most appropriate drug management for these patients?

- (A) Chloramphenicol
- (B) Ceftriaxone

- (C) Gentamicin
- (D) Penicillin

82. A 30-year-old pregnant woman came to a health center to attend her first ANC visit. On examination, the nurse identified that woman's amniotic fluid index to be 24, fetal weight is 10% greater than the gestational age, fetal heart rate is 110/min and maternal blood pressure is 90/60mmHg. She was diagnosed for amniotic fluid disorder.

What is the best characteristic that led to this diagnosis?

- (A) Fetal heart beat 110/min
- (B) Blood pressure 90/60mmHg

- (C) Increased Fetal weight
- (D) Amniotic fluid index 24

83. A 37-week pregnant mother came to a health center for her fourth ANC visit. She complains difficulty of breathing, lethargy and fatigue. On physical examination, she has pale conjunctiva and her vital signs are in a normal range. The nurse suspected anemia.

What is the most appropriate lab investigation that would confirm diagnosis?

- (A) Urine analysis
- (B) Hematocrit

- (C) Blood sugar
- (D) VDRL test

84. A pregnant mother came to a health center for second ANC visit. Her blood pressure is 120/70mmHg, pulse rate 80/min, her weight is 56kg. On abdominal examination, the nurse assessed her fundal height by using finger method and fundal height above the umbilicus allows four fingers.

What is most likely gestational age based on the fundal height?

- (A) 32 weeks
- (B) 28 weeks

- (C) 24 weeks
- (D) 20 weeks

85. A 32-week pregnant mother came to a hospital with complaints of fever and chills for two days. She has loss of appetite and joint pain. On physical assessment, her blood pressure is 100/70 mmHg, pulse rate 84/min and her body temperature is 38.5°C. Her blood film result shows positive for *falciparum* plasmodium.

What is the most likely maternal complication on this woman?

- (A) IUGR
- (B) Anemia

- (C) Post-term labor
- (D) Spontaneous abortion

76. A 43-year-old male patient comes to a hospital with complaint of penetrated skin of two days duration. He also complains hematuria and dysuria. On laboratory, there are cercariae ova in urine. Most of the time the patient has swimming habit from their local river.

What is the most appropriate management for this case?

- (A) Administration of Diclofenac
- (B) Administration of Praziquantel
- (C) Avoid swimming in water bodies
- (D) Clearing of vegetation in water bodies

77. A 34-year-old man came to a hospital from malaria endemic area with the complaints of fatigue, headache and fever. The nurse performed further assessment and diagnosed the patient with malaria.

What clinical finding is most helpful to diagnose the patient is in complication?

- (A) Fever
- (B) Myalgia
- (C) Jaundice
- (D) Headache

78. A 36-year-old female patient comes to a health institution with complaint of skin rash on her genital area for two months. She has hard symmetrical chancre but do not itch and the serological test (VDRL) is positive. The chancre was painless and disappeared within one month. Her husband tells her that he had similar case for the last six months.

What is the best clinical feature to diagnose this case?

- (A) Non itchy chancre
- (B) Chancre on genital area
- (C) Positive serological test
- (D) Hard symmetrical chancre

79. A 27-year-old patient comes to a hospital with the complaints of acute onset profuse and purulent urethral discharge, urgency, itching sensation of the urethra and burning sensation during micturition. The patient have also painless and regular ulcer on genitalia.

What is the most likely diagnosis of this patient?

- (A) Syphilis
- (B) Chancroid
- (C) Gonorrhea
- (D) Herpes genitalia

80. An old man came to a hospital and was diagnosed with rabies after dog bite. The patient has open and deep wound, paraesthesia, and tingling sensations at the site of the bite.

What is the initial management for this patient?

- (A) Suture the wound
- (B) Administer antibiotics
- (C) Administer post exposure prophylaxis
- (D) Wash the wound with soap and water

81. A group of people came to a hospital with increasing fever, dyspnea, stridor, hypoxia, and hypotension after eating raw meat prepared for the cultural celebration. Blood culture is positive for *Bacillus anthracis* bacteria.

- What is the most appropriate drug management for these patients?
- (A) Chloramphenicol
 - (B) Ceftriaxone
 - (C) Gentamicin
 - (D) Penicillin

82. A 30-year-old pregnant woman came to a health center to attend her first ANC visit. On examination, the nurse identified that woman's amniotic fluid index to be 24, fetal weight is 10% greater than the gestational age, fetal heart rate is 110/min and maternal blood pressure is 90/60mmHg. She was diagnosed for amniotic fluid disorder.

- What is the best characteristic that led to this diagnosis?
- (A) Fetal heart beat 110/min
 - (B) Blood pressure 90/60mmHg
 - (C) Increased Fetal weight
 - (D) Amniotic fluid index 24

83. A 37-week pregnant mother came to a health center for her fourth ANC visit. She complains difficulty of breathing, lethargy and fatigue. On physical examination, she has pale conjunctiva and her vital signs are in a normal range. The nurse suspected anemia.

- What is the most appropriate lab investigation that would confirm diagnosis?
- (A) Urine analysis
 - (B) Hematocrit
 - (C) Blood sugar
 - (D) VDRL test

84. A pregnant mother came to a health center for second ANC visit. Her blood pressure is 120/70mmHg, pulse rate 80/min, her weight is 56kg. On abdominal examination, the nurse assessed her fundal height by using finger method and fundal height above the umbilicus allows four fingers.

- What is most likely gestational age based on the fundal height?
- (A) 32 weeks
 - (B) 28 weeks
 - (C) 24 weeks
 - (D) 20 weeks

85. A 32-week pregnant mother came to a hospital with complaints of fever and chills for two days. She has loss of appetite and joint pain. On physical assessment, her blood pressure is 100/70 mmHg, pulse rate 84/min and her body temperature is 38.5°C. Her blood film result shows positive for *falciparum* plasmodium.

- What is the most likely maternal complication on this woman?
- (A) IUGR
 - (B) Anemia
 - (C) Post-term labor
 - (D) Spontaneous abortion

BOOKLET CODE: 012

35. A six-month-old infant with severe acute malnutrition has developed lethargy, impaired level of consciousness with the eyes slightly open. On laboratory test, blood glucose level was 40mg/dl.

What is the most appropriate nursing intervention to this infant?

- (A) Give 12.5ml 40% dextrose + 37.5ml 5% dextrose
- (B) Give 7.5ml of 40% dextrose + 37.5 distilled water
- (C) Give 12.5ml 40% dextrose + 42.5ml distilled water
- (D) Give 7.5ml of 40% dextrose + 42.5ml of 5% dextrose

36. A two-month-old infant, who was on exclusive breast feeding, is brought to a health center with the clinical features of blood stained stool as well as bruising, umbilical stump and post circumcision site bleeding.

What is the most likely cause of this infant's problem?

- (A) Vitamin K deficiency
- (B) Vitamin C deficiency
- (C) Vitamin D deficiency
- (D) Vitamin B deficiency

37. A mother brought her nine 10-month-old child to well-baby clinic. The nurse noticed that the child can say dada/mama and was able to grasp objects with thumb and forefinger. But the mother informed the nurse that the child could have gross motor developmental delay, the child's weight was 9 kg.

What should the nurse assess primarily to disprove the mother's worry?

- (A) Confirm ability of the child to crawl
- (B) Confirm ability of the child to sit independently
- (C) Confirm ability of the child to stand handling objects
- (D) Confirm ability of the child to make tripod posture

38. A 16-month-old child was brought to well-baby clinic for growth and developmental check up. The nurse in charge noticed some developmental delay based on the Freud's principle.

What could be the most appropriate nursing intervention in this case?

- (A) Advise the mother about appropriate feeding
- (B) Show the mother about how to train toileting the child
- (C) Advise the mother how to avoid aggressive behaviors of a child at this age
- (D) Recommend mother to collect learning materials to enhance intellectual development

39. A certain health center plans to do a two days immunization campaign in two selected kebelles where immunization coverage is low. Then, nurses were assigned to give vaccines for consecutive two days as it is planned.

What is the most appropriate vaccine storage to keep the potency of vaccines?

- (A) Cold box
- (B) Refrigerator
- (C) Water-packs
- (D) Vaccine carrier

BOOKLET CODE: 012

30. A four-day-old neonate, with red and swollen eye, was brought to a hospital. During examination, eyelids were sticky due to the draining pus. The mother gave birth at home and had no history of ANC follow up. The mother also informed the nurse that she lives around smoky industrial park.

What is the most appropriate management for this neonate?

- (A) Administer Ceftriaxone
- (B) Clean the eyelids using sterile normal saline
- (C) Apply 1% tetracycline ointment
- (D) Send specimen of pus prior to administer antibiotics

31. A three-day-old neonate, with a diagnosis of neonatal sepsis, has been admitted in a hospital. The infant is lethargic and has a pulse rate of 165/min, respiratory rate of 60/min and temperature of 36°C. Random blood sugar was 80mg/dl.

What is the most appropriate nursing management for this neonate?

- (A) Administer 100% oxygen
- (B) Give 5ml/kg 10% glucose
- (C) Put the baby on kangaroo mother care
- (D) Counsel the mother to continue breast feeding

32. A 24-week-old neonate, with a diagnosis of neonatal jaundice and neonatal sepsis, was admitted to a hospital. The child had difficulty of feeding and has base lines of temperature of 36.2°C; pulse rate of 120/min. Random blood sugar was 34mg/dl. The nurse wants to know the progress of the child after initial management of hypothermia.

What is the most appropriate way to determine progress of this neonate?

- (A) Offer feedings at 2-3 hours interval
- (B) Check the child if decreased body T^o
- (C) Recheck blood glucose after 30 minutes
- (D) Look for the presence of clinical sign of hypoglycemia

33. A seven-month-old infant with complaints of inability to suck breast milk and difficulty of swallowing complementary foods was brought to a health center.

What is the most important nursing diagnosis of this child?

- (A) Imbalanced nutrition
- (B) Ineffective breast feeding
- (C) Risk for fluid volume deficit
- (D) Ineffective infant feeding pattern

34. The nursing assessment from a two-year-old patient indicates that the child has oral cavity ulcer with difficulty of chewing. But, no obstruction from esophagus to the gastro-intestinal tract.

What is the most likely nursing diagnosis of this child?

- (A) Imbalance nutritional
- (B) Impaired swallowing
- (C) Risk for fluid volume deficit
- (D) Impaired oral mucosal membrane

24. A four-year-old child is brought to a health center with a complaint of cough of three days duration. On physical examination, the child has chest in drawing. His respiratory rate is 40/min, pulse rate is 90/min, and body temperature of 38°C .

What is the appropriate management for this child?

- (A) Refer urgently to hospital
- (B) Provide oral Amoxicillin for 5 days
- (C) Treat with Ampicillin and gentamycin
- (D) Counsel the mother to give fluid remedies

25. A 6-month-old HIV positive baby was admitted to a pediatric ward due to opportunistic infection (OIs) characterized by abrupt onset of fever, dry cough, tachypnea, dyspnea and cyanosis. Physical examination reveals scattered rales in the lungs.

What is the most likely infectious disease that the patient exhibits?

- (A) Pneumocystis carini pneumonia
- (B) Pulmonary Tuberculosis
- (C) Millary Tuberculosis
- (D) Severe Pneumonia

26. A nursing team has prepared a nursing care plan to provide a holistic care to HIV exposed infants. The mother has planned to stop breast feeding today. The team planned to do a rapid antibody test and appointed her for the next visit.

What is the most appropriate time to do this test?

- (A) At birth
- (B) At 6 weeks
- (C) At 10 weeks
- (D) At 12 weeks

27. A pediatric nurse started giving essential new born care by drying the baby thoroughly. During drying, the nurse observed that the child breaths spontaneously.

What is the next appropriate step the nurse should do in the case of this new born?

- (A) Cord care
- (B) Apply Tetracycline eye ointment
- (C) Initiate breastfeeding immediately
- (D) Place the baby in skin-to-skin contact

28. A five-day-old term neonate, with a diagnosis of neonatal sepsis, was admitted to a hospital. Prenatal history shows that the mother was febrile with offensive vaginal discharge before delivery. Mode of delivery was spontaneous vaginal delivery. The nurse suspects the woman with sepsis.

What is the most likely cause of this woman's problem?

- (A) Staphylococci
- (B) Escherichia coli
- (C) Haemophilus influenza
- (D) Group B streptococcus

29. A preterm baby has been admitted in the intensive care unit. The care giver reported that the neonate had mild fever and difficulty of feeding. Base line temperature was 37.8°C ; pulse rate is 120/min. Also he has pale eye lid with yellow conjunctivitis. The nurses diagnosed the newborn with icterus neonatrum.

What is the most likely criterion to diagnose this neonate's problem?

- (A) Mild fever
- (B) Pale eye lid
- (C) Difficulty of feeding
- (D) Yellow conjunctiva

19. A mother brought her a one-day-old neonate to a hospital with a complaint of inability to feed breast milk. The neonate was born at 36 weeks of gestation, has thick skin, and can grasp the nurse's palm strongly, but sucking reflex was weak. The neonate's respiratory rate is 56/min, temperature is 37 °C and pulse rate is 130/min.

What is the most likely indicator to be used to classify the child as preterm birth?

- (A) Thick skin
(B) Tachycardia
(C) Tachypnea
(D) Weak sucking reflex

20. A four-year-old child is brought to a hospital due to fever of two days duration. The child has no runny nose and mouth ulcer, but chills and loss of appetite. Her temperature is 38.5°C and respiratory rate is 55/min. The nurse diagnoses the case as pneumonia.

What additional manifestation is needed to confirm the diagnosis?

- (A) Chest in drawing
(B) Inspiratory stridor
(C) Convulsion
(D) Cyanosis

21. A seven-year-old child came to a hospital with a complaint of cough of three weeks duration. The child had weight loss and history of contact with tuberculosis patient in the last six months. The nurse wants to screen the child for Tuberculosis.

What is the most likely confirmatory screening method for this child?

- (A) Radiography
(B) Tuberculin Skin Test
(C) Clinical manifestations
(D) Acid fast bacilli test of sputum

22. A two-year-old child is brought to a hospital with a complaint of diarrhea of two days duration. The child is lethargic and unable to drink on arrival. The child also has sunken eyes. The nurse wants to classify the level of dehydration based on IMNCI protocol.

What is the most likely classification of this child?

- (A) No dehydration
(B) Dehydration
(C) Some dehydration
(D) Severe dehydration

23. A neonate was brought to a hospital with complaints of inability to breast feed and yellowish discoloration of the palms and soles. The neonate was born at 36 weeks of gestation with a birth weight of 2200 gram. Serum bilirubin level was 12 mg/dL.

What is the most appropriate nursing intervention for this neonate?

- (A) Provide exchange transfusion
(B) Put the neonate on Phototherapy
(C) Warm the infant by skin-to-skin contact
(D) Advise the mother to expose to natural light

14. A nine-year-old patient who was presented with white patches on the surface of the nail was treated with anti-fungal drugs by the nurse in accordance with the diagnosis.

What was the most likely diagnosis of this patient?

- (A) Tinea pedis
- (B) Tinea capitis
- (C) Tinea corporis
- (D) Tinea unguium

15. A two-year-old child reported to be awake from sleeping very irritable. The child had drooling and persistent cough. On examination, the child has decreased oxygen saturation and difficulty of feeding.

What is the priority nursing diagnosis of this child?

- (A) Ineffective breathing pattern
- (B) Poor body temperature control
- (C) High risk for hypoglycemia
- (D) Impaired sleep pattern

16. A three-year-old child was brought to a clinic with severe persistent cough episode associated with a high-pitched crowing sound and vomiting. The child was unimmunized and the child's elder brother has passed away with similar case.

What will be the priority nursing intervention to this child?

- (A) Isolate the child
- (B) Give cough depressants
- (C) Administer maintenance fluid
- (D) Maintain blood transmission precaution

17. An 11-year-old child who is with full thickness burn was brought to a pediatric ward. Then, the ward nurse admitted the child and planned to start fluid resuscitation in three divided doses. The nurse computed total fluid volume to be given for 24 hours using park land formula.

What is the most appropriate amount of fluid to be provided in the second eight hours of admission?

- (A) Half of fluid required in the 24 hours
- (B) Quarter of fluid required in the 24 hours
- (C) One third of fluid required in the 24 hours
- (D) Two third of fluid required during the first day

18. A five-year-old female child came to a clinic due to pain when swallowing food and fluid, and difficulty of breathing for three days. Physical examination with tongue depressor revealed the area was inflamed and edematous, erythematous and multiple petechial rashes were identified in the pharynx.

What is the most likely cause of this child's problem?

- (A) Escherichia coli
- (B) Group B streptococci

- (C) Haemophilus Group virus
- (D) Group A β -hemolytic streptococcus

65. A matron nurse has conducted supervision in the inpatient wards of a tertiary hospital and identified the level of performance standards, communicated with staff members, reviewed activities quarterly and made correction on resource mobilization. In due process, the matron nurse received complaints about his inability to bring change and make use of subordinates' expertise knowledge.

What is the most likely supervisory role missed by the matron?

- (A) Provide performance feedback to staffs
- (B) Periodic assessment of staffs performance
- (C) Work with the staff to deliver quality service
- (D) Identify standards of good performance to staffs

66. A zonal health administration monitoring and evaluation team plans to perform controlling activities in the HIV free generation project. On the process, the team wants to know the number of nurses trained on ART and VCT to plan the next training in the project implementation area.

What is the most likely monitoring result level to be used by the team?

- (A) Input
- (B) Process
- (C) Output
- (D) Outcome

67. A researcher observed that work site injuries are the common problem among workers in metal manufacturing industry. He plans to conduct research to identify the magnitude and factors associated with work site injuries among these industry workers.

What is the most likely type of study design appropriate for this study?

- (A) Cohort
- (B) Ecological
- (C) Case control
- (D) Cross sectional

68. A researcher needs to assess the proportion of alcohol drinking at high school level students. He selected male secondary school students only in order to determine the proportion.

What is the most likely type of error committed by the researcher?

- (A) Random error
- (B) Sampling error
- (C) Non-systematic error
- (D) Non-sampling error

69. A nurse researcher wants to conduct study on access to safe water supply at household level in district X. The nurse took 500 households from 10,000 households and he used systematic sampling technique to select his sample population. The 4th household was selected as first randomly.

What is the most likely consecutive sample households included in the research?

- (A) 4, 6, 8, 12
- (B) 4, 24, 44, 64
- (C) 4, 30, 56, 82
- (D) 4, 504, 1004, 1504

60. The ministry of Health certified a referral hospital in the title of the excellent infection prevention performance of the year. During the review, the ministry found best experience of nurses in following series of related actions that must be taken to prevent infection.

What was the most likely standing plan used by the hospital?

- (A) Rules
- (B) Policies
- (C) Programs
- (D) Procedures

61. The human resource department of a primary hospital prepared a document that consisted job description, position description and hiring specification of a vacant job and posted it on the organizational display board for the applicants.

What is the most likely human resource management activity used by the department?

- (A) Staffing
- (B) Selection
- (C) Orientation
- (D) Recruitment

62. A health manager at general hospital wants to decide on effectiveness of previous staff hiring procedures and plans to motivate the staff with better achievement. During the process of implementing the motivation plan, the manager wants to use benchmark to increase salary for those staff who fulfilled their duty as expected.

What is the most work performance data to be used by the manager?

- (A) Performance appraisal
- (B) Organizational analysis
- (C) Compensation record
- (D) Operational analysis

63. In a primary hospital, nurses who are working in different wards attend nursing morning session. During the session, the achievement of a nurse was presented in form of sharing for the others as "the nurse frequently updates himself and uses his time and medical supplies properly to give nursing care for the patients".

What is the most appropriate concept of management used by the nurse?

- (A) Efficiency
- (B) Information
- (C) Effectiveness
- (D) Work Relations

64. A newly appointed matron influenced the working habits of his colleagues in all nursing care provision areas by organizing all nurses to follow harmonized standard nursing procedures and approaches suitable to the hospital. As a result, the hospital won the annual hospital service award in the country by ranking first.

What is the most likely leadership character exhibited by the matron?

- (A) Copying
- (B) Imitation
- (C) Innovation
- (D) Maintenance

60. The ministry of Health certified a referral hospital in the title of the excellent infection prevention performance of the year. During the review, the ministry found best experience of nurses in following series of related actions that must be taken to prevent infection.

What was the most likely standing plan used by the hospital?

- (A) Rules
- (B) Policies
- (C) Programs
- (D) Procedures

61. The human resource department of a primary hospital prepared a document that consisted job description, position description and hiring specification of a vacant job and posted it on the organizational display board for the applicants.

What is the most likely human resource management activity used by the department?

- (A) Staffing
- (B) Selection
- (C) Orientation
- (D) Recruitment

62. A health manager at general hospital wants to decide on effectiveness of previous staff hiring procedures and plans to motivate the staff with better achievement. During the process of implementing the motivation plan, the manager wants to use benchmark to increase salary for those staff who fulfilled their duty as expected.

What is the most work performance data to be used by the manager?

- (A) Performance appraisal
- (B) Organizational analysis
- (C) Compensation record
- (D) Operational analysis

63. In a primary hospital, nurses who are working in different wards attend nursing morning session. During the session, the achievement of a nurse was presented in form of sharing for the others as "the nurse frequently updates himself and uses his time and medical supplies properly to give nursing care for the patients".

What is the most appropriate concept of management used by the nurse?

- (A) Efficiency
- (B) Information
- (C) Effectiveness
- (D) Work Relations

64. A newly appointed matron influenced the working habits of his colleagues in all nursing care provision areas by organizing all nurses to follow harmonized standard nursing procedures and approaches suitable to the hospital. As a result, the hospital won the annual hospital service award in the country by ranking first.

What is the most likely leadership character exhibited by the matron?

- (A) Copying
- (B) Imitation
- (C) Innovation
- (D) Maintenance

45. A 21-year-old woman, with no previous history of mental illness, came to a clinic complaining hearing of persistent voices commenting all her actions in the last three months. Mental status examination revealed that she has frequent derailment of speech and looks apathetic.

What is the most likely diagnosis of this patient?

- (A) Schizophrenia
- (B) Brief psychotic disorder
- (C) Schizoaffective disorder
- (D) Schizophreniform disorder

46. A known adult patient with schizophrenia and who is currently on haloperidol 5 mg/day, came for follow up visit. His attendant reports that he was doing well with this medication but started to show returning of symptoms after he resisted to take his medication. No evidence of drug side effect observed.

What is the most appropriate drug of choice for this patient?

- (A) Thioridazine
- (B) Chlorpromazine
- (C) Fluphenazine decanoate
- (D) Couple of haloperidol and Carbamazepine

47. A 19-year-old male student reports sleep disturbance, difficulty of concentrating and periods of extreme muscle tension. He is constantly worried about the health of his parents, his university study and many others since joining university eight months ago. He recognizes that his worries are often unfounded, but can't stop worrying.

What is the most likely diagnosis of this patient?

- (A) Panic disorder
- (B) Social anxiety disorder
- (C) Separation anxiety disorder
- (D) Generalized anxiety disorder

48. A 20-year-old university student was brought by the school counselor to an emergency department. He is on mood stabilizer medication for anxiety disorder. Currently, he was found sitting in the lift of the residential hall. He claims that he hears a voice talking to him despite no one is around.

What is the most likely current psychopathology of this student?

- (A) Thought disturbance
- (B) Cognitive disturbance
- (C) Emotional disturbance
- (D) Perceptual disturbance

49. A 30-year-old male patient came to a psychiatric department with a complaint of persistent low mood and insomnia for the last two weeks. He has feeling of hopelessness, worthlessness and suicidal ideation. He has no underlined medical condition. The nurse diagnosis the patient with major depressive disorder.

What is the most likely criterion used to diagnose the patient's problem?

- (A) Insomnia
- (B) Low mood
- (C) Hopelessness
- (D) suicidal ideation

30. A four-day-old neonate, with red and swollen eye, was brought to a hospital. During examination, eyelids were sticky due to the draining pus. The mother gave birth at home and had no history of ANC follow up. The mother also informed the nurse that she lives around smoky industrial park.

What is the most appropriate management for this neonate?

- (A) Administer Ceftriaxone
- (B) Clean the eyelids using sterile normal saline
- (C) Apply 1% tetracycline ointment
- (D) Send specimen of pus prior to administer antibiotics

31. A three-day-old neonate, with a diagnosis of neonatal sepsis, has been admitted in a hospital. The infant is lethargic and has a pulse rate of 165/min, respiratory rate of 60/min and temperature of 36°C. Random blood sugar was 80mg/dl.

What is the most appropriate nursing management for this neonate?

- (A) Administer 100% oxygen
- (B) Give 5ml/kg 10% glucose
- (C) Put the baby on kangaroo mother care
- (D) Counsel the mother to continue breast feeding

32. A 34-week-old neonate, with a diagnosis of neonatal jaundice and neonatal sepsis, was admitted to a hospital. The child had difficulty of feeding and has base lines of temperature of 36.2°C; pulse rate of 120/min. Random blood sugar was 34mg/dl. The nurse wants to know the progress of the child after initial management of hypothermia.

What is the most appropriate way to determine progress of this neonate?

- (A) Offer feedings at 2-3 hours interval
- (B) Check the child if decreased body T⁰
- (C) Recheck blood glucose after 30 minutes
- (D) Look for the presence of clinical sign of hypoglycemia

33. A seven-month-old infant with complaints of inability to suck breast milk and difficulty of swallowing complementary foods was brought to a health center.

What is the most important nursing diagnosis of this child?

- (A) Imbalanced nutrition
- (B) Ineffective breast feeding
- (C) Risk for fluid volume deficit
- (D) Ineffective infant feeding pattern

34. The nursing assessment from a two-year-old patient indicates that the child has oral cavity ulcer with difficulty of chewing. But, no obstruction from esophagus to the gastro-intestinal tract.

What is the most likely nursing diagnosis of this child?

- (A) Imbalance nutritional
- (B) Impaired swallowing
- (C) Risk for fluid volume deficit
- (D) Impaired oral mucosal membrane

24. A four-year-old child is brought to a health center with a complaint of cough. On physical examination, the child has chest distress during the respiratory rate is 40/min, and body temperature of 39 °C.

What is the appropriate management for this child?

- (A) Refer urgently to hospital
- (B) Provide oral Amoxicillin for 5 days
- (C) Treat with Aspirin and paracetamol
- (D) Counsel the mother to give fluid remedies

25. A 6-month-old HIV positive baby was admitted to a pediatric ward due to opportunistic infection. On physical examination, the child has chest distress, dry cough, tachypnea, dyspnea and anorexia. Physical examination reveals scattered rales in the lungs.

What is the most likely infectious disease that the patient exhibits?

- (A) Pneumocystis carinii pneumonia
- (B) Pulmonary Tuberculosis
- (C) Malaria/Tuberculosis
- (D) Severe Pneumonia

26. A nursing team has prepared a nursing care plan to provide a babies care to fully express milk. The mother has planned to stop breast feeding today. The team planned to do a rapid antibiotic test and advised her for the next visit.

What is the most appropriate time to do this test?

- (A) At birth
- (B) At 6 weeks
- (C) At 10 weeks
- (D) At 12 weeks

27. A pediatric nurse started giving essential new born care by doing the baby thoroughly. During drying, the nurse observed that the child breaths spontaneously.

What is the most appropriate step the nurse should do in absence of this new born?

- (A) Cold care
- (B) Apply Tetracycline eye ointment
- (C) Initiate breastfeeding immediately
- (D) Place the baby in skin-to-skin contact

28. A five-day-old term neonate, with a diagnosis of neonatal sepsis, was admitted to a hospital. Perinatal history shows that the mother was febrile with offensive vaginal discharges before delivery. Mode of delivery was spontaneous vaginal delivery. The nurse suspects the woman with sepsis.

What is the most likely cause of this woman's problem?

- (A) Staphylococci
- (B) Escherichia coli
- (C) Haemophilus influenza
- (D) Group B streptococcus

29. A preterm baby has been admitted in the intensive care unit. The care giver reported that the neonate had mild fever and difficulty of feeding. Basal temperature was 37.8 °C, pulse rate is 120/min. Also he has pale eye lid with yellow conjunctivitis. The nurse diagnosed the newborn with icterus neonatorum.

What is the most likely criterion to diagnose this neonate's problem?

- (A) Mild fever
- (B) Pale eye lid
- (C) Difficulty of feeding
- (D) Yellow conjunctiva

30. A four-day-old neonate, with red and swollen eye, was brought to a hospital. During examination, eyelids were sticky due to the draining pus. The mother gave birth at home and had no history of ANC follow up. The mother also informed the nurse that she lives around smoky industrial park.

What is the most appropriate management for this neonate?

- (A) Administer Ceftriaxone
- (B) Clean the eyelids using sterile normal saline
- (C) Apply 1% tetracycline ointment
- (D) Send specimen of pus prior to administer antibiotics

31. A three-day-old neonate, with a diagnosis of neonatal sepsis, has been admitted in a hospital. The infant is lethargic and has a pulse rate of 165/min, respiratory rate of 60/min and temperature of 36°C. Random blood sugar was 80mg/dl.

What is the most appropriate nursing management for this neonate?

- (A) Administer 100% oxygen
- (B) Give 5ml/kg 10% glucose
- (C) Put the baby on kangaroo mother care
- (D) Counsel the mother to continue breast feeding

32. A 34-week-old neonate, with a diagnosis of neonatal jaundice and neonatal sepsis, was admitted to a hospital. The child had difficulty of feeding and has base lines of temperature of 36.2°C; pulse rate of 120/min. Random blood sugar was 34mg/dl. The nurse wants to know the progress of the child after initial management of hypothermia.

What is the most appropriate way to determine progress of this neonate?

- (A) Offer feedings at 2-3 hours interval
- (B) Check the child if decreased body T⁰
- (C) Recheck blood glucose after 30 minutes
- (D) Look for the presence of clinical sign of hypoglycemia

33. A seven-month-old infant with complaints of inability to suck breast milk and difficulty of swallowing complementary foods was brought to a health center.

What is the most important nursing diagnosis of this child?

- (A) Imbalanced nutrition
- (B) Ineffective breast feeding
- (C) Risk for fluid volume deficit
- (D) Ineffective infant feeding pattern

34. The nursing assessment from a two-year-old patient indicates that the child has oral cavity ulcer with difficulty of chewing. But, no obstruction from esophagus to the gastro-intestinal tract.

What is the most likely nursing diagnosis of this child?

- (A) Imbalance nutritional
- (B) Impaired swallowing
- (C) Risk for fluid volume deficit
- (D) Impaired oral mucosal membrane

55. A 53-year-old male HIV positive patient was admitted in a surgical ward after BPH surgery. During admission, he was calm and cooperative. He has been on HAART for the last four years but, currently he has an additional intravenous antibiotic. One day after the operation, his condition started deteriorating as evidenced by disorientation, fluctuating level of consciousness and perceptual distortions.

What is the most likely diagnosis of this man?

- (A) Delirium
- (B) Medication-induced Neurocognitive disorder
- (C) Neurocognitive disorder due to HIV infection
- (D) Neurocognitive disorder due to another medical condition

56. A six-year-old boy was brought to a hospital with a complaint of pattern of diminished sustained attention and higher levels of impulsivity than his age mates. He had persistent problems with academic achievement. He was frequently sent to the school principal and his parents were similarly frustrated at home. During Examination, he has poor speech articulation and difficulty to wait his turn.

What is the most likely diagnosis of this child?

- (A) Autistic disorder
- (B) Conduct related disorder
- (C) Pervasive developmental disorders
- (D) Attention deficit hyperactivity disorder

57. The presidents of public university in Ethiopia addressed overall activities of the university for the graduation attendees in his opening speech. At lunch time, the president signed a community service provision project agreement with government officials.

What is the most likely managerial role played by the president?

- (A) Liaison
- (B) Leader
- (C) Figurehead
- (D) Entrepreneur

58. The chief executive officer (CEO) of a hospital has organized a task force to carry out SWOT analysis on the entire functional areas of the hospital. After gathering and analyzing data, the task force came up with evidences showing frequent shortage of drug supply, presence of poorly motivated staff, and low availability of transportation facility in the hospital.

What is the most likely SWOT analysis component observed in this hospital?

- (A) Threat
- (B) Strength
- (C) Weakness
- (D) Opportunity

59. The human resource officer of a hospital wants to estimate the number of nurses required to be employed for next year. Then, the officer consulted the chief administrator to identify criteria for the estimation and agreed to appraise the number based on the number of services in the hospital.

What is the most appropriate base for estimating the human resource required by the hospital?

- (A) Health service targets
- (B) The health needs approach
- (C) The health care demands
- (D) Human resource-to-population ratios

BOOKLET CODE: 011

72. A 32-year-old man was presented with a complaint of watery diarrhea of two days duration. The diarrhea has a sudden onset of copious. He has severe muscle pain due to frequent diarrhea. The laboratory finding shows the presence of small, comma-shaped, motile, Gram-negative bacilli on stool.

What is the most effective prevention method for this case?

- (A) Sanitation
- (B) Antibiotics

- (C) Monitor output
- (D) Parenteral vaccine

73. A 45-year-old known HIV patient came to an outpatient department with complaint of nausea, vomiting and abdominal cramp. The nurse highly suspects Giardia lamblia infection.

What is the most likely additional assessment finding that supports the nurse's suspicion?

- (A) Foul-smelling greasy stools
- (B) Pale greasy bloody diarrhea

- (C) Pale greasy mucoid diarrhea
- (D) Prolonged watery diarrhea

74. A 16-year-old female patient visits her doctor with a complaint of cough and headache of three days duration. On history, she has sudden onset fever, runny nose, myalgia, chills and sore throat. The cough is severe and protracted.

What is the priority nursing diagnosis for this patient?

- (A) Acute pain
- (B) Hyperthermia

- (C) Impaired comfort
- (D) Impaired skin integrity

75. During a seminar presentation, a graduating student presents about the mode of transmission of leprosy. But students were confused about the main method of transmission and asked the presenter for clarification.

What is the best answer to be explained by the presenter?

- (A) Through sexual contact
- (B) Through respiratory secretion
- (C) Through broken skin and drainage
- (D) Through mother to child trans placental

67. A male nurse in the operating room mentors 4th year graduating nurses and teaches the students about different zones in the operation room. He explained about the area where sterile procedures are carried out and masks are required to supplement surgical attire.

- What is the most likely zone in the operation room the nurse explained about?
- (A) Semi-restricted Area
 - (B) Unrestricted Area
 - (C) Restricted Area
 - (D) Sterile area

68. A 19-year-old female client was brought to an emergency care unit after exposure to cyanide containing by-product. The nurse initiated emergency nursing assessment to do possible interventions.

- What is the most likely common clinical manifestation the nurse may get from this patient?
- (A) High blood pressure and cyanotic
 - (B) Tachycardia and tachypnea
 - (C) Bradypnea and bradycardic
 - (D) Cyanosis and dizziness

69. A 57-year-old woman was presented to surgical emergency unit with diagnosis of multiple ribs fracture. Assessment findings showed a paradoxical chest-wall movement with inspiration and expiration.

- What is the priority nursing intervention next to immobilization?
- (A) Administering anti pain
 - (B) Monitor signs of respiratory infection
 - (C) Instructing the lie on the affected side
 - (D) Instructing the patient to take deep breaths

70. A 68-year-old man with a history of bronchiectasis is admitted for newly diagnosed heart failure. He reports that he is able to breathe easily only in a sitting position. The attending physician ordered chest physiotherapy and postural drainage.

What is the priority nursing diagnosis for this patient?

- (A) Anxiety
- (B) Activity intolerance
- (C) Ineffective airway clearance
- (D) Impaired physical comfort

71. A 25-year-old female patient was admitted to orthopedic inpatient unit with a diagnosis of dislocation at the left lower extremity. The surgeon performed immobilization and reduction to stabilize the patient and handover to the nurse for follow up.

What is the most likely instruction the nurse should notify the patient?

- (A) Do not flex the hip
- (B) Reduce range of motion
- (C) Reporting abnormal sensation
- (D) Keep the knees apart at all times