

**1, A 30 year- old prisoner presents with intermittent fever and headache of two days duration. He claims that five of his roommates had similar symptoms and were treated in the prison. blood film examination showed a splinter cell which one of the following is most appropriate antibiotic choice for this patient?**

**A, ceftriaxone B, ciprofloxacin C, procaine penicillin D, vancomycin**

**2, A 29 year- old man comes to emergency clinic with crampy abdominal pain and bloody diarrhea of one day. Stool microscop shows trophozoites which one of the following most likely cause?**

**A, ascaris lumbricoides B, entamoeba histolytica C. giardia lamblia D, trichuris trichiura**

**3, A 25 year- old male with known liver disease presents with cough, fever and weight loss for one month sputum examination for acid fast bacilli is positive 3 times.**

**Which combination anti tuberculosis treatment is most appropriate for this patient?**

**A, 2ERHZ/4HE**

**B, 2RHZ/4RH**

**C, 2SERH/6RH**

**D,2SRZE/6HE**

**4, A 55 year- old male comes to emergency OPD because of excruciating pain in his right big toe.He was diagnosed to have gouty arthritis. Which of the following is the best intial pharmacotherapy?**

**A, Ceftriaxone B, Allopurinol C, indomethacin D, Morphine**

**5, which the following is recommended for treating Chlamydia trachomatis infection?**

**A, metronidazole B, fluconazole C, doxycyline D, Gentamicin**

**6, A health officer treating a 74 years old man who was admitted with shortness of breathing and fever . his temperture on admission was 39<sup>0</sup> c.on physical examination he is sick looking and has decrease air entery at the right basal lung field, the chest x-ray shows an infiltrate at the right lower lobe. What is the most likely causative organism?**

**A, streptococcus pneumonia**

**B,Legionella pneumophila**

**C,viral pneumonia**

**D, klebsiella pneumonia**

**7, what is the treatmen of choice for mild persistent asthma?**

**A, Adrenaline**

**B, Nebulized albuterol**

**C, salbutamol**

**D,steroids**

**8, A 30 year- old male with long history of sever asthma patient to the emergency room with shortness of breath. He has been previously admitted to the hospital and was once intubated for severe respiratory distress. Which of the following physical finding indicate a life threatening asthma ?**

**A,diffuse wheeze B,PR= 110/min C,RR=26/min D, silent chest**

**9,A 62 year- old woman with heart failure develops pneumonia and large pleural effusion. Laboratory I nvestigation shows serum lactate dehydrogenase (LDH) 420mg/dl and serum protein 8g/dl .thoracentesis is performed in an effort to determine which the pleural effusion is due to HF or pneumonia.which of the following finding would indicate that the pleural effusion is due to HF?**

**A,Glucose content of 20mg/dl**

**B, LDH content of 100mg/dl**

**C, PH of 7.13**

**D, protein content of 6gm/dl**

**10, A 35 year- old pregnant woman presents with fever easily fatigability and change in mental status. On physical examination , BP=100/60mmhg, PR=108/min, RR=20/min. T° =37.5°c and has pale conjunctiva and confused. Laboratory examination reveals hemoglobin=7g/dl, random blood sugar=30mg/dl and plasmodium falciparum positive . which of the following in the initial step in the management of this patient ?**

- A, Artesunate IV**
- B, Artemether-lumefantrine orally**
- C, Blood transfusion**
- D, Glucose infusion**

**11, A 35 year- old woman presented with four episodes of spasm. Generalized rigidity and trismus after she sustained a puncture over the left foot two days ago. She is fully conscious. Which of the following is the most appropriate pharmacotherapy?**

- A, Carbamazepine B, Diazepam C, Phenobarbital D, Phenytoin**

**12, A 22 years old HIV positive man with a presumptive diagnosis of pneumocystis jiroveci pneumonia has no signs of respiratory distress. Which of the following is the most appropriate pharmacotherapy?**

**A, Fluconazole**

**B, fluconazole and prednisolone**

**C, Cotrimoxazole**

**D, Cotrimoxazole and prednisolone**

**13 A 35 year- old woman presents with fever and acute lower left – quadrant pain. She states that it began last night and began in the back and radiates to the lower left – quadrant . she also has dysuria and urinary frequency. Her body temperture is 38.9°c. There is lower left quadrant tenderness and left costovertebral tenderness . what is the most likely diagnosis?**

**A, Diverticulitis B, Volvulus C, Ovarian torsion D, pyelonephritis**

**14, A 50 year- old male with chronic kidney disease presents with swelling of his feet. Physical examination revealed BP= 150/90mmhg PR=78/min RR= 18/min and bilateral ankle edema. Laboratory studies showed BUN=62 mg/dl, serum creatinine=3.2mg/dl, serum electrolytes( $K^+$  level=5.6meq/l,  $Na^+$  = 146meq/dl) and total plasma cholesterol Level of 260 mg/dl. Which of the following drugs improves the prognosis of his disease?**

**A, Enalapril**

**B, Hydrochorthiazide**

**C, Salt restriction**

**D, simvastatin**

**15, A 22 year –old man is admitted to a burn unit after he sustained a scald burn. Three days after his admission, a bright red blood starts to ooze out through the NG-tube. He also has mild epigastric pain. Under GI endoscopy shows multiple, tiny ulcers in the corpus of the stomach. Which one of the following is the most likely diagnosis**

**A, peptic ulcer disease**

**B, gastric atrophy**

**C, acute erosive gastritis**

**D, non-erosive gastritis**

**16, a 60 year old man had cirrhosis of the liver secondary to hepatitis B for years, in the last couple of months his jaundice started to worsen and developed sleep disturbance. He also complained of vague, right upper quadrant pain since four weeks, physical examination shows a rough, nodular liver, which one is an additional useful diagnostic test for this patient?**

**A, five-hydroxy indoleacetic acid**

**B alpha-fetoprotein**

**C,carcinoembryogenic antigen**

**D,hepatitis B virus DNA**

**17.A 47 year old man comes to the hospital complaining of epigastric pain that is exacerbated by eating. A test for helicobacter pylori turned positive and endoscopy confirmed a duodenal ulcer. Which one is the most appropriate therapy?**

**A.amoxicillin,clarithromycin and omeprazole**

**B.metronidazole and anti acids**

**C.omeprazole and antiacids**

**D.rantidine,amoxicillin and metronidazole**

**18.A 40 year old woman has a one week history of jaundice and right upper quadrant abdominal pain.associated symptoms include dark urine and pale stools but no weight loss. Liver function test :**

**AST=50IU/L.,ALT=350IU/L.,bilirubin=40 $\mu$ mol/L what is most likely diagnosis?**

**A.viral hepatitis B**

**B. gallstones**

**C.alcholic hepatitis**

**D.pancreatic carcinoma**

**19. a 60 year old man who had been smoking a pack of cigarettes daily for the last twenty years presented with one year history of increasing dyspnea on exertion and accumulation dry cough. Chest examination reveals hyper resonance all over the lungs and decreased breath sounds. Which of the following is most likely diagnosis?**

**A, Bronchiectasis**

**B,Asthma**

**C,Emphysema**

**D, chronic bronchitis**

**20,A 36 year old woman came with complaints of malaise chills and fever. She also had nausea, head ache and anorexia. She reported occasional iv drugs use. On auscultation, a loud holosystolic murmur is detected. What is next best step to confirm the diagnosis?**

**A, cardiac catheterization**

**B,chest CT scan**

**C, ECG**

**D,Echocardiogram**

**21. A 60 year-old man came to the OPD with complaints of palpitation. Irritability and insomnia for two months. He is**

**taking amiodorone for cardiac arrhythmia and enalapril for hypertention. What is the most appropriate next step in management?**

- A. refer for psychiatric evalution**
- B.Measure thyroxin and TSH levels**
- C. Administer propranolol**
- D. Administer nifedipine**

**22. A 27 year- old woman complains of fever and fatigue. Non tender, enlarged cervical and axillary lymph nodes are noted. Chest x-ray shows hilar lymphadenopaty and diffuse interstitial infiltrates. Lymph node biopsy shows non – caseating ganuloma. Which drug is the most appropriate therapy?**

- A.Allopurinol**
- B.Cyclosporine**
- C.Glucocorticoid**
- D. Isoniazed**

**23. A 35 year-old known diabetic patient presented with mild hypertension and micro-albuminuria. Blood lipids and renal function tests are normal. There is evidence of left ventricular**

**hypertrophy on echocardiography. Which one is the most appropriate anti-hypertensive?**

**A.ACE inhibitors**

**B.bata blockers**

**C.Calcium channel blockers**

**D. Thiazide diuretics**

**24. which one of the following signs is associated with meningitis?**

**A. Rovsing's sign**

**B.kernig's sign**

**C.Tinel's sign**

**D. Trousseau's sign**

**25. A 52 year-old female presents to the clinic with four months of fatigue, lower extremity burning pain and numbness.laboratory test reveals Hgb=8.5g/d, MCV=118fl, WBC=4000/ $\mu$ l and platet=150,000/ $\mu$ l.what is the most diagnosis?**

**A. Aplastic anemia(normocytic)**

**B. Folate- deficiency anemia**

**C. iron-deficiency anemia**

**D. vitamin B12 deficiency anemia**

**26. A 25 year-old women is receiving packed RBC for severe anemia. She develops fever and chills fifteen minutes after initiation of transfusion. What is the next best step in the management of thi patient?**

**A. Anti-histamines**

**B. Glucocrticoids**

**C. normal saline infusion**

**D. stop transfusion**

**27. A 66 year-old diabetic man suddenly collapsed in a hospital cafeteria and quick evaluation revealed an unresponsive man with no respiratory effort and pulseless carotids. Which of the following is the most likely diagnosis?**

**A. cardiac arrest**

**B. Hypoglycemia**

**C. Non-convulsive seizure**

**D. syncope**

**28. A patient with a long standing valvular heart disease secondary to rheumatic fever is being treated for heart**

**failure. Which of the following drugs delays the progression of the disease?**

**A. Digoxin-use for increase cardiac contraction**

**B. Enalapril**

**C. Frusemid**

**D. penicillin**

**29. A 30 year- old man collapses at a wedding and is witnessed to have seizure lasting two minutes. when a paramedic arrives and calls her name and asks some questions she mumbles and no one could hear what she said. When pressure was applied to her nail bed she opened her eyes, rub her nail and push away the hand holding her nail bed. what is his Glasgow comma (gcs)?**

**A.5**

**B.9**

**C.11**

**D.12**

**30. A 60 year old man is admitted with left side hemiparesis. CT reveals a middle cerebral artery infract. What is the most significant risk factor for his condition?**

**A. Diabetes**

**C. Hypertension**

**B. hypercholesterolemia**

**D. smoking**

**31. A health officer is evaluating a patient with lymphoma. Which of the following lymph node characteristics are least likely?**

**A. Firm and rubbery**

**B. Discrete and mobile**

**C. Matted with discharging sinus**

**D. Non – tender**

**32. A 20 year old woman who has been diagnosed with pulmonary tuberculosis and started anti-tuberculosis treatment two weeks ago is found to be HIV positive on screening. On physical examination, she has pale conjunctiva and laboratory investigation shows CD4 count of 100 cells/ $\mu$ l. Which one of the following is most appropriate first antiretroviral regimen for this patient?**

**A. ABC/3TC/NVP**

**B. AZT/3TC/EFV**

**C. D4T/3TC/NVP**

**D. TDF/3TC/EFV**

**33. A 40 year old alcoholic develops productive cough and fever. Chest x-ray shows an air fluid level in a superior segment**

of the right lower lobe. which one of the following is the most appropriate chemotherapy?

A. Ceftazidime and ciprofloxacin

B. Ceftriaxone and gentamycin

C. Metronidazole and crystalline penicillin

D. vancomycin and chloramphenicol

34. A 27 year old man complains of fever, cough with rusty – coloured sputum and pleuritic chest pain. Chest examination revealed increased tactile fremitus and dullness to percussion in the right lower lung field. what is the most likely diagnosis?

A. bronchopneumonia

C. lung abscess

B. lobar pneumonia

D. pleural effusion

35. A 60 year-old man presented with sudden onset of right sided weakness on both upper and lower extremities before a day. The weakness is maximal at onset and during sleep. He is hypertensive but does not take medications what is the most likely diagnosis?

A. Hemorrhagic stroke

C. Subarachnoid hemorrhage

B. Ischemic stroke

D. Transient ischemic

36. A 60 year-old alcoholic patient presented with fever, headache and neck pain of two days duration. On

**examination body T=39°C and has neck stiffness. Bacterial meningitis was suspected but lumbar puncture was deferred because he has papilledema. Which one of the following is the best empiric therapy for this patient in addition to ampicillin?**

**A. vancomycin and chloramphenicol**

**B. ceftriaxone and vancomycin**

**C. ceftriaxone and chloramphenicol**

**D. cloxacillin and ciprofloxacin**

**37. A 50 year-old woman presented with fever, cough and chest pain for two days. Chest x-ray showed pleural effusion. Pleural fluid analysis showed: WBC=5,000/mm<sup>3</sup>, with 90% neutrophils, pH=7.2, blood culture was positive for streptococcus pneumonia. What is the next best in management of this patient besides antibiotics?**

**A. dexamethasone**

**B. chest CT scan**

**C. chest tube insertion**

**D. lung function test**

**38. A 20 year-old man presents with generalized body swelling of two months. Physical examination reveals BP=120/70 mmHg,**

**ascites and +3 pretibial edema. Protein =4.5g/24hours urine collection .which one of the most likely diagnosis?**

**A.chronic liver disease**

**C.Nephritic syndrome**

**B.Heart failure**

**D.Glomerulonephritis**

**39.Which of the following conditions causes left lower quaderant abdominal pain?**

**A.Acute appendicitis**

**C.Meckel's diverticulum**

**B.Diverticulitis**

**D.volvulus**

**40.A 32 year-old man presented with an anterior neck mass of one year duration.On examination he has a hoarse voice and a solitary 2cm x 2cm thyroid non tender swelling on the right side which has firm consistency and fixed to the underlying structures . There was no lymphadenopathy. What is the most likely diagnosis?**

**A. Toxic nodule**

**B.Thyroiditis**

**C.thyroid cancer**

**D.Goiter**

**41. A 68 year old man is admitted to hospital with a complaint of intermittent lower abdominal pain ,bowel habit changes and rectal bleeding one month duration. He has tensmus and**

**incomplete sensation of defecation and has lost weight. What is the most likely diagnosis?**

**A. amebic dysentery**

**C. sigmoid volvulus**

**B.colorectal cancer**

**D.crohn's disease**

**42.A . 55 year – old man presented to the surgical OPD with rectal bleeding of one month duration. He has no protruding mass per rectum and digital rectal examination was non-revealing. What is the next best step in managing this patient?**

**A. Barium enema**

**C.CT-Scan of the abdomen**

**B.Coloscopy**

**D.plain abdominal X-ray**

**43.A. 30 year- old man presented to the emergency surgical OPD with perianal pain and bright red rectal bleeding durin defecation. He has fever chills . perianal examination showed shallow laceration posteriorly. What is the most likely diagnosis?**

**A.Anal fissure**

**C. perianal fistula**

**B.perirectal abscess**

**D.Thrombosed hemorrhoids**

**44.A 20 year-old man was seen at the emergency department following road traffic accident of one hour duration. He has sustained trauma to the abdomen. On examination; vital signs:BP=80/50mmhg and PR=120/min.He hastier marks on**

**the abdomen and it was tender. The patient is started on iv fluids. What is the next step in the management?**

- A. abdominal CT- scan**
- B. Exploratory laparotomy**
- C. Observation**
- D. Plain abdominal x-ray**

**45. a 30 year-old man presented with fatigue and dyspepsia of three months duration. Physical examination revealed a pale conjunctiva with a palpable mass at the epigastrium that moves with respiration. What is the gold standard diagnostic modality ?**

- A. Abdominal Ultra sound**
- B. CT scan of the abdomen**
- C. Upper Gi Endoscopy**
- D. plain abdominal film**

**46. A 23 year-old farmer presented to the OPD with abdominal pain and frequent vomiting of three hours duration, on examination the abdomen is slightly distended with visible peristalsis. Plain abdominal x-ray showed a step ladder pattern air fluid level. Which of the following is the most likely diagnosis ?**

- A. caecal volvulus**
- B. Compound volvulus**
- C. sigmoid volvulus**
- D. small bowel volvulus**

**47. A 65 year – old man was seen at the Emergency OPD with abdominal pain, distension and failure to pass feces and flatus of two days duration. He had two similar attacks in the last three years and was successfully treated with rectal tube deflation. On examination BP= 90/60mmHg and PR=110/min. The abdomen was tender with absent bowel sounds. Which of the following measures should be avoided?**

**A. Analgesics B. Antibiotics C. iv fluid D. rectal tube deflation**

**48. A 45 year- old housewife presented with progressive submandibular swelling of two days duration. She visited a dentist and undergone tooth extraction a day prior to the start of her symptoms. Examination showed tense, shiny bilateral submandibular swelling with no area of fluctuation. She also has drooling of the saliva and protruded tongue. What is the diagnosis of this patient?**

**A. Ludwig's angina**

**C. Periodontal abscess**

**C. submandibular cellulites**

**D. submental abscess**

**49. A 31 year-old male patient was brought to the emergency OPD following a car accident. He was struck on his chest wall movement on the right side. Chest X-ray showed three rib**

**fracture at two sites. What is the most likely diagnosis of this patient?**

**A.Flail chest**

**B.Hemothorax**

**C. multiple rib fracture**

**D.simple rib fracture**

**50. A 20 year old man came to emergency 30 minutes after he sustained stab injury to his left posterior chest. He has shortness of breathing . He is acutely sick looking with BP=110/60mmhg , PR=90 and afebrile. has decreased air entry on the lower third of the left chest with visible air rush through the stab wound. What is the initial step in management of this patient?**

**A. Apply airtight dressing**

**B.Apply triangular dressing**

**C.Suture the wound**

**D.Take chest x-ray**

**51.A 30 year-old man came after sustaining car accident. He has trauma to his chest. Examination shows stable vital signs,point tenderness on left lateral chest with good air entry bilaterally. Ches x-ray shows fracture of the sixth and seventh**

**ribs with no haemopneumothorax. What is appropriate management for this patient?**

**A. Analgesics and antibiotics**

**B. Analgesics only**

**C. prophylactic chest tube**

**D. Refer for fixation of the ribs**

**52. A 12 year-old boy came after he fell down from a tree of four meters height. After he fell down from a tree of four meters height. After the accident he failed to move his legs and feels pain on his back. On examination, he is conscious, BP=80/45mmHg, PR=60/min. He has bruise and severe tenderness on his back. At the level of T4 and T5. Power on his lower extremities is 2/5. What is the likely diagnosis in this patient?**

**A. Bilateral femoral fracture**

**B. Hypovolemic shock**

**C. mild head injury**

**D. spinal cord injury**

**53. A 64 year-old male patient presented to the Emergency OPD after he failed to pass urine for 24 hours. He had difficulty of urination for the last one year. The previous night he took**

**seven bottles of beer with friends. On examination he is acutely sick looking in pain. Bladder is palpable 10 cm above the pubic bone and tender. What is the most likely cause for this acute urinary retention?**

**A. Benign prostatic hyperplasia**

**B. Bladder cancer**

**C. Bladder stone**

**D. Urethral stricture**

**54. A 43 year-old male patient to the Emergency OPD after being involved in a road traffic accident before a day. He has failed to urinate since the incident and is unable to move his right lower limb . on examination his vital signs are normal , there is right pubic tenderness and blood on the penile meatus. What should be the next step in management?**

**A. Decompress with percutaneous needle puncture**

**B. Do image-guided catheterization**

**C. insert a foley catheter to decompress the bladder**

**D. perform emergency suprapubic cystostomy**

**55. An 18 year-old male presented with right scrotal swelling of one day duration. He also has associated pain which started while he was asleep and woke him up. He also has fever and**

**pain during urination. On examination right testis is tender, hot to touch, swollen, and the testicular cord is not thickened. What is the most likely diagnosis?**

**A. Testicular torsion**

**B. Epididymo-orchitis**

**C. Infected hydrocele**

**D. Torsion of the testicular appendages**

**56. A 31 year-old male presented came to the OPD after he sustained trauma to his right flank. He fell down from a two story building .He complains of pain at the site and has gross hematuria. On examination, He is acutely sick looking in pain BP=110/70mmhg, PR=96/min. There is a bruise over the right flank area that is tender to touch. Which organ is most likely injured ?**

**A. kidney B. Liver C. Ureter D. Urinary Bladder**

**57. A 60 year-old female patient presented with right sided flank pain and hematuria of one month duration .she also had significant weight loss night sweats and malaise. On physical examination, she is chronically sick looking with normal vital signs, a bimanually palpable mass on the right flank . what is the most likely diagnosis?**

**A. Chronic pyelonephritis**

**B. Neuroblastoma**

**C. Renal cell cancer**

**D. wilm's tumor**

**58. A 35 year-old male patient presented to the Emergency Room after sustaining a bullet injury to his left fore arm, intact distal neurovascular examination. X-ray showed comminuted displaced radioulnar fracture. Which of the following is a temporary means of immobilization?**

**A. Apply POP cast**

**B. External fixation**

**C. Lateral arm traction**

**D. plating and screw**

**59. A 65 year-old patient sustained a fall down accident three hours prior to presenting to the Emergency Room. He fell down with the arms extended. On examination vital signs were normal with a right wrist deformity. X-ray showed displaced colle's fracture. Which one is the best management option?**

**A. Application of a POP cast**

**B. Closed reduction and casting**

**C. open reduction and plating**

**D. skin traction for few days then casting**

**60. A 20 year-old male patient came to you with right knee pain of five days duration. On examination BP=120/70mmhg PR=84/min. The knee is swollen and tender with signs of knee effusion. X-ray showed a wide joint fluid analysis revealed turbid fluid with 100,000WBC/mm<sup>3</sup>. Which of the following is the most likely diagnosis?**

- A. Acute osteomyelitis**
- B. Osteoarthritis**
- C. pyogenic arthritis**
- D. Tuberculosis of the knee**

**61. A mother brought a 20 day-old male neonate that failed to move his right extremity since five days of age after delivery at home. On examination, the right distal thigh is grossly swollen. The skin is shiny and tender. Laboratory work up showed elevated WBC count with a left shift, ESR was elevated. x-ray showed. Periosteal reaction with distortion of the soft tissue fat plane. Which of the following is the most likely diagnosis?**

- A. Acute Osteomyelitis**
- B. cellulitis**
- C. Femoral shaft fracture**
- D. Pyomyositis**

**62. A 26 year-old GIII PI woman at 20 weeks of gestation came for antenatal care followup. Previous obstetric history was unremarkable. Which of the following cardiac findings requires further evaluation?**

- A. Diastolic murmur**
- B. Grade II systolic murmur**
- C. shifting of PMI**
- D. Third heart sound**

**63. A 24 year-old GIII PI woman came with a complaint of decreased fetal movement at 33 weeks of gestation. What is the next best step in the management of this patient?**

- A. Kick chart**
- B. Biophysical profile**
- C. Strict bed rest**
- D. Immediate delivery**

**64. A 39 year-old GV PIV lady at 34 weeks gestation has come with complaint of abdominal discomfort and decreased fetal movement. On examination vital signs are normal, shiny and tense abdomen, symphysio-fundal height 38 cm, ballotable but hardly felt fetal part and fetal heart beat difficult to appreciate. Which of the following is highly likely with this patient?**

- A. Fetal neural tube defect**
- B. maternal cardiac illness**
- C. fetal renal agenesis**
- D. pre eclampsia**

**65. a 22 year-old nulliparous woman at 42+3 weeks gestational has bishops score of 10. which of the following would be the most appropriate induction method for this patient?**

**A. Amniotomy**

**B. Ergometrine**

**C. Misoprostol**

**D. Oxytocin**

**66. A 32 year-old G III P II at 39 weeks of gestation has gone into labor. She was diagnosed to have gestational diabetes during ANC follow up. Vital signs were normal, BMI being 30kg/m<sup>2</sup>. Clinical estimated fetal weight is 4.5kg she was augmented for protracted cervical dilation and entered into second stage 2Hrs back. Which of the following puts her at higher risk of having shoulder dystocia?**

**A. Fetal weight**

**B. Maternal diabetes**

**C. parity**

**D. protracted 1<sup>st</sup> stage**

**67 . A 26 year-old GII PI woman at 20 weeks of gestation visits a doctor's office for prenatal care. He counseled her for a normal well balanced diet?**

**Calcium**

**Folic acid**

**Iron**

**Magnesium**

**68. A 26 year old primigravida at 35 weeks gestation complains of headache and facial puffiness. Her BP= 160/100mmhg and her reflexes are brisk. Which urinalysis result is highly likely in this patient?**

**A. Glycosuria**

**B. Hematuria**

**C. Ketonuria**

**D. proteinuria**

**69. A 25 year-old woman GII PI presents at 35 weeks gestation for a routine ANC visit at the health center. Her blood pressure is 150/100mmhg . and her urine protein is +3. Physical examination is otherwise unremarkable. She reports headache, but no upper right quadrant pain or blurred vision . which of the following is the next appropriate management?**

**A. schedule for next visit**

**B. provide Mgso<sub>2</sub> and refer**

**C. start induction**

**D. give antihypertensive**

**70. A 38 year-old primigravida woman presents for a routine visit at 39 weeks of gestation. Her BP is persistently 140/90mmhg and her urine protein is +2. Physical examination is otherwise unremarkable, and she is completely asymptomatic. Her cervix is 2cm dilated and 90% effaced with vertex presentation at station 0 which one is the most appropriate management option?**

**A. Admission for observation**

**B. Immediate cesarean section**

**C. outpatient follow up till 40 weeks**

**D. start induction of labor**

**71.a A 34 year-old primigravida lady comes with cessation of fetal movement of four days duration at gestational age of 35 weeks. She is otherwise asymptomatic and has no medical problems. Examination shows stable vital signs, symphysio-fundal height of 34cms, no uterine contraction and absent fetal heart beat . what is the best initial step in the management of this patient?**

**A. Consider Termination**

**B. Do coagulation profile**

**C. prophylactic antibiotics**

**D. Ultrasound for confirmation**

**72. A 17 year-old girl has noted cyclic stamp, lower abdominal pain that begins on the day of her menstrual flow and lasts for 2-3 days also her menarche worsening in the last 2 years. Her periods are regular and pelvic examination is normal. Which of the following is the most likely diagnosis?**

**A. Adenomyosis**

**B. Dysmenorrhea**

**C. Endometriosis**

**D. uterine myoma**

**73. A 21 year-old GII PI mother at 24 weeks gestation complains of severe vulvar itching and cotton cheese like discharge. A wetmount slide shows no clue cells but hyphae. Which of the following is most likely diagnosis?**

**A. Bacteria vaginosis**

**B. Candida albicans**

**C. Chlamydia trachomatis**

**D. trichonomas vaginalis**

**74. A 29 year-old woman presents with vaginal pruritus associated with white curd like discharge. On examination there is erythematous macular lesion but no vesicles. Investigation shows no clue cell, negative with test. Which of the following is the treatment of choice for this case?**

**A. Azithromycin**

**B. Fluconazole**

**C. Metronidazole**

**D. Oral acyclovir**

**75. A 26 year-old GII PI woman at 26weeks of gestation came for ANC followup. She hasn't appreciated fetal movement yet. she had one episode of bleeding a month back and fetal heart beat was absent. which of the following is the most likely diagnosis?**

**A. incomplete abortion**

**B. inevitable abortion**

**C. missed abortion**

**D. threatened abortion**

**76. A 42 year-old healthy nulliparous woman has a firm, irregular shaped, 12 week- sized midline abdominal mass. Which of the following is the best initial step in the evaluation of this patient?**

**A. Abdominal x-ray**

**B CT- Scan**

**C. MRI**

**D. Ultrasound**

**77. Which of the following is NOT a component of active 3<sup>rd</sup> stage management?**

**A. controlled cord traction**

**B. Nipple stimulation**

**C. oxytocin provision**

**D. Uterine massage**

**78. Which of the following isn't a component of active 3<sup>rd</sup> stage management?**

**A. controlled cord traction**

**B. Nipple stimulation**

**C. Uterine massage**

**D. oxyticin provision**

**79. Which of the following is a component of prerequisites for forceps application?**

**A. Cervix more than eight centimeters**

**B. Gestational age beyond 37 weeks**

**C. Ruptured membrane**

**D. vertex presentation**

**80..in the active management of 3<sup>rd</sup> stage of labor, placenta failed to be delivered by controlled cord traction. Which one is the next best management?**

**A. pitocin iv drip**

**B.Fundal pressure**

**C.Manual removal of placenta**

**D. wait for spontaneous expulsion**

**81. A 32 year-old nulligravida woman came to gynecology OPD with a complaint of heavy menstrual flow of one year duration. She also gave history of constipation and pelvic heaviness and failure to conceive of four years duration. On examination the is central firm abdominal pelvic mass of 12 weeks size otherwise normal. Which of the following is this case?**

- A. Endometrial hyperplasia**
- B. Myoma**
- C. Ovarian tumor**
- D. uterine sarcoma**

**82. A PIII mother who gave birth to an alive healthy term neonte vaginally before an hour was found to be restless and pale. On physical examination BP=60/40mmhg, PR=120/min RR=24/min, T=36<sup>0</sup>C; Uterus 22 weeks and soft and active vaginal bleeding . the most appropriate initial step in the management of this patient is:**

- A . Administer crystalloids**
- B. administer intra-nasal oxygen**
- C. blood transfusion**
- D. call help**

**83. A 35 year-old GVPV mother at gestational age of weeks presented to gynecology OPD with a complaint of vaginal bleeding of four days duration. She had excessive vomiting and big for date uterus which of the following is the most likely diagnosis?**

- A. Cervical polyp**
- B. Ectopic pregnancy**
- C. incomplete abortion**
- D. molar pregnancy**

**84. Which of the following contraceptive methods is the most effective?**

- A. Combined oral contraceptive pills**
- B. femal condom**
- C. Intrauterine contraceptive device**

**D. Medroxy progesterone**

**85. A 17 year-old student came to a clinic seeking pregnancy termination service at gestational age of 10 weeks. She gave a history of sexual assault. Which of the following is the best method of termination of pregnancy for this case?**

**A. Dilatation and curettage**

**B. intramuscular methotrexate**

**C. manual vacuum aspiration**

**D. Mifepristone and misoprostol**

**86. A 22 year-old primigravida who came for a routine ANC visit tested positive for HIV. When should a health officer initiate ARV drugs for this woman?**

**A. AS soon as possible**

**B. At 20 weeks of gestation**

**C. At 28 weeks of gestation**

**D. At 36 weeks of gestation**

**87. A 20 year-old primigravida at 38 weeks was admitted in second stage of labor and stayed for 90 minutes. Fetal heart beat is raging between 80-90/min vertex presentation with the occiput direct anterior, station +3 grade one molding. No cuput and grade- II meconium stained fluid. Which one of the following is the next best in management?**

**A. Expedite delivery by vacuum right away**

**B. Refer her for cesarean section**

**C. Re-position her into the left lateral position**

**D. secure iv line and await spontaneous vaginal delivery**

**88. A para III mother who gave birth to an alive male neonate at home before an hour was brought to a hospital because of**

restlessness and protrusion of mass per vaginum. She is acutely sick looking, BP is not recordable, PR=132/min, and feeble RR=20/min. T=36°C. a globular fleshy mass with minimal bleeding protruding through the introitus. What is the most appropriate initial step in management of this patient?

- A. intra nasal oxygen 6 L/min
- B. Reposition the uterus
- C. Resuscitation with iv fluid
- D. start oxytocin drip

89. A 6 year old diabetic child admitted to a Hospital with a diagnosis of severe diabetic acidosis has been receiving normal saline and insulin for the last three hours. Which electrolyte abnormality is expected?

- A. hypomagnesemia
- B. Hyponatremia
- C. Hypokalemia
- D. Hypophosphatemia

90. A 9 month old boy is brought to a health center in a unresponsive state after repeated episodes of vomiting and diarrhea. Physical examination showed a rapid and feeble pulse, skin pinch returns very slowly. Multiple attempts to secure an iv line have failed. What is the next best step in the management of this infant?

- A. Do blood film and determine blood glucose
- B. Give ORS through NG-tube and refer
- C. obtain blood sample for complete blood count
- D. perform a lumbar puncture

**91. A 6 month old infant presented with cough and fast breathing of three days duration. Further inquiry reveals history of repeated chest infections, interruption during breast feeding and poor weight gain. Precardial examination reveals active precordium and pansystolic murmur at left lower sternal border. Which of the following most likely explains those findings?**

- A. Atrial septal defect**
- B. ventricular septal defect**
- C. patent ductus arteriosus**
- D. Tetralogy of fallot**

**92. An 8- year-old boy known cardiac patient developed low-grade fever, fatigue myalgia in the last two weeks. On examination he was found to have developed a new murmur, petechial rashes and mild splenomegaly, which one is the most likely diagnosis?**

- A. Acute rheumatic fever**
- B. Dilated cardiomyopathy**
- C. infective endocarditis**
- D. Rheumatic heart disease**

**93. which of the following vaccines should NOT be given in an infant with severe immune suppression?**

- A. Hepatitis B Vaccine**
- B. Measles Vaccine**
- C. Meningococcal vaccine**

**94. A 6 week old infant presented with high grade fever three hours after he took the first of his scheduled vaccines. On examination he is febrile and irritable otherwise there are no**

**findings in other systems. What is the most appropriate management for this infant?**

- A. Admit and observe**
- B. Initiat oral antibiotics**
- C. investigate for infection**
- D. Paracetamol suppository**

**95. A 14 –month old male child is brought to the ER with watery diarrhea of 24 hours duration. Physical examination reveals irritable infant, weight= 12 kg. sunken eye balls and eagerness to drink. What is the correct type and volume of fluid for this infant?**

- A. Ringer's lactate,300ml**
- B.0.4% Normal saline, 700ml**
- C. oral rehydration solution, 900ml**
- D. Ringer's lactate, 500ml then ORS 500ml**

**96. which of the following clinical conditions that appear after pertussis vaccination is an absolute contraindication to subsequent administration of the vaccine?**

- A. Fever and convulsion whith in 3 days**
- B. Immediate anaphylactic reaction**
- C. inconsolable cry longer than 2 hours**
- D. shock-like state after 2 days**

**97. you are evaluating an HIV exposed neonate immediately after delivery. The mather had proper ANC follow up and was managed according PMTCToption B+. which of the following is an appropriate recommendation for this new born?**

- A. Avoid breast feeding**
- B. Cotrimoxazole prophylaxis in the first weeks of life**

**C. DNA-PCR test within 48 hours**

**D. nevirapine for 6 weeks**

**98. A 3 month old infant has rhinorrhea caugh and noisy breath. Physical examination reveals RR=72/min PR=142/min and T=37.8<sup>0</sup>c he is irritable and has diffuse wheezes and chest retraction. Whate is the most likely diagnosis?**

**A. Bronchiolitis**

**B. Gastro-esophageal reflux disease**

**D. severe pneumonia**

**99. which of the following best describe clinical features of Guillain-Barre syndrome?**

**A, Ascending symmetric paralysis without sensory change**

**B. Asymmetric paralysis and sensory change**

**C. Symmetric paralysis and central nerve palsy**

**D. symmetric paralysis and sensory change**

**100. A 5 year –old boy with HIV infection is receiving Zidovudine, Lamivudine and Nevirapine. He presents with progressive difficulty of breathing for three weeks. He has no history of fever. on examination PR=108/min RR=72/min T=37<sup>0</sup>c and oxygen saturation is 70%.chest radiograph reveals bilateral diffuse alveolar disease. What is the most likely diagnosis?**

**A. Heart failure**

**B. pneumocystis pneumonia**

**C. pulmonary tuberculosis**

**D. severe bacterial pneumonia**

**101. The recommended time to start foletes supplementation for a matherwho previously had a baby boy with**

**myelomeningocele to prevent its recurrence in the next pregnancy is**

- A. At the first missed period**
- B. End of first trimester**
- C. one month before conception**
- D. third trimester**

**102. which of the following CSF profile is expected in a child with meningoencephalitis?**

- A. Leukocytosis with neutrophil predominance**
- B. Normal protein level**
- C. normal or slightly reduced glucose level**
- D. organism on Gram stain**

**103. which of the following is the correct percentage of dextrose to treat neonatal hypoglycemia?**

- A. 5%**
- B. 10%**
- C. 40%**
- D. 50%**

**104. what is the compression to ventilation ratio in resuscitating a newborn?**

- A. 3:1**
- B. 5:1**
- C. 15:1**
- D. 30:1**

**105. A 36 month-old male infant presented with high grade fever and vomiting of two days duration. He has right costovertebral angle tenderness. White cell count showed leukocytosis with left**

shift and blood film is negative for hemoparasites. What is the most likely diagnosis in this child?

- A. Pyelonephritis
- B. Hepatitis
- C. Appendicitis
- D. Lobar pneumonia

106. A 5 year-old boy sustained a penetrating injury to his right leg. Two weeks later he developed swelling of the injured leg and high fever. X-ray of the affected limb showed features of acute osteomyelitis. What is the most likely etiology in the above case scenario?

- A. *Pseudomonas aeruginosa*
- B. *Klebsiella pneumonia*
- C. *Streptococcus pyogenes*
- D. *E. coli*

107. A 2 year-old child has delay in all developmental milestones for his age. On examination he has macrocephaly, severe stunting, protruded tongue. Umbilical hernia and coarse facial features. The best initial test is to determine serum level of;

- A. cortisol
- B. Growth Hormone
- C. Thyroid stimulating hormone
- D. Vitamin D

108. Which one of the following drugs is an appropriate antidote for malathion poisoning?

- A. Atropine
- B. Flumazenil
- C. N-acetylcysteine

**D. Naloxone**

**109. A 3 year-old boy presented with a complaint of intense pruritis worse at night. On examination there are vesicular and popular eruptions at the interdigital spaces. Which one is the best treatment for thi child? Dx scabies**

**A. Acyclovir**

**B. Erythromycin**

**C. Mebendazole**

**D. Permethrin**

**110. A 2-hours term neonate is delivered by elective C/C for placenta previa. On physical examination PR=170/min, RR=72/min and oxygen saturation of 80%. What is the most likely cause of respiratory distress in this neonate?**

**A. Congenital pneumonia**

**B. Perinatal Asphyxia**

**C. Respiratory distress syndrome**

**D. Transient tachypnea of the new born**

**111. A 2 year- old child presented with difficulty of breathing of two hours. He had a breaking cough and hoarseness of voice for the last two days. On physical exam he is agitated, has audible stridor at rest and lower chest indrawing. What is the most likely diagnosis ?**

**A. Croup syndrome**

**B. Foreign body aspiration**

**C. Pertussis**

**D. Severe pneumonia**

**112. A 7 year old female patient who is on follow-up for bronchial asthma presented with shortness of breath of three**

hours. On physical examination she is conscious, able to talk in complete sentence, RR=62/min and PR=110/min. chest examination reveals diffuse expiratory wheeze. What is the initial best step in the management of this patient?

- A. Aminophylline
- B. Magnesium sulphate
- C. Oral prednisolone Epinephrine
- D. salbutamol puff

113. A 3 year-old boy is brought to the ER with high grade fever and shortness of breath of two days. On physical examination RR=62/min, T= 39°C, lower chest indrawing with bilateral basal coarse creptations. The chest X-ray showed left basal lung pneumatocele. Which of the following is the most likely etiology?

- A. Haemophilus influenza
- B. Staphylococcus aureus
- C. Streptococcus pneumonia
- D. Streptococcus pyogenes

114. A 13- year old street boy presented to the ER with high grade fever of 5 days. He has epistaxis, fatigue and headache. On physical examination heart rate 120/min, respiratory rate 32/min, T=39°C and has petechial rash on trunk. Which of the following is the best initial test for diagnosis?

- A. Blood film
- B. complete blood count

**C. CSF analysis**

**D. widal test**

**115. An 18-months old child with sever acute malnutrition is noticed to have diffuse left thigh swelling which is erythematous, hot and tender to touch. It is also fluctuating. which one of the following is the most likely diagnosis?**

**A. Cellulitis**

**B. Hip arthritis**

**C. Osteomyelitis**

**D. pyomyositis**

**116. A 2 year old child has delay in developmental milestones for his age. On physical examination he has macrocephaly, severe stunting, protruded tongue, umbilical hernia and coarse facial features. Which investigation is very helpful in managing this patient?**

**A. Cortisol**

**B. Growth hormone**

**C. TSH**

**D. vitamin D**

**117. which of the following drugs is helpful for treatment of malathion poisoning?**

**A. Flumazenil**

**B. N-acetyl cysteine**

**C. Atropine**

**D. Naloxone**

**118. A health officer was attending delivery of a post term pregnancy. Upon follow-up of the lobar there was a record of fetal tachycardia and grade III meconium stained amniotic fluid. The newborn then failed to breathe spontaneously immediately after delivery. What is the most likely cause of respiratory problem in this baby?**

- A. Respiratory distress syndrome**
- B. Congenital pneumonia**
- C. Meconium aspiration syndrome**
- D. Congenital heart disease**

**119. A 5 year old child is diagnosed with Group A beta hemolytic tonsillo-pharyngitis. Which one is a non – suppurative complication of this condition?**

- A. Otitis media**
- B. Peritonsillar abscess**
- C. Retropharyngeal abscess**
- D. Rheumatic fever**

**120 . which of the following samples give the highest yield for diagnosing visceral Leishmaniasis?**

- A. Blood**
- B. Bone marrow aspirate**
- C. Lymph node aspirate**
- D. Splenic aspirate**

**121. A 4 day female neonate is brought to helth centere with failure to open her mouth and breast**

**feed. She was well in the first three days after which the mother noticed excessive cry with stiff body posturing. The mother had no ANC follow up and delivery was at home. What is the most likely diagnosis in this neonate?**

**A. Hypocalcemia**

**B. sepsis**

**C. Meningitis**

**D. Tetanus**

**122. A 3-year-old boy came to the OPD with diarrhea. Stool examination revealed trophozoite of *Giardia lamblia*. What is the first-line drug to treat this parasite?**

**A. Tinidazole**

**B. Metronidazole**

**C. Paromomycin**

**D. Albendazole**

**123. The data collected about a body mass index of physically active population looks positively skewed. Which measure of central tendency is advisable to be used for such types of data?**

**A. Median**

**B. Mean**

**C. Variance**

**D. Range**

**124. Sampling is the procedure some members of the given population are selected as representative part of the entire population. The information collected from the sample will be generalized to;**

**A. studay population**

**B. Target population**

**C. Sampling frame**

**D. study unit**

**125. A health officer wanted to conduct a research to determine the prevalence of ANC utilization of mathers in a certain district with infinite population. The prevalence of ANC utilization frome a previous study was found to be 45.7%.what is the final sample size to address his objective?**

**A. 250**

**B. 382**

**C. 402**

**D. 459**

**126. In a study on percentage saturation of bile, information was collected on the age of patients to see whether a difference existed between the two measures. Which method of data presentation is most appropriate for this data?**

**A. Box and whisker plots**

**B. Histogram**

**C. Multiple bar graph**

**D. scatter plot**

**127. which of the following statement is true regarding hypothesis testing ?**

**A. If the P- value is less than  $\alpha$ , the null hypothesis is accepted**

**B. If the P-value is less than  $\alpha$ , the null hypothesis is rejected**

**C. Null hypothesis; there is a difference between the hypothesized true values**

**D. Rejecting the null hypothesis when it is true is type II error**

**128. The number of modes of transmission has an effect on the control or eradication of infectious disease. From this prespective. Which of the following diseases is more difficult to control?**

**A. Pulmonary tuberculosis**

**B. Relapsing fever**

**C. Syphilis**

**D. Trachoma**

**129. Prevalence(new + old case) and incidence rate(new case) are widely used as measures of morbidity, which of the following statements is true about these two measures of morbidity?**

**A. both equally measure the risk of acquiring a specific disease**

**B. follow up of a group of people is necessary to estimate incidence rate**

**C. follow up of a group of people is necessary to estimate point prevalence rate**

**D. prevalence rate tells more of the risk of acquiring a specific disease**

**130. An estimated 90 people out of 100 cases of Ebola died in the recent outbreak. What does this proportion represent?**

**A. case fatality rate**

**B. cause specific mortality rate**

**C. Crude death rate**

**D. Proportionate mortality ratio**

**131. Suppose psychosis is a rare health problem in a specific country. Researchers wanted to study design is most appropriate to assess this association?**

- A. case control study**
- B. cohort study**
- C. Cross sectional study**
- D. Experimental study**

**131. A study was conducted to investigate the effect of HIV infection on mortality among people with TB in Ethiopia. Individuals with TB were recruited from hospitals and their HIV status determined. They were then followed –up over ten years to compare mortality rates in the HIV positive group. Which study design is employed?**

- A. case- control study**
- B. Cohort study**
- C. Ecological study**
- D. Randomized Controlled**

**132. which of the following micro-nutrients enhances the absorption of iron?**

- A. Calcium**
- B. Vitamin E**

**C. Vitamin C**

**D. Zink**

**133. A significant number of under-five children were diagnosed with serious neurological problems and anemia in a certain district. Researchers found out that the families of these children were strict vegetarians. Which nutritional deficiency are the children suffering from?**

**A. Folic Acid**

**B. Vitamin B12**

**C. Vitamin B3**

**D. Vitamin B6**

**134. The dietary assessment in a certain community subsist on a diet of corn meal and boiled greens. Patients coming from this area have alternate diarrhea and constipation. They have glossitis and scaly dermatitis. What is the most likely deficiency?**

**A. Pyridoxin**

**B. vitamin E**

**C. Niacin**

**D. Vitamin k**

**135. A researcher who conducted a study on malnutrition among under five children found out a high prevalence of stunting in that community. What does this finding imply?**

- A. Occasional micronutrient deficiency**
- B. High rate of genetic growth defect**
- C. Long standing nutritional problem**
- D. High rate of acute malnutrition**

**136. A young man consumed 100 gram serving of whole-wheat spaghetti containing 10 grams of protein, 40 gram of carbohydrate and 5 grams of fat. What is the estimated content in kcal?**

- A. 205**
- B. 245**
- C. 450**
- D. 500**

**137. Which of the following vector borne diseases is transmitted by sand fly?**

- A. Filariasis**
- B. Leishmaniasis**
- C. Onchocerciasis**

#### **D. Trypanosomiasis**

**138. A researcher aims to initiate a behavioral change among individuals considering the socio-cultural factors after identifying personal characteristics of the people. The best theory or model to apply is:**

- A. Health belief model**
- B. Social cognitive theory**
- C. Theory of planned behavior**
- D. Trans-theoretical model**

**139. A health extension worker divided participants of a certain kebele into different groups to discuss the district's main health problems and provided feedback on the main findings to the whole group. What type of method is implemented?**

- A. Brain storming**
- B. panel discussion**
- C. Buzz group**
- D. case**

**140. If your aim is to bring behavioral change in hygiene and sanitation among a rural population, the best theory/ model you can use for planning the change will be:**

- A. Health belief model**
- B. Trans-theoretical model**
- C. Precede-proceed model**
- D. Theory of planned behavior**

**141. In a community dialogue some people were convinced about the importance of having a public latrine and wanted to build one in their village. But they didn't have enough resources for construction. At what communication stage did the message fail?**

- A. Acceptance**
- B. Attention**
- C. Behavioural change**
- D. understanding**

**142. several cases of acute watery diarrhea among under five children was reported in a rural village. The residents of the village use water for domestic purpose from unprotected well, which was located up gradient to a public latrine facility. Which immediate measure needs to be taken to address this priority public health problem?**

- A. Fencing the well water**
- B. Treating the well water**

**C. Abandon the well**

**D. protect the well water**

**143. A communal well water source development project was planned to be developed in a rural village. What needs to be identified for the outcome of the project to be effective?**

**A. Acceptance of the water source by the community**

**B. Cost of materials for development purpose**

**C. Electrolyte composition of the water source**

**D. Level of trace minerals in the water source**

**144. In a rural village open defecation was found to be a common practice. The district health officer planned to alleviate the problem by applying an appropriate sanitation technology. Which one is an important factor to consider in selecting appropriate sanitation technology?**

**A. Availability of latest technology**

**B. Cultural beliefs and practices**

**C. Characteristics of the waste**

**D. Technical expertise available**

**145. In health center hazardous waste disposal was found to be a critical problem. Which of the following waste disposal options is the most appropriate?**

**A. Burying**

**B. Composting**

**C. incinerating**

**D. offsite Disposal**

**146. The malaria prevalence in a certain district is increasing from time. A team has been established to develop malaria epidemic control and prevention program. What priority intervention needs to be considered to effectively limit the progress of the epidemic?**

**A. Application of outdoor residual spray**

**B. Destruction of mosquito breeding sites**

**C. Enforce usage of insect repellants**

**D. Timely treatment of cases**

**147. Nowadays, Reproductive health programs?**

**A. Demands more health professionals**

**B. Reduce access to programs**

**C. Reduce missed opportunities**

**D. systematically excludes men**

**148. Some experts argue that WHO's definition for maternal mortality has limitations. What is its limitation in measuring maternal mortality ratio?**

- A. IT disregards deaths that occur after 42 days**
- B. It misses deaths during pregnancy**
- C. It misses major causes of maternal death**
- D. It misses the time most maternal deaths occur**

**149. If an organization wants to design a strategy targeting the most important direct causes of maternal mortality globally. Which cause should get the first priority?**

- A. Abortion**
- B. Hemorrhage**
- C. Hypertension**
- D. Sepsis**

**150. If a health officer wants to know risk of maternal death do to level of fertility and risk of mathernal deathper pregnancy or per birth. Which measure of maternal mortality should he/she use?**

- A. Maternal mortality ratio**
- B. Maternal mortality rate**

**C. Life time risk of maternal mortality**

**D. Proportionate maternal mortality**

**151. Of the strategies to reduce maternal mortality, which one of the following interventions is most effective in reducing the likelihood of developing pregnancy related complications?**

**A. Family planning**

**B. Antenatal care**

**C. emergency obstetric care**

**D. Nutrition intervention**

**152. Which one is the main reason for low contraceptive use in Ethiopia?**

**A. Breast feeding**

**B. Fear of side effect**

**C. Religious beliefs**

**D. Resistance from partners**

**153. Of the strategies to reduce maternal mortality, which of the following interventions is most effective in reducing the likelihood of developing pregnancy-related complications?**

**A. Antenatal care**

**B. Emergency obstetric care**

**C. family planning**

**D. Nutritional intervention**

**154. A manager of health center wants to provide BCG vaccine for infants on Fridays. Which of the following is the best management function that the manager used?**

**A. Organizing**

**B. Planning**

**C. Controlling**

**D. Implementing**

**155. A Hospital manager allocated a budget for three different activities. Which of the following is the next step in planning?**

**A. Designing strategy**

**B. Prioritizing problem**

**C. setting objective**

**D. Situationa analysis**

**156. A zonal health Department head assigned a budget for different tasks for a fiscal year. Which decisional role did he play?**

**A. Negotiation role**

**B. Dissemination role**

**C. Resource allocation role**

**D. Monitoring role**

**157. The total population of certain locality was around 25,000. Which of the following health facilities is suitable to provide health service to that community?**

**A. District Hospital**

**B. Health center**

**C. Health post**

**D. health station**

**158. A college dean who has experts in different professions as employees left them to do their activities. What leadership style did he apply?**

**A. Autocratic**

**B. Bureaucratic**

**C. Democratic**

**D. Laissez faire**

**159. There was interpersonal conflict in the same department of a certain health facility. What was the best management option that they do?**

**A. Leave the issue for those involved**

- B. Manage the conflict after a month**
- C. Switch one of them to another department**
- D. Resolve the conflict as soon as possible**

**160. A 24 year-old woman come to your office for follow up on her depression. She had a history of suicidal attempt and severe peri-partum depression in the past. Which of the following is appropriate management?**

- A. Continue to take fluoxetine**
- B. shift to another antidepressant**
- C. Shift to antipsychotic drug**
- D. start mood stabilizer**

**161. A 26 year-old woman brought by her sister to the hospital with seven months history of unusual behavior. The patient gradually develops poor self –care, isolation and sleep herself and laughs alone. On mental status examination she has blunt affect, auditory hallucination and has no insight to her illness. Which of the following is the most likely diagnosis?**

- A. Delusional disorder**
- B. Schizoaffective disorder**
- C. Schizophreniform disorder**
- D. schizophreiform disorder**

**162. A 28 year –old man diagnosed as a case of major depressive disorder with psychotic features. He is treated with haloperidol and amitriptyline. Two months later, his mood symptoms have resolved and he is no longer psychotic. Which of the following best describes the next step?**

- A. Both drugs should be tapered and discontinued**
- B. Haloperidol should be tapered and discontinued**
- C. Amitriptyline should be tapered and discontinued**
- D. Both drugs should be continued for 6 months**

**163. A 53 year-old man comes to the Hospital complains of poor appetite, loss of pleasure in daily activities, sleeplessness, and decreased libido for 3 weeks. He is contemplating retiring, expressing concern that his poor performance in his job. On mental status examination he shows empty feeling and makes two errors in subtraction serial 3's from 20. Which of the following is the most likely diagnosis?**

- A. Bipolar disorder**
- B. Cyclothymia**
- C. Dysthymic disorder**
- D. Major depressive disorder**

**164. A 35 year-old man diagnosed to have Major depressive disorder was given 75 mg amitriptyline at bed time. On the third day he starts hearing voices which tell him that "he is bad person ". On mental status examination he feels sad and has auditory hallucinations. Which of the following is the most appropriate treatment?**

**A. Diazepam**

**B. Fluoxetine**

**C. Haloperidol**

**D. Sodium valproate**

**165. a 55 year old alcoholic is brought to the Emergency Room with a problem of irrelevant talking . Three days ago he is quit alcohol after he was counseled. On mental state examination he is disoriented to time, place and person. He has visual hallucination and agitation. There is no history of head injury or medical illness. Which of the following is the most likely diagnosis?**

**A. Alcohol induced psychosis**

**B. Alcohol intoxication**

**C. Delirium tremens**

**D. Wernicke's encephalopathy**

**166. A 55 year-old man comes to Emergency Room with acute retention of urine for 12 hours. He was suffering from depression and taking drugs for the last ten days. On examination there is distended bladder. No other physical findings. Which drug is the most likely culprit?**

**A. Amitriptyline**

**B. Fluoxetine**

**C. paroxetine**

**D. Sertraline**

**167. A patient comes to your clinic with a complaint of hearing difficulty of one month duration. There is only a set of tuning forks for hearing assessment at the clinic. Which tuning fork should be used for the assessment?**

**A. 256 Hz**

**B. 512 Hz**

**C. 102 Hz**

**D. 204 Hz**

**168. A patient comes demanding for ear wash. Which of the following would be a contraindication for ear wash?**

**A. Beads**

**B. Chronic otitis media**

**C. Cockroach**

**D. Impacted wax**

**169. A 37 year-old female presented with a cockroach in the right ear canal. She is so much in discomfort because of the movement of the insert. On examination ,the right ear has big dry central perforation of the tympanic membrane. What is the management?**

**A. Add 3% hydrogen per oxide to kill the insect**

**B. Add ethyl alcohol to kill the insect**

**C. Irrigation with warm water**

**D. Take the insect out with a small artery while alive**

**170. A 22 year-old male patient has left sided ear pain with no discharge. He has no history of trauma and physical examination revealed tenderness over the tragus. What is the most likely diagnosis?**

**A. Acute otitis externa**

**B. Acute otitis media**

**C. Cerumen impaction**

**D. chronic otitis media**

**171. During a school survey on refraction, a 6 year-old girl was found to have a refractive error of (+3) on the left eye while**

**the right eye was (+6). If the refractive error is not corrected by spectacles, which of the following is LEAST expected to happen?**

- A. Amblyopia**
- B. Floaters**
- C. poor school performance**
- D. Strabismus**

**172. what is the leading cause of visual impairment globally?**

- A. Cataract**
- B. Diabetic retinopathy**
- C. Glaucoma**
- D. Uncorrected refractive error**

**173. A 51 year-old man came to your OPD complaining he can't read contact numbers from his mobile phone since 6 months. He used to be able to read by holding the phone further away. He has no other ocular complaints. He had 6/6 distance vision on both eyes. What is the most likely diagnosis?**

- A. Age-related macular degeneration**
- B. Astigmatism**

**C. Cataract**

**D. Presbyopia**

**Answer of the above question**

**1.B**

**2.B**

**12.C22.B32.B42.B52.D62.D72.B82.D92.C102.A112.D122.B**

**3.C**

**13.D23.C33.B43.A53.A63.A73.B83.D93.B103.B113.A123.A**

**4.C**

**14.B24.C34.B44.D54.D64.B74.B84.A94.C104.D114.A124.B**

**5.C**

**15.C25.C35.A45.B55.B65.A75.C85.D95.C105.A115.B125.--**

**6.A**

**16.C26.A36.B46.D56.A66.D76.D86.A96.B106.C116.C126.--**

**7.C**

**17.A27.A37.C47.B57.A67.C77.B87.B97.D107.B117.C127.C**

**8.D**

**18.B28.A38.A48.C58.A68.D78.B88.C98.D108.A118.D128.B**

**9.D**

**19.D29.A39.D49.C59.B69.C79.D89.C99.D109.D119.B129.C**

**10.D 20.D30.B40.D50.A60.C70.A80.A90.B100.B110.—  
120.D130.A**

**131.A141.A151.B161.C171.--**

**132.C142.A152.D162.D172.A**

**133.B143.A153.A163.A173.B**

**134.C144.B154.A164.C**

**135.C145.C155.A165.--**

**136.B146.A156.B166.--**

**137.B147.A157.B167--**

**138.B148.A158.C168.B**

**139.B149159169**

**140.A150160170**