

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

HUMAN NUTRITION



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority and is essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and to the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, Leadership and Management, Ethics, and Health Promotion as outlined in the Information Booklet 2025.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey—and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Human Nutrition

1. A 50-year-old man was presented to an OPD with complaints of shortness of breath, chronic chest pain, weakness, and lack of appetite. He also reported palpitation while he used soft drinks or fatty foods. His dietary history revealed with a diet of high fatty food, burger, alcohol and soft drinks regularly. His BMI, MUAC, waist to hip ratio, and skin fold thickness were 29kg/m², 30cm, 0.96, and 0.2mm respectively. His biochemical test showed that HDL Cholesterol level of 30mg/dL. Finally, he was diagnosed with central obesity.

What is the most appropriate anthropometric measurement used to reach the diagnosis in this case?

- A. Body Mass Index
- B. Mid Upper Arm Circumference
- C. Waist to Hip Ratio
- D. Skin Fold Thickness

ANSWER: C

2. A 22-year-old male patient was presented to a clinic with a complaint of shortness of breath that worsened during heavy work and long journeys on foot. On dietary assessment, he was found to have two bottles of beer per day for four days of the week. Besides, he reported that he had been regularly consuming fatty foods. On a biochemical test, his HDL and LDL cholesterol levels were 30 mg/dL and 170 mg/dL, respectively. Then, before planning a management, the dietitian put a diagnosis for the problem.

What is the most likely diagnosis of metabolic problem in this scenario?

- A. Hyperlipidemia
- B. Hypercholesterolemia
- C. Atherosclerosis
- D. Hypertriglyceridemia

ANSWER: A

3. A 41-year-old female patient, who was operated on for gastric ulceration four months ago, came to the clinic with a chief complaint of headache and dizziness of four days duration. She had a history of insomnia, heavy duty, and lack of appetite, and she took a large amount of fluid. Her blood pressure was 100/70 mm Hg. Biochemical tests showed a sodium level of 120 mmol/L, a potassium level of 4.0 mmol/L, a calcium level of 3.2 mmol/L, and a hemoglobin level of 12 g/dL.

What is the most likely metabolic problem in this case?

- A. Fluid imbalance
- B. Electrolyte imbalance
- C. Hypertension
- D. Diabetes mellitus

ANSWER: B

4. A 27-year-old male obese patient came to a clinic for nutritional management of his weight. He had history of using fatty foods and alcohol regularly. His dietary history revealed a diet high in simple sugars, fried foods, and low in fruits and vegetables. He felt palpitation on fast diet and soft drink use. His waist-to-hip circumference ratio was 0.95 cm. On biochemical tests, he had signs of hypercholesterolemia.

What is the most likely healthy diet recommendation to be applied in this case?

- A. Decrease intake of fried foods
- B. Increase intake of simple sugars
- C. Decrease intake of fruits
- D. Decrease intake of vegetables

ANSWER: A

5. A five-month-old male infant, who showed a decreased breast milk intake for the past one week, was referred to a health center from a health extension worker. His caregiver reported that the child had not been fed well and was not gaining weight. The child appeared to be small for his age, and the nutritionist wanted to determine his nutritional status.

What is the most appropriate anthropometric index measurement to be used in this case?

- A. Weight for height
- B. Height for age
- C. Weight for age
- D. Weight for length

ANSWER: D

6. A two-year-old male child is presented with watery diarrhea. His caregiver reported that the child has lost weight. On physical examination the child has bilateral pitting edema on his leg and multiple patchy skins. The health professional suspects edematous malnutrition as a differential diagnosis and wants to assess the nutritional status of the child.

What is the most appropriate anthropometric measurement to be used in this case?

- A. Waist circumference
- B. Mid upper arm circumference
- C. Hip circumference
- D. Head circumference

ANSWER: B

7. A four-year-old female child was brought to a health center with complaints of loss of appetite and discoloration of skin. Her mother reports that the child has decreased food intake. On nutritional assessment, the nutritionist has confirmed that there is dependent bilateral pitting edema in both feet. In addition, lower legs, hands and lower arms are swollen.

What is the most likely edema grade for this case?

- A. Grade +
- B. Grade ++
- C. Grade +++
- D. Grade ++++

ANSWER: B

8. A two-year-old child was brought to a clinic with a complaint of poor growth. The care giver reports that the child has lost his appetite and lost interest in everything. On assessment, the child has a moon face appearance, has multiple patchy skin areas in his abdomen, and some fissures in his groin area. What is the most important sign needed to reach at a decision of the diagnosis?

A. Sunken eye
B. Pale conjunctiva
C. Bilateral pitting edema
D. Muscle wasting

ANSWER: C

9. A seven-month- old male child was brought to a hospital for his immunization schedule. His vital signs are stable, and he has received his immunization as scheduled. However, when the nutritionist performs a physical examination, the child has stunting.

What is the most appropriate anthropometric measurement index used in this case?

A. Length for age
B. Weight for age
C. Weight for height
D. Head circumference

ANSWER: A

10. A three-year-old male child, who had not been fed well for the past four months, was brought to a clinic. His mother reported that he had poor weight gain. He was exclusively breastfed for two months. On assessment his weight was 10 kg and his length was 85cm. The health professional wanted to assess presence of chronic malnutrition in this child.

What is the best anthropometric measurement index to be used in this case?

A. Weight for age
B. Height for age
C. Weight for Height
D. Head circumference

ANSWER: B

11. A four-year-old female child was presented to a pediatric clinic with a complaint of low appetite of two weeks duration. Her mother said that she did not gain weight for more than a year. On anthropometric assessment, the z-scores for her weight-for-height, height-for-age, and weight-for-age were -2.0 SD, -2.6 SD, and -1.9 SD, respectively.

What is the most appropriate anthropometric interpretation in this case?

A. Stunting
B. Wasting
C. Normal
D. Overweight

ANSWER: A

12. A two-year-old child was brought to a nutrition clinic with complaint of lack of weight gain. On dietary assessment, it was found out that the child had been fed on cow and formula milk starting from the neonatal period. Her weight-for-age measurement was -1.6 SD.

What is the most appropriate interpretation for this scenario?

- A. Mild wasting
- B. Mild stunting
- C. Normal
- D. Mild underweight

ANSWER: D

13. A 13-month-old child was brought to nutrition OPD by the mother, for she was concerned about the recent weight loss and loss of appetite of her child. The child was almost entirely dependent on her breast milk and rejected new food items. On nutritional examination, MUAC was 12.8cm, and weight-for-height was -1.96.

What is the most appropriate interpretation for this scenario?

- A. Mild wasting
- B. Moderate wasting
- C. Mild stunting
- D. Moderate stunting

ANSWER: A

14. A 19-month-old girl was brought to nutrition OPD by her mother with a complaint of failure to thrive. The child was alert and had good appetite; however, she appeared thin. On examination, the nutrition professional confirmed that the child had not edema, weight-for-height of a Z-score of -1.52, MUAC of 11.7cm, and hemoglobin of 12.4mg/dl. Finally, the nutrition professional diagnosed MAM and planned to treat the child as an outpatient department.

What was the most appropriate admission criterion fulfilled in this scenario?

- A. WFH
- B. MUAC
- C. Hemoglobin
- D. Age

ANSWER: B

15. A three-year-old child was brought to nutrition OPD with a complaint of recent weight loss. The child was alert and had good appetite; however, she appeared thin. On examination, the nutrition professional confirmed that the child had not edema, weight-for-height (WFH) of a Z-score of -2.52, MUAC of 12.7cm, and FBS of 110 gm/dl. She was diagnosed MAM and planned to treat the child in the outpatient department.

What were the most important admission criteria in this case?

- A. MUAC and FBS
- B. MUAC and no edema
- C. WFH and no edema
- D. WFH and FBS

ANSWER: C

16. An 11-month-old child was brought to nutrition OPD by the mother, who was concerned about the recent weight loss and loss of appetite. The child was almost entirely dependent on her breast milk and rejected new food items. On nutritional examination, MUAC was 11.8cm, and weight-for-height was a z-score of -2.5.

What is the most appropriate interpretation for this scenario?

- A. Overweight
- B. Underweight
- C. Stunting
- D. Wasting

ANSWER: D

17. A 16-month-old child was brought to a hospital with complaints of intermittent diarrhea, and loss of weight of one week duration. On clinical assessment, the child appeared to be sick but was eager to drink. There was no pertinent finding from urine and stool examination. On anthropometric assessment, MUAC was 12cm, and a z-score for weight-for height was -2.75.

What is the most appropriate interpretation in this case?

- A. Complicated severe acute malnutrition
- B. Uncomplicated severe acute malnutrition
- C. Moderate acute malnutrition
- D. Moderate chronic malnutrition

ANSWER: C

18. An 18-month-old child came to a nutritional therapy clinic due to lack of appetite for more than two weeks. On clinical assessment the dietitian found that the child had significant body fat loss. On anthropometric assessment, his weight-for-age reading was a Z score of -3.6 SD. What is the most likely interpretation of this case?

- A. Severely underweight
- B. Severely stunting
- C. Mildly underweight
- D. Mildly stunting

ANSWER: A

19. A 13-month-old female child was brought to a nutritional therapy clinic with weakness and irritation for more than two weeks. On clinical assessment, the nutritionist found that she had bilateral pitting edema and her weight-for-length measurement was a Z score of -3.8SD.

What is the most possible interpretation of this scenario?

- A. Mildly stunting
- B. Severely Wasting
- C. Severely Stunting
- D. Mildly Wasting

ANSWER: B

20. A nine-month-old female child was brought to a pediatric clinic with complaint of fever. On clinical assessment, the dietitian found that she could not swallow any form of food including breast milk. Her body temperature was 39.1 degree Celsius and MUAC measurement was 10.5cm. Finally, the dietitian admitted the child to stabilizing center.

What is the most important admission criterion for this case?

- A. MUAC result
- B. Good appetite test
- C. High grade fever
- D. Pitting edema

ANSWER: C

21. An eight-month-old child was brought to a pediatric clinic with sign and symptoms of high desire to eat, old man face, and significant loss of body fat. His body temperature was 35 degrees Celsius. He also had a MUAC measurement reading of 10.5cm.

What is the most likely nutritional problem for this case?

- A. Non-edematous
- B. Edematous
- C. Sever edematous
- D. Mild edematous

ANSWER: A

22. A six-month-old child was brought to a pediatric clinic by her mother with complaints of lack of appetite and loss of desire to play. The dietitian conducted a physical examination and found that the child was irritable, had brown hair color, and had bilateral pitting edema. The child also had a MUAC measurement reading of 13 cm.

What is the most likely nutritional problem in this case?

- A. Marasmus
- B. Kwashiorkor
- C. Marasmic-kwashiorkor
- D. Stunting

ANSWER: B

23. A nine-month-old female child has been presented to a clinic with complaints of watery diarrhea and sunken eyeballs. The mother reports that the child has poor food intake and hasn't been breastfed until the time of examination. On physical examination her MUAC is 11.2 cm, axillary temperature is 36 degrees centigrade, and respiratory rate is 28 breaths per minute. She has no signs of shock, but her skin pinch goes slowly, and she has failed the appetite test.

What is the most appropriate management program option for this child?

- A. Targeted supplementary feeding program
- B. Blanket supplementary feeding program
- C. Outpatient therapeutic program
- D. Stabilization center program

ANSWER: D

24. A two-year-old male child was brought to a clinic for his routine checkup. His mother has reported that he has a delayed developmental growth. On physical examination, he has mid upper arm circumference value of 10.9cm, and edema grade ++. His rectal temperature is 37.1 degree Celsius; he has no sign of dehydration and no other medical complication and has passed the appetite test.

What is the most appropriate management program option for this child?

- A. Outpatient therapeutic program
- B. In-patient therapeutic program
- C. Targeted supplementary program
- D. Blanket supplementary program

ANSWER: B

25. A three-year-old child was brought to a clinic with poor weight gain. On assessment, her mid-upper arm circumference value is 11.2 cm, her axillary temperature is 36.91 degrees Celsius, her blood sugar level is 51 mg/dl, and her hemoglobin level is 9 g/dl. She had no signs of shock, diarrhea, vomiting, or eye problems. Finally, the child was admitted to the stabilization center.

What is the prior most appropriate treatment for this child?

- A. Amoxicillin for 7 days
- B. 50 ml bolus of 10 percent glucose
- C. 5ml/kg of Resomal
- D. Blood transfusion

ANSWER: B

26. A two-year-old female child has been brought by her mother to a clinic with a complaint of poor feeding practice. On assessment, weight for height is a Z score of less than -3 SD, but she has no hypoglycemia, no hypothermia, and no sign of shock or any other complication. The nutritionist has provided her weekly ready-to-use therapeutic feeding ration.

What is the most appropriate additional treatment that can be given for the child in this case?

- A. Give zinc
- B. Give ReSoMal
- C. Give iron
- D. Give antibiotic

ANSWER: D

27. A twenty-six-month-old male child has been admitted to outpatient care with a MUAC of 11.9 cm and a weight of 11.5 kg. He has grade ++ bilateral pitting edema at admission. He has begun his treatment with ready to use therapeutic feeding, and by the third week, he weighs 11 kg and edema is reduced to grade +.

What is the most appropriate next step for this child?

- A. Continue his treatment at OTP
- B. Refer him to stabilization center
- C. Change his feeding from RUTF to RUSF
- D. Discharge the child from treatment

ANSWER: B

28. An eight-month-male child was referred from a health post to a primary hospital for further investigation. On examination, his mid upper arm circumference value was 11.8 cm; his vital signs were stable with no edema or sign of complication. The nutritionist then started his management by giving his biweekly ready to use supplementary feeding.

What is the most likely routine medication the nutritionist should give on admission to the child?

- A. Antibiotic
- B. Iron supplementation
- C. Vitamin A supplementation
- D. Deworming

ANSWER: D

29. A mother brought her seventeen-month-old female child to a health post for immunization. On assessment, the child's axillary temperature is 36.9 and mid upper arm circumference is 11.5 cm. The child has no vomiting, no edema and no other medical complication.

What is the most likely next initial action needed for this child?

- A. Admit her to targeted supplementary feeding
- B. Admit her to blanket supplementary feeding
- C. Admit her to outpatient therapeutic feeding
- D. Admit her to stabilization center

ANSWER: A

30. A mother has come to take a ready to use supplementary feeding ration after her two-year-old child was diagnosed with moderate acute malnutrition. The nutritionist has given her counted ration and wants to counsel her to prevent reoccurrence of the case.

What is the most likely key message that the nutritionist should inform the mother to prevent relapse?

- A. To avoid sharing of the ration with the other family members
- B. To stop breastfeeding until his ration is finished
- C. To deny offering water when the child takes the RUSF
- D. To mix RUSF with other foods to make it thick

ANSWER: A

31. A fourteen-month-old female child was admitted to a stabilization center in a primary hospital for complicated severe acute malnutrition after she failed the appetite test. The child was in stabilization phase and had begun her treatment with F-75 therapeutic milk feeding and the broad-spectrum antibiotics. The child was at risk of developing hypothermia as a complication.

What is the most appropriate measure to prevent hypothermia in this case?

- A. Encouraging skin to skin contact with the mother
- B. Stopping feeding the child at night
- C. Keeping the environment cold to prevent overheating
- D. Oral fluid resuscitation

ANSWER: A

32. A three-year-old child was admitted to the targeted supplementary feeding program based on his mid-upper arm circumference measurement two months ago. Now all the necessary investigations were done, and he was clinically well and alert with no bilateral pitting edema and no medical complications. Finally, the nutrition professional wanted to discharge him from the treatment.

What is the most appropriate discharging criterion for this child?

- A. If his MUAC is greater than or equal to 12.5 for two consecutive visits
- B. If his weight for height is less than or equal to -2 for two consecutive visits
- C. If he gains weight for 3 successive days
- D. If there is supplementary program available

ANSWER: A

33. A two-year-old child was brought to a nearby clinic by his mother for failure of thriving. The mother reported to the nutritionist that she had not breast feed him since birth. The child was dependent of cow milk. The child had also diarrhea and loss of appetite. On physical assessment, the child had pale palm which crosses the palmar creases. Laboratory investigation revealed that the child had iron deficiency anemia.

What will be the most likely sign for this child?

- A. Pale palm
- B. Loss of appetite
- C. Diarrhea
- D. Failure to thrive

ANSWER: A

34. A 40-year-old man came to a private clinic for a nutritional advice due to attention deficit, poor work productivity and poor athletic performance. Biochemical test for iron showed that the adult was anemic. The nutrition expert wanted to assess his energy and nutrient intake by determining how often he consumed iron rich foods.

What is the most likely dietary assessment technique in this case?

- A. 24-hour recall method
- B. Estimated record method
- C. Diet history method
- D. Food frequency questionnaire

ANSWER: D

35. A five-year-old child who has never been supplemented with vitamin A, presented to a hospital with poor attention span, behavioral disturbance, inability to see in dim light, and bleeding gums. The child has also history of recurrent infections. On laboratory investigation, the child is having vitamin A deficiency.

What is the most likely symptom for this deficiency?

- A. Behavioral disturbance
- B. Poor attention span
- C. bleeding gums
- D. Inability to see in dim light

ANSWER: D

36. A 45-year-old male is presented to a health center with complaint of poor vision. He has undergone surgery for a bile stone. Dietary history revealed that the man has been eating food sources like carrot, milk, cheese and butter frequently. On physical examination, there is corneal scarring and ulcerations. The laboratory investigations suggested that there is steatorrhea and vitamin A deficiency.

What is the most likely risk factor for this problem?

- A. Surgery
- B. Inadequate intake
- C. Geographical location
- D. Fat malabsorption

ANSWER: D

37. A mother of an eight-year-old child living near a high-altitude mountain brought her child to the pediatric ward with a complaint of poor academic performance and failure to thrive. On dietary history assessment, the families' staple foods were from cereals. They usually use salts stored in open containers. The mother also believed seafood and fish were harmful for health. On physical examination, the child had a slightly distended anterior neck.

What is the most likely biomarker used to detect the child's problem in this case?

- A. Plasma ferritin concentrations
- B. Plasma retinol concentrations
- C. Urinary iodine concentration
- D. Plasma transferrin concentrations

ANSWER: C

38. A 15-year-old adolescent presented to a hospital with skin rashes, poor taste acuity and multiple skin lesions. The attending physician linked the adolescent to the nutrition expert for nutritional assessment. The nutrition expert was interested in dietary assessment for zinc. The adolescent instructed to list all the foods and beverages consumed for seven consecutive days including the portion size using household measures.

What is the most appropriate dietary assessment technique for this patient?

- A. 24-hour recall method
- B. Food frequency method
- C. Diet history method
- D. Food diary method

ANSWER: D

39. A three-year-old child was brought to a pediatric outpatient department with a complaint of unable to walk, poor family socialization, poor appetite, and diarrhea of one day duration. The child had been breast fed and started complementary feeding at the age of six months. They lived in a highland, where the rain and cloud were prevailing throughout year, and were difficult to see the sunlight. On a biochemical assessment, the serum vitamin D concentration was 18ng/ml, showing vitamin D deficiency.

What is the most likely symptom or sign the child would have for vitamin D deficiency in this case?

- A. Poor family socialization
- B. Poor appetite
- C. Frequent Diarrhea
- D. Unable to walk

ANSWER: D

40. A two-year-old girl was brought to an under-five OPD with complaints of difficulty sucking breast and weight loss. On physical examination, she had a Mid Upper Arm Circumference (MUAC) of 10.5 cm and a grade one bilateral pitting edema. Her heart rate was 110 beats per minute. Her respiratory rate was 40 breaths per minute. Additionally, she had engorgement of the jugular vein, cold hands and feet, and cyanosis of the fingertips. The hemoglobin level was 3 g/dL.

What micronutrient deficiency best describes this case?

- A. Moderate iron deficiency anemia
- B. Mild iron deficiency anemia
- C. Severe iron deficiency anemia
- D. Borderline iron deficiency anemia

ANSWER: C

41. A two-year-old child from a rural area was brought to a clinic by his mother complaining of dry and scaly skin. He also had a previous history of measles, body weakness and loss of appetite. On physical examination, he had significant dryness of his abdominal and back skin. Besides, there were white, foamy-looking accumulations on the conjunctiva of his eyes.

What is the most likely nutritional diagnosis for this patient?

- A. Vitamin C deficiency
- B. Vitamin K deficiency
- C. Vitamin A deficiency
- D. Vitamin D deficiency

ANSWER: C

42. A 34-year-old pregnant woman was referred to an outpatient clinic from a rural health center with a complaint of vaginal bleeding of one week duration. The patient, who lived in one of the highland villages, said that she previously had experienced two miscarriages. She also reported that she gave birth to a child with mental retardation and hearing problem two years ago. She had no family history of intellectual disability. She always added salt to the food at the time of cooking.

What is the most likely type of micro-nutrient deficiency presented in this scenario?

- A. Selenium deficiency
- B. Zinc deficiency
- C. Iron deficiency anemia
- D. Iodine deficiency disorder

ANSWER: D

43. A two-year-old male child was brought to under-five outpatient clinics with a complaint of watery diarrhea of three days duration. On average, he had two to three diarrheal episodes per day. His food intake was reduced. He had a history of two previous diarrheal attacks in the last six months. On physical examination, he was found to have a rash around his mouth.

What is most likely micro-nutrient deficiency observed in this case?

- A. Iron deficiency
- B. Iodine deficiency
- C. Zinc deficiency
- D. Selenium deficiency

ANSWER: C

44. A one-year-old female child was brought to a health center with a complaint of frequent falling and suspected deformities on her lower extremities. Her mother also reported that her child's knees knock each other whenever she tries to stand up and walk on her own. The child was born during the rainy season. On physical examination, a rachitic rosary was found at the junction of the ribs and cartilage. When the child tried to walk, her legs looked bowed and her knees knocked frequently. Besides, the child had frontal bossing.

What is the most likely nutritional diagnosis in this scenario?

- A. Vitamin D deficiency
- B. Vitamin A deficiency
- C. Vitamin K deficiency
- D. Vitamin E deficiency

ANSWER: A

45. A five-year-old boy was brought to an under five OPD with a complaint of frequent epistaxis of two days duration. The blood was bright red. He also complained of headache and body weakness. He had previous history of the same problem a month ago. The hematocrit level was border line.

What is the first step in the management of this case?

- A. Give iron supplementation
- B. Give counseling about iron rich foods
- C. Investigate the underlying cause
- D. Give counseling about factors affecting bioavailability of iron

ANSWER: C

46. A four-year-old female child was brought to a clinic with a complaint of generalized body fatigue. She had also loss of appetite and weight loss. She came from malaria endemic area. On physical examination, both palms were pale and Mid-upper arm circumference (MUAC) was 11 cm. She showed no sign of congestive heart failure. Her hematocrit level was 11percent.

What is the most likely management in this case?

- A. Transfuse whole fresh blood at 10 ml/kg slowly over 3 hours
- B. Transfuse 5-7 ml/kg packed cells over 3 hours
- C. Transfuse whole fresh blood at 15 ml/kg slowly over 3 hours
- D. Transfuse 10-15 ml/kg packed cells over 3 hours

ANSWER: A

47. A seven-month-old male child was brought to a health center OPD with a complaint of a red and flat rash around his face. The rash spread to other parts of the body but was not associated with an itching sensation. He also had a fever and runny nose. Besides, his eyes were sensitive to light. On physical examination, his axillary temperature was found to be 37.9 degrees centigrade. Moreover, there was a red lesion on most surfaces of the skin. Finally, he was diagnosed with having a vitamin A deficiency.

What is the most likely nutritional management for this case?

- A. Give 50,000 IU of vitamin A supplement
- B. Give 100,000IU of vitamin A supplement
- C. Give 200,000IU vitamin A supplement
- D. Give 500,000IU vitamin A supplement

ANSWER: B

48. A three-year-old female child who is being treated for severe acute malnutrition (SAM) in the stabilization center (SC) of a health center recently complained of difficulty of opening the eyes. Her mother had reported that the child cries when she tries to open and blink her eyes. On physical examination, both of her eyes looked red and there is dryness of the conjunctiva and cornea.

What is the most likely first nutritional management for this case?

- A. Give vitamin A supplementation
- B. Give iron supplementation
- C. Give nutritional counseling
- D. Give zinc supplementation

ANSWER: A

49. A 35-year-old female patient came to adult OPD with a complaint of anterior neck swelling. She had difficulty of breathing and swallowing. She also had tremor and oligomenorrhea. She was living in a goiter endemic area for years. On physical examination, large multinodular anterior neck swelling was found. Besides to a surgical management, the doctor referred her for a nutritional management.

What is the most likely nutritional management to be applied by the nutritionist in this case?

- A. Advice on the importance of surgical management
- B. Advice on increasing consumption of vegetables
- C. Advice on the utilization of iodized salt
- D. Advice on increasing the consumption of cassava

ANSWER: C

50. A two-year-old child was brought to a clinic with a complaint of watery diarrhea of three days duration. The child had abdominal cramp and nausea, but no vomiting. On examination, there was no sign of dehydration. Besides, the stool examination result was normal.

What is the mostly recommended nutritional management for this case?

- A. 20mg of zinc for 10 days
- B. 10mg of zinc for 5 days
- C. 20 mg of zinc for 5 days
- D. 10 mg of zinc for 10 days

ANSWER: A

51. A two-year-old male child was brought to a pediatric OPD with a complaint of leg pain of one month's duration. The mother reported that the baby was not properly exposed to sunlight. His dietary intake showed a cereal-based diet with poor consumption of animal-source foods. He also stopped breastfeeding at the age of seven months. On physical examination, the child's legs were bowed, making it difficult to stand up and walk on his own.

What is the most appropriate management for the underlying nutrient deficiency in this case?

- A. Vitamin A supplementation
- B. Vitamin D supplementation
- C. Vitamin E supplementation
- D. Vitamin K supplementation

ANSWER: B

52. A five-year-old male child was brought to an OPD with a complaint of easily gum bleeding. He also had fatigue and malaise. On physical examination, swollen bleeding and necrotic gums were seen. Besides, there were small and red skin discolorations and easily bruising skin. He had also clogged and enlarged, hyperkeratotic hair follicles.

What is the most likely nutritional management to be applied in this scenario?

- A. Vitamin A 100-400 mg daily for three months
- B. Vitamin K 200-300mg daily for three months
- C. Vitamin E 150-400mg daily for three months
- D. Vitamin C 100-500 mg daily for three months

ANSWER: D

53. A two-year-old male child was brought to a health center with a complaint of dryness of the eye. His mother also reported loss of appetite and loss of weight. He had a history of recurrent diarrhea. He also had generalized body weakness. On physical examination, his weight for height and MUAC measurements were normal. After medical treatment, the physician referred him to a nutritionist. The nutritionist checked the child monitoring chart before providing him with vitamin A. However, the nutritionist told the mother to bring her child after one month.

What is the most likely monitoring chart information the nutritionist checked in this case?

- A. Previous vitamin A supplementation date
- B. Previous history of recurrent diarrhea
- C. Anthropometric measurement
- D. Sign and symptoms of the child

ANSWER: A

54. A five-year-old male child had a history of recurrent diarrhea and acute respiratory illnesses for the last six months. The physician managed the case and referred the child to a nutritionist for further nutritional management. The nutritionist advised the mother to feed her child zinc rich foods.

What is the most likely nutritional advice provided in this case?

- A. Eat protein rich foods
- B. Eat carbohydrate rich foods
- C. Eat high fat foods
- D. Eat high calorie foods

ANSWER: A

55. A 15-year-old girl from a mountainous village was worried by the presence of few students in her class who had anterior neck swelling. She went to a nearby clinic in her school and asked the nutritionist what she could do to reduce her risk of having the same problem. On dietary assessment, her main food has been *injera* with shiro. She has also informed him that there is a nearby river in her village where different types of fish species were found. However, she could not get any of the fish types in her dish.

What is the most likely dietary advice the nutritionist provided in this case?

- A. Improve consumption of kales
- B. Improve consumption of fish
- C. Improve consume of injera
- D. Improve consumption of Shiro

ANSWER: B

56. A 29-years-old woman gave birth to a healthy baby in the obstetrics ward of a hospital. She delivered by spontaneous vaginal delivery. There was no post-delivery complication, and the nurse brought the baby to the mother and started to advice the mother on breastfeeding.

What is the first advice about breastfeeding the nurse should provide in this case?

- A. To start breastfeeding within 1 hour
- B. To start breastfeeding within 30 minutes
- C. To start breastfeeding within 2 hours
- D. To start breastfeeding within 1 day

ANSWER: A

57. An 11-year-old girl was brought to a health center with a complaint of excessive weight gain. The mother reports that she frequently feeds the child with carbonated foods on a daily basis. The nutritionist has planned to assess the nutritional status of the child.

What is the most appropriate assessment method should be used in this case?

- A. MUAC for Age
- B. BMI for age
- C. Height for age
- D. Weight for age

ANSWER: B

58. A 64-year-old woman visited a hospital for medical check for hypertension, type-2 diabetes mellitus, and hyperlipidemia. On nutritional assessment, her BMI was 28 kg/m^2 , MUAC was 26 cm, waist to hip ratio was 0.85 and waist circumference was 94 cm. The nutritionist wants to confirm if the patient has central obesity.

Which nutritional assessment finding most likely predicts the diagnosis of central obesity in this case?

- A. BMI equals to 28 kg/m^2
- B. MUAC equals to 26cm
- C. Waist to hip ratio equals to 0.85
- D. Waist circumference equals to 94 cm

ANSWER: D

59. A 45-year-old man has visited a physician for routine medical checkup. The man is non-smoker, having long working hours in office. And even on weekends he spends most of the time in home working on his laptop. He used to be a chronic smoker. Besides, he occasionally drinks alcohol in social gatherings. His BMI is 32 kg/m^2 . The physician refers him to a nutritionist for counseling.

What is the most appropriate counseling the nutritionist would provide in this case?

- A. Alcohol consumption
- B. Cigarette session
- C. Sedentary life
- D. Comprehensive life style

ANSWER: C

60. A 45-year-old office worker has been struggling with weight gain in the last few years. He leads sedentary life with long working hours in the office and consumes sugar beverages on a daily basis. He has no known food allergy and food restriction. He wants to make changes in his life style to prevent overweight/obesity.

What is the most suitable dietary meal plan recommending to prevent overweight/obesity in this case?

- A. Low carbohydrate diet, high protein diet
- B. Low fat diet, high carbohydrate diet
- C. Balanced diet with controlled portion size
- D. Diet with packed weight-loss meals

ANSWER: C

61. A 34-year-old man visited a clinic for routine medical checkup. The man was leading life style of long hour sitting. He enjoyed eating sweet foods and animal products mostly. He occasionally was drinking alcohol in social gatherings. His BMI was 27kg/m^2 . The man was motivated for life style modification seeking a nutritionist for guidance.

What is the most appropriate counseling for this patient?

- A. Engaging in busy work days and nights stressfully to lower weight gain
- B. Encouraging to grab Obesogenic foods between regular meals
- C. Suggesting to skip lunch and compensate during dinner
- D. Recommending for low-fat diets and plenty amount of whole grains

ANSWER: D

62. A 40-year-old man had been diagnosed with obesity and received counseling on the weight management strategies. After four months, he came for a follow up visit. On anthropometry assessment, the BMI of the patient decreased from 35 to 27kg/m^2 . His MUAC decreased from 28cm to 27 cm.

What is the most appropriate anthropometry finding that shows the effectiveness of the weight management in this case?

- A. Average MUAC value
- B. MUAC of 27cm
- C. Average BMI value
- D. BMI of 27 kg/m^2

ANSWER: D

63. A 40-year-old man has been diagnosed with moderate obesity. On dietary assessment, his favorite foods were animal product-based, and he did not consider fruits and vegetables as essential nutritional components. He thought being obese was an indicator of being from a high social class. The nutritionist counseled on the overall health consequences of obesity, and the man was partially convinced to make lifestyle modifications. On a follow-up visit, the nutritionist assessed the patient's weight management effectiveness. However, his obesity was worsening to severe obesity. The nutritionist concludes what should be done to manage the case.

What is the most appropriate measure the nutritionist should take in this scenario?

- A. Take the patient to sign up for locally available gymnasium for physical activity
- B. Convince the consequences of obesity if left untreated via life modification
- C. Encourage the patient for heavy alcohol drinks in the daily base
- D. Encourage the patient to proud of his being obese

ANSWER: B

64. A 46-year-old male with cardiovascular disease and on follow up, came to a nutrition clinic for lipid profile checkup. He had undergone surgery for appendicitis a month ago. This patient was adhering to whole cow milk, meat and other fatty diet thinking that these meals would be helpful for his quick recovery of the wound and injuries faced during the operation.

What is the most appropriate nutritional assessment method used to evaluate the cholesterol level in this case?

- A. Anthropometric
- B. Clinical
- C. Biochemical
- D. Dietary

ANSWER: C

65. A 37-year-old man, who leads physically active lifestyle, working regular exercise, has come to a hospital for a medical checkup. The man is a non-smoker and drinks alcohol occasionally. He eats fruits and vegetables on a daily basis. He consumes packed foods rich of potassium and sodium most of the time. On examination, lipid profile is in the normal range, BMI was 22kg/m^2 , and blood pressure was 150/110 mmHg.

What is the most common risk factor for developing hypertension in this case?

- A. Sedentary lifestyle
- B. Alcohol drinking
- C. Eats fruits and vegetables daily
- D. Increase intake of packed food

ANSWER: D

66. A 37-year-old man works in a bar as a 'salesperson. His customers invite him to all things they use. So, he enjoys dances, meals, drinks, and others with the customers. He uses less red meat, a daily high amount of alcohol, fewer vegetables, and plenty of whole grains. One day, he notices that his body weight is increasing. He has gone to a health center for a medical checkup if the situation is associated with his weight gain. On assessment, his waist circumference is 100 cm and his BMI is 26 kg/m^2 . His blood pressure is 150/100 mm Hg. A nutritionist wants to make the patient aware of what factor causes his hypertension.

What is the most likely risk factor for developing hypertension in this case?

- A. Alcohol drinking
- B. Physical activity
- C. Vegetable and fruit consumption
- D. Whole grain consumption

ANSWER: A

67. A 44-year-old house wife mother came to a health center with complaints of thirst, hunger, and fatigue. She has family history of diabetes mellitus. On assessment, fasting plasma blood glucose was 140mg/dl; BMI was 16.5kg/m^2 , High density lipoprotein was 54mg/dl, and systolic blood pressure was 120 mm Hg. She leads active lifestyle, adheres on fruits and vegetables. Due to caring of her body image, she usually prefers to eat less carbohydrates and protein rich foods.

What is the most likely diagnosis for this scenario?

- A. Overweight
- B. Diabetic mellitus
- C. Underweight
- D. Hypertensive

ANSWER: C

68. A 46-year-old man was diagnosed with consistent blood pressure reading of 140/100 mmHg. The patient was referred to nutrition clinic for dietary management. On nutritional assessment it was found that, the patient was using minimal whole grain, less fruit and vegetables; plenty amount of meat and high sweet beverages. He also took two bottles of alcohol per day regularly. The patient was motivated to modify his lifestyle.

What is the most appropriate meal plan recommended for this case?

- A. Plentiful fruits and vegetables
- B. Plentiful red meat and sweet beverages
- C. Minimal whole grains
- D. Minimal alcohol regularly

ANSWER: A

69. A 60-year-old man came to the clinic for a routine medical checkup. He had a sedentary lifestyle and was overweight. On assessment it was found that he had a meal habit of consuming animal products commonly. On the other hand, he consumed lesser amounts of fiber-source foods. His random plasma glucose was 110 mg/dl, and hemoglobin A1C was 5.5 percent. The patient has been motivated to take intensive lifestyle management to prevent being diseased by NCD. So, a nutrition expert wanted to provide a dietary meal plan recommendation for this patient.

What is the most appropriate nutritional recommendation for this case?

- A. Plenty amount of sugar beverages, and less fruits and vegetables
- B. Plenty amount of unsaturated fat source, protein, fruits and vegetables
- C. Ketogenic diets limited proteins, starchy carbohydrates.
- D. Ketogenic diets, processed food, and fruits and vegetables

ANSWER: B

70. A 56-year-old obese woman, who leads sedentary life style, has been diagnosed with diabetes Mellitus, cardiovascular diseases, and breast cancer. She has been referred for dietary recommendation. She used to have varieties of fruits and vegetables, whole grain products and drinks large amount of water every day. She also drinks alcoholic beverages daily. She is discharging her daily duty, and she also makes regular physical activity.

What is the most appropriate dietary recommendation for this case?

- A. Consume healthy protein source foods.
- B. Consume high macronutrient food source
- C. Consume low macronutrient source foods.
- D. Consume high fat diets

ANSWER: A

71. A 56-year-old patient was diagnosed with chronic kidney failure and undergone kidney transplant. Post-kidney transplant, the dietetic patient was consulted for nutritional recommendation.

What is the most likely nutritional recommendation needed in this case?

- A. Provide tolerable unit of alcohol per day.
- B. Provide high carbohydrate diets
- C. Provide high fat diets
- D. Provide tolerable lean protein diets

ANSWER: D

72. A 16-year-old male adolescent used to consume processed and fatty foods daily. Besides, he loved playing video games for a minimum of 5 hours per day. On a nutritional assessment he had history of skipping meals, taking processed food, and plenty amount of water daily. He was taking breakfast and dinner with his family rarely. On assessment, BMI was +3.2 standard deviation, serum triglyceride, low density lipoprotein, and high-density lipoprotein was 162mg/dl, 138mg/dl and 36mg/dl respectively.

What is the most appropriate dietary counseling for this case?

- A. Encourage family meals
- B. Water intake should be restricted
- C. Breakfast should be skipped
- D. Fast food meals should be encouraged

ANSWER: A

73. A 64-year-old male patient was presented to a nutrition clinic with activity intolerance, shortness of breath and fatigue, anxiety and peripheral fluid retention. The internist diagnosed the patient that he had stage-C heart failure and provided all the medical treatments, and consulted the nutrition expert for dietary management. On nutritional assessment, he had history of irregular and high portion meal feeding habit. He was feeding any source of food. He was not worried for the contents of the food nor how the food had been cooked.

What is the most appropriate nutritional counseling in this case?

- A. Recommend small and frequent meals
- B. Recommend large and infrequent meals
- C. Recommend on adherence to high-sodium diets
- D. Recommend on smoked and processed meats

ANSWER: A

74. A 60-year-old obese and known hypertensive patient has been on hydrochlorothiazide for the last six months. She has a monthly follow up in dietetic clinic where she gets individualized dietary management for both weight loss and high blood pressure. She is always strict on her dietary plan. The dietitian planned to monitor anthropometric measurement to determine her risk of cardiovascular diseases (CVD).

What is the most appropriate index to be used in this case?

- A. Body mass index
- B. Waist circumference
- C. Hip circumference
- D. Waist to hip circumference ratio

ANSWER: D

75. A 49-year-old known chronic kidney disease (CKD) male patient undergoes Hemodialysis three times a week. The dietitian followed his dietary intake and nutritional status regularly. The dietitian recommended taking a daily protein of 1.2 g/kg of his body weight. On the next visit the dietitian found that he had edema of both legs in the last three days. On dietary assessment, he said that as he could not afford the recommended daily quantity of protein, he was consuming what was available in his home.

What is the most likely recommendation should be given to him in this case?

- A. Adherence to the diet plan
- B. Adherence to medication
- C. Adherence to physical exercise
- D. Adherence to dialysis

ANSWER: A

76. A 35-year-old pregnant woman living with HIV, who was on highly active antiretroviral therapy (HAART), came to a hospital for a follow up. The CD4 count was stable when compared with the last one and the laboratory findings showed that the lipid profile and liver function test results were within the normal range. Then the attending physician linked the patient to the nutrition unit for assessing her nutritional status.

What is the most appropriate nutritional assessment method used for this case?

- A. Body mass index (BMI)
- B. Mid upper arm circumference (MUAC)
- C. Weight for age
- D. Waist to hip ratio

ANSWER: B

77. A 40-year-old adult, who was on highly active retroviral therapy, was come to a hospital with severely wasted muscle. Laboratory investigation revealed that CD4 count was 200cells/L. The attending doctor sent to the nutrition outpatient department for understanding about the effects of nutrition on HIV progression and vice versa.

What is the most appropriate description to be provided for this patient in this case?

- A. Poor nutrition hastens HIV infection recovery
- B. Nutrient metabolism decreased by HIV infection
- C. HIV infection increases nutrient absorption
- D. HIV increases breakdown of body protein

ANSWER: D

78. A 35-year-old man was presented to a medical OPD with complaints of night sweating, weight loss and cough of 2 months duration. He had contact history with a patient known for tuberculosis. On laboratory investigation, gene expert revealed that the patient had mycobacterium tuberculosis. The attending physician linked the patient to the nutrition outpatient department for understanding about the effects of nutrition on tuberculosis and vice versa before treatment was started.

What is the most appropriate description to be provided for this patient?

- A. Tuberculosis makes malnutrition worse
- B. Malnutrition decreases the initiation of active tuberculosis
- C. Under-nutrition is not related with Tuberculosis
- D. Tuberculosis increases nutrients metabolism

ANSWER: A

79. A 35-year-old woman in her third trimester pregnancy, who came from lowland area, was presented with complaints of fatigue and headache. She was treated for malaria a week ago. She had no family history of hypertension, diabetes mellitus and goiter. On physical examination, blood pressure of 120/80mmHg, temperature of 36.7 degree centigrade, and eye examination revealed that she had white sclera and pale conjunctiva.

What is the most likely diagnosis for this patient?

- A. Vitamin A deficiency
- B. Iron deficiency anemia
- C. Iodine deficiency disorder
- D. Vitamin D deficiency

ANSWER: B

80. A 32-year-old pregnant woman came to a clinic with complaints of pallor, fatigue, weakness and behavioral disturbances. She had also frequent tapeworm infestation in the current pregnancy. She had no vision problem, no family history of hypertension, goiter, and diabetes mellitus. On biochemical assessment it was found that hemoglobin level of 10.5mg/dl, serum vitamin D of 25ng/l, and the serum plasma retinol concentration of 28micro gram/dL. Her dietary practice was characterized by poor consumption of vitamin C rich foods, fortified cereals, meat, fish, and poultry. What is the most appropriate dietary intervention for this patient?

- A. Adequate intake of vitamin D rich foods
- B. Adequate intake of vitamin A rich foods
- C. Adequate intake of iron rich foods
- D. Adequate intake of iodine rich foods

ANSWER: C

81. A 60-year-old male patient, who is on a highly active antiretroviral therapy, presented to a hospital with weight loss and loss of appetite. On physical examination, he was wasted and lost 6 percent of his baseline weight.

What is the most appropriate diet recommended to this patient?

- A. Proteins rich foods
- B. Low carbohydrate containing foods
- C. Food's rich of trans-fats
- D. Low unsaturated fat containing foods

ANSWER: A

82. A 50-year-old HIV positive patient was presented to ART clinic with angular stomatitis and dermatitis. The patient had poor dietary consumption for meat and dairy products. On biochemical assessment, hemoglobin was 14.5mg/dl, serum vitamin D was 25ng/l, and the serum plasma retinol concentration was found to be 28microgram/dL.

What is the most appropriate intervention recommended to this patient?

- A. Vitamin A supplementation
- B. Riboflavin (B2) supplementation
- C. Vitamin D supplementation
- D. Folic acid supplementation

ANSWER: B

83. A 65-year-old, who has been on HIV treatment for the last ten years, was presented to a health center with complaint of loss of appetite on her follow up. The mother had also recurrent infection, taste problems, and bruising of skins. The patient was linked to the nutrition unit for assessment of common micronutrient deficiencies.

What is the most likely nutritional assessment method used for this patient?

- A. Acute Phase Proteins
- B. Lipid profile Assessment
- C. Hematological Assessment
- D. Anthropometric Assessment

ANSWER: C

84. A 22-year-old adolescent, who was on the continuation phase of anti-tuberculosis treatment, came to a health center for follow up. On the anthropometrics assessment, the weight of the adolescent showed a decrease from the baseline measurement. Laboratory findings showed no evidence for infection. The attending physician confirmed that the patient had adherence for the treatment. The patient was referred to the nutrition clinic for nutritional counseling.

What is the most appropriate counseling for this patient?

- A. Nutritional status check for the next visit is not important
- B. Eating more and a variety of food stuffs is good for optimal health
- C. Drinking untreated water decreases the disease progression
- D. Skipping anti-tuberculosis drugs decreases the side effects

ANSWER: B

85. A 32-year-old male was diagnosed with HIV/AIDS. He has started ART and was linked to the nutrition clinic for nutritional support. He had normal body weight and was eager for nutritional counseling to maintain his body weight and the overall health.

What is the most appropriate nutritional counseling for this patient?

- A. Recommend him to take low protein foods
- B. Recommend him to consume diets rich of omega-3 fatty acids
- C. Encourage him to avoid fruits and vegetables in his diet
- D. Encourage him to drink alcohol

ANSWER: B

86. A nutrition expert planned to conduct nutrition assessment to determine the adequacy of the nutrient intake of individuals for food aid in an area where the population affected by drought, and many people were exposed to hunger.

What is the most appropriate nutritional assessment method to be applied in this case?

- A. Anthropometric
- B. Biochemical
- C. Clinical
- D. Dietary

ANSWER: D

87. A team of nutrition professionals organized by the Woreda Public Health Emergency Management (PHEM) planned to conduct a nutritional assessment to understand the emergency situation following a recent political unrest. They planned to obtain an overview of the nutritional situation, and determine areas and population groups affected by an emergency.

What is the most appropriate assessment method to be applied in this case?

- A. Rapid Survey
- B. Repeated surveys
- C. Sentinel sites surveillance
- D. Surveillance

ANSWER: A

88. A human nutrition professional has planned to conduct a rapid assessment to screen pregnant mothers for productive safety net program in a district following a prolonged drought. Reports are showing that people are suffering from hunger and many more are dying.
What is the most appropriate assessment tool to be applied in this case?
- A. Body Mass Index
 - B. Mid Upper Arm Circumference
 - C. Waist circumference
 - D. Skin fold thickness
- ANSWER: B
89. A rapid response nutrition team went to the Nile basin of Ethiopia following a sudden flood. The flood caused massive destruction of houses and livelihoods. As a result, thousands of pregnant women were crying for food aid to survive. The team conducted a rapid assessment to screen the pregnant mother for acute malnutrition.
What is the correct cut of point to diagnose acute malnutrition?
- A. Body mass index less than 18 Kg/m^2
 - B. Mid upper arm circumference less than 23 cm
 - C. Body mass index greater than 18 Kg/m^2
 - D. Mid upper arm circumference greater than 23 cm
- ANSWER: B
90. A nutrition professional went to a refugee camp where hundreds were displaced from their residences due to armed conflict in Ethiopia and majority of them were elders. During a rapid nutritional assessment, the nutritionist found that almost all of the elders had a MUAC reading of less than 18.5 cm.
What is the most likely diagnosis in this scenario?
- A. Moderate Acute Malnutrition
 - B. Chronic malnutrition
 - C. Sever acute malnutrition
 - D. Moderate chronic malnutrition
- ANSWER: C
91. A team of researchers conducted research on the nutritional status of under five-year-old children who were victims of prolonged drought. After analyzing the nutritional data, the prevalence of acute malnutrition was found to be 15 percent.
What is the most appropriate classification for this case?
- A. Acceptable
 - B. Poor
 - C. Serious
 - D. Critical
- ANSWER: D
92. The nongovernmental organization planned to provide dry ration for people highly affected by food insecurity by a sudden flood. The population was entirely dependent on general food rations as their source of food.
What estimate should be considered as minimum nutritional requirements in this case?
- A. Water
 - B. Energy
 - C. Vitamins
 - D. Minerals
- ANSWER: B

93. A group of nutrition experts identified that the prevalence of moderate acute malnutrition among under-five-year-old children was 13 percent in the eastern part of Ethiopia, where recurrent drought coupled with manmade humanitarian crises occurred. Moreover, the team confirmed that there was a high burden of infectious diseases and the households were suffering from food insecurity.

What is the most likely intervention program to be tailored in this scenario?

- A. Targeted supplementary feeding
- B. Therapeutic feeding
- C. General food ration
- D. Blanket supplementary feeding

ANSWER: A

94. A team of graduating class nutrition students was assigned to an internally displaced site to screen acute malnutrition for appropriate management. The team found that many children were with severe acute malnutrition, had medical complication, and bilateral pitted edema.

What is the appropriate management program to be applied in this case?

- A. General food ration
- B. Blanket feeding
- C. Outpatient therapeutic
- D. Inpatient therapeutic

ANSWER: D

95. A dietitian who works for a local NGO was assigned to assess the overall condition of a refugee camp. She identified that there were a diversified group of people in the camp and a significant number of individuals were complaining of diarrhea and vomiting. She planned to provide a nutritional counseling before leaving the camp.

What is the appropriate information to be included in her counseling?

- A. Hygiene and sanitation
- B. promoting breast milk substitutes
- C. Encourage food sharing
- D. Promote food selling

ANSWER: A

96. A research team organized by the Federal Ministry of Health of Ethiopia (FMOH) wanted to assess the effect of nutritional status on chronic non-communicable diseases, particularly morbidity and mortality due to cardiac diseases.

What is the most appropriate anthropometric assessment method to be applied in this case?

- A. Skin fold thickness
- B. Hip circumference
- C. Waist circumference
- D. Thigh circumference

ANSWER: C

97. A 34-year-old woman visited a nutrition unit in a private clinic for her iron supplementation on her appointment. She asked the dietitian about the highest amount that she could take daily. What nutrient intake standard could the dietitian use to provide the most appropriate recommendation?

- A. Estimated average requirements (EARs)
- B. Recommended dietary allowances (RDAs)
- C. Adequate Intakes (AIs)
- D. Tolerable upper intake levels (ULs)

ANSWER: D

98. A 13-year-old girl was brought to a hospital by her mother, who was concerned with the unusual tiredness and frequent headache. The girl had been complaining of fatigue, and she was struggling to execute the usual activities in her school and during sport activities. She had no family history of bleeding disorder or cancer. Physical examination also showed pale conjunctiva and a grade-I systolic murmur. The laboratory report indicated a hemoglobin of 7.1 mg/dl (normal 12-16 mg/dl) and a mean corpuscular volume (MCV) of 72 fl (normal 80-100 fl).

What is the most likely diagnosis for this scenario?

- A. Anemia of chronic diseases
- B. Iron deficiency anemia
- C. Sickle cell anemia
- D. Hemolytic anemia

ANSWER: B

99. A 32-year-old pregnant woman came to a hospital for prenatal check-ups in her third trimester. She experienced excessive thirst and urination in the previous few weeks. She also complained mild headache, occasional blurring of vision and unusual fatigue. During examination, her blood pressure was within the normal range; however, the fasting blood glucose exceeded the diagnostic threshold.

What is the most likely diagnosis in this case?

- A. Type 1 diabetes mellitus (T1DM)
- B. Type 2 diabetes mellitus (T2DM)
- C. Gestational diabetes mellitus (GDM)
- D. Latent autoimmune diabetes in adults (LADA)

ANSWER: C

100. A pregnant woman came to a primary health care unit with complaints of feeling weak, fatigue, and lightheadedness. During consultation, she started to disclose that she was strict on certain food items due to cultural and religious beliefs. Upon further questioning, it was confirmed that she avoided animal-source foods like meat, eggs, fish, and some fruits and vegetables. She was also worried and embarrassed to share her and her family's dietary habits.

What is the most likely description for this scenario?

- A. Food insecurity
- B. Food taboos
- C. Eating disorder
- D. Peer pressure

ANSWER: B

101. A newborn baby was sent to a pediatric neurologic unit for evaluation with his mother. He was born with a sac-like protrusion at the lower back. The mother had no ANC follow up and did not take any supplement. The MRI investigation showed incomplete formation around the second sacral spine.

What nutrient deficiency most likely explains this scenario?

- A. Vitamin B1 (Riboflavin) deficiency
- B. Vitamin B 3 (Niacin) deficiency
- C. Vitamin B6 (Pyridoxine) deficiency
- D. Vitamin B9 (Folate) deficiency

ANSWER: D

102. A two-day-old male neonate, who was born in a nearby hospital, came back to the pediatric OPD with a complaint of being unable to suck breast milk appropriately. He was the first child for the nineteen-year-old mother and her husband. She delivered vaginally without complication with a birth weight of 3400 grams. During evaluation, the baby's weight was decreased to 3100 grams. The mother reported that her nipples were engorged and tender, and she did not know how to present her breast to the suckling child.

What is the most likely sub-optimal breast-feeding practice that brought back the child to the hospital?

- A. Late initiation of breast feeding
- B. Early initiation of breast feeding
- C. Poor positioning and attachment
- D. Infrequent breast feeding

ANSWER: C

103. An 82-year-old man from a rehabilitation center complained fatigue, muscle twisting and spasm, lower back pain, and serious leg pain recently. He had a Parkinson's disease for five years and had frequent history of fracture. The bone density scan (DXA) showed a poor bone density.

What is the most common nutrient that hurried the patient's problem?

- A. Potassium
- B. Magnesium
- C. Calcium
- D. Zinc

ANSWER: C

104. A 27-year-old woman, who came from a drought-prone, food-insecure rural kebele, visited a health facility seeking medical care to help her to become pregnant. She was amenorrheic for the last year. She has never used any family planning method before. She was convinced that lack of a menstrual period was the cause of infertility. The ultrasound finding showed an adequate number of follicles on both ovaries.

What is the appropriate nutritional counseling to improve her chance of conception?

- A. Advise her on how to improve weight gain
- B. Advise her on regular exercise to reduce weight
- C. Hormonal therapy to enhance fertility
- D. Develop binge eating habit to improve body shape

ANSWER: A

105. A 39-year-old woman, who has been diagnosed for hyperlipidemia, has come to a hospital for nutritional guidance. She is really concerned about which fats and oils were recommended to her case. What is the most appropriate recommendation to this scenario?

- A. Liquid fats, like vegetable oils, are good choices
- B. Organic lipids, like butter, are good choices
- C. Solid fats, like margarine, are good choices
- D. Replacing vegetable oils with butter

ANSWER: A

106. An 18-year-old adolescent was brought to the university student's clinic accompanied by her friends. She had recurrent upper air way infection. She also complained recent weight loss and loss of appetite. She was smoking cigarette and drinking alcohol for the last two years. After receiving the medical care, the clinician consulted the nutritionist for nutritional counseling.

What is the most important nutrient nutritionist going to recommend for this client?

- A. Vitamin B1
- B. Vitamin C
- C. Iron
- D. Iodine

ANSWER: B

107. A multidisciplinary health team containing physician, dietitian, and psychologist are approaching a young lady brought to the nutrition unit. She has been diagnosed with eating disorder two weeks ago. Her parents claimed that her nutritional habit is not changing despite their efforts to feed and follow her.

What kind of nutritional counseling should the team provide in this case?

- A. Commenting on the appearance and weight gain to make her ashamed
- B. Addressing the physical, emotional and social issues
- C. Follow restrictive dietary rules and guidelines
- D. Provide calorie-dense foods to restore weight and body shape

ANSWER: B

108. A 37-year-old pregnant woman came to a nutritional care center for a routine check-up on her appointment day. She was devoted to follow nutritional advices to maintain healthy dietary habit and good pregnancy outcomes.

What is the most appropriate recommendation the nutritionist should provide in this case?

- A. Rely solely on prenatal vitamins for nutrient intake
- B. Limit carbohydrate rich foods to prevent macrosomia
- C. Ensure adequate intake of foliate, iron and vitamins
- D. Follow vegetarian diet without supplementation

ANSWER: C

109. A mother delivered a healthy baby vaginally one month ago in a hospital. She decided to consult a dietitian to ensure the nutritional needs of lactation to nourish her baby and maintain her own health.

What is the most appropriate nutritional recommendation for this woman?

- A. Avoid dairy products in her diet to prevent colic pain on the baby
- B. Ensure adequate fluids and water during breast feeding
- C. Begin one spoon of honey to improve the child's immunity at 6 months
- D. Skip meals to decrease postpartum weight gain.

ANSWER: B

110. A 13-week-old baby girl was presented to clinic with a complaint of watery diarrhea and fever of one day duration. The mother reported that she was giving her baby water and lemon to satisfy the child's thirst. During assessment, it was confirmed that the child had low weight for age.

What is the most likely nutritional counseling provided to the mother in this case?

- A. Add formula feeding to improve the child's weight
- B. Use cup to feed water and lemon
- C. Put the child on exclusive breast feeding
- D. Begin family food rich in vitamin A and zinc

ANSWER: C

111. A dietitian was consulted to provide nutritional counseling in postnatal ward for a woman who gave birth two hours ago. The dietitian noticed that the baby was not breastfed appropriately because the woman was inexperienced for child attachment.

What is the best counseling that the dietitian should provide in this case?

- A. Lying down on back while breast feeding
- B. Cradling the baby in a semi-inclined position
- C. Hold the baby's head above the breast
- D. Supporting the baby with one hand while breast feeding

ANSWER: B

112. Normal aging is associated with changes in most physiological systems, which may eventually lead to functional and structural deteriorations. This in turn results sub-optimal nutritional status in the elderly.

What is the most appropriate nutritional counseling to this age group?

- A. High calorie content diet
- B. High nutrient dense diet
- C. Low calorie content diet
- D. Daily antioxidant supplementation

ANSWER: B

113. A public health nutritionist suspected that many children were affected by protein-energy malnutrition due to existing severe food shortages in a community. In the meantime, before conducting a thorough nutritional assessment, she conducted home to home visit to look for children with visible wasting, skin changes, body swelling (edema), and hair changes.

What is the most appropriate nutritional assessment method used in this case?

- A. Anthropometric
- B. Biochemical
- C. Clinical
- D. Dietary

ANSWER: C

114. A humanitarian organization wants to provide supplementary foods for pregnant women living in a community reportedly affected by lack of food and widespread hunger. Since the available supply of supplementary foods is limited, the nutritionist conducted a nutrition assessment of all pregnant women to screen those severely affected by acute malnutrition.

What is the most preferred nutritional assessment method to apply in this case?

- A. Body Mass Index
- B. Waist Circumference
- C. Mid Upper Arm Circumference
- D. Hip Circumference

ANSWER: C

115. A rural kebele located in the highlands underwent a monthly health review meeting. One of the presenters from the health office reported a recent rise in the number of spontaneous abortions. In addition, the community nutritionist presented that 10 percent of the pregnant women in the woreda had a visible neck swelling. After discussions, the participants emphasized the need for a nutrition screening for iodine deficiency in all pregnant women.

What is the most appropriate screening tool that should be used in this case?

- A. Serum thyroxin
- B. Serum thyroid stimulating hormone
- C. Urinary Iodine
- D. Serum Iodine

ANSWER: C

116. A national food consumption survey reported that the consumption of Vitamin A rich fruits and vegetables was very low among lactating women. However, it was not known if this poor consumption levels were correlated with Vitamin A deficiency in these groups of population. Then, senior nutrition researchers conducted a large-scale study to determine the prevalence of Vitamin A deficiency in lactating women.

What is the most likely nutrition assessment tool used by the researchers for this case?

- A. Serum retinol
- B. Serum retinol-binding protein
- C. Serum carotenoids
- D. Serum retinal esters

ANSWER: A

117. A researcher studied the prevalence and determinants of child malnutrition in a *woreda*. She was particularly interested in assessing linear growth of the children. Then, the results of her study showed that 15 percent of the children in the woreda had Height for Age Z scores (HAZ) of less than -3 standard deviations (SD).

What is the most likely diagnosis in this case?

- A. Moderate wasting
- B. Severe wasting
- C. Moderate stunting
- D. Severe stunting

ANSWER: D

118. A community nutritionist in a nearby woreda conducted anthropometric assessment to estimate the effect of measles epidemic on child malnutrition levels. The assessment results showed that 21 percent of the children in the woreda had Weight for Height Z scores (WHZ) greater than -3 but less than -2 standard deviations (SD).

What is the most likely diagnosis for this scenario?

- A. Moderate wasting
- B. Severe wasting
- C. Moderate stunting
- D. Severe stunting

ANSWER: A

119. A nutrition professional conducted research on iodine deficiency disorders in a highland community. For a biochemical test a urine sample was taken from school age children. The findings indicated that many children had a median urinary iodine concentration value of 17microgram/L.

What is the most likely diagnosis for this scenario?

- A. Adequate iodine status
- B. Mild iodine deficiency
- C. Moderate Iodine deficiency
- D. Severe Iodine deficiency

ANSWER: D

120. A community nutritionist conducted home-based nutrition assessment among pregnant women in mountainous community where, agricultural cereals are the staple food along with some legumes, tubers and vegetables like cabbage and cassava. He found several pregnant women with small palpable neck swelling.

What is the most likely nutrient deficiency for this scenario?

- A. Iron
- B. Zinc
- C. Iodine
- D. Calcium

ANSWER: C

121. A community nutritionist planned to implement an immediate intervention for severe iodine deficiency in a mountainous village known to be goiter-endemic and had poor access to nutritious foods. Meanwhile, a recently conducted nutrition survey showed that 10 percent of pregnant women in the village had iodine deficiency.

What is the most appropriate nutritional intervention to be provided to this case?

- A. Provision of education on the importance of iodine
- B. Provision of fruits and vegetables for everybody
- C. Provision of iodized salt for everybody
- D. Cooking demonstration for pregnant women

ANSWER: C

122. Vitamin A deficiency (VAD) is known to be highly prevalent in a resource-limited rural woreda. The local administration persistently applied various nutrition specific intervention modalities to alleviate the problem; but it could not bring a significant change. Finally, the nutritionists from agricultural sector planned to implement nutrition-sensitive interventions that could be aligned with the specific interventions.

What is the most likely intervention approach for this scenario?

- A. Gardening of vitamin A rich fruits and vegetables
- B. Provision of large-scale Vitamin A supplements
- C. Provision of Vitamin A fortified food products
- D. Management of people affected by vitamin A deficiency

ANSWER: A

123. A nutrition researcher studied the nutritional adequacy of local complementary foods for children. One of her findings indicated that local complementary foods had low micronutrient contents as compared to the standard. Based on this finding, she recommended mothers and local producers to consider cost-effective, safe and sustainable food-based approach to address micronutrient deficiencies.

What is the best nutrition intervention that must be implemented in this case?

- A. Promotion of horticultural activities
- B. Nutrient supplementation
- C. Food fortification
- D. Dietary modification

ANSWER: C

124. A rural community was affected by a long-standing food insecurity and malnutrition. As a potential intervention, the government established a program in which they employed one family member of each household and paid them in terms of food not money.

What nutrition intervention approach best describes the government's program in this case?

- A. Food subsidy
- B. Food for work
- C. Food stamps
- D. Income generation

ANSWER: B

125. A nutrition researcher studied on the feeding practices of children in a community with high burden of protein energy malnutrition (PEM). In his findings, he concluded that more than 90 percent of the children had history of optimal breast-feeding practices and consumption of monotonous cereal-based foods only.

What is the most likely nutritional intervention applied in this case?

- A. Nutrition education on optimal breastfeeding
- B. Nutrition education on dietary diversification
- C. Supplementation of therapeutic foods
- D. Food fortification

ANSWER: B

126. A group of nutrition researchers in a rural community studied the prevalence of stunting and its determinants factor in children under the age of five years and they found that the most notable positive determinant factor was ownership of chicken by households.

What is the most likely nutritional intervention could be recommended in this scenario?

- A. Promotion of exclusive breastfeeding
- B. Promotion of water, and sanitation and hygiene
- C. Promotion of early childhood care and development
- D. Promotion of poultry production

ANSWER: D

127. A team of nutrition experts in the Minister of Health organized to assess the food availability for consumption at national level during Covid-19 pandemic. Nationally, regardless of regional differences, many people faced massive food shortage for consumption.

What is the most appropriate assessment method to be used in this case?

- A. Food account
- B. Estimated food record
- C. A 24- hour recall
- D. Food balance sheet

ANSWER: D

128. A graduating class nutrition professional planned to conduct research to assess the food insecurity status of individuals in the households from conflict affected woreda where the food crops and livelihood of the community were destroyed and looted.

What is the most appropriate assessment method to be used for this case?

- A. Food consumption survey
- B. Market survey
- C. Living Standard Measurement Survey
- D. Poverty assessment survey

ANSWER: A

129. A nutrition student conducted a research project to determine the household food insecurity status in the recurrent drought affected Woreda. After the analysis of the data, it was found that the majority of the households had food insecurity access scale score of 24 out of 27.

What is the most appropriate interpretation for this finding?

- A. Mild food insecurity
- B. Moderate food insecurity
- C. Severe food insecurity
- D. Food secure

ANSWER: C

130. A team of nutrition professionals conducted a rapid emergency assessment and they found out that many household members were suffering from food insecurity following a recurrent drought in the area. Moreover, recently it was identified that almost all members of the households went the whole day and night without eating.

What is the most appropriate information to be included in their report?

- A. The capacity of the team
- B. The socio-cultural situation
- C. The geographical situation
- D. Categories of vulnerable groups

ANSWER: D

131. Following an armed conflict, reports from the area showed that people were under severe shortage of food access with high mortality. The prevalence of acute malnutrition among children was 22 percent and there was a widespread poverty. The people were trying to cope through irreversible livelihood asset stripping.

What is the most appropriate coping strategy that explains this scenario?

- A. Insurance strategies
- B. Crisis strategies
- C. Distress strategies
- D. Non-defined strategies

ANSWER: C

132. A team of nutrition professional planned an emergency intervention in widespread, high intensity conflict area where livelihood assets were completely lost and the social systems collapsed. The prevalence of acute malnutrition among children was 32 percent. Massive starvation, death, and displacement were evident.

What coping strategy has been applied before intervention in this scenario?

- A. Insurance strategies
- B. Distress strategies
- C. Crisis strategies
- D. Survival Strategies

ANSWER: D

133. A district officers performed inspection of nutritional supply shops. The officers had immense experience in inspection. The district head noticed that the report presented by these officers in media was different. The officers also revealed information that should have been addressed by the district head. The head of the district was working to solve this problem and planning to provide a training or workshop as a solution.

What is the most appropriate component of the prepared workshop in this scenario?

- A. Code of conduct
- B. Good governance
- C. Competitive remuneration
- D. Effective communication

ANSWER: A

134. A nutrition health professional is working on a health center usually report what he did not actually doing. Even though his coworkers advised him to report what he has done only; he keeps reporting the patients managed and treated by his colleagues as if he has performed the tasks.

What is the most likely professional ethics the nutrition health professional missed?

- A. Justice
- B. Patience
- C. Veracity
- D. Compassion

ANSWER: C

135. A client presented to a private clinic with a sudden onset of acute illness. The health care provider in charge noted that the complaint that the client presented did not need any treatment as it could be resolved by it-self. As a result, the client thanked the health care worker for providing her with the correct and necessary information about her medical condition.

What is the most likely ethical principle the health care workers are applying?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-maleficence

ANSWER: C

136. Health care workers in the nutrition department are working together as a team. The team regularly share information about their client and support each other for better outcomes of patients.

What is the most likely professional practice demonstrated in this scenario?

- A. Obtaining informed consent from a patient.
- B. Maintaining the confidentiality of the client
- C. Ensuring public participation and acceptance.
- D. Providing genuine professional communication.

ANSWER: D

137. A nutritionist was planning to provide training to his coworkers on how to prevent disrespectful care. The nutritionist demonstrated with a role play. A story that portrayed characters of some health care providers showing an intentional humiliation, blaming, denying services, laughing at patients, refraining from introducing himself to the patient and not recognizing patients by their names throughout the interaction.

What is the most likely aspect of disrespectful care revealed in this scenario?

- A. Non-dignified care
- B. Non-consented care
- C. Non-confidential care
- D. Abandonment of care

ANSWER: A

138. A nutritionist working at a primary hospital accepted a patient with a warm welcome; showed kindness and provided appropriate consultation tailored to the needs of the patient. As result, the patient was satisfied with the care and adhered to the nutritional consultation.

What is the most likely principle of compassionate care revealed in this scenario?

- A. Attention
- B. Affection
- C. Acceptance
- D. Acknowledgment

ANSWER: B

139. A program coordinator assigned a health care worker to develop a plan, deliver an activity and report the performance on improvement of district level leader's awareness on food and nutrition issues. Following this, the health care worker prepared an advocacy material, provided it in a half a day workshop and documented it. Finally, they prepared the activity report.

What is the most likely performance indicator utilized to report the work result in the above scenario?

- A. Number of districts advocacy provided
- B. Number of days the advocacy provided
- C. Number of leader's advocacy provided
- D. Number of advocacy events conducted

ANSWER: D

140. A nutrition health professional was preparing a seminar on a topic related to the nutritional status of children under five globally. He was reading a number of publications globally and recorded the average daily intake of nutrition. He got two points that contradict the publications. Then, he decided to look at the procedural component of the methods of the research.

What is the most likely research method used by the professional to get the best evidence in this case?

- A. Cohort
- B. Case control
- C. Cross-sectional
- D. Interventional

ANSWER: D

141. A nutritionist has been working in a rural hospital for a long time. During discussion in the morning session, he realized that he was not updated on the most recent information and updates on nutrition due to a lack of timely trainings in the institution. However, he heard that there was an informally organized training in the nearby town where he worked.

What is the most appropriate decision to access the professional development in this scenario?

- A. Participate in the training even if it is informally organized.
- B. Request his boss to arrange him good international training.
- C. Continue updating from the morning session and coworkers.
- D. Practice with the existing skills, as he is a graduate professional.

ANSWER: A

142. A nutrition health professional working in a primary hospital was striving to be committed to work in the hospital and update his capacity to practice safely, effectively, and legally within a scope of practice. Based on self-initiated plans, he had participated in three formal trainings organized on newly adopted guidelines on nutritionist-related professional practices.

What is the most likely health care practice the nutritionist demonstrated in this scenario?

- A. Professionalism
- B. Development in Ethics
- C. Improving in career structure
- D. Continuing professional development

ANSWER: D

143. A nutritionist at a primary hospital was consulted about a patient by a health care professional from an internal medicine department. The nutritionist recalled that he examined the same patient a week before after checking his card. At the moment, he was in a hurry as the working hour was over. He asked the patient about the problem and the progress from the previous visit. He finally decided to prescribe him another medication.

What is the most appropriate action the professional should take next in this scenario?

- A. Record in a card and give a prescription.
- B. Give the prescription by the time he comes back after a week.
- C. Record it on a card and tell him to go to the referring department.
- D. Record it on a card and give the card to take to another department.

ANSWER: A

144. A health care worker scheduled an inspection of a restaurant in the town. He planned to interview food handlers and conduct physical observation. Hence, he started to prepare a checklist for the observation.

What is the most appropriate initial step to be conducted according to the standard practice of inspection?

- A. Handling of raw food.
- B. Practice of personal hygiene.
- C. Cleanliness of the tables in the dining room.
- D. Presence of obvious indoor air pollution.

ANSWER: A

145. A leader of a food security agency in the region wants to provide region-wide training for the corresponding subordinates and nutritionists working in the area. Accordingly, he wants to provide training on the topic 'strengthen the national capacity to manage natural and manmade food and nutrition emergencies in a timely and appropriate manner, including for internally displaced persons and refugees.

What is the most appropriate source used by the leader to give the training in this case?

- A. Guide to Clinical Nutrition Care.
- B. National Food and Nutrition Strategy.
- C. Health and Health-Related Institutions Regulation Training Manual.
- D. Ethiopian Guide to Clinical Nutrition Care for Children and Adults with HIV.

ANSWER: B

146. A nutrition professional intended to advocate on food safety for hotel employees. He had already conducted a quick assessment on individual and community needs for the advocacy.

What is the most appropriate next step the professional should take in this case?

- A. Implement health education programs.
- B. Plan effective health education programs.
- C. Coordinate the provision of health education services.
- D. Evaluate the effectiveness of health education programs.

ANSWER: B

147. A nutrition professional is assigned to provide advocacy on the utilization of micronutrients for youth. He is planning on the common nutritional problems in Ethiopia for the advocacy service.

What is the most likely content of the advocacy plan that the nutrition professional is intending to provide in this scenario?

- A. Promote the use of carbohydrates.
- B. Promote the use of iodized salt.
- C. Provision of calcium supplementation.
- D. Provide facility-based nutrient supplementation.

ANSWER: B

148. A researcher stated at a particular section of proposal development stages as follows: 20 percent of the school children, 10 percent of pregnant women, and 15 percent of the elders suffered from malnutrition, which causes 25 percent anemia and 10 percent stunting. Even though different activities are taken by the government to reduce the burden of malnutrition in the community, the seriousness of the issue still remains high. Hence, this study aims to identify the factors associated with the magnitude of malnutrition in the community.

Which proposal development step was most described in the above scenario?

- A. Writing the background of the study
- B. Writing the statement of problem
- C. Writing the scope of the study
- D. Writing the significance of the study

ANSWER: B

149. A nutrition graduate student submitted his proposal on the magnitude of anemia among pregnant women attending antenatal care to the college seeking ethical clearance. Then, the research review committee suggested resubmission of the proposal after incorporating the possible risks of the study and protecting the rights of respondents.

What is the most likely comment given in the above proposal development step?

- A. Purpose of the Study
- B. Summary of the study
- C. Ethical Consideration
- D. Outcome of the study

ANSWER: C

150. A human nutrition graduate student wants to assess the prevalence of goiter among reproductive-age women in a district. Accordingly, the student obtained new comments from his advisor to keep the logical flow of the questions in his first proposal.

What is the most likely comment given in the above proposal development step?

- A. Put background questions last
- B. Put sensitive questions last
- C. Put stigma-related questions first
- D. Put controversial questions first

ANSWER: B

151. A researcher wanted to assess the magnitude of factors associated with good breastfeeding practice in a city. In the literature review, the researcher noted that, as compared with the national plan, poor breastfeeding practice was observed in the city where the proposed research would be conducted.

Which proposal development step was indicated in the above scenario?

- A. The significance of the study
- B. The statement of the problem
- C. The sampling procedure of the study
- D. The data quality assurance

ANSWER: B

152. A researcher wants to provide information on the frequently observed epidemics of measles in a district. Hence, the researcher constructed his achievement as an incidence of measles among communities living in a district in 2025.

What is the most appropriate objective for the above proposal development step?

- A. To assess the incidence and predictors
- B. To assess prevalence and associated factors
- C. To assess incidence and prevalence
- D. To determine the incidence

ANSWER: D

153. A researcher conducted a study to determine the prevalence of anemia among reproductive-age women who visited public hospitals. The data were collected through history taking from the women in person, and additional explanation was provided to them when they were not clear with the questions.

What is the most likely data collection technique applied in this case?

- A. Observation
- B. Mailed questionnaire
- C. Face-to-face interview
- D. Focus group discussions

ANSWER: C

154. A graduate student of human nutrition wants to study the perception of colostrum feeding among elders in a district. The student calculated a sample size of eighteen and plans to collect the data through exploring their attitude and feeling and reported it back to him.

What is the most appropriate data collection method that will be applied in this case?

- A. In-depth Interview
- B. Observation
- C. Self-Administered Questionnaire
- D. Mail-based questionnaire

ANSWER: A

155. A student wants to show the relationship of anemia and weight of school children in a country. Accordingly, the student was commented on to consider graphs to present the relationship of anemia and weight together with texts and tables, and the student amended his proposal accordingly.

What is the most likely data presentation method that should be applied in this case?

- A. Bar
- B. Pie
- C. Graph
- D. Scatter plot

ANSWER: D

156. A human nutrition researcher wanted to determine the knowledge and associated factors of minimum dietary intake among pregnant women attending antenatal care follow-up in a hospital. Accordingly, the student suggested incorporating age, religious affiliation, monthly income, and educational status of the study participants to strengthen the first proposal to be convincing.

What is the appropriate suggested independent variable in the above proposal development step?

- A. Knowledge
- B. Educational status
- C. Socio-demographic factors
- D. Past medical history

ANSWER: B

157. A human nutrition graduate student presented his research findings on the prevalence of rickets among under-five children in a district. The study finding showed that 15 percent had rickets. Furthermore, this was significantly associated with lack of sunlight exposure in the early childhood period.

What is the appropriate interpretation of the information given in this case?

- A. 15 out of 100 children developed rickets.
- B. 75 out of 100 children developed rickets.
- C. 85 out of 100 children developed rickets
- D. 65 out of 100 children developed rickets.

ANSWER: A

158. A department head of human nutrition requested the registrar office to provide the cumulative grade report of graduate students of the department for his grade audit. Accordingly, the registrar reported the 6 students' grades as 2.2, 2.8, 2.6, 3.4, 2.3, and 3.7.

What is the appropriate data interpretation in this case?

- A. Most of the students' grades are greater than 3.0.
- B. Most of the students' grades are less than 3.0.
- C. The majority of the students' grades are equal to 3.0.
- D. Half of the students' grades are less than 3.0.

ANSWER: D

159. A 34-year-old woman presented to a clinical nutritionist with complaints of easy fatigability and lightheadedness. The physical examination revealed that she was pale. The clinician decided to put her on Hem-Up syrup for 6 months; however, the patient did not make good progress, and the clinician decided to use the recent information provided by the researcher.

Which evidence-based practice was applied in the above scenario?

- A. Best available evidence
- B. Clinical practice
- C. Patient value
- D. Patient preference

ANSWER: A

160. A 45-year-old woman presented to a clinical nutritionist with complaints of easy fatigability and lightheadedness. The physical examination revealed pale conjunctiva, and the clinician then decided to put her on Hem-Up syrup for 6 months, but the patient wanted to take a liver-rich diet to promote weight loss based on the information she heard that the level of hemoglobin will increase by 1 unit per 1 pound intake of liver. Accordingly, the nutritionist began to download all available evidence and evaluated the evidence based on the predefined checklist.

What is the most appropriate step of evidence-based practice the clinician should do?

- A. Searching
- B. Appraising
- C. Asking
- D. Acting

ANSWER: B

161. A clinical nutritionist challenged by a patient having poor progress of anemia after being treated by Hem-Up for the last two weeks. Then, the clinician decided to search the best available evidence online and got many primary articles done on it, which made him confused about utilization. Finally, his team decided to use the best article over others based on the standard.

What is the most likely study design to utilize evidence in the above scenario?

- A. Cross-sectional
- B. Case-Control
- C. Experimental
- D. Cohort

ANSWER: C

162. A human nutrition researcher wanted to determine the new onset and predictors of goiter among pregnant women who had antenatal care follow-up at a given hospital. Hence, the researcher developed his proposal and submitted it to the funding sector. Accordingly, the sector returned the proposal to the researcher since they had doubt about the achievement of the proposal.

What is the appropriate objective of the above proposal development step?

- A. Incidence and predictors of goiter
- B. Prevalence and factors of goiter
- C. Prevalence of goiter
- D. Incidence of goiter

ANSWER: A

163. A human nutrition researcher intended to conduct a study on the prevalence of calcium deficiency among adult men at a hospital. The researcher planned to enroll the study participants randomly and found that the study participants were from 18 to 64 years old in his pre-assessment study. Accordingly, the researcher considered using a descriptive summary measure that handled the effect of the outliers.

What is the most appropriate summary measure in the above proposal development step?

- A. Mean
 - B. Median
 - C. Mode
 - D. Range
- ANSWER: B

164. A nutrition professional working in a health center went to the community to conduct weekly supervision. The professional found that there was a significant problem with diversifying food in that community. She wanted to communicate to the community on how to solve the problem.

What is the most likely first role of the nutritionist to solve this problem?

- A. Thinking of the behavior that could promote mothers' preference.
- B. Finding reasons for the mother's behaviors on the feeding problem.
- C. Encouraging mothers to choose what is best suited to their preference.
- D. Promote mothers to stick to cultural feeding practices.

ANSWER: B

165. A senior health education professional working in the regional health office went to the zonal health office for annual supervision. He met the health education experts in the zone and gave an orientation on how to bring behavioral change. During his orientation he emphasized that "the health educators should find out the real health needs of people before they plan a health education session."

What is the most likely principle of SBCC emphasized by senior health education professionals?

- A. Principle of reinforcement
- B. Principle of interest
- C. Principle of participation
- D. Principle of comprehension

ANSWER: B

166. A senior nutrition expert working in a certain health center conducted a home visit in a particular community to evaluate the effectiveness of the health education that was provided by health extension workers. On his visit he observed that the mothers started preparing complementary food by using different cereals that were believed to be diversified. The senior expert asked a mother for responses on certain issues to confirm the learning objective was already achieved or not.

What is the most likely principle of SBCC addressed in this case?

- A. Principle of leadership act
- B. Principle of good human relations
- C. Principle of feedback to issues
- D. Principle of Successive Experience

ANSWER: C

167. A health education expert in a health center called a nurse working in the under-five unit for orientation on how to counsel mothers on maternal and child feeding practices to bring behavioral change. During an orientation he told a nurse, "Health education should start from the existing culture and try to change habits and practices gradually."

What is the most likely principle of SBCC addressed in this case?

- A. Health education efforts should be ambitious for change.
- B. Health education efforts should aim at small changes in a graded fashion.
- C. The effort of health education should be made for mothers to learn all together.
- D. Health education should provide opportunities for mastering changed practices.

ANSWER: B

168. An environmental health professional working in a zonal hospital plans to give orientation to all cleaners of the hospital on the management of hazardous healthcare waste. During orientation, the expert explains all the steps in the management of hazardous waste with detailed explanations, emphasizing the process of separating different categories of waste by using different color-coded waste bags or bins.

What is the most appropriate hazardous healthcare waste management step to be employed in this case?

- A. Recycling and reuse of healthcare waste
- B. Respect waste handling technique
- C. Segregation of healthcare waste
- D. Act on waste minimization

ANSWER: C

169. A 26-year-old boy came to a clinic with the chief complaint of bloody diarrhea of a month's duration. He has a history of swimming in stagnant water and became weak. Moreover, the laboratory investigation result indicated that a schistosomiasis species was found.

What is the most likely prevention mechanism to be employed for this case?

- A. Proper site selection of house
- B. Decrease needs to visit breeding site
- C. Decrease needs to visit breeding site
- D. Improve water quality and control snail population

ANSWER: D

170. An environmental health expert of a health center went for monthly supervision to his catchment area. He observed most of the households were using public latrines for excreta disposal. The expert recommended that the community should construct the cheapest type of excreta disposal system, which could be constructed using local materials and local skills. It was also considered the most common sanitation system in areas where there was no adequate water supply.

What is the most appropriate type of latrine to be recommended by an expert?

- A. Pit latrine
- B. VIP latrine
- C. Trench latrine
- D. Borehole latrine

ANSWER: A

171. An environmental health expert of a woreda was invited by the management of the cement factory existing in his locality to provide health education on environmental hazard management. The expert focused his lesson as follows: a hazard is something that is harmful to our health and further identifies the type of hazard. The expert also discussed different steps in hazard management. What is the most likely primary step to be implemented during hazard management?
- A. Hazard/risk analysis and evaluation
 - B. Establish the context and identify the hazard
 - C. Communicate and consult
 - D. Treat the hazard/risk
- ANSWER: B
172. A nutrition professional, who was working in the woreda health office, was invited to provide health education on food safety for food handlers from nearby hotels. When the professional provided a health education, he stated that delays in eating prepared food can lead to the growth and reproduction of microorganisms in the foodstuff. What is the most likely principle of food safety stated in this case?
- A. Eat cooked foods immediately after they get prepared.
 - B. Avoid contact between raw and cooked food.
 - C. Keep all kitchen surfaces and utensils meticulously clean.
 - D. Use safe water in food preparation and for washing fruits and vegetables.
- ANSWER: A
173. A nutritionist and an environmental health expert working in the woreda health office were invited by an investor, who is working in dietary production, to provide training about milk hygiene to his workers. The training incorporated modules such as application of hazard analysis and critical control point (HACCP) and income material requirements. What principle of food safety was addressed by health professionals in their training?
- A. Establishment design and facility
 - B. Control of operations
 - C. Personal hygiene
 - D. Production information and consumer awareness
- ANSWER: B
174. A health team working in the woreda health office conducted a weekly inspection in selected establishments that were located in their woreda. During an inspection the team found that the food handlers were chewing khat (chat) in the food preparation area. The health team took measures on the hotel as a result of the violation of safety regulations. What principle of food safety was violated by the hotel?
- A. Establishment design and facility
 - B. Control of operations
 - C. Personal hygiene
 - D. Production information and consumer awareness
- ANSWER: C

175. A senior nutrition professional, who was working at the woreda health office, called food handlers to his office to provide orientation on food safety. He said that food can be at risk for contamination while it is being prepared. Therefore, food handlers should wear a white gown and cape during food handling.

What is the most likely principle of food safety that was emphasized by nutritionists?

- A. Establishment design and facility
- B. Control of operations
- C. Personal hygiene
- D. Production information and consumer awareness

ANSWER: C

176. A 25-year-old man came to a hospital with complaints of abdominal cramps, headache, and vomiting. After laboratory investigation, the patient was diagnosed with food intoxication. The case was referred to the nutrition counseling unit after appropriate treatment, and then the counselor provided detailed information on possible factors that could cause food poisoning for this patient.

What is the most likely cause for the patient's illness that could be discussed during the patient's counseling?

- A. Poor personal hygiene
- B. Time-temperature abuse
- C. Consumption of poisonous plants
- D. Food prepared using contaminated water

ANSWER: C

177. According to the annual report of a health facility, waterborne diseases were among the top 10 diseases. Based on this information, the health facility planned to conduct a community health education, which focused on the prevention and control of waterborne diseases.

Which disease prevalence can be decreased after giving the health education in this case?

- A. Trachoma
- B. Malaria
- C. Schistosomiasis
- D. Cholera

ANSWER: D

178. A 47-year-old woman came to a health center with complaints of fever, abdominal pain, vomiting, and diarrhea. A health officer referred the woman to a nutritionist for counseling after conducting an appropriate examination and prescribed medication. Based on that, a nutritionist provided counseling on food storing techniques, such as avoiding placing cooked foods on plates that previously were contaminated with raw meat and eggs and keeping raw meat, poultry, and eggs away from other foods, grocery bags, and refrigerators.

What is the most likely food safety principle that the nutritionist provided for this woman?

- A. Canning
- B. Chilling
- C. Separating
- D. Blanching

ANSWER: C

179. According to a report of a woreda health office, the prevalence of schistosomiasis is very high due to open defecation and practices of the community washing in the river. To reduce the prevalence, the woreda health office organized a health education session in the public markets by using amplifiers.

What is the most likely water-associated disease addressed by the health education in this case?

- A. Water-washed diseases
- B. Water-based diseases
- C. Waterborne diseases
- D. Water-related diseases

ANSWER: B

180. A nutritionist, who was working in a particular village, found that the prevalence of foodborne diseases was very high. The organization for which the nutritionist was working assigned him to provide health education for this community to reduce the prevalence of foodborne diseases.

What is the most likely prevention mechanism to be provided for this community?

- A. Exclude food handlers with abscesses and purulent lesions.
- B. Wearing gloves when handling food
- C. Avoiding home canning and other food preservation techniques
- D. Avoid eating canned foods in damaged containers.

ANSWER: D

181. A 40-year-old female patient came to a health center with complaints of a two-day history of fever, chills, sweating, myalgia, arthralgia, headaches, anorexia, weight loss, and dry cough. After additional physical and laboratory examinations, the patient was diagnosed with brucellosis.

What is the most likely prevention mechanism to be applied for this case?

- A. Proper handling and storage of leftover foods
- B. Remove containers that can hold water.
- C. Ventilating the premises with natural openings
- D. Do not drink raw milk and milk products.

ANSWER: A

182. An eight-year-old female patient brought to a hospital with complaints of fever, malaise, anorexia, nausea, abdominal discomfort, and jaundice. Laboratory examination confirmed that the patient was affected with the hepatitis A virus.

What is the most likely prevention mechanism to be provided for this patient?

- A. Treatment of infections in food handlers such as skin and throat infections
- B. Recognizing, preventing, and controlling infections in domestic animals and pets
- C. Good sanitation, personal hygiene, hand washing, and proper disposal of feces
- D. Educate on the careful storage and use of chemicals (their storage away from foods)

ANSWER: C

183. A 10-year-old girl was brought to a hospital with a complaint of abdominal pain, diarrhea, tiredness, nausea, and vomiting. After a thorough physical and laboratory examination, the physician diagnosed the problem of the patient as trichuriasis. Finally, he referred to the nutritionist's office for health education.

What is the most likely prevention mechanism to be employed for this patient?

- A. Washing vegetables and fruits contaminated with soil before use.
- B. Improvement of working conditions for handlers of animal products.
- C. Disinfect articles in contact with the patient and soiled by discharges of the patient.
- D. Control cats and prevent them from gaining access to sandboxes and sand piles.

ANSWER: A

184. A nutritionist planned a nutrition education session for a group of athletes. The topic of the session was the number of total calories recommended for active athletes. The nutritionist intended to address the calories recommended from each macronutrient.

What is the most appropriate nutrition education to be provided in this scenario?

- A. Athletes should consume 40-55 percent of their calories from mixed nutrients.
- B. Athletes should consume 65-75 percent of their calories from carbohydrates.
- C. Athletes should consume 15-30 percent of their calories from proteins.
- D. Athletes should consume 30-35 percent of their calories from fats.

ANSWER: B

185. Many studies indicated that the adolescent female athletes, female athletes with heavy menstrual losses, athletes with meat-restricted diets, and distance runners who may have increased gastrointestinal bleeding suffered from micronutrient deficiencies.

What is the most likely nutrition education to be provided for such an athlete?

- A. Vitamin K supplementation
- B. Vitamin C supplementation
- C. Iron supplementation
- D. Iodine supplementation

ANSWER: C

186. A nutrition professional was providing counseling to a 25-year-old male endurance athlete with a weight of 60 kg on the recommended daily allowance of protein to meet the energy demand.

What is the recommended daily allowance of protein in this case?

- A. 72 to 84 g protein/day
- B. 48 to 56 g protein/day
- C. 58 to 66 g protein/day
- D. 48 to 66 g protein/day

ANSWER: A

187. A 70 kg male athlete, who had intensive and repetitive training, came to a nutritionist looking for nutritional counseling on the energy requirement to carry out the training.

What is the appropriate nutritional counseling to be provided for this athlete?

- A. CHO intake of 490-700 g/day
- B. CHO intake of 450-550 g/day
- C. Protein intake of 80-100 g/day
- D. Protein intake of 80-90 g/day

ANSWER: A

188. An active athlete came to a health facility and asked a nutritionist about the recommended intake of fat from total energy intake. Then, the nutritionist told him the recommended percentage of fat per day from total calorie intake.

What is the most appropriate nutritional counseling to be provided on daily intake of fat?

- A. To consume 30 percent or more of total energy intake
- B. To consume 30 percent or less of total energy intake
- C. To consume 35 percent or more of total energy intake
- D. To consume 35 percent or less of total energy intake

ANSWER: B

189. A health extension worker has a weekly health education plan on food hygiene and safety principles. During the provision of health education, she stated that the mothers should cover their hair and wear a clean cloth while they prepare diversified food for their children.

What principle of food safety was addressed by the health extension worker in her education?

- A. Establishment design and facility
- B. Control of operations
- C. Personal hygiene
- D. Production information and consumer awareness

ANSWER: C

190. A middle-level manager of a hospital develops a work plan that specifies what lower-level units within each division must do, how they do it, and who is in charge at each level. The plan will be used for one year and helps to achieve short-term goals.

What is the most likely type of plan developed by the manager in this case?

- A. Operational plan
- B. Tactical plan
- C. Strategic plan
- D. Contingency plan

ANSWER: B

191. Reform activities are initiated by a manager of the hospital to enhance work efficiency and employee satisfaction. The approach followed by the manager was to arrange employees' positions in a way that more senior health professionals were assigned to more comfortable corner offices and junior ones at the front desks despite their activities. After a while he started to observe a chaotic movement of clients from one office to the other and a delay in the accomplishment of some tasks.

What is the appropriate management principle recommended for the above problem?

- A. Convergence of Work
- B. Shortest decision path
- C. Management by objectives
- D. Function determines structure

ANSWER: D

192. A nutritionist working in a primary hospital approached his manager to talk about his frustration with the work assignment in the hospital. The nutritionist believes that he is under-serving the hospital because of his current ward assignment. He feels overburdened by the number of tasks assigned to him and by the fact that he is not using his specialization, which puts him under lots of pressure to carry out his tasks.

What is the most appropriate organizing action addressing the above concern?

- A. Decentralize nutritionist tasks to others
- B. Review scheme of division of labor
- C. Delegate some of the duties to others.
- D. Coordination of activities in the ward

ANSWER: B

193. An assessment conducted on the productivity of a health facility, which showed major gaps in the areas of customer service and efficient use of resources, which led to low productivity. To address the problem, the facility administration changed the organization design by regrouping similar activities, expertise, and resources within single work units.

What is the most likely organizational design implemented in this case?

- A. Team structure
- B. Functional structure
- C. Matrix structure
- D. Divisional structure

ANSWER: B

194. A referral hospital in a capital city had employees from other nations and cultural backgrounds, which was unique from the rest of the staff. Recently, the manager of the hospital started to observe disagreement among a group of professionals working in the hospital, which resulted from conflicting values.

What is the most appropriate training type recommended for this case?

- A. Interpersonal skills training
- B. Diversity training
- C. Basic literacy training
- D. Problem-solving training

ANSWER: B

195. A leader of a nutrition program decided to develop a team that would constitute members who would diagnose nutritional problems in the community from different perspectives, devise a strategy to address those problems, search for sources of funding, monitor and evaluate the progress, and promote the activities and results to the community and stakeholders.

What is the most appropriate team to conduct the above task?

- A. Cross-functional team
- B. Functional team
- C. Self-managed team
- D. Project team

ANSWER: A

196. A primary hospital administrator wanted to motivate staff in the organization. The employees complained about low salary, uncomfortable work conditions, bad policies, poor relations with supervisors, lack of recognition and reward for achievement, and slow advancement. Based on the complaint, the administrator increased the salary, revised policies, improved working conditions, and maintained good relations with supervisors. But the administrator did not observe that much change in the motivation level of the employees.

What is the most appropriate motivation strategy for the above employees?

- A. Further increase salary
- B. Rewarding high achievers
- C. Renovate the workspace
- D. Firing bad supervisors

ANSWER: B

197. A nutrition expert in a referral hospital wanted to deliver a message regarding the importance of iodization of salt to all health professionals working in the hospital. The expert wanted to select an effective communication type by considering the simplicity of the message and size of the audience. What is the most appropriate communication type helpful to the expert in this case?

- A. Organizational communication
- B. Interpersonal communication
- C. Mass communication
- D. Intrapersonal communication

ANSWER: C

198. A Leader of a private hospital was known for her easy-going attitude by her followers and she gave all of the decision-making power to the employee with minimal supervision. She exerted little control over her coworkers, leaving them to sort out their roles and tackle their work assignments on their own. Annual review meeting revealed that the hospital was inefficient in its human and other resource management, lost significant number of customers and almost at blink of bankruptcy.

What is the most likely leadership style followed by the leader in this case?

- A. Autocratic
- B. Democratic
- C. Laissez-faire
- D. Participative

ANSWER: C

199. A team leader was assigned to lead a team of nutrition experts who work on national nutrition program within a non-governmental organization. The expert teams were highly experienced in the area, capable of accomplishing their task and had high level of readiness to take every responsibility. What is the most appropriate style to lead the above team of experts?

- A. Delegating
- B. Selling
- C. Telling
- D. Participating

ANSWER: A

200. Clients and health professionals of a primary hospital were asked about their satisfaction with regard to service given in the hospital. The clients reported they were satisfied by the care they received in the hospital and they felt the leaders of the hospital considered them as their priorities. They expressed the health care professionals as compassionate and caring ones. Similarly, the health professionals mentioned that leaders of the hospital were supportive, collaborative and put their followers at front.

What is the most likely leadership style shown in the above case?

- A. Servant
- B. Transactional
- C. Democratic
- D. Charismatic

ANSWER: A