

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION (EHPLE) PRACTICE TEST BOOKLET 2025

PHYSIOTHERAPY



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH - ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, Leadership and Management, Ethics, and Health Promotion as outlined in the Information Booklet 2025.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Physiotherapy

1. A 60-year-old male patient with cervical radiculopathy of seven years duration following car accident came to physiotherapy OPD. On assessment, the patient reported limitation of multidirectional movements, severe pain that radiates to arms, muscle weakness, and difficulty of grasping objects. What is the most appropriate objective examination that has to be performed in the neck?

A. Checking hypomobility
 B. Checking hypermobility
 C. Checking reflex
 D. Checking sensation

ANSWER: A

2. A 55-year-old male patient, who has been admitted and treated at surgical ward after he sustained mild injury of neck region following car accident three years ago, came to physiotherapy clinic. On examination, the physiotherapist identified neck pain, bilateral upper extremities pain, numbness and tingling along with report of loss of balance and coordination, lack of fine motor skills and awkward gait.

What is the most likely diagnosis for this case?

A. Cervical radiculopathy
 B. Cervical myelopathy
 C. Myofascial neck pain
 D. Whiplash injury

ANSWER: B

3. A 30-year-old male patient came to a hospital with temporomandibular joint dysfunction following a motor bike accident two days ago. He was treated immediately at emergency unit with manipulation, antibiotic and anti-pain. After two weeks of appointment, he was referred to a physiotherapy clinic for physiotherapy intervention. During physical examination, the physiotherapist noticed that the patient cannot open his jaw.

What is the most appropriate long term goal of treatment in this case?

A. Reduce pain
 B. Restore range of motion
 C. Restore posture
 D. Restore normal functional activities

ANSWER: D

4. A 55-year-old female patient came to physiotherapy clinic with a complaint of temporomandibular joint dysfunction. During physical examination, the physiotherapist noticed that the patient could not open her jaw.

What is the most appropriate goal of treatment for this case?

A. Strengthening of facial muscles
 B. Restoring jaw opening and closing activities
 C. Restoring range of motion of neck
 D. Strengthening of neck and shoulder muscles

ANSWER: B

5. A 36-year-old male patient came to physiotherapy OPD with a complaint of chronic neck pain radiating to the forehead which is aggravated by neck flexion and rotation movements. On examination, tightness of neck muscles, and tingling and burning sensation were noticed. And he has undergone regular physiotherapy intervention for the past five weeks but pain could not be improved. What is the next most appropriate step in this patient's care?

- A. Prescribe strong anti-pain
- B. Refer or request other investigations
- C. Reassess and modify treatment plan
- D. Continue with the same treatment plan

ANSWER: B

6. A 45-year-old female patient came to a hospital with a complaint of severe lower back pain that radiates to the back of her legs. The pain is aggravated by standing for long period of time and lifting heavy objects. MRI revealed L4-5 disc protrusion with L-5 nerve compression.

What is the most appropriate first action to be taken by the physiotherapist in this case?

- A. Assessment of range of motion
- B. Take appropriate patient history
- C. Assessment of muscle power
- D. Design treatment plan

ANSWER: B

7. A 50-year-old male patient came to a physiotherapy clinic with a complaint of pain on the pelvic and sacroiliac joint. On examination, tenderness was noticed on both sacroiliac joints was noticed. Then, he was diagnosed with sacroiliac joint dysfunction.

What is the most commonly used special test to examine this case?

- A. Patric's test
- B. Straight leg raise test
- C. Phalen's test
- D. Apprehension test

ANSWER: A

8. A 65-year-old male patient came to a hospital with severe lower back pain characterized by sharp and shooting radiating type of pain down to the back of legs. On examination, the physiotherapist found that straight leg rising is positive.

What is the most likely diagnosis of this case?

- A. Osteoporosis
- B. Sciatica
- C. Lordosis
- D. Scoliosis

ANSWER: B

9. A 60-year-old female patient came to physiotherapy clinic with a complaint of back pain following postmenopausal osteoporosis. Due to the osteoporosis, anterior wedge fractures occurred to several vertebrae, in the upper to the midthoracic spine, causing structural scoliosis that contributed to decrease in height.

What is the most likely diagnosis of this type of kyphosis?

- A. Flat back
- B. Hump back
- C. Dowager's hump back
- D. Round back

ANSWER: C

10. A 55-year-old patient came to a physiotherapy clinic with acute low back pain radiating to both of his lower limbs following lifting heavy objects. On examination, tenderness on lumbar region was noticed.

What is the appropriate short term goal for this case?

- A. To strengthen core muscles
- B. To alleviate pain
- C. To enhance cardiovascular
- D. To restore range of motion

ANSWER: B

11. A 59-year-old female patient came to a physiotherapy clinic with a complaint of sharp lower back and left leg pain radiating to her left knee for more than six years. On examination, tenderness in bilateral lumbar, para-spinal and left gluteal muscles was noticed.

What is the most appropriate treatment goal in this case?

- A. Develop appropriate individual program at patient's gym including aerobic and strengthening components
- B. Education and instruction in home exercise program and appropriate biomechanics at work and home
- C. Achieve functional ROM of hips and lumbar spine, including full ability to squat and return to neutral
- D. Neuromuscular re-education and proprioception training for neutral spine in sitting and standing

ANSWER: C

12. A 55-year-old woman, who has seven children, came to neurosurgery OPD with a complaint of chronic lower back pain and urine incontinence. The neurosurgeon then consulted for physiotherapy intervention after excluding red flags. On examination, the physiotherapist found tenderness on lumbar region and weakness of the pelvic floor muscles.

What is the most appropriate initial treatment for this case?

- A. Lumbar traction
- B. Fitness exercise
- C. Kegel's exercise
- D. Cervical traction

ANSWER: C

13. A 45-year-old female patient came to a hospital with a complaint of chronic lower back pain that radiates to the back of her legs. The pain is aggravated by standing for long period of time and lifting heavy objects. On evaluation, there is tenderness on the lumbar region and disc protrusion with L-5 nerve compression which is confirmed by MRI. There was no improvement with physiotherapy interventions and NSAIDS.

What is the most appropriate next action in this case?

- A. Core stability exercise
- B. Refer for further investigation
- C. Prescribe strong anti-pain
- D. Strengthening exercise

ANSWER: B

14. A 45-year-old male patient came to a physiotherapy clinic with the complaint of pain on the left shoulder after he sustained fall down injury from a building two months ago. After examination, the physiotherapist confirmed that the diagnosis was rotator cuff syndrome. What is the most likely factor that affects the diagnosis this case?
- A. Impingement syndrome
 - B. Shoulder dislocation
 - C. Scapular impairment
 - D. Adhesive capsulitis
- ANSWER: B
15. A 35-year-old office worker came to a hospital with complaints of chronic pain and stiffness of the hand and wrist. The pain worsens during computer typing. On examination, there was pain and swelling on the wrist joint. A diagnosis of carpal tunnel syndrome was made. What is the most appropriate special test to be used in this case?
- A. Tinel sign test
 - B. Allen's test
 - C. Phalen's test
 - D. Froment paper sign test
- ANSWER: C
16. A one and half-year-old girl was brought to a physiotherapy OPD with a complaint of developmental delay. The physiotherapist wants to assess the child to provide a definitive diagnosis and plan of treatment. What are the optimal assessment parameters to be used to determine the child's diagnosis?
- A. Range of motion assessment, and sensory, speech, and fine motor skill evaluation
 - B. Observation for deformity, muscle power, muscle length and cognitive assessment
 - C. Gross motor, fine motor, speech and comprehensive evaluation of functional abilities
 - D. Observation for deformity, manual muscle testing, tone and sensory assessment
- ANSWER: C
17. A pediatric physiotherapist performing physical examination and postural analysis on a 12-year-old girl who had idiopathic scoliosis found out a classic rib hump on the right thoracic spine during forward bend test. The physiotherapist also noticed a lateral thoracic curvature with concavity to the left and a slight compensatory tilting in the lumbar region with concavity to the right. There was also shortening of paraspinal muscles in the left thoracic and right quadrates lumborum. What is the best possible diagnosis for this child?
- A. Left thoracic right lumbar structural scoliosis
 - B. Right thoracic left lumbar non structural scoliosis
 - C. Left thoracic right lumbar non structural scoliosis
 - D. Right thoracic left lumbar structural scoliosis
- ANSWER: D
18. A physiotherapist evaluating a 16-year-old girl with spinal deformity found right thoracic positional scoliosis. The main concern of the girl is the abnormal spine posture which she is ashamed of especially at school. The physiotherapist designing a treatment plan wants to prioritize the girl's concern. What are the most likely treatment techniques that should be used to achieve the goal?
- A. Stretching, strength training for trunk muscles and correction exercises
 - B. TENS, infrared, stretching exercises and strengthening for back musculature
 - C. US, Stretching the left trunk muscles and overall back strengthening program
 - D. TENS, therapeutic US, soft tissue release techniques and stretching exercises
- ANSWER: A

19. A man brought his four-year-old child who just started walking with a complaint of developmental delay in his motor skills. The physiotherapist did a thorough subjective and objective examination and diagnosed a motor developmental delay.

What subjective examination best determines the plan of treatment in this case?

- A. Timing of major milestones
- B. History of child delivery
- C. Sex of the child
- D. Age of the child

ANSWER: A

20. A five-year-old boy was brought to a hospital with a complaint of fall down injury to his forearm while jumping on the sofa. The orthopedician made a diagnosis of stable radio-ulnar shaft fracture and kept him on a circular pop cast that runs from elbow to mid-metacarpals. The consulting physiotherapist who assessed the boy after a week performs an objective assessment.

What is the most appropriate objective examination that can be performed in this case?

- A. Observation for digits swelling, ROM of the elbow and shoulder joints
- B. ROM for the elbow, muscle power for supinators and pronators, observation
- C. Elbow and wrist ROM, muscle power for elbow and wrist flexors
- D. Elbow, wrist, and metacarpals ROM, and sensory assessment for the forearm

ANSWER: A

21. A one-month-old male infant was referred to a physiotherapy clinic by a neurologist since he was diagnosed with obstetric brachial plexus palsy. The neonate prefers to keep his right hand in a fixed position and seems to neglect his right upper extremity. The physical therapist performed a thorough physical assessment and planned for intervention accordingly.

What is the most likely clinical presentation of the child's upper extremity?

- A. Shoulder adduction internal rotation, elbow flexion, forearm pronation, wrist and digits flexion
- B. Shoulder adduction internal rotation, elbow extension, forearm pronation, wrist and digits flexion
- C. Shoulder flexion abduction, elbow flexion, forearm pronation, wrist and digits flexion
- D. Shoulder abduction external rotation, elbow flexion, forearm supination, wrist and digits flexion

ANSWER: B

22. A five-year-old boy fall over boiling water which caused a burn injury to his chest and both upper extremities while roaming in the room. His father immediately grabbed him and took him to a hospital. The child was diagnosed with 2nd degree superficial partial thickness burn with a total body surface area of 18 percent. Active wound covering the chest, anterior aspect of the arms and the axillary area was noted on examination. The medical team admits the child for follow-up treatment performing proper diagnosis and wound care management. The physiotherapy team evaluated the child and planned a treatment.

What is the most appropriate treatment plan for day 1?

- A. Positioning for contracture avoidance and chest exercise education
- B. Stretching exercises for anterior muscle groups and slight strength training
- C. Chest exercises and cast application with elbow in extension to avoid contracture
- D. Stretching exercises, strength training and infrared treatment

ANSWER: A

23. A nine-year-old boy diagnosed with duchenne muscular dystrophy with marked joint contracture, weakness of leg and trunk muscles, impaired pulmonary function and functional ability was brought to a physiotherapy clinic for his follow-up. After performing reassessment, the physiotherapist designed a SMART treatment goal.

What is the most likely treatment goal that fits for this child?

- A. Correcting of contracture with casting
- B. Improving muscle strength to grade 5/5
- C. Treating pulmonary capacity until normal
- D. Preventing deformity and maintaining functioning

ANSWER: D

24. A 13-year-old boy was diagnosed with posterior cruciate ligament injury of the right knee. The boy plays soccer representing his school; while on training, he fall down with flexed knees. The boy has been following physical therapy for the past four months and wants to play a match in the coming week. The physical therapist planned to evaluate the child if he is in a position to play the upcoming match.

What functional outcome most likely predicts his readiness to return to the match?

- A. Full knee ROM
- B. Quadriceps strength
- C. Hamstring muscle length
- D. MRI finding of the PCL ligament

ANSWER: B

25. A 54-year-old male patient has been brought to a physiotherapy clinic with a complaint of weakness of the right side of the body and loss of balance secondary to lesion involving the cerebellum. After an intensive physiotherapy follow-up for the last three months, he is able to walk but now his main concern is the problem with maintaining standing balance. The physiotherapist wants to assess and evaluate his balance.

What is the most appropriate outcome measure that determines the balance of this patient?

- A. Timed up and go test
- B. Rivermead motor assessment scale
- C. Fatigue severity scale
- D. Visual analogue scale

ANSWER: A

26. A 35-year-old man was admitted to a medical ward with a complaint of severe headache which is not relieved by pain reducing medications. Upon chart review, the physiotherapist found that the patient was admitted eight hours ago and he has pain and stiffness in his neck that started suddenly with the headache. Laboratory report showed, increased WBC count, ESR and C-reactive protein. Lumbar puncture has been taken and results are not ready yet.

What is the best diagnosis of this patient?

- A. Brain abscess
- B. Encephalitis
- C. Brain tumor
- D. Meningitis

ANSWER: D

27. A 48-year-old patient diagnosed with post poliomyelitis syndrome was referred to a physiotherapist for a follow-up evaluation and management. The patient complained of fatigue which commonly occurs in the afternoon, weakness in the right lower limb and breathing difficulties.

What are the best exercises to treat the patient's condition?

- A. High intensity resistance training for the affected lower limb and chest exercises
- B. Chest exercises, TENS for the affected lower limb, and functional re-education
- C. Motor control re-education, sensory re-education, and chest exercise
- D. Gait training, provision of ankle foot orthosis and lumbar support

ANSWER: C

28. A 34-year-old male patient with bacterial meningitis post TBI was brought to a physiotherapy clinic with complaints of muscle weakness and limitation of neck ROM after a period of hospitalization. The physiotherapist did a thorough assessment and designed patient centred treatment plan.

What is the most appropriate subjective question that involves the patient in the treatment planning?

- A. Does your movement limitation affect your daily activities?
- B. Does your weakness interfere with your daily activities?
- C. Have you been to a physiotherapist before?
- D. What is your specific goal?

ANSWER: D

29. A 45-year-old woman presented to a physiotherapy clinic with a complaint of pain and paresthesia in her wrist and hand which worsens during the night and when typing on a computer. The pain started few months back as simple paresthesia in the hand and worsened over time. Physiotherapy examination findings show atrophy of the thenar eminence, sparing of sensation over the thenar eminence and positive tincl and phalen signs.

What is the most likely diagnosis of this case?

- A. Ulnar nerve palsy
- B. Radial nerve palsy
- C. Carpal tunnel syndrome
- D. Cervical radiculopathy

ANSWER: C

30. A 24-year-old male patient who sustained traumatic brain injury was brought to a physical therapist for rehabilitation. The patient was so desperate and he wishes at least he could help his family during diaper changing. The physiotherapy evaluation shows a grade 1+ muscle power in all upper limb muscles and 2- in lower limb muscle groups. The patient has both urinary and faecal incontinence. The physiotherapist sets rehabilitation goal while prioritizing the patient's expectation.

What is the most appropriate type of exercise to address this goal?

- A. Sensory re-education and biofeedback
- B. Bridging and trunk exercises in hook lying
- C. PNF techniques for upper and lower extremities
- D. Resisted exercises for upper and lower extremities

ANSWER: B

31. A 24-year-old ambulating man was brought to a physiotherapy clinic with a complaint of weakness in the legs. Few months ago he had sustained a car crash and was hospitalized for two months. He mentioned that the accident was bad and he initially could not move his legs. Currently, he has almost recovered fully except for the difficulty in stair climbing. A physiotherapist sets a plan to treat the weakness.

What is the best strength training that addresses his problems?

- A. Proprioceptive neuromuscular facilitation techniques
- B. Resistance training for quads and hamstrings
- C. Trunk exercises using physioball
- D. Tandom walking exercises

ANSWER: B

32. A patient was brought to a physiotherapy clinic with a complaint of progressive muscle weakness which started at the upper and lower limbs gradually resulted in difficulty of sitting and standing. The information documented in the medical chart indicated that the condition seems amyotrophic lateral sclerosis characterized by progressive degeneration of motor neurones. Upon subjective assessment, the physiotherapist was asking about the mechanism of onset, duration of onset and past medical history of the patient.

What is the most likely additional information needed to identify this case?

- A. Gender and age of the patient
- B. Social history of the patient
- C. Drug history of the patient
- D. Occupation of the patient

ANSWER: A

33. A 65-year-old male patient was admitted to a hospital with a complaint of muscle weakness. Even though multidisciplinary team was involved on the assessment, the professionals faced difficulty in understanding the case in detail. After a detail discussion and investigation, they agreed that the patient's case is not common and mostly affects males between the age of 50 and 70. The disease progressively damaged the efferent nerve cells of central nervous system and the prognosis is poor. What is the most likely diagnosis of this case?

- A. Guillain–Barre syndrome
- B. Cerebrovascular accident
- C. Motor neuron diseases
- D. Multiple sclerosis

ANSWER: C

34. A 45-year-old patient was admitted to a hospital two weeks ago with difficulty to perform basic functional activities like bed mobility, sitting and standing secondary to a motor neuron disease. The caregivers are very supportive and interested to make him independent in his daily life activities. What is the most likely long term goal for this case?

- A. Independence in bed mobility; rolling to either side, side lying to sitting
- B. Independence in daily washing and upper body dressing in sitting
- C. Independence in transfer from bed to chair and chair to bed
- D. Independence in washing and dressing in standing

ANSWER: D

35. A patient was brought to a physiotherapy clinic with a complaint of progressive neurological diseases with bilateral upper and lower limb weakness. He was in a physiotherapy follow-up for the last two months by doing strengthening exercises and functional activities to treat muscle weakness. Even though the improvement seems promising, the physiotherapist planned to evaluate the prognosis of the patient.

What is the best tool to evaluate the prognosis of this patient?

- A. Nottingham sensory assessment scale
- B. Medical research council scale
- C. Modified Ashworth Scale
- D. Motor Assessment Scale

ANSWER: B

36. A 35-year-old male patient was brought to a physiotherapy clinic with a difficulty of standing and walking due to stroke. Upon physical examination, the physiotherapist asked the patient to flex his left upper limb in sitting position and the patient tried to generate a force but was not able to fully flex but when the patient is at sideline position, movement becomes easy and can flex his elbow fully. What is the most appropriate objective assessment for this case?

- A. Active range of motion
- B. Passive range of motion
- C. Muscle strength
- D. Muscle tone

ANSWER: C

37. A patient was admitted to a hospital with a complaint of sudden onset of left side body weakness which resulted in difficulty of walking and standing. The patient's past medical history indicated that he was smoker, alcoholic and physically inactive. The physical examination result indicated that there is spastic left side hemiparesis and sensory loss, in which upper extremity and face are mainly involved than the lower extremity.

What is the most likely diagnosis of this case?

- A. Right middle cerebral artery occlusion
- B. Right anterior cerebral artery occlusion
- C. Left anterior cerebral artery occlusion
- D. Left middle cerebral artery occlusion

ANSWER: A

38. A patient with central nervous system lesion was brought to a physiotherapy clinic with a complaint of right side body weakness, loss of sensation and shoulder pain. Upon physical examination, the therapist noticed that there is spasticity on the affected side and upper limb is in shoulder adduction, elbow and wrist with a flexed position. The physiotherapist planned to treat the spasticity and related complications.

What is the best management for this case?

- A. Sustained stretching
- B. Strengthening exercise
- C. Bilateral arm training
- D. Active exercise

ANSWER: A

39. A 40-year-old male patient has been brought to a physiotherapy clinic with a history of left side hemiplegia due to central nervous system disorder; difficulty of standing and walking a month before. He was taking intensive physiotherapy intervention to improve muscle strength, sensory re-education and gross motor activities. After a month of physiotherapy follow-up, the patient returned with complaint of loss of balance and coordination called dysmetria. The physiotherapist planned to evaluate the prognosis of patient coordination.

What is the most appropriate means to evaluate the prognosis of this case?

- A. Examining awareness of movement
- B. Examining the joint position sense at rest
- C. Examining by drawing a circle or figure eight
- D. Examining in recognize the form of objects by touch

ANSWER: C

40. A 50-year-old patient was brought to a physiotherapy clinic with complaints of muscle weakness, headache, visual disturbance, and numbness which progressively led her to difficulty of standing and walking. On the subjective assessment, the patient explained that there was a family history similar to this case. Imaging result indicated that there was inflammation and demyelination that resulted in slow neural transmission of the nerves which are affected by the immune system.

What is the most likely diagnosis of this case?

- A. Amyotrophic lateral sclerosis
- B. Guillain-Barre syndrome
- C. Multiple sclerosis
- D. Parkisons diseases

ANSWER: C

41. A 70-year-old female patient came to a physiotherapy clinic with complaints of weakness of extremities and tremor on her left hand. Upon examination, the physiotherapist found muscle weakness on the left upper and lower limb with a muscle manual testing of 2 out of 5 and 3 out of 5 respectively. The gait pattern of the patient is trendelenburg, with a strong lateral lean to the left side of her body.

What is the most appropriate treatment plan for this case?

- A. To increase the hip abductors muscle strength from 2 to 3 within two weeks
- B. To increase the hip flexors muscle strength from 2 to 3 within two weeks
- C. To increase the quadriceps muscle strength from 2 to 3 within two weeks
- D. To increase the hip adductors muscle strength from 2 to 3 within two weeks

ANSWER: A

42. A patient with a neurodegenerative disorder came to a physiotherapy clinic with a history of loss of sitting balance, inability to transfer and move from sitting to standing position. A physiotherapist was providing intervention directed at promoting postural control focusing on static control, weight bearing, balance and coordination of the patient. After a month physiotherapy follow-up, the therapist noticed that the patient can move independently with a support of forearm crutch. However, there is foot drop and ankle instability in the gait analysis.

What is the most appropriate evaluation of this case?

- A. The prognosis is good, need aggressive balance and coordination exercises
- B. The prognosis is good , no need of additional treatments and discharge
- C. The prognosis is good, needs additional ankle foot orthosis devices
- D. The prognosis is good , needs additional walking cane

ANSWER: C

43. A 40-year-old patient was brought to a physiotherapy clinic with a complaint of muscle weakness, numbness and tingling sensation on the extremities. He was admitted to a hospital for a month with unknown cause of the disease. Based on the subjective assessment, the physiotherapist planned to assess sensation especially the cortical sensory of the patient.

What is the most appropriate objective assessment for this case?

- A. Two-point discrimination
- B. Movement awareness
- C. Light touch
- D. Vibration

ANSWER: A

44. A 52-year-old patient was admitted to a hospital with guillain–barre syndrome. He had a five days history of progressive weakness of upper and lower extremities which progressively led to breathing difficulty. Upon physical examination, the physiotherapist found that there is muscle weakness on extremities which ranges from 1 to 3 in manual muscle testing. However, the sensory part is relatively less affected and reflex is preserved.

What is the most likely diagnosis of this case?

- A. Acute inflammatory demyelinating polyneuropathy
- B. Acute motor and sensory axonal neuropathy
- C. Acute motor axonal neuropathy
- D. Miller-Fisher syndrome

ANSWER: C

45. A 30-year-old male patient was admitted to a hospital with a peripheral nervous system disorder. He had a seven days history of progressive weakness of upper and lower extremity which initially starts distally. Upon physical examination, the therapist found that a muscle manual testing (MMT) of lower extremities muscle group is 2/5 and 1/5 for upper extremities muscle group, and there is numbness and tingling sensation in the extremities. The physiotherapist planned to regain the strength of the muscles.

What is the best management for this case?

- A. Resisted exercise
- B. Passive exercises
- C. Free active exercise
- D. Mobilizing exercise

ANSWER: C

46. A 50-year-old football player was brought to a physiotherapy clinic with a complaint of right knee joint pain and swelling due to fall down injury while he was in a game yesterday. The patient walks with the support of his two friends and cannot bear weight on the affected lower limb. The physiotherapist planned to take subjective history on the status of the pain with movement and rest.

What is the most relevant subjective assessment for this case?

- A. Asking about the type and intensity of pain
- B. Asking whether the pain varies with time of day
- C. Asking about the aggravating and easing factors
- D. Asking whether the pain is constant or intermittent

ANSWER: C

47. A 28-year-old patient was referred to a physiotherapy clinic with a complaint of right knee joint pain and swelling due to fall down injury while he was running. Upon physical examination, there was pain with a visual analog scale of 6/10, and tenderness and swelling on the anterior side of knee joint. X-ray investigation revealed no fracture on the site. Based on the symptoms and examination result, the physiotherapist suspected that there might be anterior cruciate ligament injury.

What is the most appropriate objective assessment tool to be used for this case?

- A. McMurray's tests
- B. Posterior draw test
- C. Lachman's test
- D. Varus stress test

ANSWER: C

48. A 67-year-old man who had a history of outstretched fall down injury seven months ago presented with a complaint of pain, swelling, and limited movement of his affected part. The radiological finding revealed an impacted type of fracture on the affected part.

What is the most likely diagnosis of this patient?

- A. Supracondylar fracture
- B. Shaft of humerus fracture
- C. Proximal humerus fracture
- D. Humerus neck fracture

ANSWER: D

49. A 30-year-old man who had history of motor vehicle accident seven months ago came to physiotherapy clinic with complaint of difficulty of performing overhead activities. Upon examination, he had 50 degree right shoulder flexion and 10 degree shoulder extension.

What is the most appropriate SMART goal for this patient?

- A. To improve 50 -70 degree shoulder flexion in a week
- B. To improve 50 -70 degree shoulder flexion within two months
- C. To improve 50-130 degree right shoulder flexion within three days
- D. To improve 50-130 degree right shoulder flexion within a month

ANSWER: D

50. A 28-year-old man was presented with severe pain, swelling, and paresthesias symptoms on the left thigh region due to bullet injury. The radiological finding showed that he has a closed transverse femoral shaft fracture. Orthopedic surgeon managed the fracture with skeletal traction. The physiotherapist was consulted and planned to retard muscle atrophy and maintain strength of the quadriceps muscles.

What is the most appropriate physiotherapy intervention for this case?

- A. Eccentric exercise
- B. Isometric exercise
- C. Isokinetic exercise
- D. Concentric exercise

ANSWER: B

51. A 22-year-old man had history of inhalational burn injuries on his face, hand, chest, and anterior part of upper extremities a year ago. Upon examination, the patient had full thickness burn and 30 percent TBSA. In addition to this, the patient noticed shortness of breath and burning kind of pain on the burnt part of his body.

What is the most likely positive prognostic indicator of this case?

- A. Age of the patient
- B. Location of burn
- C. Degree of burn
- D. Type of burn

ANSWER: A

52. A 50-year-old woman who is an osteoarthritis patient came to physiotherapy outpatient clinic three weeks back. She was complaining of morning stiffness, swelling and pain on carpal, tarsal and phalangeal joints in the morning. Moreover, she had a relaxed feeling and decrement of pain feeling during the day times.

What is the most appropriate assessment for this patient?

- A. Easing and aggravating factors
- B. History of present complaint
- C. Chief complaint
- D. Medical history

ANSWER: A

53. A 40-year-old woman came to physiotherapy outpatient clinic with a complaint of morning stiffness, bilateral symmetrical symptoms and swelling of PIP, MCP and MTP joints which is getting worse through time since a year ago. The radiological findings showed bony erosion and localized decalcification of joints.

What is the most likely diagnosis of this patient?

- A. Osteoarthritis
- B. Rheumatoid arthritis
- C. Ankylosing spondylitis
- D. Juvenile idiopathic arthritis

ANSWER: B

54. A 50-year-old woman came to a physiotherapy clinic with history of chronic left knee joint pain. The pain developed gradually since the last five years. On examination, the physical therapist found knee extensor muscle weakness which was 4 out of 5 in MMT grading.

What is the best appropriate treatment plan for this condition?

- A. Manual strength exercise of left knee extensor with 3 set and 10 repetitions
- B. Mechanical strength exercise of knee extensor with 3 set and 10 repetitions
- C. Manual resistance exercise of knee extensor with 3 set and 10 repetitions
- D. Strengthening exercise of knee extensor with 3 set and 10 repetitions

ANSWER: A

55. A 53-year-old married woman came to physiotherapy outpatient clinic with a diagnosis of chronic rheumatoid arthritis on her hip and knee joints. She is presented with symptoms of pain, morning stiffness and weakness on her joints. On examination, physiotherapist found weakness on knee extensor and hip flexor muscles. Accordingly, the physiotherapist planned to improve the muscle adaptation and functional performance of the patient.

What is the most appropriate treatment for this case?

- A. Strength exercise
- B. Functional splint
- C. Dry heating pad
- D. Hydrotherapy

ANSWER: A

56. A 57-year-old woman who works as a nurse for the last 25 years came to a physiotherapy outpatient clinic with a diagnosis of right knee osteoarthritis for the last 12 years. She was following a physiotherapy treatment for the last three years. However, the physiotherapist noticed a gradual worsening of symptoms with severe pain and restricted mobility with a difficulty to perform her daily routines.

What is the most appropriate evaluation for the prognosis of this patient?

- A. The prognosis is poor and advise her to discontinue her work
- B. The prognosis is poor and advise her to request leave
- C. The prognosis is poor and recommend her for joint replacement
- D. The prognosis is poor and advise her to continue physiotherapy follow-up

ANSWER: C

57. A 50-year-old female patient who sustained a fall down injury a month back came to physiotherapy clinic. She has complaints of pain, paresthesias, and numbness on the medial nerve distribution of the right upper limb. In addition to this, she had nocturnal pain and pain during activities. The symptoms worsen gradually and especially when she is engaged in some form of activity. Radiological investigation confirmed that there is no any fracture sign.

What is the relevant subjective assessment described in this case?

- A. Personal history
- B. Medical history
- C. Present history
- D. Past medical history

ANSWER: C

58. A 35-year-old man who had a fall down injury on the right shoulder joint was treated surgically. After surgery, the patient's complaint is decreasing range of motion in conjunction with pain. Upon examination, the physiotherapist found signs of warmth at the site, erythema, and effusion.

What is the most likely diagnosis of this patient?

- A. Stiffness
- B. Infection
- C. Frozen shoulder
- D. Impingement syndrome

ANSWER: B

59. A 43-year-old lady came to physiotherapy outpatient clinic with a diagnosis of rheumatoid arthritis three years ago. She has noticed morning stiffness, pain and restricted mobility on her knee and elbow joints. On examination, physical therapist found five degree lag for knee and elbow joint extension range of motion exercise.

What is the most appropriate goal of this patient?

- A. Improve right knee extension 175-180 degree within three months
- B. Improve right knee extension 175-180 degree within two weeks
- C. Improve knee extension to full range of motion within a day
- D. Improve knee extension 175-180 degree within a week

ANSWER: B

60. A 30-year-old man who is a tennis elbow player for the last 10 years came to physiotherapy outpatient clinic after surgical repair of his right shoulder rotator cuff tear. Upon physical examination, the physiotherapist found pain, limited range of motion, and weakness on the affected side.

What is the most appropriate treatment for the restricted movement at this stage?

- A. Pendulum exercise
- B. Resistance exercise
- C. Shoulder active ROM
- D. Electrical stimulation

ANSWER: A

61. A 30-year-old patient who was admitted in a hospital with a diagnosis of type II superior labral anterior posterior (SLAP) lesion of the left upper limb was treated surgically. Immediately after the surgery, the physiotherapist is consulted to treat the patient who positioned his affected limb in a sling.

What is the most appropriate management for this patient until two week post surgery?

- A. Perform overhead activities
- B. Avoid active elbow flexion
- C. Rotator cuff isotonic strengthening
- D. Posterior glide in the plane of scapula

ANSWER: B

62. A 50-year-old male patient with a diagnosis of tennis elbow was treated surgically two weeks ago. The splinting and suture were removed after a week. He has full shoulder, forearm, and hand mobility with elbow active range of motion (AROM) of 50 degree to 120 degree. His pain remains 4/10 through VAS even at rest.

What is the most appropriate intervention based on the prognosis of this patient?

- A. Ultrasound
- B. Link to physician
- C. Grade II mobilization
- D. Anti-pain medication

ANSWER: B

63. A 29-year-old office worker pregnant woman came to a physiotherapy clinic with a complaint of pain in her lower back which radiates to posterior thigh and leg that minimally interferes with her daily life and compromises her ability to work in the office. Her pain is 6 in visual analogue scale and aggravates by prolonged sitting. She denies a history of trauma.

What is the most appropriate subjective assessment to be asked in this case?

- A. Family history
- B. Working posture
- C. Past medical history
- D. Associated medical problem

ANSWER: B

64. A 25-year-old third trimester pregnant woman came to a physiotherapy OPD with complaints of pain in her right wrist joint, burning and tingling sensation in the palmar aspect of lateral three fingers and lateral half of the 4th finger. She is currently diagnosed with gestational DM and hypertension.

What is the most likely diagnosis for this case?

- A. Carpal tunnel syndrome
- B. Cubital tunnel syndrome
- C. Guyons tunnel syndrome
- D. Pronator teres syndrome

ANSWER: A

65. A 25-year-old 3rd trimester pregnant woman came to a physiotherapy OPD with a diagnosis of carpal tunnel syndrome. After assessment, a physiotherapist wants to prescribe orthosis for the client. What is the most appropriate orthosis in the management of this patient?

- A. Elbow orthosis
- B. Wrist hand orthosis
- C. Shoulder elbow orthosis
- D. Shoulder elbow wrist orthosis

ANSWER: B

66. A 25-year-old female who is diagnosed with carpal tunnel syndrome had follow-up physiotherapy intervention for the past one month. Previously she had wrist pain and sensory symptoms but no motor involvement. After 10 sessions of physical therapy intervention, she started to complain difficulty of doing household activities and being unable to hold objects by the right hand.

What is the most appropriate next step management for this patient?

- A. Advise the patient to discontinue the therapy
- B. Advise the patient to do activities with left hand
- C. Advise the patient to stop house hold activities
- D. Send the patient for Nerve conduction test

ANSWER: D

67. A 40-year-old female patient came to a physiotherapy OPD with a complaint of pelvic girdle pain, urinary incontinence, pressure and heaviness in the pelvis. She was diagnosed with pelvic organ prolapse.

What is the most appropriate assessment that needs to be asked in relation to past medical history?

- A. Number of children
- B. Smoking history
- C. Previous hysterectomy
- D. Family history of pelvic organ prolapse

ANSWER: C

68. A 38-year-old woman on her 15th postpartum day came to a hospital with complaints of difficulty to control her urine during coughing, laughing, sneezing and while she was lifting heavy objects. She denies history trauma and she had a normal vaginal delivery with a normal labour duration.

What is the most likely diagnosis?

- A. Urinary urgency and frequency
- B. Stress urinary incontinence
- C. Urge urinary incontinence
- D. Mixed urinary incontinence

ANSWER: B

69. A 35-year-old postpartum patient with urinary incontinence was linked to physiotherapy OPD. The patient complains urinary incontinence which manifested while she engages herself in heavy home activities, laughing, sneezing and coughing.

What is the most appropriate management for this patient?

- A. Pelvic floor muscle strengthening exercise
- B. Pelvic floor muscle stretching exercise
- C. Ultrasound for pelvic floor muscle
- D. Infrared (IRR) for pelvic floor muscle

ANSWER: A

70. A 32-year-old female patient, who underwent C-section and is on follow-up physiotherapy interventions, complained increased pain, redness, swelling, bleeding and drainage from the incision site on the 10th day of follow-up assessment.

What is the most appropriate next step in the management of this patient?

- A. Send the patient to physician
- B. Advice to continue the intervention
- C. Discharge the patient with home advice
- D. Increase the intensity of exercise

ANSWER: A

71. A 72-year-old male client came with a diagnosis of osteoarthritis of the hip and depression. He has history of back pain with chronic lumbar spine degeneration. He lost his daughter and was unable to lead an independent life due to pain. He walks with walker/stick, better in morning, slower on stairs and uneven surfaces, and sometimes pain may be a limiting factor.

What is the most appropriate subjective assessment included under personal history of this patient?

- A. Age
- B. Assistive devices
- C. Family situation
- D. Back pain

ANSWER: C

72. A 75-year-old male client came to physiotherapy OPD with a diagnosis of dementia and history of hypertension, ischemic heart diseases, and nocturnal confusion. He has mild visual lose, and is immobile and wheelchair dependent.

What is the most appropriate outcome measurement tool to assess the patient cognition?

- A. Barthel index
- B. Perceived exertion rate
- C. Mini mental state examination
- D. Functional ambulation classification

ANSWER: C

73. An 85-year-old female patient came with complaints of lower extremity pain and difficulty in outdoor movement. After initial assessment, the client claims that she has feeling of sadness, hopelessness and lose of interest to do activities and she prefers to spend her time sitting at home the whole day. The family also complained that she has difficulties of concentrating, low self-esteem, decreased motivation, apathy, and excessive guilt and those symptoms get worse mainly in the morning.

What is the most likely diagnosis for this case?

- A. Dementia
- B. Delirium
- C. Depression
- D. Hallucination

ANSWER: C

74. An 80-year-old male patient who previously was diagnosed and treated for Parkinson disease came to a physiotherapy OPD with complaints of inability to move independently; full assistance is required for all hygienic, dressings and transfer needs. He is unable to feed himself and requires full assistance with positioning, sitting and set-up of meals.

What is the most appropriate assistive device for this patient?

- A. Cane
- B. Crunch
- C. Walker
- D. Wheelchair

ANSWER: D

75. A 73-year-old patient with complaints of decreased mobility and balance is following physiotherapy intervention. The patient uses cane for walking outside home and he has fear of falling and is depressed because he is not fast enough while walking. After one month of physiotherapy intervention, a physiotherapist plans to examine the progress of the patient walking speed.

What is the most appropriate evaluation tool for this patient?

- A. Timed 'Up and Go'
- B. Berg Balance Scale
- C. 10 meter walk
- D. Barthel Index

ANSWER: C

76. A 75-year-old man is complaining of left-sided chest pain radiating to his axillary area with the medical diagnosis of non Q-wave anterior myocardial infarction. The pain appeared about an hour ago. Pulse rate is 112/min and in sinus rhythm. Blood pressure is 145/100mmHg. He has no history of diabetes, hypertension and known cholesterol problems. He has a habit of smoking 4 cigarettes a day and up to a half a pack daily in the past.

What is the most likely subjective component that can be taken as a cardiac risk factor for the patient?

- A. Chief complaint
- B. Medical diagnosis
- C. Vital sign
- D. Personal history

ANSWER: D

77. A 32-year-old former athlete presented with a complaint of chest pain. The pain description reveals pressure, heaviness, squeezing, tightness, and burning like pain around the substernal, jaw, shoulder, back and arm. The pain is aggravated by training in the field and emotional stress for the duration of 5 -15 minutes and relieved by rest.

What is the most likely explanation based on the given clinical manifestation?

- A. Angina pectoris
- B. Stable angina
- C. Unstable angina
- D. Anginal equivalents

ANSWER: A

78. A 34-year-old female patient was presented with a complaint of left-sided chest pain radiating to her axillary area. The chest discomfort was described as if somebody sat on her chest. The pain started about an hour ago. Serial EKG showed ST segment depression in the precordial leads. Serial troponin I showed an increase to 10mg/L at 6 hours and 20mg/L at 18 hours. Pulse rate is 112/min and in sinus rhythm. Blood pressure is 145/100mmHg. She has a past medical history of herpes zoster infection, prediabetic and cholesterol with a BMI of 27kg/m², smokes 6 cigarettes a day and up to a half a pack a daily in the past.

What is the most likely diagnosis?

- A. Myocardial infarction
- B. Chronic heart failure
- C. Valvular heart diseases
- D. Cardiomyopathy

ANSWER: A

79. A 49-year-old male patient came to physiotherapy clinic with a complaint of right leg pain. There is redness and swelling throughout the foot and ankle distal to the knee that has developed over the last three days. The patient reports no trauma and complains of a deep ache in the calf musculature.

What is the most appropriate initial treatment?

- A. Refer to physician for further examination
- B. Elevate the lower extremity and apply an ice modality
- C. Instruct on range of motion exercises and begin a home exercise program
- D. Instrument assisted manual soft tissue mobilization to decrease the swelling

ANSWER: A

80. A physiotherapist is treating and following a patient for cardiac rehab for the past two months. He started treating the patient since day three after admission in the hospital. The client is following an exercise programme over a period of time and the physiotherapist noticed the patient is stressed and planned to reduce patient stress levels and clear thought processes.

What is the most appropriate management for a good prognosis of the patient?

- A. Relaxation exercises
- B. Regular exercise
- C. Getting into low level training
- D. A reduced score on the exercise scale

ANSWER: A

81. A 55-year-old man diagnosed with chronic asthma presented with complaints of chest pain, shortness of breath, coughing, and wheezing. He was immediately admitted to a medical ward to manage the significant decrement of saturation of oxygen. On the 4th day of admission, he was referred to physiotherapy clinic to consult for pulmonary rehabilitation. During physiotherapy subjective assessment, the therapist asked him for the presence of chronic diseases. What is the most likely subjective assessment component exercised in this case?

- A. Onset of the condition
- B. Mechanism of the condition
- C. Social history
- D. Associated medical history

ANSWER: D

82. A 75-year-old man came to a hospital complaining of increasing fatigue and shortness of breath for a year which progressively worsened over the last three years and it becomes tough when he is in yard work, and shaving. He smoked 2 packs per day from age 13 to 65. On physical examination, a high-pitched crackle during inspiration was noted. Chest x-ray showed small lung volumes, interstitial pattern on lung field, poorly defined cardiophrenic angles bilaterally and small cardiothoracic index.

What is the most appropriate diagnosis for this case?

- A. Asthma
- B. Cystic fibrosis
- C. Restrictive lung disease
- D. Chronic obstructive pulmonary disease

ANSWER: C

83. A 57-year-old male cigarette smoker who had chronic stable obstructive pulmonary disease visited physician with a complaint of coughing, expectoration, and dyspnea. He was managed with anti-cholinergic drugs, and then consulted for pulmonary rehabilitation.

What is the most appropriate physiotherapy management for the dyspnea?

- A. Huffing and coughing
- B. Manual Percussion
- C. Pursed lip exercise
- D. Riding bicycle

ANSWER: C

84. A 35-year-old male patient who had chronic bronchitis for the last two years came to a hospital. He was under physiotherapy follow-up program for the last two months with a management of deep breathing exercise. Two weeks before, the symptom got worse, and the therapist decided to stop and refer the patient back to the physician for further investigation. On review examination, he was diagnosed with pulmonary tuberculosis and started antibiotic and anti-TB drugs. A month later, he started a physiotherapy program with another session.

What is the most appropriate physiotherapy treatment for good prognosis of productive cough?

- A. Breathing exercises
- B. Pursed lip exercise
- C. Huffing exercise
- D. Riding bicycle

ANSWER: C

85. A 58-year-old male cigarette smoker patient diagnosed with chronic obstructive pulmonary disease came to physiotherapy OPD for pulmonary rehabilitation. He had a follow-up program of three times in a week for the last three weeks. Upon re-examination, he feels fine apart from dyspnea for the last one week.

What is the most appropriate positioning to improve prognosis of this case?

- A. Supine positioning
- B. Prone positioning
- C. Upright positioning
- D. Relaxation positioning

ANSWER: D

86. Physiotherapy undergraduate students were trained about how to manage patients in ICU. Their instructor taught them about the indication and contraindication of prone position which is frequently used for patients in acute medical and surgical condition admitted in ICU.

Which case is the most likely contraindication for prone position?

- A. Early acute respiratory distress syndrome
- B. Acute lung insufficiency
- C. Cerebral hypertension
- D. Pulmonary edema

ANSWER: C

87. A 63-year-old male patient was admitted to an ICU with complaints of shortness of breath and pleuritic chest pain. He had got car accident a month ago and sustained comminuted fracture at both distal tibio-fibular bones. The fracture was managed with external fixation and he prefers to lie down in supine position for the last one month. On examination, his vital signs were temperature of 36.8 degree centigrade, pulse rate of 85/min, respiratory rate of 26/min, blood pressure of 110/60mmHg, and SPO2 of 84 percent. Duplex ultrasonography test is positive.

What is the most likely diagnosis for this case?

- A. Pulmonary emboli
- B. Pulmonary edema
- C. Pleural Effusion
- D. Lung abscesses

ANSWER: A

88. A 28-year-old female who sustained a car accident was immediately admitted to ICU with a diagnosis of open head injury and fracture on right distal humeral bone. Surgical procedure was done for the head injury and right humeral fracture. She was assisted with mechanical ventilation. Upon examination, all her vital signs and other parameters were controlled. After auscultation physiotherapy assessment, the therapist found accumulation of secretion in apical segments of both upper lobes. Therefore, the physiotherapist planned to drain this accumulated secretion.

What is the most appropriate positioning technique for this case?

- A. Supine with raising the head of the bed 45 degree with fatted on the foot of the bed
- B. Prone with raising the head of the bed 45 degree with fatted on the foot of the bed
- C. Supine with elevation of the foot of the bed 45 degree
- D. Prone with elevation of the foot of the bed 45 degree

ANSWER: A

89. A 15-year-old male patient was admitted to an ICU eight weeks ago. He sustained quadriplegia secondary to spinal cord injury with a fracture of C3 to C8 vertebral bone. On presentation, he was unconscious and undergone spinal surgery. He was put under mechanical ventilator and consultation for physiotherapy intervention was made. After two weeks of follow-up, the therapist together with ICU team staff, undertaken observation and found bedsores on the ischial tuberosity.

What is the most appropriate prognostic physiotherapy intervention needed for this case?

- A. Breathing exercise
- B. Positioning
- C. Spirometry
- D. Percussion

ANSWER: B

90. A 28-year-old male soccer player was admitted to emergency unit. He lost his consciousness while he was playing soccer in the field. There was no any contact injury with the player or with the ball. He was taken to emergency unit immediately but it was impossible to save his life. On his previous file; he had no history of chronic medical problems.

What condition is considered as a risk for sudden death from the point of fitness perspective?

- A. Cardiovascular conditions
- B. Musculoskeletal conditions
- C. Neurological conditions
- D. Pulmonary conditions

ANSWER: A

91. A 36-year-old female marathon runner came to physiotherapy center with an intent to start physical fitness program. She undertakes a regular medical and physical fitness check-up by physician. She was referred to physiotherapy for therapeutic exercises which emphasizes the improvement of vital capacity of lung.

What is the most appropriate fitness training techniques applied for this case?

- A. Muscular endurance
- B. Muscular strength
- C. Flexibility exercise
- D. Aerobic exercise

ANSWER: D

92. A 25-year-old male combatant has been following his physical fitness program in a physiotherapy OPD. The designed exercise by the therapist was targeted on body building within three months. After SOAP assessment, the therapist realized that he is in need of more supplementary foods to achieve the plan.

What nutrient item is most likely recommended for the combatant to take?

- A. Carbohydrate
- B. Vitamin
- C. Protein
- D. Lipid

ANSWER: C

93. A nine-year-old male student had got a fall down injury while he was playing soccer. He was admitted to orthopedic ward for surgical procedure of fracture on his right distal and shaft of humeral bone. He was referred to physiotherapy OPD with difficulty of extending his wrist joint and holding things by his right hand.

What is the most likely structure injured in this case?

- A. Peroneal nerve
- B. Radial nerve
- C. Medial nerve
- D. Ulnar nerve

ANSWER: B

94. A 25-year-old male soccer player who had a complaint of pain and limping while he walks visited physiotherapy OPD. Upon examination, the therapist instructed the patient to stand with his right leg and observed the position of the patient's ASIS, PSIS, and iliac crests.

What is the most likely special test used for this case?

- A. Hip scouring test
- B. Trendelenburg
- C. Thomas test
- D. Ober's test

ANSWER: B

95. A 26-year-old female soccer player has got contact injury on her right knee joint. She fall down with right foot in plantar flexion position; right knee in slightly flexion and moving into full extension and external rotation. She complained swelling soon after the injury. On examination, Lachman test was positive whereas anterior and posterior drawer tests were negative.

What is the most likely diagnosis of this case?

- A. Anterior cruciate ligament sprain
- B. Posterior cruciate ligament sprain
- C. Medial collateral ligament sprain
- D. Lateral collateral ligament sprain

ANSWER: A

96. A 25-year-old male basketball player was admitted to orthopedic ward for the complaint of swelling and difficulty to stand due to contact injury on his right ankle. His MRI investigation showed a complete calcaneal tendon rupture of his right ankle. After a tendon repair procedure, he was referred to physiotherapy for further rehabilitation.

What is the most appropriate immediate physiotherapy care at this stage?

- A. Non weight bearing gait training using crutches
- B. Strengthening exercise of the quadriceps muscle
- C. Stretching exercise of the calcaneous tendon
- D. Strengthening exercise of the calf muscle

ANSWER: A

97. A 28-year-old female volleyball player visited a medical OPD with a diagnosis of patellar tendonitis of right knee joint. She had complaint of pain when jumping, ascending and descending the stairs. After her case was managed by steroids, she was referred to physiotherapy clinic for rehabilitation.

What is the most appropriate physiotherapy technique for this case at this stage?

- A. Running on treadmill
- B. Stair climbing exercise
- C. Running on rough surface
- D. Straight leg raise exercise

ANSWER: D

98. An eight-year-old male student got a fall down accident with outstretched hand during soccer match. He was admitted in orthopedic ward with a fracture of left distal humeral bone with elbow dislocation. His case was managed with reduction and sling. He was referred to physiotherapy for wrist drop rehabilitation.

What is the most appropriate preventive strategy in this case?

- A. Encourage passive movement at left wrist joint
- B. Encourage passive movement at left elbow joint
- C. Immobilize movement at left elbow joint
- D. Immobilize movement at left wrist joint

ANSWER: A

99. A 15-year-old boy was referred to a physiotherapy rehabilitation center with transfemoral amputation done six months ago secondary to snake bite. He reported that, he feels as if his amputee limb is alive hence he stands to walk suddenly and feels pain. He is also worried about accepting the overall situations; is future life in maintaining friendship and continuing his education after grade 12. What is the most likely condition that has to be addressed before prescribing device in this case?

- A. Phantom limb pain
- B. Fear of putting weight of stump
- C. School drop stress
- D. Psychosocial issues

ANSWER: D

100. An 80-year-old man has been amputated through his right proximal femur 3 weeks ago and presented to physiotherapy OPD. Upon his medical chart review, the physiotherapist found that the amputation was due to complication of leprosy and sensory loss at the distal part of residual limb. He also experienced weakness and numbness in both lower extremities.

What is the most advisable mobility device for this patient?

- A. Wheelchair
- B. Transfemoral prosthesis
- C. Axillary crutches
- D. Walking frames

ANSWER: A

101. A 36-year-old male driver sustained a left tibiofibular comminuted fracture following road traffic accident 20 days ago. He was managed with external fixation and referred to a physiotherapy clinic. On examination, the physiotherapist found left foot drop and inability to put weight on left foot. The patient is optimistic and eager to ambulate and return to his job.

What is the most appropriate early mobility device to ambulate in this case?

- A. Foot orthotics and elbow crutch
- B. Knee Ankle Foot Orthotics and elbow crutch
- C. Knee orthotics and elbow crutch
- D. Ankle Foot Orthotics and elbow crutch

ANSWER: D

102. A 37-year-old female secretary has experienced paraplegia 20 days ago secondary to GBS. Physiotherapy was consulted once her medical conditions were stabilized. Upon examination, the physiotherapist found poor trunk control test, bilateral lower limbs muscle power 2/5 on manual muscle testing and she had foot drop (MMT equal to 1). Her family has no any other income except her salary. She is informed that she will lose her job unless she gets back within a month.

What is the most appropriate early mobility device that would help her?

- A. Wheelchair
- B. Axillary crutch
- C. KAFO
- D. AFO

ANSWER: A

103. A 24-year-old college student came to physical rehabilitation center with complaint of pain and gait deviation. He has been using left trans-femoral prosthesis for the first time for the last eight months. He reports that he feels as if his limb does not exist. On examination, the physiotherapist found normal stump shape, good gluteus muscle function of the residual limb and properly fitted stump with the socket. No problem was reported on the device by orthopedic technician but he stopped using the device after two days of returning home. Thus, he is looking for a new prosthesis to go back to school.

What is the most appropriate initial action before prescribing a new device in this case?

- A. Check phantom pain and plan for re-fitting
- B. Check the muscle power and plan strengthening exercises
- C. Check an alignment of prosthesis and make adjustments
- D. Check ROM and plan post fitting training with old prosthesis

ANSWER: A

104. A 64-year-old man has been following a physiotherapy services for post stroke left side hemiplegia and elbow contracture because of severe spasticity four months ago. On SOAP assessment, the physiotherapist found improvement with passive ROM exercises of elbow and pain from VAS 7/10 to 5/10 within two weeks.

What is the most appropriate orthotic device to be prescribed in this case?

- A. Rigid shoulder abduction orthotics
- B. Non-articulated elbow immobilization orthotics
- C. Shoulder immobilization orthotics
- D. Articulated adjustable forearm orthotics

ANSWER: B

105. A six-year-old boy came to physiotherapy OPD with difficulty of standing and walking since birth secondary to cerebral palsy. He is still crawling by dragging his both lower limbs to play with his neighbourhood kids and to move indoor and outdoor. On physical examination, the physiotherapist noticed spasticity on bilateral lower extremities and has episodes of a frequent fall while trying to stand up. No sensory problem finding and has normal motor functions in upper limbs.

What is the most appropriate device to help him to stand?

- A. Para podium
- B. Hip Control Hip Orthoses
- C. Knee Ankle Foot Orthotics
- D. Ankle Foot Orthotics

ANSWER: B

106. A 49-year-old security officer suffered a left long trans-femoral amputation after miss fired self gunshot. After six months of amputation he was referred to rehabilitation center prosthesis. The officer is excited to be independent and eager to get back to his job. Upon examination, the physiotherapist's evaluation of stump assessment revealed minimal edema, little redness seen after training, and residual limb change in size. The residual limb muscle power of hip flexor, abductor, extensor and adductor are fair.

What is the most likely prescribed prosthesis in this case?

- A. Temporary trans-femoral prosthesis
- B. Temporary trans-tibial prosthesis
- C. Definitive trans-femoral prosthesis
- D. Definitive trans-tibial prosthesis

ANSWER: A

107. A 14-year-old girl revisited a physical rehabilitation center with a complaint of pain and gait deviation. She has been using left trans-femoral prosthesis for the last five months. Upon physical examination, there is blister at distal residual limb, poorly fitted stump with the socket. Thus, she is looking for new prosthesis to go back to junior school.

What is the most likely evaluation to be considered to change the prosthesis?

- A. Age
- B. Gender
- C. Educational status
- D. Nature of amputation

ANSWER: A

108. A 62-year-old man was referred to rehabilitation center after seven months of right trans-tibial amputation for prosthesis fitting. He is known diabetic mellitus and hypertensive patient for the last 10 years. He has completed gait training indoor and smooth surface outdoor sessions and fitted with trans-tibial prosthesis. He is discharged back to home with appointment scheduled after one month. What are the most likely domains to be checked in the next visit for this patient?

- A. Skin irritation, cosmetic and gait
- B. Cosmetic, ischial seat and gait
- C. Cosmetic, stump length and gait
- D. Stump length, pain and gait

ANSWER: A

109. A 45-year-old man came to physiotherapy clinic complaining of constant dull aching pain in his lower back and sharp shooting pain in his right leg and occasional burning sensation in the anterior aspect of his right thigh. He had undergone lumbar discectomy and laminectomy at multiple level lumbar spines of L3/L4, L4/L5 and L5/S1 for the severe spinal stenosis two months ago. He is worrying about his current health condition and the situation is threatening his marriage and spiritual life.

What is the most possible explanation of the impact of pain in this case?

- A. Bio-psychosocial Model
- B. Medical model
- C. Behavioral model
- D. Spiritual model

ANSWER: A

110. A 55-year-old man came to physiotherapy OPD complaining of constant dull aching pain in the neck region with sharp shooting pain down his left arm. MRI results revealed that C6/C7 cervical degenerative disc disease. Upon examination, muscle strength of left elbow extensor and wrist extensor are fair but reduced triceps tendon reflex.

What is the most possible explanation of the source of suffering in this patient?

- A. Neuropathic pain
- B. Emotional pain
- C. Physiologic pain
- D. Spiritual pain

ANSWER: A

111. A 17-year-old girl, who experienced incomplete spinal cord injury (ASIA-B) of T10 20 days ago by catastrophic car accident, came to a physiotherapy clinic. She was brought by wheelchair being immobilized by a thoraco-lumbar corset. When examination is started, the girl only showed the site injury where she feels pain. She looks depressed because of her health situation and life encounters. What is the most appropriate assessment tool to determine the intensity of the pain?

- A. Numerical rating scale
- B. Oswestry disability index
- C. Short form 36
- D. Ronald Morris disability index

ANSWER: A

112. A 58-year-old woman, who was diagnosed with breast and cervix cancer, was brought to pain and palliative care ward with complaint of constant burning with VAS 9/10. She reported urine incontinence, weakness of right upper limb, and difficulty of dressing herself. Her pain makes her emotionally unstable, worried about her family life and blames her God for letting her to be a cancer patient.

What is the most likely early treatment plan in this case?

- A. Family education
- B. Pain reduction
- C. Improve muscular power
- D. Improve function

ANSWER: B

113. A 60-year-old farmer who has been recently diagnosed with multiple myeloma is undergoing combined radiotherapy and chemotherapy in the hospital. Since admission, he has been complaining of pain in the back, right shoulder, and left groin. He has been evaluated thoroughly using MRI and bone scan, which revealed wedge fractures of thoracic spine (T7 and T8) and osteolytic regions in the right humerus, ribs, and right femur. His medication regimen includes acetaminophen, tramadol, and morphine sulfate. The attending hematologist is having problems with his pain control. Physiotherapist was consulted and through his medical notes, found that he is on antibiotics for left lower lobe pneumonia.

What is the most appropriate immediate action to be taken for this patient?

- A. Recommend a dorso thoraco-lumbar brace
- B. Apply trans-cutaneous electrical stimulation
- C. Use percussion technique
- D. Mobilization of thoraco-lumbar spine

ANSWER: A

114. A 78-year-old man was diagnosed two months ago with advanced carcinoma of the pancreas. He has extensive bony metastasis, and his main problem is severe, intolerable abdominal and lower back pain. The physician has consulted physiotherapist for an intervention, since he planned to assess for a possible neurolytic celiac plexus block. The physician is doing so in order to keep the pain under control. The patient has expressed his desire to spend the last days of his life with his family. Review of his medical records showed that his liver function tests are abnormal and his clotting is slightly deranged. Other medical problems include a 4 cm size aortic aneurysm, COPD, and hypertension. What is the most likely poor prognostic indicator in this case?

- A. COPD
- B. Pain
- C. Bony metastasis
- D. Degenerative changes of the spine

ANSWER: C

115. A 58-year-old woman who has been complaining of lower back pain and severe groin pain following cervical cancer radiotherapy and chemotherapy sessions was referred to physiotherapy clinic. The physiotherapist started treatment targeting moderate fecal incontinence, pelvic organ prolapse and chronic urinary incontinence and constant burning pain 9/10 for the past four weeks. After weeks of rehabilitation, she reported pain reduction to 7/10 and she was sent back home with her regular appointment.

What is the most likely good prognostic result in a maintenance phase in this case?

- A. Improve pain reduction
- B. Improve pelvic organ prolapse
- C. Improve fecal incontinence
- D. Improve urine incontinence

ANSWER: A

116. A 45-year-old patient came to a physiotherapy clinic with complaints of pain and progressive limitation of ROM in the shoulder joint of two months duration. At its early stage, the pain gets worse at night and the stiffness limited the abduction and external rotation of the shoulder. Currently, the patient complains pain at all times and all the movements of shoulder joints are severely restricted.

What is the most appropriate electrotherapy modality to be used in this case?

- A. Interferential therapy
- B. Ultraviolet radiation
- C. Infrared radiation
- D. Therapeutic Ultrasound

ANSWER: A

117. A 57-year-old male patient with left hemiplegia after stroke attack of four months duration was presented to physiotherapy OPD. He also reported difficulty of lifting his left shoulder and arm and performing over head activities. Upon physical examination, the left humeral head was displaced caudally and shoulder abductors and extensors were weak (2/5 on MMT).

What is the most appropriate electrotherapy modality to be used in this case?

- A. Neuromuscular electrical stimulation
- B. Functional electrical stimulation
- C. Neuromuscular electrical stimulation
- D. Trans-cutaneous electrical nerve stimulation

ANSWER: B

118. A 25-year-old man sustained a complete rupture of his right Achilles tendon five weeks ago while playing football. The tendon was surgically repaired one week later. He has been referred for physiotherapy with complaints of mild discomfort at the surgical incision site that worsens while walking. Plaster of Paris Cast was removed yesterday and he has been instructed to walk, bearing weight as tolerated, wearing a heeled boot. The objective examination has revealed reduced dorsiflexion ROM (-10 degree), mild swelling, tenderness, and redness in the area of the surgical site and atrophy of the calf muscles on the right.

What is the most appropriate modality to be used in this scenario?

- A. Ultraviolet Radiation
- B. Transcutaneous electrical nerve stimulation
- C. Infrared therapy
- D. Cryotherapy

ANSWER: D

119. A 37-year-old male patient with the complaint of frozen shoulder after immobilization due to right proximal humeral fracture three months ago has come to physiotherapy clinic. The physiotherapist has evaluated and found both active and passive ROM of shoulder abduction and external rotation excessively limited and prescribed shockwave diathermy before stretching exercise and mobilization techniques.

What are the most appropriate therapeutic frequencies and wavelengths to be used in this case?

- A. 54.24 MHz and 30 m
- B. 40.68 MHz and 2 m
- C. 27.12 MHz and 11 m
- D. 13.56 MHz and 22 m

ANSWER: C

120. A 45-year-old male patient came to a hospital with unilateral heel pain. He reports pain that gets worse in the morning and inability to bear weight on the foot while getting up from bed. A lateral view x-ray revealed calcaneal spur. A physiotherapist planned to apply short wave diathermy for the heel pain.

What is the mostly likely placement of the electrodes to be used?

- A. Mono-planar technique
- B. Contra-planar technique
- C. Coplanar-technique
- D. Crossfire technique

ANSWER: A

121. A 35-year-old athletic woman came to sport physiotherapy clinic with a complaint of left ankle joint sprain while playing tennis table. The physiotherapist noticed that there is pain and gross edema on her left ankle joint and cryotherapy was his plan of care. The physiotherapist also noted that she has a peripheral vascular disease.

What is the most likely precaution that has to be considered before applying Cryotherapy in this case?

- A. Oedema
- B. Ankle sprain
- C. Peripheral vascular disease
- D. Being female

ANSWER: C

122. A 19-year-old footballer came to physiotherapy OPD with a complaint of left knee extensor weakness. Upon physical examination, left knee extensor muscle strength was found to be 2/5 on MRC scale. The plan of the physiotherapist was to use Faradic current stimulation for muscle re-education.

What is the most appropriate procedure to apply Faradic current stimulation in this case?

- A. Place electrodes on muscle belly with high frequency setting
- B. Place electrodes on motor point with low frequency setting
- C. Place electrodes on muscle belly with low intensity setting
- D. Place electrodes on motor point with high frequency setting

ANSWER: B

123. A 50-year-old woman was referred to physiotherapy clinic with an edema over her left lower extremity. After taking appropriate assessment, the physiotherapist sets plan to apply galvanic current to reduce the edema.

What is the appropriate principle of the galvanic current application in this case?

- A. Continuous current over edematous area with low intensity setting
- B. Interrupted current over edematous area with low intensity setting
- C. Continuous current over edematous area with low frequency setting
- D. Interrupted current over edematous area with low frequency setting

ANSWER: C

124. A 40-year-old male patient came to physiotherapy OPD with mild to moderate lower back pain that radiates to his left buttock and lateral thigh of five weeks duration. On MRI investigation, mild postero-lateral disc bulge at L5-S1 was found which caused left L5-S1 radiculopathy. His pain is aggravated by sitting for more than 30 minutes and relieved by walking or lying down. Otherwise, he has no numbness, tingling and weakness of the lower extremities. Upon physical examination, lumbar spine active range of motion is significantly restricted in forward bending and left side bending, both exacerbate left lower back and lower limb pain.

What is the most appropriate equipment to be used in this case?

- A. Mechanical traction
- B. Physio ball
- C. Lumbo-sacral corset
- D. Pelvic belt

ANSWER: A

125. A 69-year-old male patient came to physiotherapy OPD with right shoulder pain and difficulty to perform overhead daily activities. He has sustained a fall down injury two months ago. He underwent surgical repair of torn rotator cuff muscles. Upon evaluation of the right side shoulder, his muscle power was 2/5 on MRC scale.

What is the most appropriate exercise modality in this case?

- A. Shoulder press
- B. Bench press
- C. Shoulder pulley
- D. Floor dumbbell press

ANSWER: C

126. A 55-year-old woman, who has sustained a rupture of right anterior cruciate ligament two weeks ago, came to physiotherapy OPD. Two days ago, she underwent successfully completed an intra-articular reconstruction of ACL. The surgeon recommended her right knee joint flexion and extension range of motion to be initiated.

What is the most appropriate mechanical equipment to be used immediately after the procedure?

- A. Leg press with hyper extended knee joint
- B. Continuous passive ROM for knee joint
- C. Intermittent knee joint traction
- D. Compression therapy for knee joint

ANSWER: B

127. A 34-year-old man with bilateral lower limb weakness since one month ago was linked to physiotherapy unit. He was admitted to acute medical care unit and sent for chest x-ray that revealed normal. Currently, he is ambulating using two axillary crutches with left foot dragging on floor. On observation, the patient's left quad and calf appear to be atrophied than the right. The physiotherapist planned to do a baseline muscle strength record for tracking rehabilitation progression.

What is the most appropriate tool to measure muscle strength of knee flexors in this case?

- A. Oswestry Disability Index (ODI)
- B. Leg press
- C. Handheld dynamometer
- D. Quad bench

ANSWER: C

128. A 39-year-old man with complaint of lower cervical pain following cutting a lot of trees seven days ago was linked to physiotherapy OPD. The pain was unremitting pain in the midline of the lower cervical and across the upper trapezius area bilaterally. His active ROM was restricted and painful at the end of the ROM in all planes, and overpressure increases the pain.

What is the most likely contraindication to cervical traction?

- A. Unremitting neck pain
- B. Active range of motion restriction
- C. Pain in passive range of motions
- D. Pain at end of range of motion

ANSWER: A

129. A 45-year-old male carpenter has undergone surgical management of left ankle fracture four months ago. After removal of ORIF, he was linked to physiotherapy OPD with left ankle joint stiffness, weakness, balance while standing and walking. A physiotherapist has incorporated wobble board in the rehabilitation program.

What is the most likely purpose of incorporating the wobble board in the rehabilitation program?

- A. Reduce ankle joint edema and pain
- B. Enhance balance and proprioception
- C. Strengthen the ankle joint musculature
- D. Improve ankle joint range of motion

ANSWER: B

130. A 47-year-old man came to physiotherapy OPD with a diagnosis of right shoulder impingement syndrome. He has been working as a carpenter for the last 20 years. He has experienced pain gradually which is limiting him from doing activities that require movement above his shoulder level due to the pain. Upon physical examination, shoulder abductor and flexor muscle power was found 3/5 on MRC and Hawkin's and Neer test was positive.

Which is the most appropriate equipment that requires greater precaution in this case?

- A. Shoulder pulley
- B. Shoulder press
- C. Dumbbell floor press
- D. Arm curl

ANSWER: B

131. A 21-year-old male college student has sustained a traumatic cervical spinal cord injury 25 days ago. He had surgery of internal fixation and decompression using a right iliac bone graft to stabilize the fracture site. The surgeon linked him to physiotherapy clinic immobilizing with Philadelphia collar with consultation notes. Upon initial physical examination, he was able to extend his elbows, but could not flex his fingers of both hands and flex and extend his knees, and could not stand. The physiotherapist plan was to initiate locomotor training as early intervention. What is the most appropriate equipment to retrain walking in this case?

A. Parallel bar
B. Roller walker
C. Weight supported treadmill
D. Axillary crutch

ANSWER: C

132. A physiotherapist measured the range of motions (ROM) of 10 patients who had elbow contracture which they developed following the mass car accident injury. The ROM measurements taken are 165 degree, 140 degree, 125 degree, 125 degree, 90 degree, 145 degree, 125 degree, 125 degree, 90 degree, and 165 degree. What is the mean of the ROM measurements taken in the above scenario?

A. 139.5 degree
B. 129.5 degree
C. 125.0 degree
D. 165.0 degree

ANSWER: B

133. Top ten graduate batch physiotherapy students' thesis research score were 87, 90, 91, 81, 91, 80, 85, 76, 82, and 81. What is the range of students' thesis research score in the above scenario?

A. 17
B. 16
C. 15
D. 14

ANSWER: D

134. Randomly selected nine graduate batch physiotherapy students' thesis research exam scores were 83, 78, 83, 85, 96, 88, 78, 84, and 86 percent.

What is the mean of exam scores in the above scenario?

A. 84.45
B. 83.55
C. 84.55
D. 83.45

ANSWER: C

135. A physiotherapist conducted a study to determine the magnitude of clubfoot among newborns delivered in a rural health centers in 2023. After employing the required methodological procedures, the report showed that 25 had clubfoot among the 500 newborns during the study period.

What is the magnitude of clubfoot in this scenario?

- A. 20 percent
- B. 25 percent
- C. 10 percent
- D. 5 percent

ANSWER: D

136. A physiotherapist conducted a study and determined the number of newly diagnosed autistic disorders among 5000 under five-age children in 2023. After employing the required methodological procedures, the number of newly diagnosed autistic disorders children was 25 among children at risk in 2023.

What is the rate of newly diagnosed autistic disorders per 1000 in the above scenario?

- A. 5
- B. 10
- C. 15
- D. 20

ANSWER: A

137. A physiotherapist conducted a research among children with autistic disorder selected through systematic random sampling technique from an autism center. After the required sample size was calculated, data were collected by watching and recoding the behavioural acts they showed that characterize their main behaviours.

What is the most likely data collection tool used in this scenario?

- A. Observation
- B. Self-administered interviews
- C. Postal and telephone interviews
- D. Focus group discussions

ANSWER: A

138. A physiotherapist who is working at a hospital reported to the medical director of the hospital that 10 female and 15 male new clients with low back pain joined the hospital for physiotherapy service.

What type of frequency measure is used in the above scenario?

- A. Rate
- B. Ratio
- C. Count
- D. Proportion

ANSWER: C

139. A physiotherapist conducted a study on the new onset of lower back pain among patients with disc prolapse. The physiotherapist collected and analyzed the data after the study participants were followed for a month. Finally, 1.2 per 1000 (95 percent CI; 8, 12) low back pain was observed.

What is the most likely measure of lower back pain used in the above scenario?

- A. Prevalence
- B. Incidence
- C. Relative risk
- D. Odds Ratio

ANSWER: B

140. A physiotherapist, who is working at a hospital, reported the number of days that patients with lower back pain attend their physiotherapeutic service to the medical director of the hospital. The report of the number of days were as follows; 2, 3, 4, 2, 5, 4, 5, and 3.

What is the most appropriate measure of central tendency in the above scenario?

- A. Mean
- B. Median
- C. Variance
- D. Standard deviation

ANSWER: A

141. A head of physiotherapy department of a hospital receives a report indicating the number of days that patients with low back pain attend their physiotherapeutic service. The report was indicated as follows; 2, 3, 4, 2, 3, 4, 5, and 3.

What is the median of the patients' attendance in the above scenario?

- A. 2
- B. 3
- C. 4
- D. 5

ANSWER: B

142. A researcher wanted to determine the level of knowledge among female university students towards stroke. Thus, the researcher invited selected students to the hall and distributed the questionnaire simultaneously to all students to fill and return the questionnaire to him.

Which data collection technique is appropriate in the above scenario?

- A. Self-Administered
- B. Interviewer Administered
- C. Focused-Group Discussion
- D. Mailed questionnaire

ANSWER: A

143. An instructor who is working at a medical school follows a teaching approach which appropriately matches with real-life needs. The instructor realized that this type of approach will help the learners to acquire competence in using essential information in the decision making process for the analysis and solution of the problems.

What most appropriate educational method was used in the above scenario

- A. Problem-based learning
- B. Case-based learning
- C. Lecture-based learning
- D. Adult learning

ANSWER: A

144. A physiotherapist provided health education for patients with lower back pain. However, the physiotherapist realized that the patients did not get the necessary information and decided to re-educate them via a method that involves them through imitation, observation, feedback, analysis, and conceptualization.

What is the most appropriate educational principle used in the above scenario?

- A. Case-based
- B. Role play
- C. Interactive lecture
- D. Group discussion

ANSWER: B

145. A physiotherapist is conducting a research on the level of satisfaction of patients visiting outpatient physiotherapy department. He communicated to research participants that the research participants can discontinue or refuse to participate in the research at any time during the process.

What is the most likely ethical principle applied in this scenario?

- A. Justice
- B. Consent
- C. Beneficence
- D. Non-maleficence

ANSWER: B

146. A senior physiotherapist is working as a head of clinical department in a tertiary level hospital. He planned to make a short presentation about the Ethiopian health and health related institution standards and food standards for his staff in one of the afternoon sessions as part of an in-service training program.

What is the most likely standard the professional employed in this scenario?

- A. Company standards
- B. Voluntary standards
- C. Obligatory standards
- D. International standards

ANSWER: C

147. An assistant physiotherapy practitioner practicing at a district hospital while making a morning round came across a patient with total sensory loss of lower limb and he decided to make some positioning and posture adjustment to prevent pressure sore.

What is the most appropriate preventive action to be followed in this case?

- A. Allow clothes that are either too tight or too loose fitting
- B. Avoid clothes that are either too soft or comfortable
- C. Avoid clothes with thick seams and buttons in areas of pressure
- D. Allow clothes with thick seams, buttons, or zippers in areas of pressure

ANSWER: C

148. While a senior physiotherapist was in due course to conduct a physical examination for a 20-year-old woman who presented to an outpatient unit, he ensured the patient's privacy to undress and dress her body.

What is the most likely ethical principle demonstrated in this scenario?

- A. Dignity
- B. Equality
- C. Empathy
- D. Sympathy

ANSWER: A

149. A junior physiotherapist is working in outpatient physiotherapy department of a busy tertiary hospital. While providing care, the patients shouted and complained on the physiotherapist. However, the physiotherapist realized that the patients are under enormous stress and pain; as a result, he forgave them and continued his task.

What is the most likely ethical principle practiced in this scenario?

- A. Openness
- B. Empathy
- C. Sympathy
- D. Confidentiality

ANSWER: B

150. A 72-year-old female patient, who has undergone post-right transtibial amputation secondary to arteriosclerotic gangrene a day before, came to a physiotherapy clinic. On physical examination, she has independent rolling to left and to the right. She can also independently do supine-to-sit and return. What would be the most appropriate focus of a prosthetic training program for this patient?

- A. Teach this patient to go up and down steps
- B. Teach the patient how to cross an obstacle
- C. Start initial balance training program
- D. Evaluate the stump length

ANSWER: C

151. A physiotherapist working in a busy outpatient clinic received a referral of a 70-year-old man who recently had a stroke. This patient has limited mobility and is struggling with activities of daily living. The patient's daughter insists on intensive physiotherapy because she is concerned about her father's safety. However, the patient is hesitant about aggressive therapy.

What is the most appropriate ethical approach to be followed in this scenario?

- A. Prioritize client's autonomy and respect his wishes
- B. Prioritize client's concerns and push for proposed treatment
- C. Prioritize the well-being of the client regardless of principle
- D. Engage in open communication with both the client and his attendant

ANSWER: D

152. A 49-year-old male patient is referred to a prosthesis and orthosis center for fitting services. A physiotherapist is working as a member of the orthotic clinic team composed of a physician, physical therapist, and orthotist. After a detailed patient assessment, a decision was made to ensure an optimum match between the patient's biomechanical and psychological requirements and an appropriate orthosis capable of performing its intended function.

What is the most appropriate step that should be followed by the orthotic team?

- A. Prescription, evaluation, training on donning of the orthosis
- B. Training on donning of the orthosis, prescription, and evaluation
- C. Evaluation, training on donning of the orthosis, prescription
- D. Prescription, evaluation, training on donning of orthosis

ANSWER: A

153. A 50-year-old man who is a long truck driver has sustained car accident. He is referred to a physiotherapy clinic with a diagnosis of soft tissue injury to the right lateral side of upper arm.

What is the most appropriate initial step to be followed in this scenario?

- A. Ensure collaboration with the co-workers
- B. Obtain informed consent prior to intervention
- C. Start assessing and providing services to the patient
- D. Keep appropriate medical information about the treatment

ANSWER: B

154. A multidisciplinary team involves professionals from different disciplines who approach patient care from their own perspectives. All members of the multidisciplinary team hold separate individual consultations and meet regularly to discuss shared goals and patient findings.

What is the most likely explanation that shows the contribution of the team in this case?

- A. Integrates single discipline approaches into a single consultation
- B. Integrates multiple discipline approach into a multiple consultation.
- C. Assessment, intervention, and goals are conducted in the absence of the patient
- D. The team maps out prognosis, generate new ideas, and optimize patient care

ANSWER: D

155. A 35-year-old patient is referred to a physiotherapy clinic with a medical diagnosis of right side hamstring muscle strain. A junior physiotherapist working in the clinic made a complete assessments and evaluation for the patient and kept the records electronically.

What is the most appropriate component of record to be included in this case?

- A. Problem list
- B. Outcome measure
- C. Patient diagnosis
- D. Associated medical problems

ANSWER: C

156. A 45-year-old woman came to a physiotherapy clinic with a complaint of non-specific low back pain. She has three daughters and two sons and lives on the 3rd floor of government rented apartment. Her husband is the main earning member of the family.

What most appropriate physiotherapy process has to be followed in this context?

- A. Assess, evaluate, modify, implement, discharge
- B. Evaluate, modify, implement, discharge, assess
- C. Assess, plan, implement, discharge, evaluate, modify
- D. Assess, plan, Implement, evaluate, modify, discharge

ANSWER: D

157. A senior physiotherapist working in tertiary level hospital has received complaint from a concerned physician on the basis of specification of physiotherapy intervention that has been provided for chronic lower back pain.

What is the most appropriate component of documentation and registration that has to be included to solve the complaint?

- A. Clearly document which health service was provided at each client visit
- B. Separate and document the billing and financial records for each practice
- C. Communicate effectively with the client regarding the differences in health provider roles performed
- D. Separate appointment books and/or distinct days and times for providing each service

ANSWER: A

158. A new junior physiotherapist joined well organized physiotherapy team of a tertiary hospital. The team provided him a job description, advised him to have respectful, open, clear, and honest contact in all professional interactions, converse effectively with clients to promote their understanding of proposed services, and identify potential barriers to effective conversation and make a reasonable effort to address these barriers.

What is the most appropriate physiotherapy practice standard to be exercised?

- A. Standard of client-centered care
- B. Standard of competence
- C. Standard of communication
- D. Standard of collaborative practice

ANSWER: C

159. A 60-year-old female hypertensive patient came to a physiotherapy clinic with the complaint of being physically inactive. A physiotherapist working in the clinic assessed her body composition which was 80kg weight and 1.60m height.

What is the most likely health problem of this patient?

- A. Under weight
- B. Appropriate weight
- C. Over weight
- D. Obesity

ANSWER: D

160. A 58-year-old male hypertensive patient came to a health facility with complaints of blurred vision and pain at occipital area. After complete physical and laboratory examination, the physiotherapist noticed that the blood pressure of the patient is 160/100mmHg.

What is the appropriate blood pressure level category of this measurement?

- A. Normal diastolic pressure
- B. Pre-hypertension
- C. High hypertension grade 1
- D. High hypertension grade 2

ANSWER: D

161. A physiotherapist is planning a community-level health education at selected *Kebele* of a town. His aim is introducing the new idea about diabetes mellitus and making patients with diabetes mellitus to be familiar with signs, symptoms and complications of the disease.

What is the most likely health education approach to be used in this scenario?

- A. Informing
- B. Persuading
- C. Educating
- D. Entertaining

ANSWER: A

162. A physiotherapist working in rehabilitation center is planning public health emergency preparedness of poliomyelitis outbreak at a community level.

What is the most likely primary prevention of emergency preparedness?

- A. Detect victim
- B. personal hygiene
- C. Immunization
- D. Short-term counseling

ANSWER: C

163. A 29-year-old woman came to orthopedic clinic with complaints of chronic neck pain and weakness. Her history showed that she spends a most of her time sitting at the computer. On assessment, she developed a postural dysfunction.

What is the most likely ideal alignment of this patient?

- A. Head tilted or rotated
- B. Thoracic spine is slightly concave posterior
- C. Humeral head is posterior to acromion
- D. Thoracic spine is slightly convex posterior

ANSWER: D

164. A 30-year-old male teacher presented to physiotherapy OPD with ankylosing spondylitis. Physical examination shows a flattening of the lumbar lordosis, increased thoracic kyphosis and generalized hypo-mobility in all spinal movements as well as restricted chest expansion.

What is the most likely principle that has to be taken into account at early phase of exercise program?

- A. Postural correction programs
- B. Range of motion exercise programs
- C. Strengthening exercises programs
- D. Aerobic exercise programs

ANSWER: A

165. A physiotherapist, who is working as community based rehabilitation focal person, is screening a 50-year-old male patient during community level assessment to identify the appropriate rehabilitation measures. The patient is known diabetic patient and the right trans-tibial amputation was done before two months following a massive road traffic accident.

What is the appropriate prevention counseling to be provided to this patient?

- A. Primary prevention
- B. Secondary prevention
- C. Tertiary prevention
- D. Injury prevention

ANSWER: C

166. A 25-year-old male athlete came to a clinic. He is a known chronic diabetic mellitus patient and he is on anti-diabetic medication. He needs advice on prevention of disability related to hypoglycemia before doing exercise.

What is the appropriate preventive measure of this problem?

- A. Administer insulin less than one hour before exercise
- B. Administer insulin more than one hour before exercise
- C. If blood glucose greater than 100mg/dL, ingest 15g or more carbohydrate
- D. If blood glucose less than 250mg/dL, delay exercise and measure urine ketones

ANSWER: B

167. A 22-year-old football player came to a clinic with the complaint of right knee joint pain and inability to bear weight with the affected extremity. After physical examination, a physiotherapist notices that he has anterior cruciate ligament injury.

What is the most appropriate preventive strategy in this case?

- A. Decrease muscle activation
- B. Decrease landing forces
- C. Increase dynamic knee valgus
- D. Increase varus and valgus moments

ANSWER: B

168. A 40-year-old male office worker, who plays tennis once per week and likes to walk daily, came to a health facility with a complaint of pain in the midportion of his right Achilles' tendon. His pain has been gradually worsening over the past three months, and the care provider confirmed that, he has ankle instability.

What is the most likely functional disability factor in this case?

- A. Arthrokinetic restriction
- B. Synovial changes
- C. Degenerative changes
- D. Strength deficit

ANSWER: D

169. A 66-year-old office worker, presented with lumbar pain due to an acute disc protrusion at L4/5. Examination result reveals pain in early flexion and positive slump and straight leg raise on the left lower extremity but has no neurological sign.

What is the most likely health promotion and safety of this domain?

- A. Assess specific risks and barriers
- B. Recognize the need for continuity of treatment
- C. Develop treatment plans based on best evidence
- D. Evaluate clinical situations where standard treatment recommendations

ANSWER: A

170. A senior rehabilitation expert conducts training for health care professionals working in disability rehabilitation centers. The rehabilitation expert explained that there are domains of Knowledge required for Disability Management (DM) Professionals. Based on the discussion of the expert, it is incumbent that clinicians who wish to expand their role into DM need to possess working knowledge that typically resides in non-medical domains. These domains include legal, human resources, technology, risk management, ethics, management, insurance, reimbursement, labor relations, business, and accreditation.

Which is the most likely explanation of the ethical domain?

- A. Patient abandonment
- B. Treatment protocols
- C. Workers' compensation
- D. Medicare fraud and abuse

ANSWER: A

171. A final year graduating class physiotherapy students conducted internship program in a community about how to increase community awareness on disability management. The students describe the basic concepts about disability management and the basic difference between disability management and case management.

Which is the most appropriate explanation about disability management?

- A. Focus on population
- B. Focus on individual
- C. Focus on high-cost or catastrophic
- D. Applied to acute conditions

ANSWER: A

172. A manager of a tertiary hospital wants to achieve planned goals of the year. To be in line with the goals, he arranged a weekly meeting. In each meeting he motivates, inspires, and encourages the employees.

What is the most likely managerial function exercised in this scenario?

- A. Planning
- B. Organizing
- C. Controlling
- D. Leading

ANSWER: D

173. In a newly established organization the board members conducted election and assigned a person with a job description which emphasizes the legitimate right to make decisions, give orders, and allocate resources to achieve the desired outcomes.

What component of organizing is undertaken by the board?

- A. Accountability
- B. Responsibility
- C. Authority
- D. Delegation

ANSWER: C

174. A chief clinical director of a tertiary hospital had a meeting with the head of department of physiotherapy and received formal complaint of extremely high number of daily patient flow in the OPD. Hence, the department faces challenges to accomplish different daily departmental activities. On his informal assessment, the director interviewed and observed that there was high flow of patients and long waiting time among patients. In addition he observes that a single physiotherapist is performing the whole task in the unit.

What most appropriate management function should be improved?

- A. Planning
- B. Organizing
- C. Staffing
- D. Leading

ANSWER: C

175. A medical director of a general hospital had signed a memorandum of agreement for clubfoot management with orthopedic specialist, physiotherapist and orthotics units. In the agreement, 320 patients are planned to receive club foot management service within six months. To ensure the effectiveness of the planned activity the medical director continuously checks the monthly performance against the plan and took corrective actions when necessary. He ensured that the plans remained on track periodically.

What is the most appropriate management function exercised by the director?

- A. Organizing
- B. Controlling
- C. Staffing
- D. Leading

ANSWER: B

176. A newly appointed leader of a national physiotherapy association conducted a general meeting to share his transformational strategic plan of the next five years. In the meeting, he is friendly, approachable and exhibits consideration for the status, wellbeing and needs of members.

What most appropriate leader behavior is shown in this case?

- A. Directive leadership
- B. Supportive leadership
- C. Participative leadership
- D. Achievement oriented leadership

ANSWER: B

177. A senior physiotherapist working as a unit head of a general hospital received an award for being the best performer of physiotherapy activities in the hospital. The physiotherapist has an outstanding knowledge and skill on how to perform activities. Additional to his academic achievements, he also periodically updates his skills with trainings and short term courses.

What most appropriate managerial skill does the physiotherapist possess?

- A. Technical Skills
- B. Practical skill
- C. Human Skill
- D. Conceptual Skill

ANSWER: A

178. A district level hospital manager planned to start physiotherapy service in the coming years. The service acquires different human resources, infrastructures and finance. To start the service, he revised the strategic plan document as an initial step. The established team as a second phase wants to assess the organization's current capacity, availability of resources, organizational limitations, external enabling and disabling factors to start the service.

What most appropriate strategic planning tool should the team use?

- A. Stakeholder analysis
- B. SWOT analysis
- C. Situational analysis
- D. Problem analysis

ANSWER: B

179. On the fourth quarter of the 1st year of a project implementation, the project manager assessed the progress of the project by comparing the plan against the achievement. The manager identified major areas of improvement. He then called meeting to key project staff members and made further discussion to take corrective measures for the identified gaps.

What is the most likely management function exercised by the manager?

- A. Monitoring
- B. Evaluation
- C. Reporting
- D. Progress

ANSWER: A

180. The physiotherapy association hired an evaluator who will work on evaluation of a project which ended up before two years. The main aim of the project was improving the life of physically disabled women in the community. The evaluator assessed the effectiveness to the beneficiaries through different mechanism. He measured the quality of life of physically disabled women who were the primary beneficiaries and founded a greater improvement from the baseline data. Additionally the women's family and society as a whole showed a great improvement from the baseline state. Finally the team recommended scaling up the project encompassing wide areas.

What most appropriate type of evaluation is implemented by the evaluator?

- A. Impact evaluation
- B. Input evaluation
- C. Outcome evaluation
- D. Process evaluation

ANSWER: A

181. A senior manager of a primary hospital planned to conduct performance evaluation on physiotherapy service across all wards including outpatient department. He worked out the plan that shows when and how to evaluate the service. He then developed objectives and methods to perform the evaluation activity.

What most appropriate evaluation step should the manager follow next?

- A. Report
- B. Application
- C. Intervention
- D. Data collection

ANSWER: D

182. A senior physiotherapist planned to open undergraduate physiotherapy program in one of the re-known universities in 2025. In line with other necessary activities, he designed a need assessment for relevance of opening the program from the community and stakeholders.

What is the most appropriate purpose of the need assessment for the above case?

- A. Assess current situation
- B. Assess progress towards goals and objectives
- C. Assess program output
- D. Assess impact of the program

ANSWER: A

183. Federal ministry of health was preparing a five years strategic plan with staff and stake holders. After the discussion the team formulated list of goals regarding health care services that have to be achieved in the upcoming years.

What most appropriate goal can be stated for the above case?

- A. To eradicate protein-energy malnutrition among risky target populations
- B. To increase the caloric intake among infants by 30 percent in the district by year three
- C. To reduce prevalence of anaemia in pregnant women by 30 percent in Ethiopia by 2015
- D. To improve the quality of HIV counselling services at ANC clinics in the district

ANSWER: A

184. A manager working on monitoring and evaluation was in the process of identifying appropriate performance indicators depending on five necessary criteria. Therefore he selected one important performance indicator that allows consistent assessment over time and among different observers.

What most appropriate criterion is used to select the performance indicator?

- A. Clarity
- B. Reliability
- C. Validity
- D. Measurability

ANSWER: B

185. A 40-year-old-male construction worker came to physiotherapy clinic complaining of right shoulder pain that lasted for one week. He spent the weekend painting his house. Since then, he reported pain in the anterior and lateral aspect of his right shoulder. He described the pain in his shoulder as a deep dull aching but sharp during active elevation of his arm. He had difficulty of sleeping on his right shoulder at night. Upon physical examination Neer's test was positive and tenderness of the greater tubercle along the bicipital groove was evident.

What is the most appropriate diagnosis for this case?

- A. Bursitis
- B. Biceps tendinitis
- C. Impingement syndrome
- D. Cervical degenerative disk disease

ANSWER: C

186. A 35-year-old female cleaner was presented to a physiotherapy outpatient department with complaints of paresthesia on the thumb, index, and middle fingers. Prior to the onset of paresthesia she was working on a laundry machine. She also claimed a history of repetitive trauma on her hand. Upon physical examination, the physiotherapist noticed positive result of Phalen's test.

What is the most likely diagnosis of this case scenario?

- A. Carpal tunnel syndrome
- B. DeQuervain's tenosynovitis
- C. Cubital tunnel syndrome
- D. Radial nerve entrapment

ANSWER: A

187. A 37-year-old female patient came to physiotherapy outpatient department with complaint of consistent numbness and paresthesia of right forearm. The pain worsens with pronation and supination of hand or grip-related activities and she experienced gradual weakness of the distal forearm muscles. On examination there was weakness of thenar intrinsic and median innervated extrinsic muscles.

What is the most likely diagnosis of this case?

- A. Radial nerve entrapment
- B. Median nerve entrapment
- C. Ulnar nerve entrapment
- D. Posterior interosseous nerve entrapment

ANSWER: B

188. A 65-year-old male patient who has been diagnosed with adhesive capsulitis came to a tertiary level hospital. He complained shoulder pain, stiffness, and restriction of motion. Also, he mentioned difficulty of overhead activities and inability to perform day to day activities. He had a history of diabetes mellitus and sedentary lifestyle. After investigation, the patient was referred to a physiotherapist for consultation.

What is the most likely primary goal of therapeutic intervention for this case?

- A. Introducing moderate-vigorous intensity exercise
- B. Grade III or IV joint-play mobilization
- C. Progressive resistance strengthening exercise
- D. Pain free functional range of movement exercise

ANSWER: D

189. A 53-year-old male construction worker came to a tertiary level hospital with complaint of increasing pain and discomfort on his right shoulder. An MRI investigation result for continued symptoms, showed abnormal supraspinatus tendon with probable rotator cuff tear. Rotator cuff tear repair was done. Then, he was referred to physiotherapy for consultation on his 3rd of post operative week.

What is the most appropriate therapeutic management to be done in this case?

- A. Pendulum exercise
- B. Resistant band exercise
- C. Carrying and transferring objects
- D. Free weight exercise

ANSWER: A

190. A 60-year-old female patient came to physiotherapy clinic with complaint of spontaneous onset of shoulder pain, which is often severe and disrupting sleep pattern for 20 weeks duration. She often rests the arm and notes a little bit relief of pain, but increased stiffness with rest. Upon assessment, she had restricted range of motion in a characteristic pattern of loss of external rotation, internal rotation, and abduction. She was under physiotherapy follow up receiving TENS, range of motion in the pain-free range.

What is the most appropriate prognostic phase for this case?

- A. Freezing phase
- B. Stiffing phase
- C. Thawing Phase
- D. Frozen phase

ANSWER: A

191. A 45-year-old male instructor came to physiotherapy clinic with complaint of right anterior shoulder pain when overthrowing or wrenching of his arm. He was treated with Rest, Ice, Compression, Elevation (RICE) protocol and was reassessed. Upon reassessment, apprehension and posterior capsule tests were negative, but the pain was still consistent. After thorough re-evaluation, isolated capsular and labral tests were found to be positive.

What is the most appropriate prognostic decision to be made for this case scenario?

- A. Conservative management
- B. Modification of intervention
- C. Referral for further evaluation
- D. Rest and splint

ANSWER: C

192. A 32-year-old male came to physiotherapy clinic with complaint of anterolateral hip pain, aggravated by prolonged sitting, leaning forward, and pivoting motions. A CT scan result showed an abnormal articulation of the acetabular rim and the proximal femur. On physical exam, there was limited range of motion due to pain. The flexion, adduction, and internal rotation tests were positive.

What is the most likely diagnosis of this case?

- A. Osteoarthritis of the hip
- B. Iliotibial band syndrome
- C. Femoral neck stress fracture
- D. Femoroacetabular impingement

ANSWER: D

193. A 55-year-old male who has been diagnosed with osteoarthritis came to physiotherapy clinic. He claims persistent knee pain at both weight bearing and rest. His pain is frequently associated with tightness, morning stiffness less than 30 minutes, and often awakens at night. However, he gets relief after motion, his knee becomes painful and stiff again once the weight-bearing tolerance of the joint is exceeded by prolonged standing or walking.

What is the most appropriate systemic cause for being diagnosed with osteoarthritis?

- A. Nutritional deficiency
- B. Joint malalignment
- C. Repetitive micro trauma
- D. Obesity

ANSWER: A

194. A 52-year-old male merchant who was diagnosed with rheumatoid arthritis came to physiotherapy clinic with complaint of morning stiffness in and around bilateral elbow, knee, and ankle joints. The x-ray investigation confirmed typical of rheumatoid arthritis on hand, wrist and knee joints. Upon physical examination, he experienced limited mobility and signs of inflammation including pain, redness, swelling, and warm joints.

What is the most likely therapeutic goal to be set to improve independence in this case?

- A. Passively maximize sufficient range of motion for all joints
- B. Maximize joint stability to decrease biomechanical stress on all affected joints
- C. Promote patient training in activities related to daily living
- D. Educate family and caregivers to support the patient in activities related to daily living

ANSWER: C

195. A 20-year-old male patient came to physiotherapy clinic with a moderate medial knee ligament injury or Grade-II Sprain while taking shower two days ago. He has acute pain. He had no history of associated medical illness and trauma. On examination, knee joint pain, swelling, and tenderness are noticed.

What is the most appropriate treatment plan to be done in this case?

- A. Progressive exercise techniques and functional activities
- B. Grade I and II mobilization
- C. Avoid varus stress
- D. Protection to allow healing

ANSWER: D

196. A 30-year-old female bank officer presented to physiotherapy clinic with complaint of right lateral ankle pain a day after an injury. The pain gets worse when she attempted to walk long distance, and climb up and down stairs. Upon physical examination, the physiotherapist noticed significant swelling and tenderness over the involved ligaments and soft tissues.

What is the most important treatment plan to be done in this case scenario?

- A. Functional bracing
- B. Weight bearing
- C. Protect, rest, ice compression, elevation
- D. Prolonged use of walking boot or cast

ANSWER: C

197. A 48-year-old traffic police man came to physiotherapy clinic with complaints of heel pain while walking on hard surface and climbing up and down stairs of four months duration. He is obese weighing 95 kilogram. An x-ray investigation revealed right plantar calcaneal spur. On examination, there is localized tenderness at the plantar fascial attachment of the calcaneus, in the medial arch area and in the abductor hallucis muscle. Also there is limited range of motion of the great toe and ankle dorsiflexion.

What is the most important initial therapeutic management to be done in this case?

- A. Ice massage, rest, and anti-inflammatory medication
- B. Concentric loading exercises for the gastrocnemius and arch musculature
- C. Open kinetic chain activities and neuromuscular activities
- D. Grade III-IV joint mobilizations and hip adduction exercises

ANSWER: A

198. A 71-year-old, obese woman came to a physiotherapy clinic with complaint of right hip pain. She feels the pain when hip is internally rotated, and morning stiffness spans less than an hour. She received isometric and isotonic exercise, as well as proprioceptive and balance training. However, she bothered about the escalating nature of the pain during standing and prolonged walking.

What is the next most appropriate prognostic approach to be done on this case?

- A. Neuromuscular and functional training exercise for lower extremities
- B. Home exercise programs including exercise booster sessions
- C. Walking and stationary bicycles
- D. Static to dynamic balance activities

ANSWER: A

199. A three-year-old female child who has been diagnosed with cerebral palsy came with her mother to a physiotherapy clinic. The child experienced upper and lower limb weakness, poor vision, and tightness to lengthen her extremities. On assessment, there is limited range of motion both in upper and lower limb joints and ankle joint contracture.

What is the most likely motor type subjective marker for this abnormal developmental milestone?

- A. Spastic
- B. Hemiplegic
- C. Diplegic
- D. Quadriplegic

ANSWER: A

200. A six-month-old male infant who has been diagnosed with spinal defect came with his mother to physiotherapy clinic. Upon examination, an external sac is noticed on his back at the lumbar region. Also he lost his mobility in all settings.

What is the most likely clinical manifestation to be seen on this case?

- A. Absence of bilateral motor and sensory function below the level
- B. Absence of bilateral motor and sensory function above the level
- C. General body weakness
- D. Swallowing difficulty

ANSWER: A