



Pediatrics and Child Health Nursing

Time left 2:25:44

Question 1

Not yet answered

Marked out of 1.00

[Flag question](#)

A woman gave birth to a 1200 gram baby in her 36th week of gestation. The baby was admitted to an intensive care unit immediately after birth. What is the most likely classification of this newborn?

- ☐ a. Very low birth weight
- ☐ b. Very preterm
- ☐ c. Low birth weight
- ☐ d. Extremely low birth weight

Time left 2:23:21

Question 2

Not yet answered

Marked out of 1.00

[Flag question](#)

A school age child was presented to a pediatric clinic with a chief complaint of cough of five days duration. On assessment, the child had maculo-papular rash, mouth ulcer, poor appetite, history of recurrent pneumonia and diarrheal diseases. The child was diagnosed to have vitamin A deficiency disorders. What is the most likely cause of vitamin A deficiency in this case?

- ☐ a. Measles infection
- ☐ b. Recurrent pneumonia
- ☐ c. Poor appetite
- ☐ d. Recurrent diarrhea

Time left 2:20:07

Question 3

Not yet answered

Marked out of 1.00

[Flag question](#)

A two-year-old child is brought to a health center due to diarrhea of six days duration. The child has night blindness, bitot's spots, and corneal ulceration. The child has also weight for height less than -3 Z-score, MUAC less than 11.5 cm, and bilateral pitting edema. A nurse plans to treat vitamin A deficiency as recommended by IMNCI protocol. What is the most appropriate management for this child?

- ☐ a. Give two doses on day 1 and 2.
- ☐ b. Give single dose RUTF, since RUTF has vitamin A
- ☐ c. Give three doses of Vit A on day 1, 2 and 15
- ☐ d. Give single dose on day 1 if child didn't get Vitamin A in the last one month.

Time left 2:19:29

Question 4

Not yet answered

Marked out of 1.00

[Flag question](#)

A pediatric nurse researcher conducted a study on adherence level of HIV medication among children who are on ART follow up. The researcher selected 400 study participants from 4000 children on ART who are on follow up from the registration logbook by taking an assumption of all children on ART follow up have similar characteristics. What is the most likely sampling technique used by the researcher?

- ☐ a. Multistage sampling
- ☐ b. Simple random sampling
- ☐ c. Cluster sampling
- ☐ d. Stratified sampling

Time left 2:18:11

Question 5

Not yet answered

Marked out of 1.00

[Flag question](#)

A four-year-old child was presented to pediatric emergency room with a complaint of scald burn of chest and abdomen of two hours duration. On examination, the child weighs 20kg, and has full thickness burn of 20 percent of total body surface area. A Physician ordered maintenance fluid in addition to emergency fluid. What is the total fluid estimate required to manage this child for the first eight hours?

- ☐ a. 1,550ml
- ☐ b. 1,650ml
- ☐ c. 1,250ml
- ☐ d. 1,450ml

Time left 2:17:32

Question 6

Not yet answered

Marked out of 1.00

[Flag question](#)

A Four-year-old child was brought to an emergency department of a hospital after sustaining head injury of one hour duration. On assessment, the child has history of one episode of seizure and two episodes of vomiting, loss of consciousness, had nasal bleeding, slow and irregular breaths and dilatation of the pupils. The nurse assessed, traumatic brain injury and stabilized him in a recommended position. What is the next most appropriate next nursing intervention for this child?

- ☐ a. Check pupil reaction every 4 hours
- ☐ b. Seek medical attention
- ☐ c. Awaken twice during the night
- ☐ d. Assess pain and give analgesics

Question 7

Not yet answered

Marked out of 1.00

[Flag question](#)

A nurse who works in a pediatric ICU is challenged by shortage of human resource, work overload, increasing death of new born, malfunctioning equipments and overcrowded environment. The nurse frequently requested a hospital's management to solve the issues, but it did not act on it, rather they focused on grand projects. As these conditions become worsen, the nurse gradually exhibited low self-esteem, cynicism, and total negativism to his work. What is the most likely term that most likely describes current status of the nurse?

- ☐ a. Stress
- ☐ b. Burnout
- ☐ c. Absenteeism
- ☐ d. Turnover

Question 8

Not yet answered

Marked out of 1.00

[Flag question](#)

A woman brought 11-month-old infant to a nearby health center for immunization. A nurse evaluated the developmental milestones and appreciated that the gross motor, fine motor and cognitive developmental levels are normal. However, the child has delayed language development. What is the most likely indicator of delayed development in this child?

- ☐ a. Failure to speak understandable sentences
- ☐ b. Saying "dada" and "mama" for anyone
- ☐ c. Saying dada for his/her dad and mama for his/her mom
- ☐ d. Failure to speak two word sentence

Question 9

Not yet answered

Marked out of 1.00

🚩 Flag question

A four year-old child has been brought to a well baby clinic for growth assessment. The anthropometric measurements have been taken and revealed that; MUAC=13.5cm, BMI for age =3.1 Z-score, weight for age=2.5 Z-score and height for age =0.9 Z-score. What is the most appropriate nursing diagnosis in this case?

- ☐ a. Obesity
- ☐ b. Risk for stunting
- ☐ c. Risk for overweight
- ☐ d. Overweight

Question 10

Not yet answered

Marked out of 1.00

🚩 Flag question

A 10-year-old child was presented to emergency room with a complaint of skin injury secondary to human bite and treated as an outpatient. On the initial assessment, the child had a 2cmx3cm lacerated wound on his right hand. On the first day, a nurse cleaned and sent home with a follow-up schedule. After two days, the child came for follow-up. What is the most likely indicator of good progress of the child's condition?

- ☐ a. Absence of wound discharge
- ☐ b. No sign of wound inflammation
- ☐ c. Closure of the wound
- ☐ d. No sign of systemic infection

Question 11

Not yet answered

Marked out of 1.00

🚩 Flag question

A school age child is brought to a health center due to a complaint of frequent fight with colleagues at school. The class room teacher told to his family that the child is impulsive and hyperactive in class. Accordingly, the mother suspected for behavioral disorder and brought the child to a health center. A nurse examining the child, and suspected attention deficit hyperactivity disorder. What is the most likely additional symptom that supports the nurse's diagnosis?

- ☐ a. Lack of concentration
- ☐ b. Delay in language development
- ☐ c. Poorly integrated verbal communication
- ☐ d. Strike out at people

Question 12

Not yet answered

Marked out of 1.00

🚩 Flag question

A 34-week-old-newborn with a birth weight of 2000 gram was brought to a hospital 30 minutes after birth in a health center. The newborn has is not able to suck the breast and a temperature of 36.3 degree centigrade. The respiratory rate was also 62 breaths per minutes. What is the most likely risk factor for the newborn's problem?

- ☐ a. Hypothermia
- ☐ b. Born in health center
- ☐ c. Low birth weight
- ☐ d. Prematurity of organs

Question 13

Not yet answered

Marked out of 1.00

🚩 Flag question

A woman who gave birth at 36 weeks of gestation brought her two- day-old newborn due to fast breathing of one day duration. The newborn had severe chest indrawing, jaundice, irritability and heart rate of 200 beats/minute. Finally, the neonate is diagnosed with newly onset neonatal sepsis. What is the most appropriate antibiotic in this case?

- ☐ a. Ampicillin and Gentamycine
- ☐ b. Amoxicillin and corticosteroids
- ☐ c. Corticosteroids and vancomycine
- ☐ d. Ceftriaxone and vancomycine

Question 14

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A school of nursing wants to assess student's knowledge, skill and attitudes towards patient care domain. They selected 6 clinical procedures and every student has performed all the selected tasks. Without performing all the tasks, students would not be promoted to the next academic program. What is the most appropriate assessment method for the above scenario?

- ☐ a. Structured feedback report
- ☐ b. Log books
- ☐ c. Structured practical examination
- ☐ d. Clinical rounds

Question 15

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A woman came to a hospital and gave birth of a 2000 gram baby at her 35 weeks of gestation. On examination, the baby was hypoxic and had increased peristalsis, passage of meconium, and relaxation of anal sphincter. The baby was diagnosed as meconium aspiration syndrome. What is the most appropriate finding that justifies this diagnosis?

- ☐ a. Hypoxia
- ☐ b. Relaxation of anal sphincter
- ☐ c. Passage of meconium
- ☐ d. Increased peristalsis

Question 16

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A woman gave birth a 2300 gram baby at 38 weeks of gestation at a health center . The baby's temperature and breathing rate were 35.5 degree centigrade and 40 breaths/minute, respectively. What is the most appropriate nursing intervention for this case?

- ☐ a. Apply cord care
- ☐ b. Provide resuscitation
- ☐ c. Keep the newborn warm
- ☐ d. Bathing the neonate

Question 17

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A woman brought her four-month old infant to a nearby health center with complaints of crying and irritability for the last 24-hours. The pediatric nurse suddenly moved the infant towards top of the bed and the child startled. What is the most likely reflex type exhibited by this child?

- ☐ a. Moro reflex
- ☐ b. Rooting reflex
- ☐ c. Grasping reflex
- ☐ d. Suckling reflex

Question 18

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

In a community where there is a belief that holy water can be the treatment of HIV/AIDS because HIV/AIDS is the result of sin. Hence, taking ART drugs wouldn't have any improvement for people living with HIV. What is the most likely determinate of health behavior in the above scenario?

- ☐ a. Enabling factor
- ☐ b. Predisposing factor
- ☐ c. Influencing factor
- ☐ d. Reinforcing factor

Question 19

Not yet answered

Marked out of 1.00

🚩 Flag question

A mother brought a three-year-old child to a health center due to generalized body rash which manifested in the facial area. Besides, the child has fever, runny nose and a temperature of 39 degree centigrade and respiratory rate of 36 breaths per minute. Then, the nurse managed the child's problems accordingly. What is the most important indicator of good prognosis after treating this child?

- ☐ a. Normal body temperature
- ☐ b. Total clearance of body rash
- ☐ c. Increased respiratory rate
- ☐ d. Absence of runny nose

Question 20

Not yet answered

Marked out of 1.00

🚩 Flag question

A health professional wants to conduct screening program on elementary school for the disease of scabies. Each school members were assessed for the disease based on case definition without specific reference to their individual risk of having or developing the disease. What is the most appropriate type of screening?

- ☐ a. Selective screening
- ☐ b. Mixed type of screening
- ☐ c. Opportunistic screening
- ☐ d. Mass screening

Question 21

Not yet answered

Marked out of 1.00

🚩 Flag question

A pediatric nurse working in a general hospital registered 10 neonates weight (in kg) who was born in a single day. The weights of the neonates are 2.8, 4.0, 3.5, 3.6, 4.1, 2.5, 2.8, 3.8, 2.9 and 3.0. What is the mean weight of the neonates born in the hospital?

- ☐ a. 4.1
- ☐ b. 2.8
- ☐ c. 3.0
- ☐ d. 3.3

Question 22

Not yet answered

Marked out of 1.00

🚩 Flag question

A 50-year of old man came to a health center with a complaint of persistent cough for more than 3 weeks. He has productive cough and positive for AFB. In addition, he has loss of weight, chest pain and night sweats. What is the most appropriate control mechanism TB disease in the above scenario?

- ☐ a. Even though it has minimal benefit, educate the public about adequate nutrition
- ☐ b. Isolation is very important, when individuals are at risk for the disease.
- ☐ c. Avoid overcrowding and health education on mode of transmission of the disease
- ☐ d. Immunization of the patient with measles

Question 23

Not yet answered

Marked out of 1.00

🚩 Flag question

A four-year-old child was admitted to a hospital with severe acute malnutrition and provided appropriate treatments. On monitoring, weight for height greater than -2 - Z-score, weight for age greater than -2 Z-score and height for age of =-2 Z-score after two weeks of treatments. What could be the most reliable criteria to discharge this child?

- ☐ a. Weight for age greater than -2 Z-score with no edema
- ☐ b. Weight for height greater than -2 Z-score with no edema
- ☐ c. MUAC = 12.5 cm with no edema
- ☐ d. Height for age greater than -2 Z-score with no edema

Question 24

Not yet answered

Marked out of 1.00

🚩 Flag question

A pediatric and child health nursing department wants to assess the nutritional status of the infants in a district. A nurse collected weight measurement, length measurement and head circumference of nine infants in the district. What is the most appropriate nutritional assessment method for the above scenario?

- ☐ a. Clinical
- ☐ b. Dietary
- ☐ c. Anthropometric
- ☐ d. Biomedical

Question 25

Not yet answered

Marked out of 1.00

🚩 Flag question

A four-year-old child was brought to a hospital with a complaint of unconsciousness of one hour duration. On examination, the child had snoring sound and secretions drooling from his mouth. What nursing diagnosis has to be prioritized of this child?

- ☐ a. Risk for aspiration
- ☐ b. Impaired gas exchange
- ☐ c. Ineffective airway clearance
- ☐ d. Risk for airway blockage

Question 26

Not yet answered

Marked out of 1.00

🚩 Flag question

A woman spontaneously gave birth after 30 hours of labour initiation at 38 weeks of gestation. The baby has signs of meconium aspiration and gasping with a respiratory rate of 20 breaths/minute. The mother had no medical problem during pregnancy. The pediatric nurse diagnosed perinatal asphyxia. What is the most likely cause of asphyxia for this newborn?

- ☐ a. Intrapartal causes
- ☐ b. Antepartal causes
- ☐ c. Prenatal causes
- ☐ d. Postpartal causes

Question 27

Not yet answered

Marked out of 1.00

🚩 Flag question

A Pediatric nurse is monitoring the immunization data for a given population in X Health Center. The total catchment population of the health center was 2000 and the annual infant target population was found to be 60. What is the estimated monthly target infant population for this health center?

- ☐ a. 5
- ☐ b. 12
- ☐ c. 15
- ☐ d. 6

Question 28

Not yet answered

Marked out of 1.00

🚩 Flag question

A five-years-old child was presented to emergency room with a complaint of Bee sting before one hour. On assessment, the child had shallow breathing, generalized body swelling, grunting and lethargic. Otherwise the child had no history of chronic illness. What is the most appropriate prioritized nursing intervention for this child?

- ☐ a. Provision of IV fluid
- ☐ b. Monitoring of vital signs
- ☐ c. Provision of diuretics
- ☐ d. Management of airway

Question 29

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A three-week-old infant presented to hospital with a complaint of diarrhea of one week duration. Upon assessment for hydration status, skin pinch goes back slowly, and the infant moves only when stimulated. The infant also has restlessness, bloody stool, and poor suckling. Based on the results, the nurse classified the infant problems as dysentery using an IMNCI protocol. What was the most likely finding to confirm this classification?

- ☐ a. Blood in stool
- ☐ b. Decreased skin turgidity
- ☐ c. Minimal fluid intake
- ☐ d. Restlessness

Question 30

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A child with cerebral palsy was admitted to a pediatric ward for rehabilitation. A nurse diagnosed impaired physical mobility related to neuromuscular impairment and started to give care to correct the problem with regular evaluation of the treatment responses. What is the most likely indicator of good prognosis to the nurse's management?

- ☐ a. Child start to play with colleagues
- ☐ b. Child demonstrates active muscle movement
- ☐ c. Child remains injury free
- ☐ d. Child has adequate mobility to perform ADLs

Question 31

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A health facility received 300 doses of DTP-HepB-Hib vaccine in a 1 dose vial in January. During monthly reporting, 150 children were found to be recorded as immunized. They had 40 doses as a start balance on 1 January and by 31 January their stock level was 70 doses. What is the percentage of-vaccine wastage in this facility?

- ☐ a. 30 percent
- ☐ b. 50 percent
- ☐ c. 44 percent
- ☐ d. 60 percent

Question 32

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A three-year-old child with severe acute malnutrition was admitted to the stabilization center and put on F-75 therapeutic food via NG-tube and appropriate antibiotics. The child had failed appetite test and bilateral pitting edema at admission. However, after seven days the NG-tube was removed because the child was able to take F-75 orally. A nurse planned to transfer the child to transition phase after one week of admission. What could be the most likely patient condition to transfer him to transition phase?

- ☐ a. No medical problems
- ☐ b. Edema starts to subside
- ☐ c. Weight gain of 10 g/kg/day
- ☐ d. The child finishes all F-75 milk easily

Question 33

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A nine-month-old infant was brought to a child health clinic for immunization. On assessment, a nurse noticed poor positioning and breast attachment with late initiation of complementary feeding. The lab result showed; Hct of 30percent and Hgb of 10mg/dl. The infant has also some palmar palor with body temperature of 37.9 degree centigrade. What is the most likely classification of this child according to IMNCI?

- ☐ a. Anemia
- ☐ b. Very Severe anemia
- ☐ c. No Anemia
- ☐ d. Severe Anemia

Question 34

Not yet answered

Marked out of 1.00

🚩 Flag question

A five-year-old child is referred from a health center to a hospital due to complicated severe acute malnutrition. On assessment, the child has limpness, lethargic, and unconsciousness with extracted eye-lids later after admission. Blood glucose level=42 mg/dl, T0= 35.8 degree centigrade, and RR=32 breaths/minute. What nursing intervention has to be prioritized in this case?

- ☐ a. Put the child on 100 percent oxygen
- ☐ b. Provide F-75 via NGT
- ☐ c. Provide 50 ml 10 percent dextrose via NGT
- ☐ d. Keep the child covered and put away from draughts

Question 35

Not yet answered

Marked out of 1.00

🚩 Flag question

A woman gave birth to a term bay in a hospital. The newborn developed yellowish discoloration of the skin and sclera six days after delivery and the baby treated for jaundice. However, the baby is brought back to hospital after a year due to hearing loss. What is the most likely complication developed in this neonate?

- ☐ a. Stage 2 acute bilirubin encephalopathy
- ☐ b. Stage 3 acute bilirubin encephalopathy
- ☐ c. Chronic bilirubin encephalopathy
- ☐ d. Stage 1 acute bilirubin encephalopathy

Question 36

Not yet answered

Marked out of 1.00

🚩 Flag question

A nurse working in pediatric ward inserted NG-tube for a child. Immediately after insertion, she documented the procedure. While documenting procedures and other care provided, the nurse used many lines and fluid to make the error unreadable. She used common abbreviations with clear and readable hand writing and signature on the correction. What documentation element was most likely mal-practiced?

- ☐ a. Documentation timing
- ☐ b. Error removal
- ☐ c. Vocabulary use
- ☐ d. Signature around error

Question 37

Not yet answered

Marked out of 1.00

🚩 Flag question

A newborn is brought to a neonatal intensive care unit with a complaint of poor breast feeding. The nurse found auxiliary temperature of 30 degree centigrade, pulse rate of 100 breaths/minute and respiratory rate of 36 breaths/minute. Consequently, the newborn was diagnosed for hypothermia. What is the most likely hypothermia classification of this newborn?

- ☐ a. Mild hypothermia
- ☐ b. Extremely severe hypothermia
- ☐ c. Severe hypothermia
- ☐ d. Moderate hypothermia

Question 38

Not yet answered

Marked out of 1.00

🚩 Flag question

A woman brought her twelve-month-old-child to a health center with a complaint of failure to stand independently; but, the child is able to creep and can say two-syllable sounds. The child also tries to drink from cup and rarely waves bye-bye. The height for age of the child resides at the 90th percentile on WHO growth chart. What nursing intervention has to be prioritized for sthis child?

- ☐ a. Counseling the mother to train the child about how to feed self
- ☐ b. Counseling the mother to encourage and train the child to stand
- ☐ c. Counseling the mother to encourage crawling and creeping
- ☐ d. Counseling the mother to bring the child to take the child to rehabilitation center

Question 39

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A six-year-old child was presented to a health center with a complaint of diarrhea of three days. The child had history of frequent episodes of diarrhea and the child is from a family of low socioeconomic status. There is also lack of health care service in the community where the child is born. The child is also found to have growth faltering. What is the immediate cause of growth faltering in this child?

- ☐ a. Poor health care service
- ☐ b. Social insecurity
- ☐ c. Frequent attack by diarrhea
- ☐ d. Low economic status of the family

Question 40

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

An epidemic investigation team got a report from health extension worker about unexpected high occurrence of malaria cases in a district. The team tried to compare the observed and the expected case of malaria occurrence at the district by considering seasonal variation. What steps of epidemic investigation most likely describes the above scenario?

- ☐ a. Prepare for field work
- ☐ b. Verifying the diagnosis
- ☐ c. Describe the epidemic with respect to time, place, person
- ☐ d. Verifying the existence of an epidemic

Question 41

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A two-year-old child was brought to a hospital with swelling of feet of four days duration. The child has recent weight gain due to edema, weight for age less than -3 Z-score, MUAC less than 11.5 cm. The child had no medical complication and passed the appetite test. The child did not take vaccine and a nurse planned to provide measles vaccine along with treatments for malnutrition. What is the most appropriate schedule to give measles vaccine for this child?

- ☐ a. Give at the 1st week of treatment
- ☐ b. Give at the 4th week of treatment
- ☐ c. Give at the 6th week of treatment
- ☐ d. Give at the 3rd week of treatment

Question 42

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A Pediatric Nurse was caring a five-year-old child. The child was brought to pediatric emergency with the diagnosis of poisoning as a result of carbon monoxide inhalation before one hour. On assessment, the child looks confused had headache, visual disturbances, irritability, nausea, pallor and cyanosis. What is the most appropriate initial nursing intervention for this child?

- ☐ a. Secure IV line for rehydration
- ☐ b. Put the child in coma position
- ☐ c. Determine blood gas level
- ☐ d. Put on humidified 100 percent oxygen

Question 43

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A 15-year-old boy with a diagnosis of meningitis is requested to have x-ray and lumbar puncture. However, the boy refused x-ray procedure. A nurse in charge said to the boy, if you don't go through the diagnostic procedures I will give you an injection and leave you without any care. What is the most likely legal liability exercised by the nurse in charge?

- ☐ a. Malpractice
- ☐ b. Negligence
- ☐ c. Battery
- ☐ d. Assault

Question 44

Not yet answered

Marked out of 1.00

[Flag question](#)

A two-year old child with birth weight of 2500 gram and birth length of 50 cm was presented to a health center due to poor food intake. A nurse measures the weight and height of the baby and concluded that the child has no problem in growth. Assessment findings; weight= 9.5 kg, height=102cm, weight for age= +1 Z-score, weight for height = +1 Z-score. What most appropriately indicates normal growth in this child?

- ☐ a. weight for height = +1 Z-score
- ☐ b. weight for age= +1 Z-score
- ☐ c. weight= 9.5 kg
- ☐ d. height=102cm

Question 45

Not yet answered

Marked out of 1.00

[Flag question](#)

A medical director of a private hospital planned to conduct clinical audit. Through the audit process, the director provided orientation for newly organized clinical audit team to improve the quality and effectiveness of clinical care. Lastly, the director emphasized that selection of the right person with the right skills was the overall purpose of the audit. What step of clinical audit plan is best explained in the above scenario?

- ☐ a. Delivering of audit fieldwork
- ☐ b. Determining the audit topic
- ☐ c. Identifying available resources
- ☐ d. Involving potential stakeholders

Question 46

Not yet answered

Marked out of 1.00

[Flag question](#)

A mother brought a one-month-old neonate with a complaint of poor weight gain. On assessment, the nurse found that poor attachment and poor positioning were major problems. The weight for age was less than -2 Z-score. The nurse classified the case as underweight and advised the mother to give home care for the infant and breastfeed often as per the need, and appointed the mother for follow up. What is the next most appropriate schedule for this infant?

- ☐ a. After 14 days
- ☐ b. After 5 days
- ☐ c. After 2 days
- ☐ d. After 7 days

Question 47

Not yet answered

Marked out of 1.00

[Flag question](#)

A researcher wants to conduct a research on knowledge, attitude and practice towards neonatal intensive care unit management protocol among junior nurses. In the previous study, the level of good knowledge among junior nurses was 40 percent with 95 percent CI and marginal error of 4 percent. What is the sample size required for this research?

- ☐ a. 384
- ☐ b. 864
- ☐ c. 576
- ☐ d. 368

Question 48

Not yet answered

Marked out of 1.00

[Flag question](#)

A child, who was presented to a health center with eye pain, lethargy, or behavioral changes, was assessed to have visual impairment. The child was admitted to a hospital, but become anxious to the unfamiliar environment. What is the most appropriate expected outcome of this condition?

- ☐ a. Child's coping strategy is improved
- ☐ b. The child accustom with the new environment
- ☐ c. The child gets free of visual impairment a day after treatment
- ☐ d. Child verbalizes comfort with environment

Question 49

Not yet answered

Marked out of 1.00

🚩 Flag question

A four-year-old child is brought to a hospital due to suspected growth and developmental retardation. On assessment, the child can hop, balance on each foot, can draw a person, and is able to identify the color of objects. The child is also able to brush teeth without any help, but cannot understand the opposite of things. What is the most likely nursing recommendation for this child?

- ☐ a. Treat the child for cognitive developmental delay
- ☐ b. Treat the child for gross motor developmental delay
- ☐ c. Inform the mother that the child is developing normally
- ☐ d. Treat the child for fine motor developmental delay

Question 50

Not yet answered

Marked out of 1.00

🚩 Flag question

A neonate is going to have surgical repair for tracheo-esophageal fistula. A surgeon requested a nurse in charge to provide consent form to the neonate's mother about the neonate's care. What is the most likely important question that the nurse will forward to the mother?

- ☐ a. Do you know that there are alternative options other than surgery?
- ☐ b. Did the surgeon have told you that your child is going to have operation?
- ☐ c. Did you understand the procedure for which they are giving consent?
- ☐ d. Did the surgeon have told you the possible harms related to the procedure?

Question 51

Not yet answered

Marked out of 1.00

🚩 Flag question

A three-week-old neonate was brought to under-five OPD with umbilical swelling of three days duration. The infant has cough, red and warm umbilicus with a body temperature of 36.7 °C, and respiratory rate of 56 breaths per minute. The neonate has no danger signs. A nurse wanted to classify the problem based on IMNCI protocol. What is the most likely classification of the newborn's problem?

- ☐ a. Pneumonia
- ☐ b. Very severe disease
- ☐ c. Local infection
- ☐ d. Local bacterial infection

Question 52

Not yet answered

Marked out of 1.00

🚩 Flag question

A nine-month-old infant was brought to a well-baby clinic for growth and development assessment. On examination, a nurse appreciated that the child cannot crawl/creep and is not able to speak a two-word sentence. The baby is also not able to stand and walk and then the nurse appreciated developmental delay. What is the most likely indicator of delayed development in this child?

- ☐ a. Not able to walk
- ☐ b. Not able to creep/crawl
- ☐ c. Not able to stand
- ☐ d. Not able to speak two-word sentence

Question 53

Not yet answered

Marked out of 1.00

🚩 Flag question

A preterm infant with complaints of poor feeding and cyanosis is admitted to a hospital. On assessment, the baby had jaundice, tachypnea, lethargy and hypotonia. Laboratory findings revealed excessive protein in the urine and decreased sodium excretion with peripheral venous blood and HCT of 78 percent. Hence, partial volume exchange transfusion was ordered for this neonate. What is the ultimate goal of this intervention?

- ☐ a. Removing 5ml of blood at a time
- ☐ b. Removing 20ml of blood at a time
- ☐ c. Removing 15ml of blood at a time
- ☐ d. Removing 10ml of blood at a time

Question 54

Not yet answered

Marked out of 1.00

🚩 Flag question

A 35-year-old woman came to a health center with complaints of sudden onset of vomiting and bloody diarrhea. She has also abdominal cramp and fever. When a nurse takes history, she told that she ate unrefrigerated food prepared before a day. What is the most appropriate control mechanism for the above scenario?

- ☐ a. Temporarily exclude food handlers with purulent lesion
- ☐ b. Keep cleanliness of the hands
- ☐ c. Cover the wounds found on the skin
- ☐ d. Reduce food handling time of food

Question 55

Not yet answered

Marked out of 1.00

🚩 Flag question

A six-year-old child was presented to a hospital with complaints of agitation and changes in level of consciousness of two days duration. On assessment, the child has history of frequent fever, headache and general malaise and history of street dog bite three weeks before. What is the most appropriate management for this case?

- ☐ a. Treating with broad spectrum antibiotics
- ☐ b. Administration of rabies vaccine
- ☐ c. Provision of antipsychotic drugs
- ☐ d. Provision of symptomatic management

Question 56

Not yet answered

Marked out of 1.00

🚩 Flag question

A school age child was presented to a pediatric nurse with visible wasting and diarrhea for more than two weeks duration. On assessment, the child had weight for age is less than -2 Z-scores, and anorexia due to poor taste for foods. A nurse diagnosed diarrhea and suspected that the child might have zinc deficiencies. What is the most important clinical finding that supports the diagnosis of diarrhea in this child?

- ☐ a. Anorexia due to poor taste
- ☐ b. Weight for height z-scores
- ☐ c. Weight for age z-scores
- ☐ d. Visible wasting

Question 57

Not yet answered

Marked out of 1.00

🚩 Flag question

A nurse working at immunization unit of a Health Center wants to store the following vaccines in the refrigerator; hepatitis B vaccine, Rota vaccine (liquid) vaccine, inactivated poliovirus vaccine, and haemophilus influenza type b (liquid) vaccine. What is the most likely vaccine that has to be stored away from the freezing point threshold?

- ☐ a. Inactivated poliovirus vaccine
- ☐ b. Rota vaccine (liquid) vaccine
- ☐ c. Haemophilus influenza type b (liquid) vaccine
- ☐ d. Hepatitis B vaccine

Question 58

Not yet answered

Marked out of 1.00

🚩 Flag question

A nurse works in a hospital which has only one dialysis machine. Two children who need dialysis came to the hospital at the same time. The nurse in charge raised question of ethical dilemma and gathered relevant data to come up with a solution for the ethical dilemma. What is the next most appropriate step to be taken by the nurse in this case?

- ☐ a. Value clarification
- ☐ b. Identifying course of action
- ☐ c. Negotiation plan
- ☐ d. Problem verbalization

Question 59

Not yet answered

Marked out of 1.00

🚩 Flag question

A two-year-old child was presented to a pediatric OPD with a complaint of inability to understand simple instructions like 'roll the ball', 'wear shoe'. The child plays with colleagues rarely and runs properly. Also, the child was unable to pay attention to simple stories and songs. What is the most likely nursing diagnosis for this child?

- ☐ a. Delayed physical growth
- ☐ b. Delayed gross motor development
- ☐ c. Delayed fine motor development
- ☐ d. Delayed child development

Question 60

Not yet answered

Marked out of 1.00

🚩 Flag question

A woman brought her 18-day-old infant with a complaint of big head. The nurse assessed and found full and tense fontanel, markedly separated cranial sutures, broad forehead and the head circumference was at 95th percentile. Skull x-ray showed; widening of sutures, prominent convolutional markings on the inner part of the skull and erosion of the sella turcica. What is the most likely diagnosis of the infant?

- ☐ a. Encephalocele
- ☐ b. Hydrocephalus
- ☐ c. Anencephaly
- ☐ d. Brain tumor

Question 61

Not yet answered

Marked out of 1.00

🚩 Flag question

A 4-year-old boy was admitted to a pediatric medical ward for the management of severe pneumonia. A nurse in charge, did not give medications correctly as per the standard of the hospital. In the morning shift, another nurse noticed that the medication given was below the standard. However, the morning shift nurse did not do anything to correct the malpractice. What legal role the nurse in the morning shift most likely failed to accomplish?

- ☐ a. Provider of service
- ☐ b. Citizen
- ☐ c. Employee or contractor for service
- ☐ d. Contractual obligation

Question 62

Not yet answered

Marked out of 1.00

🚩 Flag question

A nursing service manager in a comprehensive specialized hospital is well-known in his public relations, staff benefits, advocacy, activity in different national meetings, and being figurehead of the organization. What managerial role is practiced by the nurse service manager in the above scenario?

- ☐ a. Decisional Role
- ☐ b. Informational role
- ☐ c. Developmental role
- ☐ d. Interpersonal role

Question 63

Not yet answered

Marked out of 1.00

🚩 Flag question

A seven-year-old child was brought to a primary hospital with complaints of fever and shortness of breath. A physician ordered blood sample to be taken from the child, but the nurse faced difficulty of getting appropriate vein though the veins are visible at different sites. What most likely code of ethics should be considered in this case?

- ☐ a. The nurse provides services with respect for human dignity
- ☐ b. The nurse assumes accountability for individual nursing judgments
- ☐ c. The nurse safeguards the client's right to privacy
- ☐ d. The nurse fails to maintain competence in nursing

Question 64

Not yet answered

Marked out of 1.00

🚩 Flag question

An investigator wants to assess the prevalence of leprosy among children in a district with a sample size of 576. Before starting the actual data collection, the data collectors administered the questioner for 29 respondents in another district. Finally, the investigator corrected unclear and ambiguous words from the questionnaire. What is the most likely data quality assurance technique used by the researcher?

- ☐ a. Pre-test
- ☐ b. Pilot study
- ☐ c. Training
- ☐ d. Supervision

Question 65

Not yet answered

Marked out of 1.00

🚩 Flag question

A child developed measles case and a pediatric nurse reported this to an epidemic investigation team. In the following days, all children who had contact with the child developed measles and the measles it became an outbreak in a district. What type of outbreak most likely represents this scenario?

- ☐ a. Propagated outbreak
- ☐ b. Continuous source outbreak
- ☐ c. Intermittent source outbreak
- ☐ d. Point source outbreak

Question 66

Not yet answered

Marked out of 1.00

🚩 Flag question

A four-year-old boy was admitted to a pediatrics ICU. The boy needs emergency surgery. However, his parents are not around to give consent. A nurse worried about getting valid informed consent. What is the most likely reason for lack of valid consent?

- ☐ a. The timing to have surgery
- ☐ b. The capacity of the boy to give consent
- ☐ c. Means to get voluntary consent
- ☐ d. The ability of the nurse to give adequate information

Question 67

Not yet answered

Marked out of 1.00

🚩 Flag question

A nurse working at the immunization unit of a health center was handling vaccines. The nurse record, monitor the temperature based on cold chain standard of the refrigerator and check the vaccine vial monitor(VVM) regularly. The nurse received measles vaccine supply from the woreda health office. What is the most appropriate storage area in the refrigerator for measles vaccine?

- ☐ a. Middle compartment
- ☐ b. Upper compartment
- ☐ c. Freezing compartment
- ☐ d. Lower compartment

Question 68

Not yet answered

Marked out of 1.00

🚩 Flag question

A 2000 gram baby was born at 36 weeks of gestation. On assessment, the baby suckles breast poorly and the respiratory rate, oxygen saturation and axillary temperature were 62 breaths per minute, 80 percent, and 36.2 degree centigrade respectively. What is the most appropriate nursing intervention for this baby?

- ☐ a. Administer oxygen to correct the respiratory rate and desaturation
- ☐ b. Position newborn to facilitate drainage of secretions
- ☐ c. Put the baby on trophic feeding till the respiratory rate improves
- ☐ d. Put the baby on kangaroo mother care

Question 69

Not yet answered

Marked out of 1.00

🚩 Flag question

A 24-month-old child was brought to under-five OPD with a complaint of discharge from the ear for seven days duration. At the time of arrival, the child cries and pulls out her ear. On assessment, pus draining from the child's ear was observed and there was a swelling and tenderness behind the ear. What is the most effective management of this child according to IMNCI?

- ☐ a. Give Amoxicillin for five days
- ☐ b. Treat with topical quinolone ear drops for two weeks
- ☐ c. Give first dose of Ceftriaxone
- ☐ d. Dry ear by wicking

Question 70

Not yet answered

Marked out of 1.00

🚩 Flag question

A nine-year-old child came to emergency OPD after car accident over the left femoral bone of one hour duration. On assessment, the child has active bleeding, shallow respiration, temperature of 36.2 degree centigrade, RR of 35 breaths per minute, PR of 88 beats per minute and cold extremity. Radiological finding confirmed fracture of the femoral bone. What nursing diagnosis has to be prioritized for this child?

- ☐ a. Deficient fluid volume
- ☐ b. Risk for post-trauma syndrome
- ☐ c. Ineffective breathing
- ☐ d. Risk for infection

Question 71

Not yet answered

Marked out of 1.00

🚩 Flag question

A medicine and health science college in a university is popular in producing competent and qualified health professionals for the last 15 years. In this academic year, the college signed an agreement to work on a staff development and technology transfer with a foreign university. What component of SWOT is taken into account when signing an agreement?

- ☐ a. Strength
- ☐ b. Weakness
- ☐ c. Threats
- ☐ d. Opportunity

Question 72

Not yet answered

Marked out of 1.00

🚩 Flag question

A 2000 gram neonate was born at 35 weeks of gestation is vigorously and was admitted to an intensive care unit. On assessment, the baby is found to have difficulty of breathing secondary to meconium aspiration syndrome. What is the most appropriate nursing intervention for this neonate?

- ☐ a. Providing oxygen
- ☐ b. Providing resuscitation
- ☐ c. Providing tracheal suctioning
- ☐ d. Providing routine intubation

Question 73

Not yet answered

Marked out of 1.00

🚩 Flag question

A researcher wants to study the effect of F-100 nutritional program on malnourished patients. There is a promising possibility for patients to recover from the disease, but there is also a condition that will make it worse. The average recovery time of malnutrition for patients is 3 weeks. What is the most appropriate alternative hypothesis for the above scenario?

- ☐ a. F-100 nutritional program will not be given for patients with malnourished
- ☐ b. F-100 nutritional program effect on average time of recovery is by chance
- ☐ c. F-100 nutritional program has no any difference on the average time of recovery
- ☐ d. F-100 nutritional program results in difference on the average time of recovery

Question 74

Not yet answered

Marked out of 1.00

🚩 Flag question

A six-year-old child came to a pediatric emergency ward after sustained abdominal trauma because of car accident six hours ago. On assessment, the child has distended abdomen, dry mucus membrane, pallor, cold extremity, has no external bleeding, blood pressure of 60/40mmhg, heart rate of 100 beats/min, temperature of 36 degree centigrade, weight of 20 kg and capillary rifling greater than 3 seconds. What is the most appropriate nursing intervention for this child?

- ☐ a. Normal saline 400ml bolus fluid
- ☐ b. Dextrose in normal saline 400ml bolus fluid
- ☐ c. Dextrose in normal saline 500ml bolus fluid
- ☐ d. Normal saline 500ml bolus fluid

Question 75

Not yet answered

Marked out of 1.00

🚩 Flag question

A seven-day neonate who was admitted to a hospital has extremely low chance of survival and that even if he has the chance of living, he will probably be severely handicapped. A pediatric nurse explained his view in terms of utilitarian theory when requested for professional opinion on the management plan; and stating whatever it is, to do a best work for everyone even in the expense of the neonate's life. What ethical principle best supports the attitude of the nurse?

- ☐ a. Non-maleficence
- ☐ b. Justice
- ☐ c. Autonomy
- ☐ d. Beneficence

Question 76

Not yet answered

Marked out of 1.00

🚩 Flag question

A nurse attending labor assessed a newborn immediately after birth and appreciated that the baby cries well with a normal respiratory rate. Then, the nurse dried the baby and put on the mother's abdomen after confirming the normal breathing pattern of the baby. What is the next most appropriate step in the management of this newborn?

- ☐ a. Apply Chlorohexidine on the cord
- ☐ b. Give vitamin intramuscular K
- ☐ c. Initiate breast feeding
- ☐ d. Give TTC eye ointment

Question 77

Not yet answered

Marked out of 1.00

🚩 Flag question

A two-week-old-infant presented to a health center with a complaint of not feeding well of one day duration. Upon assessment, she has fast breathing, severe chest indrawing, and high body temperature of 38.6°C. The child moves only when stimulated. The nurse wanted to decide an optional management of the neonate based on IMNCI protocol. What is the best optional management for the neonate's problem?

- ☐ a. Advise mother to give home care for the infant
- ☐ b. Treat the child with amoxicillin
- ☐ c. Treat to prevent low blood sugar
- ☐ d. Refer urgently to hospital

Question 78

Not yet answered

Marked out of 1.00

🚩 Flag question

An eight-year-old child is brought to a hospital with a complaint of chronic skin rash. On assessment, the child has photosensitive dermatitis and signs of depressive psychosis. A nurse suspected nutritional deficiency though the anthropometric indicators are in normal range. What is the most likely nutritional deficiency related to the nurse's diagnosis in this case?

- ☐ a. Thiamin deficiency
- ☐ b. Niacin deficiency
- ☐ c. Folate deficiency
- ☐ d. Riboflavin deficiency

Question 79

Not yet answered

Marked out of 1.00

🚩 Flag question

A mother brought her three-day-old newborn to an emergency OPD with a complaint of swelling over her infant's head in the past three days. On assessment, the infant was delivered by assisting episiotomy and vacuum, none tender swelling of the head that does not cross the suture line, no fractured bone and no symptoms of jaundice. What is the most appropriate nursing intervention for this infant ?

- ☐ a. Reassure the mother as it will resolve after 3 week
- ☐ b. Start the prescribed antibiotics immediately
- ☐ c. It needs surgical consultation for intervention
- ☐ d. It needs to be aspirated by Needle

Question 80

Not yet answered

Marked out of 1.00

🚩 Flag question

A term baby is born from a mother who had HIV/AIDS, tuberculosis, and hepatitis C infection during pregnancy. On top of that, the mother was also on chemotherapy as a treatment of cancer. The mother wanted to feed breast milk, but the nurse opted formula feeding. What is the most likely condition that gave rise to formula feeding choice for this infant?

- ☐ a. Mother had HIV/AIDS
- ☐ b. Mother had Hepatitis c infection
- ☐ c. Mother had tuberculosis
- ☐ d. Mother being on chemotherapy

Question 81

Not yet answered

Marked out of 1.00

🚩 Flag question

A 10-year-old-child was admitted with brain injury was gradually developed into unconsciousness. On monitoring, the nurse noted a temperature of 35 degree centigrade, pulse rate of 88 beats/min and respiratory rate of 28 breaths/min , random blood sugar of 95 mg/dl and blood pressure of 110/70mmHg. What is the most appropriate parameter that indicates the poor prognosis of this case?

- ☐ a. Blood pressure
- ☐ b. Pulse rate
- ☐ c. Respiratory rate
- ☐ d. Temperature

Question 82

Not yet answered

Marked out of 1.00

🚩 Flag question

A four-year-old child was brought to a health center with complaint of cough of three days duration. The child has oxygen saturation of 89 percent, fast breathing and convulsions on examination. The clinician wants to classify the child's problems based on IMNCI. What is the most likely classification of the child's problem?

- ☐ a. Severe infection
- ☐ b. Severe Pneumonia
- ☐ c. Cough or cold
- ☐ d. Pneumonia

Question 83

Not yet answered

Marked out of 1.00

🚩 Flag question

A six-year-old child was brought to a hospital with complaints of fatigue and pale conjunctiva for two weeks duration. On examination, the child had low hemoglobin level (Hgb less than 10 g/dl), underweight, and pale palm. The morphologic findings implied hypo-chromic and hypo-cystic anemia. The folate and vitamin B-12 levels of the child were in normal range. What could be the most likely cause of anemia in this case?

- ☐ a. Iron deficiency
- ☐ b. Parasitic infection
- ☐ c. Megaloblastic disease
- ☐ d. Hemolysis of RBCs

Question 84

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A nurse in delivery unit of a health center attended the delivery of 1600 gm newborn who was born at 36th weeks of gestation. The newborn was actively sucking and had a good muscle tone. What initial management should be provided for this child according to IMNCI?

- ☐ a. Start Kangaroo mother care
- ☐ b. Refer urgently with the mother to Hospital
- ☐ c. Appoint follow up visit at 6th hour
- ☐ d. Give expressed breast milk by cup

Question 85

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A pediatric nurse, who is a member of epidemic management team, assigned to investigate cholera epidemic that occurred in a district. The epidemic management team focused its activity on limiting the number of cases and geographical spread of cholera, shortening the duration of the outbreak and reducing fatalities due to cholera. What public health management phase most likely describes the above scenario?

- ☐ a. Disaster recovery
- ☐ b. Disaster response
- ☐ c. Disaster mitigation
- ☐ d. Disaster preparedness

Question 86

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A nine-year-old child was admitted to a burn center because of sustaining fire burn of the face and chest. The child was given an eight hour emergency fluid. On follow-up assessment, the nurse found that the child had regular breathing, sloughed skin on the chest, erythematous skin and edematous face. What is the most appropriate nursing diagnosis of this child?

- ☐ a. Impaired tissue integrity
- ☐ b. Risk for impaired airway clearance
- ☐ c. Risk for skin integrity
- ☐ d. Impaired gas exchange

Question 87

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A 57-month-old child is brought to a pediatrics OPD with a complaint of cough of two weeks duration. On assessment, the child has lethargy, bilateral pitting edema, and MUAC of 11 cm. Vital signs shows; respiration rate is 46 beats/minute, Temperature is 39 degree centigrade, and pulse rate is 80/minute. What is the priority nursing intervention in this case?

- ☐ a. Admit the child to pediatric ward and administer oxygen
- ☐ b. Refer the child for management of SAM in OTP
- ☐ c. Admit the child to stabilization center and treat complications
- ☐ d. Admit the child to stabilization center and initiate therapeutic feeding

Question 88

Not yet answered

Marked out of 1.00

🚩 [Flag question](#)

A mother brought her premature newborn delivered at 35 weeks of gestation who weighs 2000 grams. The infant was not sucking properly. Vital Signs shows; RR=70 breaths/minute, T0= 37.5 degree centigrade, PR=100 beats/minute. The mother was in a state of confusion on how to feed her infant and asked a nurse on how to help her infant. What is the most appropriate nursing intervention in this case?

- ☐ a. Educate the mother about the importance of good attachment and initiate direct feeding
- ☐ b. Provide expressed breast milk via cup
- ☐ c. Provide expressed breast milk through NG tube
- ☐ d. Counsel the mother to give formula milk till the baby is able to suck the nipple

Question 89

Not yet answered

Marked out of 1.00

[Flag question](#)

A primi gravida mother came to antenatal care clinic on her 37th weeks of gestation and a nurse asked her about child feeding practice preparedness. What is the initial information the mother to be asked?

- ☐ a. The values of timely initiation of breast feeding
- ☐ b. The nutrient contents of breast milk
- ☐ c. Possible feeding options
- ☐ d. The effect of maternal food to the nutritional quality of the breast milk

Question 90

Not yet answered

Marked out of 1.00

[Flag question](#)

A 15-year-old child was brought to pediatric emergency after motor vehicle accident over both lower extremities. On assessment, child has history of severe bleeding, blood pressure of 80/40mmhg, RR of 28/min, PR of 90/min, temperature of 36 degree centigrade, Hg of 5g/dl and shallow breathing. What could be the most likely diagnosis of this child?

- ☐ a. Hypovolemic shock
- ☐ b. Obstructive shock
- ☐ c. Anaphylactic shock
- ☐ d. Septic shock

Question 91

Not yet answered

Marked out of 1.00

[Flag question](#)

A result section of a published article showed that the prevalence of perinatal asphyxia among neonate admitted at intensive care unit was found to be 25 percent with 95 percent CI; 20-28. No ANC follow up (AOR=2.5, 95 percent CI; 2- 4.5), stunting (AOR=3.5, 95 percent CI; 2.5-6.8), preterm birth (AOR= 4.3, 95 percentCI; 3-8.5), and low birth weight (AOR=2.8, 95 percentCI; 1.8-7.4) were associated factors of perinatal asphyxia among neonates admitted at intensive care unit in a hospital. What is the most appropriate research objective in the above scenario?

- ☐ a. To determine the prevalence of perinatal asphyxia and understand factors of perinatal asphyxia among neonates admitted at intensive care unit
- ☐ b. To assess the prevalence and associated factors of perinatal asphyxia among neonates admitted at intensive care unit
- ☐ c. To study the prevalence of perinatal asphyxia and associated factors among neonates admitted at intensive care unit
- ☐ d. To examine the potential reasons of perinatal asphyxia and its prevalence among neonates admitted at intensive care unit

Question 92

Not yet answered

Marked out of 1.00

[Flag question](#)

A Six-year-old child came to an emergency OPD with a complaint of barking cough OF one day duration. On examination, she had audible stridor at rest, supra sternal and sternal retraction at rest but on fever and agitation. What is the most likely diagnosis of this child?

- ☐ a. Acute spasmodic laryngitis
- ☐ b. Bacterial tracheitis
- ☐ c. Impending respiratory failure
- ☐ d. Acute laryngotracheobronchitis

Question 93

Not yet answered

Marked out of 1.00

[Flag question](#)

An asphyxiated newborn is brought to a newborn corner for resuscitation. A nurse provided positive pressure ventilation for 30 seconds and reassessed the baby. The reassessment findings were labored breathing with a heart rate of 70 beats per minute as well as poor muscle tone. What is the next most appropriate step in the management of this newborn?

- ☐ a. Reposition and continue bag and mask ventilation
- ☐ b. Start chest compression with continued bag and mask ventilation
- ☐ c. Start plasma volume expander with bag and mask ventilation
- ☐ d. Provide medications that can improve the heart rate

Question 94

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse is taking care of a girl who grief due to lose of her mother. During their conversation, the girl says, I am very upset after I lost her; I have difficulty while speaking to stranger. the nurse replied that I see that you find it difficult when you are taken to new persons. What therapeutic communication techniwues is most likely exercised by the nurse?

- ☐ a. Seeking of clarification
- ☐ b. Provision of general lead
- ☐ c. Paraphrasing
- ☐ d. Being specific and tentative

Question 95

Not yet answered

Marked out of 1.00

 [Flag question](#)

A ten-year-old child was brought to an emergency OPD after sustaining fire injury over posterior trunk and lower extremities. On assessment, the child had PR of 96 beats/min, RR of 38 breaths/min and temperature of 36.4 degree centigrade. The child also had decreased urine output, dry skin, unable to move the lower extremity, and sustained deep partial-thickness burn. Nearly 20 percent of total body surface area was affected. What is the most likely complication that this child is suffering from?

- ☐ a. Shock
- ☐ b. Infection
- ☐ c. Disabling contractures
- ☐ d. Pulmonary edema

Question 96

Not yet answered

Marked out of 1.00

 [Flag question](#)

A head nurse of a neonatal intensive care unit is passionate and role model, even at the time of crisis. The head nurse tackled issues which may lead to discouragement among his co-workers and most of the time the head nurse considers problems as an opportunity. What effective leadership quality is manifested by the head nurse in the above scenario?

- ☐ a. Generosity
- ☐ b. Perseverance
- ☐ c. Optimism
- ☐ d. Balance

Question 97

Not yet answered

Marked out of 1.00

 [Flag question](#)

A school of nursing in a university wants to appoint a new dean for the school. There were two applicants who have been department heads in different program of the school. Even though, both applicants have better chance to win the position than others, they end up with conflict and physical fight. As a result, the director and committee members screened them to be out of the competition. What conflict resolution strategy is applied in the above scenario?

- ☐ a. Win-win
- ☐ b. Win-lose
- ☐ c. Lose –lose
- ☐ d. Lose-win

Question 98

Not yet answered

Marked out of 1.00

 [Flag question](#)

A child with multiple, intense, disproportionate, or irrational worries, often about future events is brought to a pediatric clinic. After thorough evaluation by the nurse, the child is diagnosed to have generalized anxiety disorder (GAD). What should be the most appropriate priority nursing intervention?

- ☐ a. Refer the child for psychotherapy
- ☐ b. Provide antipsychotic drugs
- ☐ c. Give advice to the family to support the child
- ☐ d. Provide cognitive behavioral therapy

Question 99

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A six-week-old infant was brought to an immunization clinic for the second follow-up visits. On assessment, the infant is stable and looks well, no significant reaction to the previous dose except mild skin lesion. A nurse provided the appropriate vaccination and counseled the mother to bring the infant for the next schedule. What is the most appropriate schedule for this child?

- ☐ a. After four weeks
- ☐ b. After five weeks
- ☐ c. After three weeks
- ☐ d. After six weeks

Question 100

Not yet
answered

Marked out of
1.00

🚩 [Flag question](#)

A school of nursing wants to launch generic oncology nursing undergraduate program. In order to make the learner focused, a committee member was determining the content and learning methods. The curriculum includes knowledge, effectiveness and cognitive learning outcomes in measurable terms that easily understandable by the learners. What is the most appropriate step of curriculum development for the above scenario?

- ☐ a. Problem identification
- ☐ b. Educational strategy
- ☐ c. Targeted need assessment
- ☐ d. Forming goals and objectives

[Previous page](#)

[Finish attempt ...](#)