

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

SURGICAL NURSING



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH • ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, Leadership and Management, Ethics, and Health Promotion as outlined in the Information Booklet 2025.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Surgical Nursing

1. A 35-year-old patient, who has a history of digoxin therapy, is scheduled to undergo abdominal surgery. The nurse responsible for preparing the patient for surgery got the results of preoperative investigations ready. The serum potassium level is 3.2mmol/L. (NR: 3.5mmo/L – 4.5mmol/L). What is the most appropriate immediate action regarding this result?

A. Administer potassium supplement
B. Inform the surgical team about the result
C. Encourage dietary intake of potassium
D. Proceed with the surgery, as it does not matter

ANSWER: B

2. A male patient scheduled for vascular surgery is preoperatively investigated for routine tests. Considering the case, a nurse planned to add a diagnostic procedure. What special diagnostic procedure is most appropriate for this patient?

A. Electrocardiogram
B. Doppler ultrasound
C. X-ray
D. Echocardiography

ANSWER: B

3. A male patient, who has undergone extensive surgery on his upper back a week ago, is forced to have another emergency surgery for appendicitis.

What is the most likely condition that puts this patient at risk?

A. Anxiety related to perioperative events
B. Perioperative positioning injury
C. Impaired tissue integrity
D. Excessive bleeding

ANSWER: B

4. A 69-year-old male patient came to the emergency department with severe abdominal pain. After evaluation, bowel perforation was diagnosed. Significant medical history includes controlled hypertension and smoking for 50 years. A nurse is preparing the patient for surgery.

What is the most appropriate nursing intervention to be included in the plan of this patient's care?

A. Providing patient education
B. Training deep breathing and coughing exercises
C. Increasing the dose of antihypertensive drugs
D. Encouraging mobility and active body movement

ANSWER: B

5. A male patient was confused when the physician told him that his health condition was not good. Besides, the physician left the patient by telling him that he may benefit only from surgical intervention. The patient does not know about his problem, is still confused and full of fear, and does not know what procedure will be done. But he decided to undergo surgery.

What nursing intervention is most appropriate for this patient in this case?

A. Tell the patient not to fear for anything
B. Provide preoperative patient teaching
C. Facilitate family time for the patient
D. Encourage him to tell you his concerns

ANSWER: B

6. A nurse while preparing a patient for surgery identified a potential problem “Risk for hypothermia related to frequent exposure and the potential effect of induction agents”.
What nursing intervention is most appropriate for this patient’s problem?
- A. Draping the surgical site with heavy blanket
 - B. Administering warmed IV fluids
 - C. Boiling the materials to be used
 - D. Ignoring it as it is normal
- ANSWER: B
7. During the discussion before a surgery, the surgeon told the nurse that the surgery would be extensive, and a large amount of fluids and blood products should be ready. The surgeon also reminded the nurse not to forget to check the presence of Yankauer, Frazier, Poole, or Baron in the set.
What is the most important purpose of the surgeon’s action in this case?
- A. Viewing
 - B. Suctioning
 - C. Dilating
 - D. Probing
- ANSWER: B
8. A known diabetic patient is going to undergo hip replacement surgery. The patient also has controlled hypertension.
What finding of the outcome evaluation is most appropriate to proceed with this procedure?
- A. Blood pressure is slightly high
 - B. Blood glucose level in normal range
 - C. Slight decrease in respiratory effort
 - D. Depressed coughing
- ANSWER: B
9. A student nurse working in a highly contaminated and crowded surgical ward assumes that he has developed an infection while practicing at the ward. The staff nurse advised the student that he would be free from infection if he performed different precautions.
What is the most appropriate precaution that a student should perform in this case?
- A. Frequent hand washing with an antimicrobial
 - B. Using a waterless cleansing agent
 - C. Avoiding jewelry such as rings and bracelets
 - D. Wearing specially laundered or disposable clothing
- ANSWER: A
10. A nurse is working as a surgical scrub nurse for a patient undergoing surgery. During the procedure, the patient developed massive bleeding, and the nurse noted that there was too much blood in and around the operating room table and changed his gloves frequently during the procedure.
What is the most likely standard precaution that the nurse should follow?
- A. Management of puncture injuries
 - B. Prevention of puncture injuries
 - C. Use of protective barriers
 - D. No touching of mucous membranes
- ANSWER: C

11. A nursing student practicing at a surgical ward has frequent contact with patients due to the higher number of cases. He plans to practice hand hygiene more frequently, but he has dryness and skin irritation problem. The nursing instructor advised the student to perform techniques that cause less dryness and skin irritation.

What is the most likely recommended technique in this case?

- A. Alcohol-based hand rub
- B. Soap with water
- C. Ash with water
- D. Hydrogen peroxide

ANSWER: A

12. A fourth-year nursing student who is assigned to operating room (OR) told to remove mask or respirator after leaving the patient's room and close the door. The student asks the scrub nurse about the purpose of removing respirator after leaving the patient's room.

What is the most likely response of the scrub nurse in this case?

- A. It helps for proper disposal of the mask or respirator
- B. It helps in preventing mask or respirator from contamination
- C. It helps in keeping the privacy of the patient
- D. It helps in preventing contact with airborne disease

ANSWER: D

13. A nurse on duty at post-anesthetic care unit of a hospital received a postoperative patient and planned to start postoperative assessment.

What is the most appropriate initial step in the assessment of this case?

- A. Hydration and fluid status
- B. Mobility and activity level
- C. Airway and circulatory adequacy
- D. Urinary elimination status

ANSWER: C

14. A nurse assigned to a postoperative care unit planned to assess a patient on immediate arrival to the post-anaesthetic care unit. On evaluation, fast breathing, cyanosis of the lips, gums, and tongue, rapid, weak, and thready pulse with narrowing pulse pressure, low blood pressure, and concentrated urine were observed.

What is the most likely cause of this patient's problem?

- A. Positional change
- B. Effect of surgery
- C. Blood loss
- D. Hypoventilation

ANSWER: C

15. In postoperative assessment of a 23-year-old adult patient, a nurse noticed fast breathing, hypotension, hyperthermia, pus-like secretion from the wound, and an abrupt change in his mental status. What is the most likely nursing diagnosis of this patient?

- A. Surgical site infection
- B. Respiratory complications
- C. Cardiac complications
- D. Neurological complication

ANSWER: A

16. A 45-year-old patient who underwent thoracic surgery has been transferred to post post-anesthetic care unit. The patient has difficulty of hearing, and he has a history of being allergic to latex. What is the most appropriate immediate goal of postoperative care for this patient?

- A. Ensure correct closure of the incision
- B. Maintain ventilation and oxygenation
- C. Collect preoperative patient information
- D. Conduct a quick history taking and physical examination

ANSWER: B

17. A PACU nurse encourages a patient to take deep breaths, cough, and use the incentive spirometer every three hours.

What is the most likely reason for the nurse to encourage these actions?

- A. Because it is a routine postoperative action
- B. To drain surgical site secretions
- C. To clear pulmonary secretions
- D. To treat postoperative pneumonia

ANSWER: C

18. When a nurse assessed a patient who was admitted to the PACU eight hours back, the nurse noticed high body temperature and chills.

What is the most appropriate immediate action for this patient in this case?

- A. Notifying the primary care physician
- B. Administering antipyretics
- C. Applying cold compresses
- D. Administering antibiotics

ANSWER: A

19. Gas chemical sterilization is familiar in a hospital. A nurse arranged items to be sterilized and put them into the sterilizer. When the nurse checks it for adequacy, the colour of Ethylene oxide sensitive indicator has changed.

What does this change most likely indicate?

- A. Sterilization has occurred
- B. The item was exposed to ethylene oxide
- C. Wrong assembling of equipment
- D. Sterility defect in the wrapper

ANSWER: B

20. A nurse working in the minor OR of a hospital immersed all the used instruments in a Chlorine solution for 10 minutes. Then ordered a junior nurse to proceed with the processing, and the junior nurse started to clean them using a brush.

What is the most appropriate next step after the junior nurse finishes this task?

- A. Immerse them in another disinfectant
- B. Rinse them with pure water
- C. Boil them at a high temperature
- D. Load them into the autoclave

ANSWER: B

21. A nurse was discussing a discharge plan with a patient who was operated for benign prostatic hyperplasia. After the discussion, the nurse assessed some components to check his readiness for discharge.

What finding most likely changes the decision to discharge the patient?

- A. Orientation to place, events, and time
- B. Urine output less than 30mL/h
- C. Nausea and vomiting under control
- D. Minimal pain

ANSWER: B

22. A 33-year-old male patient presented to a clinic with a history of eye discomfort. His eye has been tearing for three hours, and he cannot open his eye due to the pain. On physical examination, the foreign body has stuck to the eye surface.

What is the most likely examination technique used in this case?

- A. Percussion
- B. Palpation
- C. Inspection
- D. Auscultation

ANSWER: C

23. A 30-year-old man complains of severe nasal bleeding an hour after admission to the hospital. The nurse observed that the patient had a subconjunctival haemorrhage and gingival bleeding.

What is the most likely cause of nasal bleeding in this case?

- A. Systemic cause
- B. Local cause
- C. Nasal sniffers
- D. Nasal dryness

ANSWER: A

24. A 62-year-old man presented with a history of blunt trauma, a blow from a fist. He has bruising around the orbit and maxillary area, and limitation of upward gaze.

What is the most likely priority nursing diagnosis in this case?

- A. Risk for bleeding
- B. Risk for corneal injury
- C. Risk for dry eye
- D. Risk for shock

ANSWER: B

25. A 40-year-old man presented with a history of epistaxis. He has an anterior nasal injury and severe haemorrhage.

What nursing diagnosis receives the highest priority in this case?

- A. Risk for infection related to injury
- B. Risk for shock related to bleeding
- C. Impaired gas exchange related to injury
- D. Risk for aspiration related to bleeding

ANSWER: B

26. A 52-year-old male patient presented with complaints of pain, mild fever, and ear drainage lasting two weeks. He has no history of vomiting and nausea.
What is the most likely priority plan of care for this patient?
- A. Prevent membrane perforation
 - B. Educate parents about risk factors
 - C. Prepare the patient for surgery
 - D. Prevent the causative agents
- ANSWER: A
27. A 52-year-old male patient presented with a history of tympanic membrane perforation on the left ear. He had Ciprofloxacin ear drops, 0.3 percent, 5ml, 3 drops twice daily for two weeks. However, he could not recover, and the initial topical antibiotic therapy failed.
What is the most appropriate next step in treatment in this case?
- A. Repeat initial antibiotic therapy
 - B. Culture-directed topical therapy
 - C. Culture-directed systemic therapy
 - D. Change the antibiotic therapy
- ANSWER: C
28. A 23-year-old male patient was admitted to the recovery room after he underwent nasal fracture repair under general anesthesia. On assessment, he has regained his consciousness and is oriented to time, place, and person.
What is the most appropriate prior nursing intervention for this patient?
- A. Reassure the patient
 - B. Deep breathing exercise
 - C. Reassess the level of consciousness
 - D. Schedule for discharge
- ANSWER: B
29. A 56-year-old female patient presented to a clinic with a complaint of inadvertent insertion of cereal in the ear two hours ago. On assessment, the nurse found out that she has history of ear discharge. Physical examination confirmed the presence of a foreign body in the middle ear.
What is the most appropriate action for the nurse in this case?
- A. Refer the patient
 - B. Irrigate the ear
 - C. Manual removal
 - D. Install olive oil
- ANSWER: A
30. A nurse is preparing a 36-year-old adult patient who is to undergo mastoidectomy. While counseling the patient, the nurse stressed the importance of visiting the health center when he feels sick and keeping follow-up appointments to avoid complications.
What is the most likely outcome that indicates good understanding of the patient in this case?
- A. Remains free of discomfort or pain
 - B. Demonstrates no signs or symptoms of infection
 - C. Remains free of injury and trauma
 - D. Exhibit the importance of scheduling
- ANSWER: D

31. A 45-year-old male patient with a diagnosis of obstructed bronchus is admitted to the surgical ward due to acute exacerbation of wheezing, cough, and decreased oxygen saturation. The nurse suspected the diagnosis of chronic obstructive pulmonary disease (COPD) and wanted to confirm it by percussion of the patient's chest.

What percussion note would support the nurse's diagnosis?

- A. Dull
- B. Resonant
- C. Hyper-resonant
- D. Flat

ANSWER: C

32. A 55-year-old patient came to the hospital with a history of an untreated chronic productive cough and moderate amounts of foul-smelling sputum. On assessment, the patient has an accumulation of thick and purulent fluid within the pleural space.

What is the most likely cause of the patient's problem in this case?

- A. Hematogenous infection
- B. Bacterial infection
- C. Iatrogenic causes
- D. Lung abscess

ANSWER: D

33. A 19-year-old female patient with a history of heart disease and lung abscess came to the hospital with complaints of chest pain and cough. On physical assessment, the patient has altered chest excursion and bluish discoloration of the lip and oral mucosa. She has also tachycardia, tachypnea, crepitations, and pleural effusion; but, distended neck veins, raised JVP, hepatomegaly, ascites, and leg edema were not found. The attending nurse developed a plan of care to improve the symptoms and signs through the administration of furosemide and oxygen.

What is the most likely nursing diagnosis in this case?

- A. Risk for disturbed sleep pattern
- B. Ineffective breathing pattern
- C. Impaired skin integrity
- D. Impaired physical mobility

ANSWER: B

34. A 70-year-old patient with a history of blunt chest trauma came to the surgical OPD for a regular follow-up. On assessment, the patient has difficulty elevating the upper body, decreased gastrointestinal motility, difficulty swallowing, and difficulty with sputum. The attending nurse wants to formulate a nursing diagnosis to develop a nursing plan for the care of this patient.

What is the most likely nursing diagnosis of this patient?

- A. Risk for aspiration
- B. Risk for fall
- C. Risk for electrolyte imbalance
- D. Risk for fluid volume deficit

ANSWER: A

35. A 30-year-old female patient with a previous history of nasal polyps came to a surgical OPD with complaints of severe headache and sore throat. Nursing assessment revealed that the patient has tenderness around the eyes and on both sides of the nose. The nasal mucosa has also increased redness, swelling, and exudates.

What is the most likely nursing diagnosis for this patient?

- A. Ineffective breathing pattern
- B. Impaired verbal communication
- C. Hyperthermia
- D. Acute pain

ANSWER: D

36. A 35-year-old patient who has undergone surgery for lung abscess came to the surgical OPD before his appointment date. On assessment, the patient has difficulty of elevating the upper body, abdominal discomfort, and an ineffective cough.

What is the priority care plan for this patient?

- A. Identify the causative factors
- B. Assist in self-care
- C. Promote wellness
- D. Provide ventilatory support

ANSWER: A

37. A 39-year-old male patient who underwent surgery for a lung mass was admitted to a surgical ward for full recovery. On assessment, his vital signs are within the normal range, and he has balanced input and output, improved airway, and breathing.

What is the most appropriate nursing intervention for this patient?

- A. Prepare the patient for discharge
- B. Encourage a diet that is high in protein and calories
- C. Give IV antimicrobial therapy for three weeks
- D. Plan for health promotion and wellness

ANSWER: A

38. A 60-year-old male patient came to the hospital on his appointment date to undergo surgery on the anterior part of the left lung for the management of a lung tumour. The surgery is expected to take eight hours, and the patient was on chemotherapy to shrink the tumor size. The nurse suspects a complication related to the procedure due to the previous chemotherapy and plans to assess for the occurrence of complications in pre-operatively and immediately after the surgery.

What is the most likely assessment the nurse should do in this case?

- A. Signs of infection
- B. Signs of fluid volume deficit
- C. Skin integrity
- D. Patient's pain level

ANSWER: C

39. A 44-year-old patient who underwent emergency tracheostomy for acute airway obstruction at the minor ward came to the recovery room for recovery. Upon arrival, the patient looks stable, the tube is in correct placement, and there is no sign of respiratory distress.

What is the most appropriate nursing action in this case?

- A. Perform a respiratory assessment at least hourly
- B. Chest x-ray to determine proper placement of the tube
- C. Provide post-operative prophylaxis for 48 hours
- D. Give health education about stoma care

ANSWER: A

40. A 35-year-old male patient diagnosed with atelectasis was referred to a surgical OPD to undergo postural drainage. On assessment, he looks fatigued, complains of pain around the chest, and has difficulty of sleeping in head head-down position.

What is the most appropriate nursing intervention for this patient?

- A. Put the patient in a head-down position
- B. Position the patient on side side-lying
- C. Provide anti-pain medication and put on a head-down position
- D. Ask the patient to speak out during the procedure

ANSWER: B

41. A 54-year-old patient came to the hospital with complaints of thick sputum production and dyspnea. On assessment, the patient is irritable and has decreased breathing. A nurse has planned to perform chest physiotherapy.

What is the most appropriate first step of nursing intervention in this case?

- A. Administering anti-pain drugs
- B. Positioning the patient
- C. Conducting physical examination
- D. Demonstrating the procedure

ANSWER: B

42. A 44-year-old male patient presented to a clinic with complaints of pleuritic-type chest pain, fatigue, tachypnea, cough, and purulent sputum. After completion of the assessment, the nurse educated the patient about taking medications as prescribed, performing breathing exercises, ensuring adequate fluid and dietary intake, and avoiding smoking, alcohol, and excessive activity.

What is the most likely expected outcome in this case?

- A. Maintained weight
- B. Remaining oriented and being aware of surroundings
- C. Exhibited no complications
- D. Adhered to the therapeutic regimen

ANSWER: D

43. A 32-year-old male patient with a history of Peritonsillar abscess was admitted to the surgical ward. After three days of surgical intervention, like debridement and daily wound care, the nurse planned to reassess the patient. During evaluation, his vital signs are within the normal range, and he has no fever, hypotension, or deterioration of cognitive status.
What is the most likely nursing outcome gained in this case?
- A. Relief of pain and discomfort
 - B. Absence of complications
 - C. Utilizes strategies to prevent infections
 - D. Adequate level of knowledge
- ANSWER: B
44. A 24-year-old male patient came to OPD with complaints of dyspnea, restlessness, and orthopnea of five days duration. After assessment, a physician made a diagnosis of left-sided heart failure.
What is the most likely symptom that confirms the diagnosis in this case?
- A. Blood-streaked or bloody sputum
 - B. Thick mucoid sputum
 - C. Pink frothy sputum
 - D. Copious purulent sputum
- ANSWER: C
45. A 34-year-old diabetic male patient came to a clinic with a diagnosis of deep vein thrombosis (DVT). The patient asked the nurse to tell him the main cause of his problem.
What is the most appropriate answer for the patient's query?
- A. Decreased release of clotting factor
 - B. Decreased activation of platelets
 - C. Decreased fibrin production
 - D. Decreased extremity muscle activity
- ANSWER: D
46. A 20-year-old female patient came to an emergency OPD with complaints of severe watery diarrhea and vomiting for the last two days. On examination, her skin turgor goes back very slowly, and she has sunken eyeballs.
What is the most likely nursing diagnosis in this case?
- A. Excessive fluid volume
 - B. Risk for fluid volume deficit
 - C. Fluid volume deficit
 - D. Risk for fluid volume imbalance
- ANSWER: C
47. A 32-year-old male patient is admitted to a medical ward with a diagnosis of heart failure. On nursing assessment, the attending nurse noted jugular vein distention, enlarged liver and spleen, dependent edema, distended abdomen, and swollen hands and fingers.
What is the most likely nursing diagnosis of the patient?
- A. Risk for fluid volume deficit
 - B. Risk for constipation
 - C. Risk for impaired body Image
 - D. Risk for impaired mobility
- ANSWER: D

48. A 16-year-old male patient presented with sudden fainting, sunken eyeballs and lethargy. On assessment, the patient's weight had decreased by 6 percent of his usual weight, and capillary refill took more than 3 seconds. The nurse intended to include the type of fluid to be given in nursing care plan.

What is the most appropriate fluid that should be administered to the patient?

- A. 0.9 percent normal saline
- B. 5 percent dextrose in 0.45 percent normal saline
- C. 5 percent dextrose in Ringer's lactate
- D. 10 percent dextrose in water (D₁₀W)

ANSWER: A

49. A 30-year-old patient presented to the OPD with chest discomfort for the last three days. On assessment, the patient has tachycardia, hypotension with systolic blood pressure less than 90mmHg (30mmHg less than the patient's baseline), decreased urine output, cold and clammy skin with poor peripheral pulses, agitation, restlessness, and confusion. The physician diagnosed the case as cardiogenic shock.

What is the most appropriate nursing care plan in this case?

- A. To administer angiotensin-converting enzyme inhibitors
- B. To administer once-a-day beta blockers
- C. To administer normal saline
- D. To administer morphine and oxygen

ANSWER: D

50. A 54-year-old male patient presented with sharp pain on the anterior chest, nausea, and vomiting. The patient rated the intensity of the pain as 10 on a 0-to-10 pain intensity scale. On physical examination, increased blood pressure, absent peripheral pulse and decreased level of consciousness were noted. What is the priority nursing intervention to prevent emergency harm in this case?

- A. Check for airway patency
- B. Administer drugs to decrease blood pressure
- C. Monitoring input and output
- D. Administer antibiotics to prevent infection

ANSWER: B

51. A patient with acute decompensated heart failure is receiving IV Dobutamine. After administration of the medication, the nurse wanted to monitor the patient's response to the drug.

What is the most likely expected effect of the drug in this case?

- A. Raised heart rate
- B. Dilated renal blood vessels
- C. Increased heart contractility
- D. Increased systemic vascular resistance

ANSWER: C

52. A 40-year-old male patient came to an emergency OPD with complaints of excessive sweating, hunger, tremor, and some episodes of seizure. On examination, his temperature is 39 degrees centigrade, pulse rate is 110/min, respiratory rate is 26/min, blood pressure is 130/98mmHg, weight is 60kg, and there are dilated pupils. The blood glucose level of the patient was 30mg/dl. The nurse intended to give 10 percent dextrose in water.

What is the glucose infusion rate that the nurse should include in the nursing care plan?

- A. 0.1mg/kg/min
- B. 0.2mg/kg/min
- C. 0.4mg/kg/min
- D. 0.6mg/kg/min

ANSWER: A

53. A 23-year-old female patient developed brain swelling or hyponatremic encephalopathy following excessive volume fluid therapy. The health care providers are very concerned that the condition is caused by the infused fluid.

What is the most likely responsible fluid in this case?

- A. 0.9 percent NaCl (normal saline)
- B. Lactated Ringer's solution
- C. 0.45 percent NaCl (normal saline)
- D. 5 percent dextrose in water (D5W)

ANSWER: D

54. A 44-year-old male patient was subjected to excessive blood loss during the operation. On the immediate post-operative period, a nurse is planning to evaluate the patient's fluid status after rehydration therapy has been instituted.

What is the most likely appropriate clinical indicator that the nurse can use?

- A. Blood pressure and respiratory rate
- B. Skin turgor and capillary refill
- C. Pulse rate and urine output
- D. Heart rate and cardiac output

ANSWER: C

55. A 20-year-old female patient presented to an emergency room with abdominal pain of one day duration. She described the pain as it was generalized and, currently become localized to the right lower quadrant. During physical examination, the patient experienced pain in the left lower quadrant on palpation of the right lower quadrant.

What is the most appropriate finding for this examination?

- A. Murphy sign
- B. Obturator sign
- C. Rovsing sign
- D. Psoas sign

ANSWER: C

56. A 40-year-old male patient presented to a surgical OPD with complaints of nausea and vomiting. He also experiences intermittent epigastric pain and a feeling of fullness after eating. A physician suspected a diagnosis of sliding hiatal hernia.

What is the most likely symptom that confirms the diagnosis in this case?

- A. Jaundice
- B. Heartburn
- C. Supra-pubic pain
- D. Stomatitis

ANSWER: B

57. A 38-year-old woman was admitted to a hospital for severe diarrhea and abdominal pain. Laboratory results revealed decreased hematocrit, hemoglobin, albumin, and protein levels; however, she has an elevated WBC count and ESR. The nursing care provider suspects that she has Crohn's disease.

What additional sign may confirm the diagnosis?

- A. Diminished bowel sounds
- B. Abdominal pain of 2 on a 0-10 pain intensity scale
- C. Low potassium level
- D. Hypervolemia

ANSWER: C

58. A 62-year-old male patient presented to a surgical ward with a diagnosis of peritonitis. The nurse started collecting the patient's data and performing a physical examination of the abdomen.

What is the most classic symptom that the nurse expects during nursing assessment?

- A. Rigid abdomen
- B. Increase passage of flatus
- C. Hyperactive bowel sound
- D. Increased urine output

ANSWER: A

59. A 35-year-old male patient was brought to the emergency department with severe abdominal pain. After an extensive assessment, he was diagnosed with a primary peritonitis. The patient asked the nurse to tell him the cause of his disease.

What is the most appropriate response of the nurse?

- A. It occurs due to a spontaneous bacterial infection
- B. It occurs after perforation of abdominal organs
- C. It occurs as a result of penetrating injury
- D. It occurs due to autoimmune causes

ANSWER: A

60. A 25-year-old male patient presented with a complaint of difficulty of swallowing for the last three months. He reported that initially, he felt a sensation of food sticking in the lower portion of his esophagus. He gets relieved after he intentionally vomits the food to avoid discomfort produced by prolonged distention of the esophagus by food that will not pass into the stomach. The patient also reported heartburn that was independent of the amount and time of eating.

What is the most likely medical diagnosis of this patient?

- A. Gastroesophageal reflux disease
- B. Achalasia
- C. Esophageal cancer
- D. Chronic Gastritis

ANSWER: B

61. A 55-year-old man came to a primary health care unit with a complaint of bright red bleeding on defecation. On assessment, the patient reported that it was associated with severe pain, which was relieved by avoiding excessive straining while defecating. Anorectal examination revealed prolapse outside the anal canal which can be reduced spontaneously.

What is the most likely medical diagnosis of this case?

- A. First-degree Hemorrhoid
- B. Second-degree Hemorrhoid
- C. Third-degree Hemorrhoid
- D. Fourth degree Hemorrhoid

ANSWER: B

62. A nurse is caring for a 50-year-old male patient who presented with a several-month history of heartburn after meals. He reports that these episodes last about an hour after eating, and are much worse if he lies down for sleep after a meal, and relieved if he takes fluids or antacids and maintains an upright position. Most of the time, he also experiences pain with difficulty of swallowing, abdominal discomfort, nausea, flatulence, belching, and bloating.

What is the most likely medical diagnosis of this patient?

- A. Gastroesophageal reflux disease
- B. Esophageal cancer
- C. Hiatal hernia
- D. Peptic ulcer disease

ANSWER: A

63. A 44-year-old patient presented with abdominal pain, low-grade fever, and diarrhoea that occurs five to six times per day. On assessment, abdominal tenderness and pallor were noted. Endoscopic examination revealed a cobblestone appearance on the colon.

What is the most likely medical diagnosis of this patient?

- A. Crohn's disease
- B. Ulcerative colitis
- C. Colorectal Cancer
- D. Diverticulitis

ANSWER: A

64. A nurse is taking care of a 65-year-old woman who was diagnosed with peptic ulcer disease after she presented with fatigue, severe epigastric pain associated with meals, and coffee ground emesis. She was treated with omeprazole and sucralfate; however, she did not respond to this therapy.

What is the most likely priority nursing diagnosis in this case?

- A. Risk for self-care deficiency
- B. Risk for infection
- C. Risk for imbalanced energy
- D. Risk for bleeding

ANSWER: D

65. A 40-year-old female patient with known peptic ulcer disease was admitted to a surgical ward after she presented with complaints of epigastric pain that awakens her at midnight and a feeling of abdominal fullness without a meal.

What is the most likely priority nursing problem in this case?

- A. Disturbed sleep pattern
- B. Altered thought process
- C. Fluid volume deficit
- D. Impaired mobility

ANSWER: A

66. A nurse is developing an instruction session for a 65-year-old patient with constipation. On assessment, he has not received any education regarding his health problems previously. What is the most likely instruction that the nurse should include in this case?

- A. Using a diet that is low in bulk
- B. Decreasing fluid intake
- C. Administering an enema once a day
- D. Allowing sufficient time for evacuation

ANSWER: D

67. A 39-year-old man who presented with complaints of fatigue, epigastric pain, and coffee ground emesis was admitted to a surgical ward with a medical diagnosis of peptic ulcer disease. After extensive nursing assessment, a diagnostic label of 'acute pain' and a related factor of 'excessive gastric acid secretion' was identified. Based on these data, a nurse wanted to establish a nursing intervention. What is the priority nursing intervention in this case?

- A. Administer anti-pain like aspirin for the acute pain
- B. Avoid oral medication, as the patient has to be scheduled for surgery
- C. Administer an antacid to relieve pain secondary to an ulcer
- D. Administer non-steroidal anti-inflammatory drugs for the pain

ANSWER: C

68. A 40-year-old woman was admitted to a surgical ward after she presented with bright red bloody vomiting last night. She reported that she has a history of gastritis and takes medication for it without a prescription. However, she could not provide the name and therapeutic benefits of the medication she had taken.

What is the most appropriate priority nursing intervention in this case?

- A. Filling the knowledge gap
- B. Prescribing medications
- C. Reviewing patient chart
- D. Consulting senior nurse staff

ANSWER: A

69. A nurse is taking care of a 39-year-old patient who has undergone gastrectomy three days ago. While developing a nursing care plan, the nurse wanted to include nursing interventions for the prevention of dumping syndrome.

What is the most likely appropriate nursing intervention in this case?

- A. Encourage the patient to eat high high-carbohydrate diet
- B. Instruct the patient to reduce fluid intake with meals
- C. Assist the patient to assume high flower position
- D. Assist the patient to ambulate following the meal

ANSWER: B

70. A nurse is preparing a nursing care plan to provide nursing care for a 50-year-old male patient who was admitted to a surgical ward with upper gastrointestinal bleeding. What is the most likely priority nursing intervention for this patient?
- A. Inserting an IV line for fluid resuscitation
 - B. Identifying the underlying cause and source of bleeding
 - C. Preparing the patient for surgery
 - D. Monitoring hemoglobin and hematocrit level
- ANSWER: A
71. A cleansing enema is ordered for a 55-year-old patient who is admitted to a surgical ward for the management of fecal impaction. The nurse took consent and instructs the patient to have an appropriate position during the procedure. What is the most appropriate position in this case?
- A. Left Sims' position
 - B. Right Sims' position
 - C. Left lateral with raised head of bed
 - D. Right lateral with raised head of bed
- ANSWER: A
72. A 55-year-old male farmer is admitted to a surgical ward for surgical intervention of large bowel pathology. A nurse is administering a large-volume cleansing enema to prepare a patient for intestinal surgery. Once the enema solution is introduced, the patient complains of severe abdominal cramping. What is the most appropriate prior nursing intervention in this case?
- A. Elevate the head of the bed and reposition the rectal tube
 - B. Elevate the head of the bed and modify the amount of solution
 - C. Lower the solution container and check the flow rate
 - D. Remove the rectal tube and reinsert the tube after the symptom disappears
- ANSWER: C
73. A 67-year-old man was admitted to a surgical ward with a diagnosis of diverticulitis. A nasogastric tube (NGT) insertion is ordered in response to associated severe vomiting and abdominal distention. During insertion of the NGT, the nurse wanted to confirm whether the NGT is in the stomach or not by measuring the pH value of the gastric content. What is the most likely indicator that the NGT is in the correct position?
- A. pH value below 3
 - B. pH range of above 8.5
 - C. pH range of 7.0
 - D. PH value above 7.9
- ANSWER: A
74. A 64-year-old male patient who is admitted to a surgical ward complains of acute abdominal discomfort. He has a history of chronic constipation and disturbance in the usual defecation patterns and habits. He has no hemorrhoids. What is the most appropriate nursing intervention to be performed for this patient?
- A. Advise the patient to do simple exercise
 - B. Position the client in semi-sitting to provide gravity
 - C. Fleet mineral oil retention enema
 - D. Advice to increase fluid intake
- ANSWER: C

75. A nurse giving care for a 34-year-old male patient, who is on continuous gavage, found that the gastric residual volume in the morning after eight hours is 210ml.

What is the most appropriate nursing action for this patient?

- A. Stop feeding to reduce the risk of regurgitation
- B. Instruct the patient to begin oral intake of clear liquids
- C. Add solid foods and remove the tube
- D. Ensure that the feeding tube is in the stomach

ANSWER: A

76. A 34-year-old male patient came to the emergency department with a complaint of severe lower abdominal pain of three hours duration. He reported that the pain originates in the renal area, radiating anteriorly and downward toward the testes. His temperature and blood pressure were 39.3 degrees centigrade and 90/50mmHg, respectively. Blood chemistry and a 24-hour urine test for measurement of calcium showed that there was an elevation of calcium and uric acid levels.

What is the most important physical examination to be done in this scenario?

- A. Palpation
- B. Percussion
- C. Auscultation
- D. Inspection

ANSWER: A

77. An 84-year-old male patient came to an emergency department with a complaint of urgency and difficulty of urination. The nurse immediately inserted a Foley catheter and eliminated the urine from the distended bladder. After seven days, the patient complained of burning on urination, urinary frequency, and hematuria.

What is the most likely causative microorganism of the most likely diagnosis?

- A. Escherichia coli
- B. Klebsiella
- C. Pseudomonas
- D. Staphylococcus

ANSWER: A

78. A 42-year-old male patient who underwent open ureterolithotomy presented with bleeding from the incision site. A nurse assessed and noticed that the urine output was 300ml/24 hours and the blood pressure was 80/50mmHg.

What is the most appropriate nursing diagnosis in this scenario?

- A. Deficient fluid volume related to active fluid loss manifested by low blood pressure
- B. Dehydration related to active bleeding as manifested by decreased blood pressure
- C. Deficient Fluid Volume and bleeding related to insufficient fluid intake
- D. Risk for deficient fluid volume and infection-related insufficient fluid

ANSWER: A

79. A 73-year-old male patient came to an emergency department with complaints of urinary retention and urgency of eight hours duration. Finally, the physician ordered the emergency nurse to catheterize the patient and schedule him for surgery after two weeks.

What is the most appropriate nursing diagnosis in this scenario?

- A. Impaired urinary elimination related to disruption of bladder innervation
- B. Risk for impaired urinary elimination and incontinence related to mechanical obstruction
- C. Infection related to the disease process as manifested by patient response
- D. Acute pain related to irritation of the bladder

ANSWER: A

80. A 37-year-old male patient presented with severe pain and swelling of the scrotum. He reported recent scrotal trauma in a football game. In physical examination, the right testicle is tender and positioned upper than the left testicle. Doppler ultrasound revealed that there is no blood flow to the injured testicle.

What is the most likely medical diagnosis of this case?

- A. Epididymitis
- B. Hydrocele
- C. Testicular cancer
- D. Testicular torsion

ANSWER: D

81. A 27-year-old male patient who underwent surgery for testicular cancer is complaining of severe pain and anxiety due to the removal of his testicles. On assessment, his vital signs were within the normal range.

What is the priority nursing diagnosis in this scenario?

- A. Acute pain related to the presence of a surgical incision
- B. Anxiety related to situational crisis
- C. Disturbed body image related to culture
- D. Hopelessness related to stress

ANSWER: A

82. A surgical ward nurse is providing postoperative care for a patient with a urethral stricture. The nurse does each activity based on Maslow's hierarchy of basic human needs.

What is the most likely reason for the nurse to follow this approach?

- A. To make accurate nursing diagnoses
- B. To establish priorities of care
- C. To communicate concerns more concisely
- D. To integrate science into nursing care

ANSWER: B

83. A 28-year-old female patient came to an emergency room with a history of severe lower abdominal pain and hematuria of three hours duration. Blood chemistry and a 24-hour urine test for measurement of calcium showed an elevation of calcium level.

What is the appropriate initial step in the management of this patient?

- A. Giving analgesic agents
- B. Limit sodium intake to 3 to 4g/day
- C. Avoid intake of oxalate-containing foods
- D. Drink two glasses of water at bedtime

ANSWER: A

84. A 65-year-old man came to a surgical OPD, and a diagnosis of BPH was made. The physician planned to admit the patient and do a prostatectomy.

What is the most appropriate initial step the nurse should do before the procedure?

- A. Administer the prescribed medication
- B. Inform the patient about the effects of prostate surgery
- C. Measure urine input and output
- D. Perform a nursing care plan for the patient

ANSWER: B

85. A 65-year-old male patient came to an emergency room with complaints of frequent of urination, urgency, and nocturia. On physical examination, the bladder is distended. The duty nurse decided to catheterize the patient.

What is the most appropriate initial step in this patient's care?

- A. Explain the procedure
- B. Prepare the equipment
- C. Wash hands with soap
- D. Wear sterile gloves

ANSWER: A

86. A 38-year-old female patient came to an emergency department with an electrical burn of two hours duration. The emergency nurse catheterized the patient.

What is the most appropriate nursing measure to prevent the catheter from slipping out?

- A. Inflate the balloon of the catheter
- B. Insert the catheter 3-5 inches after withdrawal of urine
- C. Instruct the patient to hold the catheter by her hand
- D. Properly secure the catheter to the thigh

ANSWER: A

87. A 34-year-old male patient presented with severe lower abdominal pain, nausea, and vomiting of three hours duration. On assessment, his body temperature and blood pressure were 39.5 degrees centigrade and 90/50mmHg, respectively. The nurse gave the patient Paracetamol 500mg PO BID.

What is the most likely expected outcome?

- A. Decrease in vomiting after a few hours
- B. Rise in blood pressure to 100/60mmHg after four hours
- C. Pain relief after few hours
- D. Reduction of body temperature to normal

ANSWER: D

88. A nurse providing postoperative care for a testicular cancer patient. On assessment, the nurse found that the patient has chronic weight loss. Then, the nurse diagnosed the condition of the patient as “imbalanced nutrition less than the body requirement”.

What is the most likely expected outcome after two weeks of nursing intervention?

- A. Improved nutritional status
- B. Improved appetite
- C. Improved symptoms
- D. Gain 2.5kg weight

ANSWER: D

89. An adult patient is admitted to a surgical ward with complaints of fatigue, lethargy, cold intolerance, and weight gain. After a complete physical examination and laboratory investigations were done, a diagnosis of hypothyroidism was made. The physician kept him on thyroxin tablet. The nurse working in the surgical ward also conducted a nursing assessment.

What is the most likely sign detected on physical examination in this case?

- A. Buffalo hump
- B. Hoarseness of voice
- C. Exophthalmia
- D. Tremors

ANSWER: B

90. A 20-year-old male patient, with a history of palpitation, fatigue, and headache, was admitted to a surgical ward for surgery. On physical examination, the nurse identified that he has high blood pressure and tachycardia. After laboratory investigations and radiologic examination, the patient was diagnosed with pheochromocytoma.

What is the most likely gland tumor that caused this condition?

- A. Thyroid gland
- B. Adrenal medulla gland
- C. Parathyroid gland
- D. Pancreatic gland

ANSWER: B

91. A patient with Type-2 DM who was on oral hypoglycemic medications came to a follow-up clinic after two years of discontinuing medication. On evaluation, severe hyperglycemia, neuropathy, and a foot ulcer were noted. The physician asked him the reason for discontinuing his follow-up. The patient stated that he was totally healed after taking the hypoglycemic medication, except that he had developed the above-mentioned problems currently.

What is the most likely nursing diagnosis of this patient?

- A. Knowledge deficit related to the disease condition as evidenced by discontinuation of medical follow-up
- B. Acute pain related to the disease condition, as evidenced by the actual patient's level of pain measurement scale
- C. Ineffective airway clearance related to disease condition as evidenced by increased respiratory rate
- D. Risk for nutritional imbalance related to disease condition

ANSWER: A

92. A 35-year-old patient with a tumor of adrenal medulla experienced headache, palpitations, sweating, and severe hypertension. Laboratory investigation revealed a high level of catecholamine.

What is the most appropriate medical diagnosis for this patient?

- A. Pheochromocytoma
- B. Thyrotoxicosis
- C. Cushing Syndrome
- D. Addison's disease

ANSWER: A

93. A 25-year-old female patient with a history of thyroidectomy came to an outpatient department with complaints of fatigue and lethargy, weight gain, cold intolerance, dry skin, and deepening of the voice. On examination, brittle hair, coarse voice, peri-orbital edema and pulse rate of 60/min were noted. Laboratory investigation indicated low T4 and high TSH level.

What is the most likely medical diagnosis of this patient?

- A. Hypothyroidism
- B. Thyrotoxic storm
- C. Cushing syndrome
- D. Creatinism

ANSWER: A

94. A 55-year-old male patient came to an outpatient department with complaints of frequent vomiting and severe left upper quadrant pain. His general condition was sick-looking, and he was shouting by stating the severity of the pain. He has dry mucus membrane, sunken eye, and blood pressure of 80/50mmHg. Serum amylase and lipase levels are high.

What is the priority nursing diagnosis of this patient?

- A. Risk for infection related to severe pain
- B. Pain related to disease condition as evidenced by patient verbalization and shouting
- C. Activity intolerance related to disease condition as evidenced by fatigue
- D. Disturbed sleep pattern related to disease condition

ANSWER: B

95. A 40-year-old male patient, who is diagnosed with acute pancreatitis, is admitted to a surgical ward with severe left upper quadrant pain. On evaluation, he is looking sick and is shouting by stating the severity of the pain. His blood pressure was 90/60mmHg, pulse rate was 100/min, and respiratory rate was 16/min.

What is the most priority nursing care plan needed for this patient?

- A. To improve respiratory status
- B. To relieve pain
- C. To maintain normal breathing
- D. To inform the patient about adherence

ANSWER: B

96. A patient with acute pancreatitis is admitted to emergency department with severe left upper quadrant abdominal pain and nausea. General condition of the patient was sick-looking and he was complaining of severe of the pain. His blood pressure, pulse rate and respiratory rate were 90/60mmHg, 100/min, and 16/min, respectively, with a pain score of 7/10.

What is the priority nursing intervention for this patient?

- A. Preventing re-infection or spread of infection
- B. Relieving the pain
- C. Maintaining normal body temperature
- D. Improving nutritional status

ANSWER: B

97. A male patient who is diagnosed with pheochromocytoma is admitted to a surgical ward for surgery. He had a history of palpitation, fatigue, and headache. On physical examination, he has high blood pressure and tachycardia.

What is the most priority nursing intervention likely for this case?

- A. Maintaining adequate cardiac function
- B. Decreasing risk of infection
- C. Maintaining normal body temperature
- D. Maintaining nutritional status

ANSWER: A

98. A female patient is admitted to the recovery room after adrenalectomy was done and she became severely sick. Her blood pressure was 60/20mmHg, pulse rate was 120/min, respiratory rate was 24/min, and she had a weak pulse and extreme weakness. An additional intravenous line was secured, and normal saline was given fast and the patient was kept on 100 percent oxygen. Thirty minutes later, the patient showed improvement with blood pressure of 100/60mmHg, pulse rate of 85/min, and respiratory rate of 16/min.

What is the most likely expected outcome of nursing intervention for this patient?

- A. Maintain normal body temperature
- B. Experiences increased comfort
- C. Experiences relief from anxiety
- D. Maintain adequate cardiac output

ANSWER: D

99. A 45-year-old patient with a history of cholecystectomy came to the surgical department with a complaint of cold intolerance.

What is the most likely skin layer affected?

- A. Epidermis
- B. Dermis
- C. Subcutaneous tissues
- D. Sebaceous gland

ANSWER: C

100. A 30-year-old patient came to a dermatology clinic with complaints of itching and a wound on the trunk. On examination, there were red raised patches on the skin over the lower back and buttocks with plaque-type lesions. A diagnosis of psoriasis was made, and 10 percent of the body surface area was involved.

What is the most likely classification of Psoriasis in this case?

- A. Mild
- B. Severe
- C. Extensive
- D. Moderate

ANSWER: B

101. A 45-year-old woman came to a clinic with a complaint of small masses in the genital and perineal area. She had a history of multiple sexual partners. After physical examination and laboratory investigations, a physician made a diagnosis of warts and treated her with cryosurgery.

What is the most likely cause of the wart in this case?

- A. Trichomonas vaginalis
- B. Human papilloma virus
- C. Exposure to ultraviolet
- D. Genetic factors

ANSWER: B

102. A young female patient came to an emergency OPD with a second-degree burn. Upon evaluation, her head, neck, and chest was injured. She was restless and had difficulty in breathing. On physical examination, she has poor skin turgor, sunken eye, and oxygen saturation of 70 percent.

What is the most likely nursing diagnosis of this patient?

- A. Ineffective breathing related to disease condition as evidenced by low O₂ saturation
- B. Knowledge deficit related to injury as evidenced by treatment refusal
- C. Risk for infection related to the disease condition
- D. Fluid volume deficit related to injury as evidenced by low blood pressure

ANSWER: A

103. A 20-year-old female patient came to a hospital with complaints of an itching sensation and redness on her lower legs. A dermatologist diagnosed her with eczema and ordered medications for it. A duty nurse also taught her how to adhere to the ordered medications. However, the patient is worried about how to attend her college education with this situation.

What is the most likely nursing diagnosis in this case?

- A. Risk for infection related to the disease condition
- B. Risk for impaired nutritional intake related to nausea and vomiting
- C. Poor self-esteem related to disease condition as evidenced by verbalization
- D. Knowledge deficit related to disease condition, as evidenced by looking for assistance

ANSWER: C

104. A 10-year-old child was brought to a hospital with complaints of fever, poor appetite, itching, and pain on his hand and fingers, which worsens in the evening. On examination, there was a lesion on his hands and fingers, his body temperature was 39 degrees centigrade, and his pulse rate was 130/min. Finally, he was diagnosed with scabies.

What is the most priority nursing diagnosis for this patient?

- A. Impaired skin integrity related to disease condition as evidenced by lesion
- B. Acute pain related to itching as evidenced by pain on the hands and fingers
- C. Risk for infection related to itching and lesion
- D. Hyperthermia related to disease condition as evidenced by high temperature

ANSWER: D

105. A 17-year-old female patient presented to a hospital emergency department with flame burn injury while she was cooking. She arrived at the hospital emergency department after two hours of the injury. The burn affected the right anterior forearm and chest. On examination, she is conscious and oriented, her pulse rate is 110/min, respiratory rate is 28/min, temperature is 36.7 degrees centigrade, and blood pressure is 90/70mmHg.

What is the most appropriate immediate nursing care plan for this case?

- A. Prevent infection
- B. Administer IV fluid
- C. Avoid contracture
- D. Provide wound care and debridement

ANSWER: B

106. A 47-year-old woman who is working in the movie industry undergone cosmetics surgery to avoid wrinkles on her face and restore her youthful appearance. She has no history of previous hospitalization. Her vital signs are within the normal range.
What is the most appropriate nursing intervention for this patient?
- A. Decide the discharging arrangement
 - B. Maintain adequate nutrition
 - C. Undergo debridement
 - D. Perform radiologic test
- ANSWER: B
107. A patient was admitted to a surgical ward with a diagnosis of appendicitis. During the surgical procedure, a surgeon found a collection of pus. Initially, the skin was left unclosed, and a surgical nurse was assigned to sew it later by appointment.
What is the most likely wound healing approach applied in this case?
- A. Primary intention
 - B. Secondary intention
 - C. Tertiary intention
 - D. Fourth intention
- ANSWER: C
108. A patient came to a hospital with complaints of swelling and discharge on the right lower leg for one one-week duration. Initially, the patient tried to manage the wound at home. On examination, the wound has purulent discharge, foreign debris, and devitalized tissues.
What is the most appropriate wound cleanser to be used in this case?
- A. Wet gauze
 - B. Hydrogels
 - C. Saline
 - D. Iodine solution
- ANSWER: C
109. A 25-year-old male patient came to an emergency department with a complaint of a sharp injury to his forearm while he was arranging the materials in a shop. The injured part of the skin was stitched and he was discharged with an appointment. On his appointment day, a nurse evaluated the effectiveness of the wound healing process.
What are the most likely findings expected from the evaluation in this case?
- A. Absence of drainage and foul around the wound
 - B. Decrease redness and swelling around the wound
 - C. Presence of warmth and pain around the wound
 - D. Resolution of bruising around the wound
- ANSWER: C
110. A 63-year-old woman was admitted to a hospital with a diagnosis of femoral fracture, and currently she is complaining of pain in the affected site. Skin traction was done to keep the alignment. She had a history of Gastritis.
What is the most appropriate anti-pain medication for this patient?
- A. Diclofenac
 - B. Ibuprofen
 - C. Celecoxib
 - D. Naproxen
- ANSWER: C

111. A 35-year-old male patient presented to a hospital with a head injury. Upon presentation, the patient was unconscious and had slight weakness in his left hand. During physical examination, the nurse instructed the patient to flex and extend his hand.

What is the most likely test checked by the nurse?

- A. Motor
- B. Sensory
- C. Reflex
- D. Coordination

ANSWER: A

112. A 67-year-old male patient presented to a hospital with complaints of severe headache and blurred vision. He has no history of trauma or any head injury. He had a history of hospitalization for skin infection three months before and was treated with IV antibiotics. On physical examination, he was negative for Kernig and Brudzinski signs.

What is the most appropriate nursing diagnosis for this patient?

- A. Acute pain related to the disease process is defined by patient verbalization
- B. Ineffective breathing related to immobility, evidenced by difficulty breathing
- C. Risk for poor skin integrity associated with mobility restriction
- D. Enhanced body image associated with surgery

ANSWER: A

113. A 60-year-old patient was brought to an emergency department by a policeman after he was found lying down on the street close to the main road. On physical examination, he is unconscious and unresponsive to shouts. He moves his extremities. His pulse rate is 100/min, respiratory rate is 30/min, temperature is 36.2 degrees centigrade, and blood pressure is 90/70mmHg. There is laceration on the scalp and right forearm.

What is the most likely medical diagnosis of this patient?

- A. Head injury
- B. Fracture of the forearm
- C. Splenic rupture
- D. Spinal cord injury

ANSWER: A

114. A 28-year-old female patient presented to a hospital with a fall down injury from a building under construction. After 24 hours of admission, the patient developed breathing difficulties. Physical examination revealed the absence of rectal tone and bradycardia in the presence of hypotension. No abnormality was detected on the ribs.

What is the most likely medical diagnosis of this patient?

- A. Hospital-admitted pneumonia
- B. Spinal cord injury
- C. Chest injury
- D. Abdominal injury

ANSWER: B

115. A nurse is providing care for a male patient who is admitted with the diagnosis of head injury and has a history of heavy bleeding. His pulse rate was 110/min, respiratory rate was 28/min, temperature was 35.8 degrees centigrade, and blood pressure was 80/60mmHg. What is the most appropriate nursing care plan in this scenario?
- A. Plan to reduce the pain within 30 minutes
 - B. Plan to stabilize the patient's vital signs within an hour
 - C. Plan to provide psychological support and calm him within one hour
 - D. Plan to prevent the dissemination of infection within one hour
- ANSWER: B
116. A nurse who is assigned to a surgical ward is providing nursing care for a patient with a diagnosis of spinal cord injury. The patient cannot provide self-care as he has developed quadriplegia due to the disease condition. The assigned nurse needs to provide skin care. What is the most appropriate nursing intervention for this patient?
- A. Protect the bony prominence
 - B. Apply talcum powder
 - C. Perform a hot bath
 - D. Assist Crunch walks
- ANSWER: A
117. A 60-year-old patient was admitted to an emergency department with the diagnosis of head injury with subdural hematoma. The surgeon assessed the general condition of the patient and told to the duty nurse to transfer him to the operating room (OR) for surgery to evacuate the hematoma. While transferring the patient to the OR, the patient's family came and asked the nurse about the situation and what was going on to the patient. What is the most appropriate nursing intervention in this scenario?
- A. Keeping the family first rather than the patient
 - B. Explains the procedure in layman's terms
 - C. No need to ask them to repeat the information back
 - D. Do not allow further questioning
- ANSWER: B
118. A patient was admitted to a surgical ward with a diagnosis of radioulnar bone fracture. Closed reduction was done. The assigned nurse in the ward has provided him with anti-pain after she assessed the overall patient condition. What is the most likely expected outcome in this scenario?
- A. Complete elimination of pain
 - B. Reduced pain to a tolerable level
 - C. Minimized use of pain medication
 - D. Avoided chronic pain
- ANSWER: B

119. An 18-year-old male patient and a 62-year-old male patient were diagnosed with hip fractures and admitted to the surgical ward for inpatient care. The type of fracture for the 18-year-old patient was comminuted and that of the 62-year-old patient was spiral. After a monthly follow-up with similar treatment and nutrition, the 62-year-old patient healed earlier than that of the 18-year-old patient. What is the most important factor that facilitated early fracture healing in this case?
- A. Age of the patient
 - B. Type of fracture
 - C. Blood supply
 - D. Bone union
- ANSWER: B
120. A 90-year-old male patient presented to a hospital with the complaints of limited range of motion and morning joint stiffness, which has lasted at least one hour for the last three months. On assessment, the nurse identified that he has pain on both knee joints and limited daily activity. What is the most likely medical diagnosis of this patient?
- A. Rheumatoid arthritis
 - B. Gout arthritis
 - C. Septic arthritis
 - D. Osteoarthritis
- ANSWER: A
121. A 60-year-old male patient who presented with complaints of pain and tenderness on his left arm was admitted to the orthopedic ward. Tenderness and decreased motility were revealed by physical examination. Laboratory and diagnostic studies evidenced that the patient has osteomalacia. What is the most likely laboratory finding that indicates the patient's condition?
- A. Low level of serum calcium
 - B. Low level of urine phosphors
 - C. Low level of ALP
 - D. Low level of urine calcium
- ANSWER: A
122. A 40-year-old male patient presented to a surgical OPD complaining of numbness on his right lower hand after he sustained a motorcycle accident two days ago. The nurse wanted to test the adequacy of the blood supply to the hand from the radial and ulnar arteries and the arcade between them. What is the most likely technique to be used in this case?
- A. Allen's test
 - B. Tinel's test
 - C. Phalen's test
 - D. Finkelstein's test
- ANSWER: A

123. A 35-year-old woman came to outpatient department with a complaint of swelling in the wrist joints of both hands. The swollen joints are tender and sensitive to touch. Laboratory diagnostic tests indicated negative rheumatoid factor, 11mg/dl uric acid level (normal value: 2.7 mg/dl-8.5 mg/dl), and WBC count of 11,000/mm³.

What is the most likely medical diagnosis of this case?

- A. Gout arthritis
- B. Osteoarthritis
- C. Inflammatory arthritis
- D. Rheumatoid arthritis

ANSWER: A

124. An 18-year-old young sportsman came to a surgical outpatient department with complaints of pain, tenderness, and loss of function at the right knee. The injured part also became deformed and short. After complete history taking and physical examination, a surgeon on duty diagnosed the patient with right knee dislocation.

For what condition is the patient at greater risk?

- A. Avascular necrosis
- B. Poor skin integrity
- C. Local tissue infection
- D. Ecchymosis

ANSWER: A

125. A 32-year-old female footballer presented with a sprained ankle following a sports injury. She was anxious about missing the upcoming football match and concerned about the repair of her injury.

What is the primary nursing diagnosis of this patient?

- A. Anxiety
- B. Impaired physical mobility
- C. Ineffective coping
- D. Acute pain

ANSWER: A

126. A 70-year-old male patient with a tibial fracture following a falling accident was admitted to the surgical ward. After evaluation, he underwent surgical repair and was immobilized in bed. He complained of discomfort at his right lower back and difficulty changing position.

What is the priority risk diagnosis in this case?

- A. Risk for impaired physical mobility
- B. Risk for impaired skin integrity
- C. Risk for disuse syndrome
- D. Risk for fluid volume deficit

ANSWER: B

127. A patient was admitted to orthopedics ward with a diagnosis of fracture. The orthopedic surgeon wanted to differentiate the type of fracture by radiologic imaging. And, the result of x-ray showed the presence of several fractured bone fragments.

What is the most likely fracture type revealed from the radiologic result?

- A. Open type fracture
- B. Green stick fracture
- C. Closed type fracture
- D. Comminuted fracture

ANSWER: D

128. A 55-year-old woman presented to a general hospital with left-sided hip joint pain. On assessment, she characterized her pain as worsening with activities and getting relief at rest. She also complained of joint stiffness at the morning for about 30 minutes. X-ray result showed the presence of narrowing spaces between hip joints.

What is the most likely medical diagnosis of this patient?

- A. Septic arthritis
- B. Osteoarthritis
- C. Rheumatoid arthritis
- D. Inflammatory arthritis

ANSWER: B

129. A 39-year-old male patient came to an emergency department after sustaining a fighting injury. During assessment, the emergency nurse found that the patient had pain, bleeding, swelling, and an open fracture on his upper right hand. After controlling the bleeding and providing painkillers, the patient was admitted to the surgical ward for further care.

What is the priority nursing care plan for this patient?

- A. Promote healing
- B. Relieve pain
- C. Prevent infection
- D. Reduce edema

ANSWER: C

130. A 60-year-old woman who had a history of kidney failure came to an adult outpatient department with a complaint of pain in the distal radius. The proximal femur was diagnosed with osteomalacia. On physical examination, skeletal deformities like spinal kyphosis and bowed legs give her an unusual appearance.

What is the most likely body system to be given an emphasis during the developing caring plan?

- A. Urinary function
- B. Gastrointestinal function
- C. Endocrine function
- D. Skin function

ANSWER: A

131. A 28-year-old male patient presented to an emergency department with falling injury. The doctor on duty sent him for x-ray, which revealed a right tibial fracture. The health care team planned to immobilize the fractured part by skeletal traction.

What is the most likely goal of immobilization?

- A. Maintain alignment
- B. Correct defects
- C. Promote physical mobility
- D. Alleviate pain associated with the fracture

ANSWER: A

132. A 60-year-old woman was admitted to orthopedic ward with a diagnosis of osteoporosis. At the orthopedic ward, the nurses provided all the necessary care, and bone density showed improvement. However, the patient was considered at risk for fracture due to her age.

What is the most effective advice to be given to the patient?

- A. Identify calcium and vitamin D rich foods
- B. Identify environmental hazards that lead to falls
- C. Provide vitamin D and calcium supplementation
- D. Engage the patient in a daily body exercise

ANSWER: B

133. A 42-year-old female patient came to the emergency department after sustaining a car accident and was admitted to the surgical ward. She was complaining of severe left lower leg pain, bleeding, and edema. Upon physical examination and diagnostic imaging, a contaminated higher-grade open fracture was diagnosed.

What is the priority nursing intervention to promote wound healing?

- A. Leave the wound open for wound drainage
- B. Control pain by providing opioids
- C. Ice pack application to control edema
- D. Monitor temperature at a normal interval

ANSWER: A

134. A 50-year-old male patient came to a general hospital with complaints of pain and stiffness in the thigh. He also reported difficulty in daily activities such as going to work. On examination, there was a swelling and tenderness with motion in the thigh muscle. After the administration of pain painkiller, a nurse planned to reduce the swelling.

What is the most appropriate intervention in this case?

- A. Immobilize with a splint
- B. Elevation of the affected leg
- C. Administer analgesics
- D. Apply a cold pad

ANSWER: B

135. A 28-year-old male patient who presented to an emergency department with a complaint of pain after sustaining a traumatic injury on the right forearm was admitted to the surgical ward. He also had swelling and tenderness over the injured arm. Diagnostic studies revealed the presence of a closed fracture. The nurse planned to immobilize the affected site.

What is the most appropriate device for immobilization?

- A. Braces
- B. Splints
- C. Fiberglass cast
- D. Plaster cast

ANSWER: B

136. A 38-year-old male patient came to an emergency department with a complaint of severe pain in the trunk and lower extremities after sustaining car accident. X-ray scan revealed fractures to both legs and the trunk. Surgical care team immobilized the patient with cast. What is the most likely type of cast used to immobilize the patient?
- A. Body cast
 - B. Hip spica cast
 - C. Short leg cast
 - D. Long leg cast
- ANSWER: B
137. A 56-year-old male patient who presented to an emergency department with severe vertebral injury following a gunshot was admitted to the surgical ward. The X-ray result revealed a vertebral fracture. Health care team immobilized the injured area. What is the most likely type of cast applied for the patient?
- A. Spica cast
 - B. Body cast
 - C. Shoulder spica cast
 - D. Hip spica cast
- ANSWER: B
138. A 44-year-old female patient came to a hospital with complaints of chills, high-grade fever, rapid pulse, tachypnea, and general malaise since 30 days after forearm surgery. A nurse provided holistic nursing care, like room ventilation, oxygen administration, and medication administration. After two hours of nursing intervention, the nurse observed a body temperature of 36.5 degrees centigrade, a respiratory rate of 18/min, and a pulse rate of 75/min. What is the most likely expected treatment outcome in this case?
- A. Complications are controlled
 - B. Infection is prevented
 - C. Body mobility is improved
 - D. Vital signs are monitored
- ANSWER: D
139. An 80-year-old male patient who developed gangrene on his left lower leg was amputated. The patient was worried and became anxious about his lost leg. He feels hopeless in his future life due to enforced dependency of being one-legged. What is the most likely expected treatment outcome the nurse should expect?
- A. Accept the changed body image and integrate changes into lifestyle
 - B. Become mobile within limitations imposed by amputation
 - C. Have reduction or absence of pain
 - D. Have no evidence of skin breakdown
- ANSWER: A
140. A mother brought a one-year-old child who has a lip abnormality, which shows disruption of the muscles of the upper lip, and has a unilateral cleft lip. What is the most likely disrupted muscle in this case?
- A. The nasolabial and bilabial muscle rings are on below and upper lip
 - B. The nasolabial and bilabial muscle rings are on two sides
 - C. The nasolabial and bilabial muscle rings are on one side
 - D. The nasolabial and bilabial muscle rings are on the lower lip
- ANSWER: C

141. In the operating room, a surgical nurse is taking care of a child who is scheduled for cleft lip repair. The nurse has used excessive draping, and the patient begins to sweat. What is the most likely nursing diagnosis in this case?
- A. Risk for perioperative hypothermia related to overwarming
 - B. Risk for perioperative hyperthermia related to overwarming
 - C. Risk for perioperative contamination related to overwarming
 - D. Risk for electrosurgical burn related to overwarming
- ANSWER: A
142. A three-month-old child is admitted to a post-anesthetic care unit after cleft lip repair was done. During handover, the circulator nurse reported the age of the patient, type of procedure performed, and the child's vital signs which was unstable. After an hour of admission to the post-anesthetic care unit, the nurse observes swelling of the tissue around the surgical site. What is the most appropriate immediate plan of care for this child?
- A. Give an analgesic
 - B. Initiate sedation
 - C. Give antibiotics
 - D. Control hemorrhage
- ANSWER: D
143. After circumcision was done for a 15-year-old child, the nurse had informed him and his family to keep the area of the operation dry for 24 hours, to keep the area clean, and to not use any antiseptic cream or ointment. What is the most likely additional information that would facilitate wound healing in this case?
- A. Do exercise for about 4 to 6 weeks per day
 - B. Avoid any sport for 4 to 6 weeks
 - C. Apply traditional ointments for 6 weeks
 - D. Do strenuous activity for 6 weeks
- ANSWER: B
144. An 18-year-old child was presented to a peripheral health facility with a complaint of penis swelling. On physical assessment, he has chronic paraphimosis, and the foreskin swells and tightens around the penis. The nurse decided that it needed emergency treatment. What is the most likely immediate nursing intervention in this case?
- A. Refer to the district hospital
 - B. Perform circumcision
 - C. Put back foreskin in place
 - D. Cold compress
- ANSWER: A
145. A mother brought a 12-year-old male child to a health center for circumcision. A surgical nurse working in minor operation room has finished collecting all the necessary materials for the procedure and started the procedure by 1 percent plain lidocaine without epinephrine injection through the skin near the base of the penis. What is the most likely priority step to be applied in this case?
- A. The preputial opening should be stretched
 - B. Homeostasis is achieved, usually by needle-tip electro-surgery
 - C. A circumferential incision is made at the reflection of the foreskin
 - D. The wound is approximated using individual fine absorbable sutures
- ANSWER: A

146. A mother brought her eight-month-old male neonate to a hospital for circumcision. A surgical nurse working in a minor operation room is preparing to perform the procedure.

What is the most priority surgical skill to apply safe circumcision?

- A. Understanding of penile anatomy
- B. Choice of appropriate suture material
- C. Gentle tissue handling
- D. Administering local anesthesia

ANSWER: A

147. A nurse is providing care for an infant in the pediatric ward who is scheduled for club foot repair. During this time, the nurse instructed the mother about nothing by mouth on the day of surgery preoperatively, including the time interval for clear liquids, breast milk, and cow milk.

What is the most likely positive outcome anticipated in this case?

- A. Demonstrated continued breastfeeding up to the time of surgery
- B. Demonstrated nothing by mouth four hours preoperatively
- C. Demonstrated nothing by mouth six hours preoperatively
- D. Demonstrated continued cow milk up to time of surgery

ANSWER: B

148. A 25-year-old female patient presented with hopelessness, inability of falling asleep, fatigue and loss of energy, recurrent suicidal ideation, and not being well concentrated on daily activities for more than 14 days. She was isolated from her friends and families. She was tearful even during interview. However, she has no other psychological and medical conditions.

What is the most likely diagnosis in this case?

- A. Anxiety disorder
- B. Major depression disorder
- C. Panic anxiety disorder
- D. Social anxiety disorder

ANSWER: B

149. A 29-year-old male patient presented with a sudden de-realization, sensations of shortness of breath or smothering, fear of losing control or going crazy and sometimes he feels as he was detached from self which lasts for one hour. He has no other psychological and medical problems.

What is the most likely diagnosis in this case?

- A. Agoraphobia
- B. Obsessive-compulsive disorder
- C. Panic attack disorder
- D. Social anxiety disorder

ANSWER: C

150. A 30-year-old female patient presented with a loss of interest in her daily activities for the last 21 days. She has lost weight, inability to fall asleep, decreased energy, and feelings of worthlessness for the last three weeks. She was indecisive nearly every day and had recurrent suicidal ideation. The health care providers confirmed that she has no other psychological or medical conditions. During the development nursing care plan, the attending nurse wants to establish a treatment goal. What is the most likely treatment goal in this case?
- A. To restore self-esteem
 - B. To promote the patient's impaired coping
 - C. To promote the patient's spiritual distress
 - D. To promote helplessness
- ANSWER: A
151. A 28-year-old male patient presented with complaints of palpitations, sweating, dyspnea, chest pain, and dizziness. On assessment, the client has sudden derealization, sensations of shortness of breath or smothering, fear of losing control or going crazy, and sometimes he feels as he is detached from himself, which lasts for one hour. The client has no other psychological or medical problems. What is the most appropriate nursing action in this case?
- A. Teaching ways to interrupt escalating anxiety
 - B. Encouraging verbalization of aspects of life not within the client's control
 - C. Reassuring that he has used adaptive activities to maintain anxiety at a manageable level
 - D. Inspiring him to accept aspects of life out of his/her control
- ANSWER: A
152. A 25-year-old female patient presented with recurrent excessive distress, persistent and excessive worry, and repeated complaints of headaches and stomachaches whenever she remembered the death of her beloved by a tragic car accident six months ago. Since then, she is also been fearful and wants to be alone. She frequently visits the cemetery to change flowers. She had no such behavior before the accident. What is the most appropriate priority nursing action in this case?
- A. Administer paroxetine 20mg every morning
 - B. Encourage her to demonstrate use of adaptive coping strategies
 - C. Motivate her to develop a structured future life plan
 - D. Discourage her to stop dysfunctional grief
- ANSWER: A
153. A 38-year-old female patient presented with more than 14 days' history of hopelessness, difficulty falling asleep, fatigue and loss of energy, recurrent suicidal ideation, and not being able to concentrate on her daily activities. She is tearful even during the interview. She was isolated from her friends and family as she prefers to be alone. She has no other psychological or medical conditions. What is the most likely expected outcome after the treatment?
- A. Interacts willingly and appropriately with others
 - B. Risk for suicide related to depressed mood
 - C. Dysfunctional grieving related to real or perceived loss
 - D. Disturbed thought processes related to withdrawal into the self.
- ANSWER: A

154. During a community level nutritional screening program, a nurse identified a 40-year-old man with decreased muscle mass, complains weakness in the lower extremity and has poor functional performance of daily activity.
What is the most appropriate nutritional assessment method to be used in this case?
- A. Calf circumference
 - B. Waist circumference
 - C. Head circumference
 - D. Mid-arm muscle circumference
- ANSWER: A
155. A nutrition monitoring and surveillance focal person at a *Woreda* health office is preparing a plan to conduct a community-level screening program by using the nutrition care process.
What is the most likely priority action in this case?
- A. Collecting nutrition-related history
 - B. Identifying nutrition-specific problems
 - C. Measuring and monitoring changes in the client's nutrition
 - D. Identifying and implementing effective individual nutritional interventions
- ANSWER: A
156. A nutrition monitoring and surveillance focal person of a *Woreda* health office is planning to conduct a campaign of community-level nutritional screening to identify iron deficiency among the population in the catchment area.
What is the most appropriate priority action in this case?
- A. Iron supplementation through the primary health care system
 - B. Iron fortification through industry-based food processing
 - C. Iron-related nutritional education and promotion
 - D. Assessment of iron nutrition status
- ANSWER: D
157. A senior surgical nurse working in a health center is performing a screening program to identify the nutritional status of individuals who are living in selected *Kebele* of the catchment area of the health center. During assessment, the nurse wanted to identify the screened population based on their risk for malnutrition.
Which groups of people are most likely susceptible to malnutrition?
- A. Older people
 - B. Pregnant women
 - C. Infants and adolescents
 - D. People with neurological diseases
- ANSWER: C
158. During community-level nutritional screening, a 40-year-old man presented with complaints of fatigue and depression of five days duration. On assessment, the nurse identified that the person has a history of bleeding, bruising, slow recovery from infections, and poor wound healing.
What is the most likely nutritional deficiency the patient developed in this case?
- A. Vitamin D
 - B. Vitamin C
 - C. Vitamin E
 - D. Vitamin B₁
- ANSWER: B

159. A nurse is planning to provide prison health education to develop an understanding of the target population problem, develop a relationship with the population, and create awareness by using a comprehensive health education model.

What is the most appropriate phase of health education planning applied in this case?

- A. Program initiation
- B. Need assessment
- C. Goal setting
- D. Programming

ANSWER: A

160. A sanitarian nurse in the surgical ward was managing the overall waste handling and disposal of waste that contains human tissues or fluids, organs, body parts, fetuses, and unused blood products.

What is the most likely category of health care waste in this case?

- A. Pathological waste
- B. Infectious waste
- C. Cytotoxic waste
- D. Non-infectious waste

ANSWER: A

161. A nurse is providing care for a 23-year-old male patient who came to a hospital with a sudden onset of copious and watery diarrhea. On assessment, the nurse identified that he has a history of occasional vomiting and muscle pain. On physical examination, the skin pinch goes back slowly. Finally, the patient was diagnosed with cholera, and the health care intervention was ongoing at a separate center.

What is the most effective method of preventing of spread of the disease?

- A. Antibiotic prophylaxis
- B. Control through sanitation
- C. A parenteral vaccine of whole killed bacteria
- D. Breastfeeding in endemic areas protects infants from disease

ANSWER: B

162. A director of a health center is preparing a monthly plan to conduct public health communication as health education is one of the components of primary health care.

What the most priority principles in the application of health education?

- A. Promoting Community Involvement and Self-Reliance
- B. Enhancing Decision-Making Skills at the Local Level
- C. Harmonizing National and Local Plans
- D. Facilitating Inter-Sectoral Action

ANSWER: A

163. A nurse working in a health center is providing public health education on environmental hygiene at selected *Kebele* in the catchment area. During the health education, the nurse used objective displays of information, including photographs and posters.

What is the most appropriate health communication method used in the above case?

- A. Visual Communication
- B. Verbal Communication
- C. Written Communication
- D. Non-Verbal Communication

ANSWER: A

164. A surgical nurse is providing public health education to enhance community participation towards the prevention of cholera. The nurse wants to find a powerful mood setter communication method to meet the objective.
What is the most appropriate health communication method to be used in this case?
- A. Music
 - B. Panels
 - C. Models
 - D. Newsletters
- ANSWER: A
165. A team of middle-level managers came together to develop an annual plan and to translate the broader plan of the health sector into measurable objectives. This plan was intended to specify what the lower-level units within each division must do, how they would do it, and who was in charge at each level.
What is the most likely type of planning to be developed in this case scenario?
- A. Tactical planning
 - B. Strategic planning
 - C. Operational planning
 - D. Contingency planning
- ANSWER: A
166. A nurse working in a hospital was newly assigned as head nurse and wanted to manage a problem that already existed. The head developed a plan on how to manage the problem.
What is the most likely type of planning developed in this case?
- A. Inactive planning
 - B. Reactive planning
 - C. Preactive planning
 - D. Proactive planning
- ANSWER: B
167. A nurse working in a *Woreda* health office has a variety of roles in the office, including giving emphasis to group process, gathering information, and giving feedback.
What is the most likely characteristic that indicates the nurse as a leader?
- A. Giving feedback
 - B. Gathering information
 - C. Seeking variety of roles
 - D. Emphasizing group process
- ANSWER: D
168. While two nurses were walking around, they started to argue about the difference between a leader and a manager. The nurse who said leaders are different from managers tried to convince by mentioning that leaders are planners, monitors, role models, and evaluators.
Which character of a leader mentioned in this case is most appropriate?
- A. Planners
 - B. Monitors
 - C. Evaluators
 - D. Role models
- ANSWER: D

169. A surgical ward head nurse identified that there is a shortage of nurses in the ward and requested the nursing director to assign some nurses from the medical ward. The nursing director assigned three nurses to the surgical ward based on the department's request. The surgical ward head nurse introduced the assigned nurses to patients, oriented them about the ward, and facilitated a working environment. What is the most likely skill of the nurse explained in this case scenario?
- A. Organizing
 - B. Planning
 - C. Staffing
 - D. Leading
- ANSWER: C
170. A department head of a surgical ward in a district hospital wanted to prepare a report by using Microsoft Excel. However, the head could not do that due to a lack of skill in Microsoft Excel. Then, the department head did it manually on paper. What is the most likely skill deficiency explained in this case?
- A. Communication skills
 - B. Conceptual skill
 - C. Technical skill
 - D. Human skill
- ANSWER: C
171. A human resource manager of a hospital recruited five new employees. The manager provided job description and took to respective department. With the immediate manager of the department, the employees were assigned to specifically identified jobs. The manager facilitated the working area and gave a desktop computer to all employees recruited. What is the most likely skill of the manager explained in this case scenario?
- A. Organizing
 - B. Controlling
 - C. Planning
 - D. Leading
- ANSWER: A
172. A health care institution that can serve 250,000 populations, which has 50 beds for admission, 30 technical and 20 non-technical staff, was built in the northeast of Ethiopia. What is the most likely type of health care institution built in this case (Ethiopian context)?
- A. Primary health care unit
 - B. District hospital
 - C. Regional hospital
 - D. Specialized hospital
- ANSWER: B
173. Ministry of Health organized a supervision team to evaluate tertiary prevention mechanisms of health institutions in Ethiopia. The team identified the hierarchies of the institutions and selected the appropriate institution. And then, the team directly went to the identified institutions. What is the most likely institution they have to choose in this case?
- A. District hospital
 - B. Regional hospital
 - C. Specialized hospital
 - D. Primary health care unit
- ANSWER: C

174. A surgical ward head nurse was a person-oriented supervisor. The head nurse controls the workers based on needs. The head nurse has done nursing rotation past few weeks. At that time, the head nurse assigned the workers considering their training and working style. The head nurse's concern during supervision was the worker's personality.

What is the most likely characteristic that shows the head nurse is a person-oriented supervisor?

- A. Concerns of workers' personality
- B. Seeking quantity and quality of the task
- C. Controlling the workers based on their needs
- D. Assignments of the workers considering their training

ANSWER: A

175. A researcher planned to assess the effects of the use of motorcycle helmets on head injuries during motorcycle accidents. Then, the researcher recruited and interviewed individuals who were treated for head injury in the emergency rooms, and those who were treated for other injuries due to motorcycle accidents.

What is the most likely study design employed for the above scenario?

- A. Cohort
- B. Case control
- C. Experimental
- D. Cross-sectional

ANSWER: B

176. Groups of researchers planned to study the effect of problem-based learning on academic performance among medical students. In the study, medical students were randomly assigned to universities that employed problem-based learning and universities that did not employ problem-based learning. At the completion of the medical education program, the academic performance of the students from both universities was assessed by taking the cumulative average grade, proportion of passing rate at the licensing exam, and by results of oral exams, which were administered centrally. What is the most likely study design used in this scenario?

- A. Cohort
- B. Case control
- C. Experimental
- D. Cross-sectional

ANSWER: C

177. A researcher has planned to conduct a study to assess the knowledge and attitudes of first-year university students of a public university in Ethiopia regarding human immunodeficiency virus (HIV). The researcher decided to collect data at the time of their first arrival at the University. As a result, the researcher selected and recruited every other student who arrived at the university during the study period.

What is the most likely sampling technique employed in this scenario?

- A. Cluster sampling
- B. Stratified Sampling
- C. Systematic sampling
- D. Simple random sampling

ANSWER: C

178. A researcher has planned to conduct a study to determine the magnitude and associated factors of turnover of nurses who are working in different health facilities of a city. For this study, nurses were recruited from selected hospitals and health centers of the city. Then, the researcher took a proportional sample size of nurses from the selected hospitals and health centers independently. What is the most likely sampling technique employed in this scenario?
- A. Cluster sampling
 - B. Convenience sampling
 - C. Stratified Sampling
 - D. Multistage sampling
- ANSWER: C
179. During an investigation of a cholera outbreak, a health professional identified six individuals with signs and symptoms of cholera. The health professional collected the following incubation period in days reported from each patient as: 3, 2, 3, 4, 5, 1. What is the mean incubation period of cholera in days for this data set?
- A. 1
 - B. 3
 - C. 4
 - D. 5
- ANSWER: B
180. A nurse reported the hemoglobin levels of pregnant women who came to the Antenatal Care (ANC) clinic for regular ANC services on a day. Then the nurse screened five pregnant women for further medical consultation, and their corresponding hemoglobin measurements in g/dl were: 12, 13, 11, 14, and 15. What is the mean hemoglobin value for this dataset?
- A. 12 gm/dl
 - B. 13 gm/dl
 - C. 14 gm/dl
 - D. 15 gm/dl
- ANSWER: B
181. A nurse reported the systolic blood pressure measurements of pregnant women who came to the Antenatal Care (ANC) clinic for regular ANC services on a day. During screening, systolic blood pressure measurements in mmHg of five pregnant women were: 120, 130, 110, 140, and 150. What is the median systolic blood pressure value for the above scenario?
- A. 120 mmHg
 - B. 130 mmHg
 - C. 140 mmHg
 - D. 150 mmHg
- ANSWER: B

182. A researcher has planned to study the attitudes of male partners towards long-acting family planning methods. For this study, the researcher considered two ANC clinics in different hospitals. Then, the researcher tried to recruit the male participants whenever they came to the clinic, accompanied their partner.

What is the most likely sampling technique applied in this case?

- A. Quota sampling
- B. Purposive sapling
- C. Snowball sampling
- D. Convenience sampling

ANSWER: D

183. A physician, who is assigned to teach surgical nursing students, provided a brief overview and presented the steps and procedures of urinary catheterization, which will be done for patients admitted in a surgical ward. Following that, the students tried to perform urinary catheterization by using anatomic models.

What is the most likely teaching method employed in this case?

- A. Role play
- B. Case study
- C. Demonstration
- D. Facilitated practice

ANSWER: C

184. An instructor has assessed the performance of final year nursing students on how to perform a pelvic examination. For evaluation purposes, the instructor lists out the detailed steps of the procedure and the criteria for evaluating the performance of the students. Accordingly, the performance of each student was evaluated.

What is the most likely assessment method used by the teacher in this case?

- A. Evaluated assessment
- B. Diagnostic assessment
- C. Norm-referenced assessment
- D. Criterion-referenced assessment

ANSWER: D

185. A lecturer, who is assigned to give a course for third-year nursing students, has planned to give an hour-long lecture regarding pregnancy-induced hypertension. The lecturer has set this objective “At the end of this lecture, students should try to identify the signs and symptoms of pregnancy-induced hypertension”.

What is the most likely objective type presented in the above scenario?

- A. Knowledge objective
- B. Skill objective
- C. Attitude objective
- D. Psychomotor objective

ANSWER: A

186. A professor has set this objective for a given course: “After completing this course, try to assess, classify, and treat a sick child in an effective and integrated manner.”
What most appropriate information is missing in the above course objective?
A. When to demonstrate the knowledge or performance
B. Who will demonstrate the competency
C. What will be demonstrated
D. How well the performance must be demonstrated
ANSWER: B
187. A 30-year-old male patient came to a clinic for a check-up. A nurse provided him with the necessary information about HIV and the benefits of HIV testing. But he refused to be tested despite the HIV counseling. The nurse told him to come at any time if he changes his mind.
What is the most likely ethical principle the nurse respected in this case?
A. Nonmaleficence
B. Beneficence
C. Autonomy
D. Justice
ANSWER: C
188. A nurse is providing care for a terminally ill patient who is in need of an urgent blood transfusion. The patient refuses the blood transfusions due to his religion.
What is the most appropriate next action to be taken by the nurse?
A. Inform the family and give the decision to them
B. Give the blood because it is important for survival
C. Discuss its benefit by respecting patient’s need
D. Withhold the blood because of the patient’s need
ANSWER: C
189. A nurse in a hospital usually commits maltreatment of children if they are not cooperating with the treatment. The head nurse heard a rumor on this and developed a strict regulation regarding child abuse and made it every health care provider’s obligation to report suspected child abuse to protect children.
What is the most likely ethical principle the head nurse is trying to correct in this case?
A. Nonmaleficence
B. Beneficence
C. Autonomy
D. Justice
ANSWER: A
190. A surgeon ordered a nurse to prepare a 28-year-old male patient for surgery. The patient asked the nurse about the qualifications of the physician who is going to perform the procedure. The nurse told him the procedure would be performed by a physician who specialized in general surgery.
What is the most likely ethical principle the nurse respected in this case?
A. Beneficence
B. Fidelity
C. Justice
D. Veracity
ANSWER: D

191. A 40-year-old known asthmatic patient with a complaint of shortness of breathing and a 36-year-old male patient with a complaint of acute diarrhea came to the emergency ward at the same time. There was only one nurse to provide service at a time. The nurse first considered the patient with shortness of breathing based on the severity of their condition.
What is the most likely ethical principle the nurse used for the decision in this case?
- A. Beneficence
 - B. Autonomy
 - C. Veracity
 - D. Justice
- ANSWER: D
192. A researcher conducted a study on the prevalence of surgical site infection and its associated factors among patients who undergone surgery in a hospital. The researcher reported the exact data without manipulation of the data.
What is the most likely ethical principle best adhered by the researcher in this case?
- A. Autonomy
 - B. Fidelity
 - C. Justice
 - D. veracity
- ANSWER: B
193. A nurse is providing care for an 18-year-old female patient in the surgical ward who was admitted following a cesarean section. On the afternoon shift, the nurse discovered that a physician has prescribed an unusually large dosage of a medication.
What is the most appropriate action of the nurse in this case?
- A. Refuse to administer the medication
 - B. Administer the medication
 - C. Notify the prescriber
 - D. Call the pharmacist
- ANSWER: C
194. A nurse is providing care for a patient who presented with a complaint of acute abdominal pain. The nurse did not report the client's complaint to the health care team, which resulted in rupture of the appendix. Finally, the patient died.
What is the most likely error committed by the nurse in this case?
- A. Negligence
 - B. Assault
 - C. Battery
 - D. Defamation
- ANSWER: A
195. A nurse administered an injection of Tramadol, as per the physician order, to a client with a complaint of pain. The nurse failed to document the site and mode of the injection on the patient card.
What is the most likely tort committed by the nurse in this case?
- A. Malpractice
 - B. Assault
 - C. Battery
 - D. Defamation
- ANSWER: A

196. A 19-year-old female patient who was admitted to a surgical ward refused to take oral medication. The nurse threatens the patient with an injection. Finally, the nurse gave the injection without the client's consent.

What is the most likely tort the nurse has been guilty in this case?

- A. Negligence
- B. Assault
- C. Battery
- D. Defamation

ANSWER: C

197. A 22-year-old male patient came to the surgical ward after colorectal surgery. He was feeling severe pain at the incision site and wanted to get anti-pain. But the nurse was out from the ward for lunch.

What is the most likely characteristic of respectful care breached in the case?

- A. Abandonment of care
- B. Non-consented care
- C. Non-confidential care
- D. Non-dignified care

ANSWER: A

198. A nurse is preparing a patient with advanced cervical cancer for surgery. The patient was feeling hopeless due to fear of a poor prognosis of the treatment. The nurse said, "I am sorry; I understand that it really is difficult for you."

What is the most likely compassionate characteristic of the nurse this case?

- A. Empathy
- B. Sympathy
- C. Motivation
- D. Sensitivity

ANSWER: B

199. A nurse is providing care for a 26-year-old male patient diagnosed with bowel obstruction. A surgeon ordered the nurse to prepare the patient for surgery. The patient is worried about his pain. The nurse, feeling concerned about his pain, tried to alleviate suffering of the patient.

What is the most likely compassionate characteristic of the nurse in this case?

- A. Empathy
- B. Sympathy
- C. Motivation
- D. Sensitivity

ANSWER: A

200. A five-year-old girl who has lost her mother came to an emergency ward with a complaint of a headache. During assessment, the nurse observed that the patient feels loneliness due to the death of her mother. The nurse, feeling sorry for the girl described her condolences.

What is the most likely compassionate characteristic applied by the nurse in this case?

- A. Empathy
- B. Sympathy
- C. Motivation
- D. Sensitivity

ANSWER: B