

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

**PUBLIC
HEALTH**



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority, essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, leadership and management, ethics, and Health Promotion as outlined in the Information Booklet 2024.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies, including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey—and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Public Health

1. A 36-year-old man, who was recently diagnosed with HIV infection, presented to the outpatient department with a three-month history of painless neck swelling over the posterior cervical region. He also has a low-grade fever, anorexia, and significant weight loss. He was repeatedly treated with antibiotics but showed no improvement. On examination, there is a non-tender 4 cm by 5 cm mass with whitish discharge. A laboratory investigation result showed an ESR of 50 mm/hour.

What is the most likely diagnosis for this case?

- A. Lymphoma
- B. TB lymphadenitis
- C. Pyogenic lymphadenitis
- D. Reactive lymphadenitis

ANSWER: B

2. A 34-year-old man with a known history of HIV infection presented to an emergency department with complaints of shortness of breath, dry cough, and fever of ten days' duration. He reported that he discontinued his HAART two years ago. On examination, the pulse rate is 120 beats/minute, the respiratory rate is 28 breaths/minute, the temperature is 37.8 degrees centigrade, and oxygen saturation is 86 percent. On lung auscultation, bilateral crackle sounds are heard. His investigation result showed that the CD4 count is 120/microliter, and the chest x-ray shows faint bilateral interstitial infiltrates.

What is the most likely diagnosis for this patient?

- A. Bacterial pneumonia
- B. Idiopathic interstitial pneumonia
- C. Pulmonary Tuberculosis
- D. Pneumocystis jiroveci pneumonia

ANSWER: D

3. An 18-year-old young man from a malarial area presented to an emergency department with complaints of high-grade fever, chills, and rigor of six days' duration. He also has associated symptoms of fatigue and dizziness. On examination, pulse rate is 128/minute, temperature is 38.9 degrees centigrade, and he has pale and pallor conjunctivae. His hemoglobin level is 7 mg/dl, his random blood sugar level is 140 mg/dl, and the blood film result shows ring-forming trophozoites of blood parasites.

What is the most likely evidence that indicates a severe sign of malaria for this patient?

- A. Pulse rate of 128/min
- B. Hemoglobin of 7 mg/dl
- C. Temperature of 38.9 degrees centigrade
- D. Random blood sugar of 140 mg/dl

ANSWER: B

4. A 45-year-old prisoner is brought to the emergency department for complaints of a high-grade fever and severe headache for the last three days. He also reported having severe chest pain and a dry cough. Prisoners from the same prison also have similar complaints. Laboratory test results showed elevated Weil-felix titer.

What is the most likely mode of transmission of the infection in this case?

- A. Feco-oral
- B. Mosquito bite
- C. Infected lice
- D. Infected tick

ANSWER: C

5. A 34-year-old man, who has been having dyspeptic symptoms for the past three years and on antacids, presented to the emergency unit with sudden severe upper abdominal pain that rapidly became generalized. On examination, the respiratory rate is 20 breaths/minute with shallow breathing, and there is generalized board-like rigidity of the abdomen. Erect chest x-ray shows free air beneath the diaphragm.

What is the most likely diagnosis for this patient?

- A. Perforated PUD
- B. Acute Cholecystitis
- C. Acute Gastritis
- D. Acute Pancreatitis

ANSWER: A

6. A 26-year-old man came to a health center with a complaint of yellowish discoloration of his eyes for a week. He has also anorexia and fatigue for the last few weeks. He admits to have had casual unprotected sexual practice with multiple partners. On examination, there is right upper quadrant tenderness, and the liver edge is palpable. Laboratory test result shows ALT of 280 U/ml and AST of 260 U/ml and hemoglobin of 14 g/dl.

What is the most likely diagnosis for this patient?

- A. Viral Hepatitis
- B. Acute Pancreatitis
- C. Acute Cholecystitis
- D. Chronic liver disease

ANSWER: A

7. A 26-year-old health officer sustained a needle stick injury while he was trying to take a blood sample from a Hepatitis B positive patient, admitted for the management of chronic liver disease. What is the next best step in the management of the health officer?

- A. Provide Hepatitis B Immunoglobulin
- B. Screen for Hepatitis B
- C. Provide Hepatitis B vaccine
- D. Start antiviral drugs

ANSWER: C

8. A 46-year-old man came to an emergency department with complaints of a productive cough and a low-grade fever of one week duration. He also reported to have loss of appetite and mild weight loss. He is a known alcoholic, and two weeks ago, he was admitted for alcohol overdose. On examination, pulse rate is 90 beats/min, respiratory rate is 24 breaths/minute, and crackles sounds are heard on auscultation of the right lower lung field. Gram staining of the sputum shows mixed bacterial flora. Chest x-ray demonstrates consolidation in the right lower lobe.

What is the most likely diagnosis of this patient?

- A. Bronchiectasis
- B. Pulmonary TB
- C. Aspiration pneumonia
- D. Necrotizing Pneumonitis

ANSWER: C

9. A 36-year-old female patient came to an emergency department with complaints of progressive left pleuritic chest pain, fever and shortness of breath for three three-week duration. She was diagnosed with HIV infection a week before. On examination, pulse rate is 90 beat/minute, respiratory rate is 22 breaths /minute, and temperature is 37.8 degrees centigrade. The left lower lung field is dull to percussion and has no air entry. On investigation, WBC count is 11,000 cells/microliter. Pleural fluid analysis showed WBC count of 900/microliter with 65 percent lymphocytes. Pleural fluid cytology was unremarkable.

What is the most likely cause of the effusion in this case?

- A. Pleural TB
- B. Mesothelioma
- C. Pulmonary embolism
- D. Parapneumonic effusion

ANSWER: A

10. A 34-year-old female patient came to an emergency OPD with complaints of progressive worsening of shortness of breath and wheezing over the past one month. She has a history of asthma since childhood, but notes that her symptoms have been progressively worsening over the past month despite adherence to her asthma medications. She described that the symptoms are ongoing on a daily basis and every night. On examination, pulse rate is 90 beats /minute, respiratory rate is 28 breath/minute, and temperature is 36.9 degrees centigrade. There is diffuse bilateral wheezing on auscultation of both lung fields.

What is the most likely diagnosis for this patient?

- A. Mild persistent asthma
- B. Mild intermittent asthma
- C. Severe persistent asthma
- D. Moderate persistent asthma

ANSWER: C

11. A 42-year-old male patient came to a health center with a complaint of progressively worsening bilateral leg swelling of one month duration. He also has shortness of breath with exertion. On examination, blood pressure is 160/90 mmHg, pulse rate is 90 beats/minute, respiratory rate is 18 breaths/minute, and fine crackles are heard on the base of both lung fields on auscultation. ECG shows left axis deviation. Echocardiogram demonstrates left ventricular wall hypertrophy with an ejection fraction of 50 percent and mild left atrial dilatation.

What is the most likely cause of heart failure for this patient?

- A. Valvular heart disease
- B. Infective endocarditis
- C. Ischemic heart disease
- D. Hypertensive heart disease

ANSWER: D

12. A 52-year-old male patient came to an emergency department with complaints of a three-week history of gradually worsening shortness of breath and fatigue. He has a history of rheumatic heart disease during childhood for which he underwent mitral valve replacement. On examination, pulse rate is 106 beats/minute and temperature is 38.6 degrees centigrade. A mechanical click was heard at the apex on cardiac auscultation, and tipped splenomegaly was detected during abdominal palpation. WBC count is 14, 000/micro liter, hemoglobin is 11 mg/dl, platelet count is 90,000/microliter, and ESR is raised.

What is the most likely diagnosis for this patient?

- A. Infective endocarditis
- B. Acute rheumatic fever
- C. Rheumatic heart disease
- D. Acute respiratory distress syndrome

ANSWER: A

13. A 50-year-old man visited a clinic for elevated blood pressure, which was detected during a screening campaign. On examination, his seated blood pressure is 155/95 mmHg. He was investigated for his basic metabolic panel, and the results were unremarkable.

What is the most likely diagnosis for this patient?

- A. Prehypertension
- B. Stage I hypertension
- C. Stage II hypertension
- D. Isolated systolic hypertension

ANSWER: B

14. A 12-year-old male child was brought to an emergency department with the complaints of tea-colored urine and swelling in his legs and face of three days duration. His mother reports that he had a sore throat and fever two weeks ago. On examination, his blood pressure is 130/90 mmHg. Urinalysis demonstrates Proteinuria (2+), hematuria(2+), and RBC casts. Serum Creatinine is 1.4 mg/dl (slightly elevated), and ASO titer is elevated.

What is the most likely diagnosis for this patient?

- A. Nephrotic syndrome
- B. Acute pyelonephritis
- C. Acute kidney injury
- D. Acute glomerulonephritis

ANSWER: D

15. A 34-year-old pregnant woman came to an emergency department with complaints of high grade fever, chills and left flank pain of two days duration. She also has nausea and three episodes of vomiting of ingested matter. On physical exam, she has costovertebral angle tenderness and urinalysis shows positive leukocyte esterase and pyuria.

What is the most appropriate initial management strategy applied for this patient?

- A. Intravenous fluid and analgesics
- B. Treat with oral antibiotics as an outpatient
- C. Admission for intravenous antibiotics and supportive care
- D. Admit and follow without intervention until the urine culture result arrives

ANSWER: C

16. A 28-year-old female patient came to the outpatient department with complaints of recurrent headaches over the past few months. She describes the headache as throbbing and pulsatile, typically located to one side of the head. She also has associated nausea, vomiting, and sensitivity to light and sound. She has no any other neurologic complaint.

What is the most likely diagnosis for this patient?

- A. Sinus headache
- B. Cluster headache
- C. Tension headache
- D. Migraine headache

ANSWER: D

17. A 38-year-old man presented to the outpatient department with a history of joint pain in his hands and wrists over the past few months. He reported that the joints are stiff in the morning and get improve as the day progresses. On physical examination, the patient has swollen and tender metacarpophalangeal, proximal interphalangeal, and wrist joints. Laboratory test result shows elevated ESR and CRP.

What is the most likely diagnosis for this patient?

- A. Gout arthritis
- B. Osteo arthritis
- C. Septic arthritis
- D. Rheumatoid arthritis

ANSWER: D

18. A 21-year-old male patient came to the outpatient department with the complaints of polyuria and polydipsia over the past two weeks. He feels more fatigued than usual and has lost some weight despite having good appetite. He denies any history of trauma or recent illness. On examination, pulse rate is 98 beats/minute and mild dehydration is noted. The health officer decides to investigate the patient in line of diabetes mellitus.

What is the best diagnostic test for this patient?

- A. Hemoglobin A1C
- B. Oral glucose tolerance test
- C. Random blood glucose
- D. Two-hour plasma glucose test

ANSWER: A

19. A 26-year-old female patient came to an emergency department with the complaints of nausea, vomiting and abdominal pain of one day duration. Over the last few days, she has been having the urge to drink water too much but does not feel quenched. On physical examination, her pulse rate is 108 beats/minute, respiratory rate is 28 breaths/min with deep and shallow breathing. She has dry mouth and buccal mucosa. Random blood glucose measurement indicated a blood glucose level of 450 mg/dl and urine is positive for glucose and ketone.

What is the most likely diagnosis for this patient?

- A. Hyperglycemia
- B. Acute pancreatitis
- C. Diabetic ketoacidosis
- D. Hyperosmolar hyperglycemic state

ANSWER: C

20. A 34-year-old female patient came to a health center with complaints of fatigue and shortness of breath with exertion for two months. She also has occasional dizziness, hair loss and brittle nails. On examination, she appears pale and has pale conjunctivae. Her hemoglobin level is 9 g/dl, and MCV is 70 fl.

What is the most likely diagnosis for this patient?

- A. Hemolytic anemia
- B. Iron deficiency anemia
- C. Megaloblastic anemia
- D. Sideroblastic Anemia

ANSWER: B

21. A 33-year-old Gravida IV, Para III pregnant woman presented to a health center with the complaint of progressively worsening fatigue and dizziness of three months duration. She also has shortness of breath during exertion. Her previous pregnancies were uncomplicated and ended up with smooth delivery. Otherwise, she has no other recent illnesses. On examination, she appears pale and has pale conjunctivae. On investigation, her hemoglobin level is 9 g/dl, MCV is 70 fl and serum ferritin level is low.

What is the most likely contributing factor to the patient's presentation?

- A. Blood loss
- B. Inadequate iron intake
- C. Decreased iron absorption
- D. Increased metabolic demand

ANSWER: D

22. A 54-year-old female patient, who has been on follow-up at a cardiac clinic for the past 10 years, was brought to an emergency unit with a complaint of shortness of breath and chest pain. She had been diagnosed with chronic rheumatic valvular heart disease. On physical examination, vital signs were; pulse rate of 132beats/minute, respiratory rate of 33breaths/minute with atmospheric oxygen, temperature of 36.5 degrees centigrade, and blood pressure of 80/40mmHg. What is the most likely cause of shock in this patient?

A. Septic
B. Hypovolemic
C. Neurogenic
D. Cardiogenic

ANSWER: D

23. A 45-year-old male patient presented to an emergency room with a complaint of a productive cough of two days duration. He also has an associated fever, shortness of breath and altered mentation of same duration. On physical examination, his vital signs showed: pulse rate of 134beats/minute, respiratory rate of 34breaths/minute with 3 liters of Intranasal oxygen, blood pressure of 70/40mmHg, temperature of 38.9 degrees centigrade. Despite being resuscitated with 30ml/kg crystalloid, which made the inferior vena cava full, and other one-hour bundle management, the shock could not be recovered.

What is the most appropriate next step in the management of this patient?

A. Initiate steroid
B. Initiate vasopressor
C. Change antibiotics
D. Add more fluid resuscitation

ANSWER: B

24. A four-year-old male child incidentally ingested multiple tablets of Isoniazid after which he was immediately taken to an emergency room.

What is the most appropriate antidote drug to be given to this child?

A. Naloxone
B. Thiamine
C. Pyridoxine
D. Sodium bicarbonate

ANSWER: C

25. A 41-year-old female patient was admitted for elective surgery after she was diagnosed with symptomatic cholelithiasis. While cholecystectomy was being performed, there was minimal spillage of bile.

What is the most likely type of surgical wound for this patient?

A. Clean the contaminated wound
B. Contaminated wound
C. Clean wound
D. Dirty wound

ANSWER: A

26. A 40-year-old male, diabetic patient, came to an emergency room on his 15th post operation day after abdominal surgery with a complaint of wound site pain. On abdominal examination, there are erythematous skin, local edema, and tenderness with minimal pus oozing from the wound. Abdominal ultrasound is unremarkable, and the surgeon diagnosed the condition as a superficial surgical site infection.

What is the most appropriate management of this patient?

- A. Local exploration with wound care
- B. Re-laparotomy for Intra-abdominal collection
- C. Broad-spectrum PO antibiotics
- D. Broad-spectrum IV antibiotics

ANSWER: A

27. A 35-year-old female patient presented to the surgical OPD with a complaint of anterior neck swelling associated with cold intolerance, loss of mood, weight gain, and sleepiness of 5 years duration. Her serum TSH (thyroid stimulating hormone) is 10mU/L (High), neck ultrasound shows cystic degeneration and FNAC result indicated a colloid goiter.

What is the most likely diagnosis of this patient?

- A. Hyperthyroidism
- B. Hypothyroidism
- C. Hashimoto's thyroiditis
- D. Suppurative thyroiditis

ANSWER: B

28. A 30-year-old female patient, who is a known psychiatric and on lithium, came with a complaint of new development of anterior neck swelling. The patient asked for preventive measures that help to avoid the occurrence of goiter.

What best strategy that can be implemented to prevent the occurrence of goiter?

- A. Avoiding cassava and cabbage diets
- B. Regular consumption of iodized salt
- C. Regular consumption of seawater foods
- D. Avoiding the use of anti-thyroid drugs

ANSWER: B

29. A 25-year-old lactating mother came to an emergency unit with a complaint of right breast swelling associated with severe pain, chills, rigor, and high-grade fever of one week duration. On breast examination, there was a tender, boggy mass on her breast with erythematous skin. Right breast ultrasound revealed an anechoic collection of likely an abscess. Her WBC count was 11,000cells/microliter.

What is the most appropriate initial management for this patient?

- A. Needle aspiration
- B. Incision and drainage
- C. Broad-spectrum IV antibiotics
- D. Broad-spectrum PO antibiotics

ANSWER: B

30. A 32-year-old male patient came to an emergency room with a complaint of the sudden onset of severe epigastric abdominal pain associated with vomiting of ingested matter of three hours duration. In the past, he had a repeated history of treatment for dyspeptic symptoms that worsen with alcohol intake. On physical examination, blood pressure of 90/60mmHg, pulse rate of 120 beats/minute, and temperature of 38 degrees centigrade. Erect chest radiograph revealed air under the diaphragm.

What is the most likely diagnosis of this patient?

- A. Acute exacerbation of peptic ulcer disease
- B. Acute alcohol induced pancreatitis
- C. Acute exacerbation of gastritis
- D. Perforated peptic ulcer disease

ANSWER: D

31. A 15-year-old male patient presented to an emergency room with a complaint of crampy abdominal pain of one day duration. The pain started around his umbilicus and later shifted to the right lower quadrant. He also has associated anorexia, nausea, two episodes of vomiting of ingested matter and low-grade fever. On abdominal examination, there was right lower quadrant direct and rebound tenderness.

What is the most likely diagnosis of this patient?

- A. Acute appendicitis
- B. Mesenteric lymphadenitis
- C. Acute ileitis
- D. Acute gastroenteritis

ANSWER: A

32. A 50-year-old male patient came to an emergency unit with a complaint of failure to pass stool and flatus associated with colicky abdominal pain of one day duration. On physical examination, his vital signs are in the normal range. On abdominal examination, he has a grossly distended abdomen with visible peristalsis, and hyper-tympanic note but no tenderness. Digital rectal examination showed a ballooned-out empty rectum and no blood on the examining finger. Abdominal radiograph revealed dilated large bowel loop. His WBC count is 9000cells/microliter.

What is the most appropriate initial management for this patient?

- A. Manual rectal tube deflation
- B. Sigmoid resection and colostomy
- C. Sigmoid resection and anastomosis
- D. IV fluid resuscitation and antibiotics

ANSWER: A

33. A 45-year-old male, known diabetic patient presented to an emergency unit with a complaint of painful swelling on his buttock associated with chills and rigor of three days duration. On physical examination, there is tender, boggy mass on the right lateral part of anal orifice with erythematous skin.

What is the most likely diagnosis of this patient?

- A. Pilonidal sinus
- B. Perianal fistula
- C. Perianal abscess
- D. Thrombosed haemorrhoid

ANSWER: C

34. A 46-year-old male patient presented to surgical OPD with a complaint of peri-anal itching sensation and protruding mass per rectum of two months duration. On physical examination, he has BMI of 30Kg/meter square. Digital rectal examination revealed grade two internal hemorrhoids. What are the best preventive strategies to halt the progress of his hemorrhoid?
- A. Dietary modification and avoid straining
 - B. Topical analgesia and physical exercise
 - C. Topical steroid and weight loss
 - D. Haemorrhoidectomy and Seitz bath
- ANSWER: A
35. A 38-year-old male patient came to an emergency unit with a complaint of failure to pass urine of one day duration. He had a history of treatment for recurrent gonorrhea. On physical examination; he has a distended urinary bladder to the level of umbilicus. Trial of catheterization failed even with small sized catheter. Abdominal and pelvic ultrasound revealed distended urinary bladder, and prostate size of 40ml.
- What is the most likely cause of urinary retention in this patient?
- A. Urethral stricture
 - B. Benign prostatic hyperplasia
 - C. Bladder neck contracture
 - D. Neurogenic bladder
- ANSWER: A
36. A 35-year-old female patient was brought to an emergency unit after she sustained road traffic accident. On examination, her vital signs were; blood pressure of 80/50mmHg, pulse rate of 120beats/minute, respiratory rate of 30 breaths/ minute, oxygen saturation of 92 percent and random blood sugar was 90mg/dl.
- What is the best initial treatment for this patient to prevent secondary brain injury?
- A. Dextrose 40 percent IV push to increase her blood sugar
 - B. Fluid resuscitation to increase blood pressure
 - C. Put her on face mask oxygen to increase saturation
 - D. Sedate her to decrease her respiratory rate
- ANSWER: B
37. A 12-year-old male child was brought to a surgical OPD with complaints of right leg swelling, pain, limping and fever of one week duration after he sustained falling down injury. On physical examination, there was a warm and swollen right leg with point tenderness. Leg radiograph showed periosteal elevation and leg ultrasound shows soft tissue oedema.
- What is the most likely diagnosis of this patient?
- A. Acute osteomyelitis
 - B. Cellulitis of the leg
 - C. Acute pyomyositis
 - D. Tibiofibular fracture
- ANSWER: A
38. A 32-year-old male patient came to an emergency unit with complaints of severe leg pain and numbness of 24 hours duration. Long leg circular cast was applied for him two days back for his right leg closed tibio-fibular fracture. After thorough physical examination, compartment syndrome was diagnosed.
- What is the most appropriate initial management for this patient?
- A. Remove the cast
 - B. Give strong analgesic
 - C. Do fasciotomy
 - D. Refer to orthopedician
- ANSWER: A

39. A 25-year-old male patient came to an emergency unit after he sustained a fighting injury to his chest with a stick. He complains of left side severe chest pain and shortness of breathing, with difficulty speaking a full sentence. On physical examination, he has blood pressure of 80/50mmHg, pulse rate of 130beats/minute, respiratory rate of 36breaths/minute, and oxygen saturation of 75 percent. There is tracheal shift to the right and an absent air entry on his left side chest with hyper-resonant percussion note.

What is the most likely diagnosis for this patient?

- A. Haemo-pneumothorax
- B. Flail chest with multiple rib fractures
- C. Simple pneumothorax
- D. Tension pneumothorax

ANSWER: D

40. A 20-year-old male patient came to surgical OPD with a complaint of right elbow joint deformity. On physical examination, his right elbow joint was at full flexion but there is limitation during extension and redundant skin with old scar on the anterior surface of the joint was seen. He had a history of flame burn injury to the site six months back.

What is the most likely diagnosis of this patient?

- A. Post burn keloid
- B. Elbow joint ankylosis
- C. Post burn contracture
- D. Elbow joint dislocation

ANSWER: C

41. A 50-year-old male patient, who is a known hypertensive patient for many years, came to an emergency unit with complaints of insidious onset, severe crampy pain and numbness of his left leg of one week duration. On physical examination, his leg is cold, mottled and sensation is lost. Dorsalis pedis and posterior tibialis pulses are absent but the femoral pulse is palpable. He has no history of claudication.

What is the most likely diagnosis of this patient?

- A. Acute Limb ischemia
- B. Chronic limb ischemia
- C. Deep vein thrombosis
- D. Wet gangrene of the leg

ANSWER: A

42. A five-year-old male child was brought to an emergency unit with complaints of repeated cough, neck and throat pain, fever and shortness of breath of four days duration. On chest auscultation, there was coarse crepitation and wheeze over the right lower third of posterior lung field. Four days back, the family noticed that he had choking incident. On chest radiograph, there was haziness over right lower third of lung field.

What is the most likely diagnosis of this child?

- A. Hyperactive airway disease
- B. Basal pneumonia
- C. Epiglottitis
- D. Aspiration pneumonia

ANSWER: D

43. A 20-year-old primigravid woman came to the health center with a complaint of missed menses of one month duration. Associated with this, she also has a history of nausea and vomiting of one week duration. Vital signs at presentation: blood pressure of 110/60, pulse rate 110 beat/minute and respiratory rate 20 breath /minute.

What is the best initial investigation for this patient?

- A. Urine HCG
- B. PAP smear
- C. Abdominal x ray
- D. Abdominal ultrasound

ANSWER: A

44. A 26-year-old woman came to health center for antenatal care visit after eight months of amenorrhea. The working health officer evaluates the women and advises her on place of delivery, saving money and items needed for birth.

What is the most likely addressed antenatal care goal?

- A. Prevention of communicable diseases.
- B. Health promotion and disease prevention.
- C. Birth preparedness and complication readiness planning.
- D. Early detection and treatment of complications and existing diseases.

ANSWER: C

45. A 23-year-old gravida two Para one woman, who is amenorrhic for the past three months, presented with complaints of vaginal bleeding with passage of tissue and lower abdominal pain of one day duration. Vital sign at presentation: blood pressure of 70/40 mmHg, pulse rate of 120 beat/minute, respiratory rates of 20 /min and temperature 36.6 degrees centigrade. On vaginal examination; cervix three cm dilated and blood on examining finger. On investigation urine HCG is positive.

What is the best initial management for this patient?

- A. Transfuse cross-match blood
- B. Manual vacuum aspiration
- C. Fluid resuscitation
- D. Antibiotic

ANSWER: C

46. A 30-year-old gravida three Para two woman, who is amenorrhic for the past two months, presented with a complaint of spot vaginal bleeding followed by lower abdominal pain of 12 hours duration. Vital sign at presentation were blood pressure of 110/60 mmHg, pulse rate of 96 beat/minute, respiratory rate of 20/min and temperature 36.5 degree centigrade. On vaginal examination, cervix was closed and blood on examining finger. Urine HCG positive and obstetric ultrasound shows intra uterine pregnancy with positive cardiac activity.

What is the most likely diagnosis for this patient?

- A. Missed abortion
- B. Inevitable abortion
- C. Incomplete abortion
- D. Threatened abortion

ANSWER: D

47. A 25-year-old gravida four Para three woman presented to health center with a complaint of dark red vaginal bleeding associated with crampy lower abdominal pain of four hours duration. On investigation, obstetric ultrasound shows alive fetus with gestational age of 34 weeks with anterior placental, which is three cm above the level of internal cervical orifice.

What is the most likely diagnosis for this woman?

- A. Abruptio placentae
- B. Placenta accreta
- C. Placenta previa
- D. Vase previa

ANSWER: A

48. A 25-year-old primigravida woman, who is amenorrhic for the past seven months, came to health center with a complaint of gush of fluid per vagina of one day duration. Otherwise, she has no fever, abdominal pain and other co-morbidity. Vital signs are within normal range and on Leopold examination, she has 31 weeks sized gravid uterus. On speculum examination, there is pooling of fluid from cervix.

What is the most likely diagnosis for this woman?

- A. Preterm labor
- B. Threatened abortion
- C. Abruptio placenta
- D. Premature rupture of membrane

ANSWER: D

49. A 20-year-old primigravida woman, who is amenorrhic for the past nine months, came to health center with a complaint of pushing down pain of five hours duration. At presentation, vital signs were blood pressure of 120/70 mmHg, pulse rate of 120 beat/minute, respiratory rate of 24 /min and temperature 36.7 degree centigrade. On vaginal examination, cervix is four cm dilated with full effacement. She has two contractions within ten minutes which stayed from 35 to 40 seconds.

What is the most likely diagnosis for this patient?

- A. Latent first stage of labor
- B. Active first stage of labor
- C. Second stage of labor
- D. Third stage of labor

ANSWER: A

50. A 30-year-old gravida three and Para two woman, who is amenorrhic for the past nine months, came to health center with a complaint of pushing down pain of six hours duration. On vaginal examination, cervix was six cm dilated and membrane ruptured and clear. She has three contractions within ten minutes which stayed from 20 to 30 seconds. After four hours of evaluation, cervical dilation is eight cm with three contractions each stayed from 30 to 35 seconds.

What is the most appropriate next step in the management of this patient?

- A. Continue labor follow up
- B. Augment labor with Oxytocin
- C. Deliver with cesarean section
- D. Deliver with vacuum assisted delivery.

ANSWER: B

51. A 21-year-old gravida four, Para two and abortion one woman, who is amenorrhic for the past nine months, came to health center with a complaint of pushing down pain of seven hours duration. At presentation vital signs were stable. On vaginal examination, cervix six cm dilated, station zero and membrane ruptured and clear. She has two contractions within ten minutes. After four hours of stay at health center she gave birth to 3600 gram male alive neonate and the health officer is applying controlled cord traction to deliver the placenta.

What is the most likely stage of labor for this woman?

- A. Latent first stage of labor
- B. Active first stage of labor
- C. Third stage of labor
- D. Second stage of labor

ANSWER: C

52. A 25-year-old woman gave birth to a 4200 gram male alive neonate with APGAR score of eight and nine at first and fifth minutes of life respectively at health center after 13 hours of labor. Her vital signs were blood pressure of 120/70 mmHg, pulse rate of 110 beat/minute and respiratory rate of 20 breath/minute.

What is the next most appropriate management for this woman?

- A. Oxytocin ,uterine massage and misopristol
- B. Oxytocin, controlled cord traction and misopristol
- C. Oxytocin, controlled cord traction and uterine massage
- D. Uterine massage ,controlled cord traction and ergometrine

ANSWER: C

53. A 25-year-old gravida three and Para two woman presented to health center with a complaint of pushing down pain of four hours duration. At presentation, her vital signs were within normal range .On Leopold examination, she has 36 weeks sized gravid uterus and on vaginal examination; cervix fully dilated, station negative one and position left mentum transverse.

What is the most likely presentation of fetus in this case?

- A. Brow
- B. Face
- C. Breach
- D. Vertex

ANSWER: B

54. A 35-year-old gravida three and Para two woman, whose gestational age is 39+6 week, came to health center with complaints of pushing down pain and passage of fluid per vagina of seven hours duration. At presentation, vital signs were within normal range. FHB is 182 beat/minute and on vaginal examination, cervix is six cm dilated and there is pulsatile mass palpate at cervical opening alongside of the presenting part.

What is the most likely diagnosis for this case?

- A. Cord prolaps
- B. Footling breach
- C. Cord presentation
- D. Shoulder presentation

ANSWER: A

55. A 35-year-old para-three mother gave birth to a three-kilogram male alive neonate after 14 hours of onset of labor. One hour after delivery, Oxytocin ten IU was given, placenta delivered by controlled cord traction and uterine massage done. Two hour after delivery, the midwife noticed massive vaginal bleeding. On vaginal examination, there is bleeding from cervical opening; otherwise no vaginal or cervical lacerations. On abdominal examination, floppy hypotonic uterus. Vital signs were; blood pressure of 80/40 mmHg, pulse rate of 125 beats per minute, respiratory rate 24 breaths per minute, and temperature 36.7 degrees centigrade.

What is the most appropriate preventive measure for this patient?

- A. Active management of third stage of labor
- B. Oxytocin injection before fetal delivery
- C. Shorten labor by augmentation
- D. Cesarean delivery of the baby

ANSWER: A

56. A 24-year-old mother presented to a health center with complaints of fever, foul smelling vaginal discharge and lower abdominal pain of three days duration. She gave birth vaginally to a male baby two weeks back at home. Upon presentation, vital signs were blood pressure of 90/60 mmHg, pulse rate of 120 / min, respiratory rate of 20 breaths per minute, and temperature 39 degree centigrade. On vaginal examination, there is cervical motion tenderness.

What is the most appropriate management for this patient?

- A. Iv ampicillin and gentamycin
- B. Po ampicillin and metronidazole
- C. Iv ampicillin an Metronidazole
- D. PO cephalaxine and Metronidazole

ANSWER: A

57. A 20-year-old woman presented to health center with a complaint of absence of menses for the past four months. Before four months, her menses was regular which comes every 28 days. Six months back, myomectomy was done for large sub mucosal myoma.

What is the most likely diagnosis for this patient?

- A. Asherman syndrome
- B. Ovarian dysgenesis
- C. Premature ovarian failure
- D. Polycystic ovarian syndrome

ANSWER: A

58. A 20-year-old girl presented to health center with complaints of purulent, malodorous, thin vaginal discharge of four days duration associated with burning, pruritus, dysuria, frequency and dyspareunia. She has a history of multiple sexual contacts. Speculum examination shows "strawberry cervix".

What is the most likely diagnosis for this patient?

- A. Trichomoniasis
- B. Neisseria gonorrhoea
- C. Chlamydia trachomatis
- D. Bacterial vaginosis

ANSWER: A

59. A 20-year-old girl came to a health center after five days of having forceful unprotected sexual intercourse with unknown offender while she was on the way to her home.

What is the most appropriate emergency contraceptive for this girl?

- A. Copper intra uterine device
- B. Combine oral contraceptive
- C. Levonorgestrel tablet
- D. Mifepristol tablet

ANSWER: A

60. A 34-year-old woman presented to health center with a complaint of unable to conceive for the past one year and six months despite adequate frequency of sexual intercourse with her husband. Before her marriage, she had multiple sexual partners and treated for sexually transmitted diseases multiple times. Her menses come every 28 days regularly.

What is the most appropriate preventive measure for this patient?

- A. Avoid any sexual intercourse before marriage
- B. Avoid early marriage before 20 years old
- C. Give awareness about condom usage for females only
- D. Give awareness about safe sex to teenagers and adults

ANSWER: D

61. A 65-year-old Para five woman came to a health center with a complaint of postmenopausal vaginal bleeding for the past four months. Vital signs at presentation were within normal range. Transvaginal ultrasound shows endometrial thickness of 10mm.

What is the next appropriate management for this patient?

- A. Hysteroscopy
- B. Endometrial biopsy
- C. Papanicolaou smear
- D. Saline infusion sonography

ANSWER: B

62. A 70-year-old Para eight woman presented to a health center with complaints of pelvic pressure, heaviness in the vagina, and feeling of sitting on a ball of four months duration. Associated with this, she has also history of urinary urgency, frequency, incontinence and protrusion of mass per vagina of the same time duration.

What is the most likely diagnosis for this patient?

- A. Pelvic organ prolapse
- B. Nabotian cyst
- C. Cervical myoma
- D. Cervical cancer

ANSWER: A

63. A neonate born from a 23-year-old primigravida mother, at gestational age of 40 weeks plus 6 days through cesarean section with birth weight of 4600gm. The mother had regular and smooth ANC follow up.

What is the classification of the newborn based on the gestational age?

- A. Term
- B. Preterm
- C. Post term
- D. Post date

ANSWER: A

64. A well-feeding male neonate after birth, who was born at home by traditional birth attendant, was brought by his mother to a health center on the 3rd day after birth with complaints of crying excessively and developed progressive difficulty in sucking and swallowing, diminished movements and stiff body while the mother touches him.

What is the most likely diagnosis for this neonate?

- A. Early onset neonatal sepsis
- B. Congenital syphilis
- C. Neonatal tetanus
- D. Birth trauma

ANSWER: C

65. A six-year-old unvaccinated child presents with repetitive cough of two-week duration. The cough resulted in loss of his breath and he vomits after the cough since the onset of the disease. The complete blood count shows raised white blood cells with marked lymphocyte predominance and CXR is pending.

What is the most appropriate management for this patient?

- A. Start Anti TB despite CXR result
- B. Admit him and give Vitamin A
- C. Start age adjusted Azithromycine
- D. Put on intranasal oxygen and observe

ANSWER: C

66. A five-year-old male child was brought to a pediatric emergency room after he loses his consciousness. The mother states that he has watery diarrhea for the last four days at least five to six episodes per day. The mother added that she gave him coffee mixed with honey but it did not cease. On physical examination, the eyeballs are sunken, and the skin pinch goes back very slowly.

What is the most likely classification of dehydration for this case?

- A. No Dehydration
- B. Some dehydration
- C. Moderate dehydration
- D. Severe dehydration

ANSWER: D

67. A ten-year-old female child, who was on HAART for the last seven years, presented with complaints of dry cough, shortness of breath, and chest pain of four days duration. She also complains of a white lesion on her tongue. On physical examination, blood pressure and temperature are within normal ranges, pulse rate is 128 beat/minute, and respiratory rate is 28 /min. She is also diagnosed with severe acute malnutrition.

What is the most likely WHO clinical HIV/AIDS staging of this patient?

- A. Stage I
- B. Stage II
- C. Stage III
- D. Stage IV

ANSWER: D

68. A seven-year-old child presented with complaints of abdominal pain, loss of appetite and occasional diarrhea of ten-day duration. The child walks by barefoot while being out of school. On physical examination, the conjunctivae and the palms are pale. The CBC reveals hemoglobin of 6.2 gm/dl and MCV 58 percent.

What is the most likely diagnosis for this patient?

- A. Amebiasis
- B. Ascariasis
- C. Hookworm
- D. Giardiasis

ANSWER: C

69. A male child came to a hospital with complaints of bloody diarrhea, tinnitus, loss of appetite and weight loss and he was diagnosed with hookworm infestation. The complete blood count was done and result shows hemoglobin level of 4.1gm/dl.

What is the most effective management for this patient?

- A. Single dose albendazol is sufficient
- B. Albendazol and vitamin B12
- C. Albendazol and transfusion
- D. Mebendazole and oral folic acid

ANSWER: C

70. A seven-year-old child, who had a history of repeated pharyngitis, brought to a hospital with complaints of migratory type of joints pain and firm, painless, non-pruritic, mobile swellings on the elbow joints and the scalp. On physical examination, temperature is 36.9 degrees Celsius and the pulse rate is 122 breaths/min. There is systolic murmur at apex and most of the joints are tender and limited on range of motions. All laboratory investigations are pending. What is the most likely diagnosis for this child?
- A. Ischemic heart disease
 - B. Acute rheumatic fever
 - C. Infective endocarditis
 - D. Dilated cardiomyopathy
- ANSWER: B
71. A two-year-old child was brought to a health center in the middle of the night after he developed hoarseness of sound, stridor during agitation and a barking type of cough. He has runny nose and sneezing since the last one week. On examination, temperature of 37.9 degree centigrade, no other pertinent findings. What is the most likely diagnosis for this child?
- A. Bacterial trachitis
 - B. Viral croup
 - C. Epiglottitis
 - D. Bronchiolitis
- ANSWER: B
72. A three-year-old child was brought to a hospital with complaints of worsening of cough and shortness of breath of six-day duration. The mother visited nearby clinic for the above complaints and treated with unspecified syrup but no improvements. On physical examination, temperature is 38.9 degree Celsius and respiratory rate is 45 breath/minute. On chest examination, there is chest indrawing, use of accessory muscles and coarse crepitation on auscultation. What is the most likely diagnosis for this child?
- A. Common cold
 - B. Severe pneumonia
 - C. Pneumonia
 - D. No pneumonia
- ANSWER: B
73. A ten-year-old female child came to a health center with complaints of reddish urine color change and facial swelling of five days duration. She had impetigo one month ago which was relieved without treatment. On physical examination, her blood pressure is Stage II hypertension range for her age and there is bilateral pitting edema on the lower extremity. Urine analysis shows RBC 3+ but urine protein is negative and serum Creatinine level is 1.9 mg/dl. What is most likely diagnosis for this child?
- A. Hereditary nephritis
 - B. Minimal change disease
 - C. Post streptococcal glomerulonephritis
 - D. Systemic lupus erythematosus nephritis
- ANSWER: C

74. An eleven-year-old male child came to a health center with complaints of high-grade fever and neck stiffness of 10 days duration. There is a similar illness in the vicinity since the last two weeks. For the above complaints, he visited nearby clinic and got treated with oral antibiotics for seven days but there is no improvement. On physical examination, the temperature is 38.7 degree Celsius and meningeal signs are positive. CSF analysis and culture is pending.

What is the most appropriate step in the management of this patient?

- A. Reassure and continue the oral antibiotics
- B. Wait till the CSF analysis and culture arrived
- C. Start broad spectrum intravenous antibiotics
- D. Start dexamethasone and wait for CSF results

ANSWER: C

75. A mother was brought her two-year-old child for growth and monitoring follow up at health center. The health professional noticed paler of the palms and conjunctiva. The mother feeds him cow's milk since his four-month of age. The complete blood count shows hemoglobin of 7.6gm/dl.

What is the most likely explanation for his signs and symptoms?

- A. Folate deficiency anemia
- B. Iron deficiency anemia
- C. Vitamin B12 deficiency
- D. Anemia of dilution

ANSWER: B

76. A four-year-old child was brought by his mother after he developed bilateral leg swelling. Since the last one-month, he had intermittent watery diarrhea at least three episodes per day and poor intake of food of a three-week duration. For the above complaints, he got treated with unspecified oral medication and ORS, but no improvement. On physical examination, the vital signs are within the normal ranges and anthropometric measurement shows weight-for-height is less than -3 SD. He has bilateral grade 3 pitting edema.

What is the most likely classification of the malnutrition in this case?

- A. Moderate acute malnutrition
- B. Severe acute malnutrition
- C. Mild acute malnutrition
- D. Chronic malnutrition

ANSWER: B

77. An eleven-month-old infant was brought to a health center by his parents after having repeated respiratory tract infection in the last two-months and concerning bowing of the legs while he tries to walk. He spends the whole day with care giver at home since two-months of age. On physical examination, the cranial bones are soft and while examining the ribs, the infant feels like beads. Wrist x-ray shows cupping of the distal radius and ulna.

What is the most effective preventive method for this case?

- A. Universal administration of vitamin K at birth for all infant
- B. Adequate sun light exposure starting from two-weeks of birth
- C. Administering multivitamin starting from two-weeks of birth
- D. Administration of multivitamin for the mother during pregnancy

ANSWER: B

78. A mother was brought her child to a well-baby clinic. The child transfers the mother's phone hand to hand, stares on the mother when she took the phone from his hands and can sit without support but cannot walk.

What is the estimated age of the child for this case?

- A. 9-months
- B. 12-months
- C. 6-months
- D. 13-months

ANSWER: C

79. A six-year-old child was brought by pedestrians' after he sustained road traffic accident to his right leg while crossing the road. On physical examination, he is able to communicate, his vital signs; pulse rate is 122 beat/minute and blood pressure is within the normal range. Chest is clear with good air entry bilaterally. There is massive bleeding from his right thigh with signs of femoral fracture.

What is the most appropriate initial management for this patient?

- A. Transfuse with cross-matched blood
- B. Put on oxygen with face mask
- C. Apply pressure to control bleeding
- D. Catheterize the patient to monitor urine

ANSWER: C

80. A-five-year old child was brought to a hospital with complaints of dry cough and shortness of breathing of four-hour duration. He had a history of similar illness three-months ago which has been treated with salbutamol puff and discharged after improvement. On physical examination, temperature is 36.7 degree Celsius, pulse rate is 110/minute, respiratory rate of 45/minute. On chest examination, there is diffuse wheeze on the upper two-third of the lung fields bilaterally and rash over the extensor surfaces of the arms.

What is the most likely diagnosis for this child?

- A. Tracheobronchomalacia
- B. Foreign body aspiration
- C. Gastroesophageal reflux
- D. Bronchial asthma

ANSWER: D

81. A 20-year-old college student came to a clinic with a complaint of the right ear discomfort upon waking from a nap in the dormitory. On physical examination, he is in pain, and a live cockroach can be seen moving in the right external auditory canal.

What is the most appropriate initial step of management in this case?

- A. Remove the roach live with forceps
- B. Refer to ENT center
- C. Instill oil to kill the roach
- D. Ear irrigation

ANSWER: C

82. A 48-year-old female patient, who was on follow-up for type-2 diabetes mellitus, went to a clinic with complaints of having left ear itching and thin watery ear discharge of 5 days duration. She has no fever. Ear examination shows the left ear filled with black and white debris and minimal watery discharge.

What is the most likely diagnosis for this patient?

- A. Acute bacterial otitis externa
- B. External canal tumour
- C. Foreign body in the ear
- D. Otomycosis

ANSWER: D

83. A 20-year-old male patient came to an emergency unit with a complaint of nasal bleeding following a punch to the face by an assailant 30 minutes ago. On physical examination, he has no gross nasal deformity; he has profuse active bleeding from the right nostril.

What is the most appropriate first step of management in this case?

- A. Adrenaline soaked pack
- B. Refer to ENT
- C. Vaseline gauze pack
- D. Watchful waiting

ANSWER: C

84. A 40-year-old male patient who is on follow-up at the cardiac clinic and takes furosemide, metoprolol, warfarin, and enalapril, came with bilateral nasal bleeding of 2 hours duration.

Which one of the medications is most likely responsible for epistaxis in this case?

- A. Furosemide
- B. Warfarin
- C. Metoprolol
- D. Enalapril

ANSWER: B

85. A 17-year-old male patient came to a clinic with complaints of sore throat, pain during swallowing, fever, and myalgia of four days duration. On physical examination, he has bilaterally inflamed grade 2 palatine tonsils covered with exudates.

What is the definitive management for this patient?

- A. Analgesics
- B. IM dexamethasone
- C. PO amoxicillin
- D. IV fluids

ANSWER: C

86. A four-year-old female patient, on treatment for acute bacterial rhinitis, was taken to a clinic with progressively worsening snoring and dysphagia. She is febrile, tachycardic, and has symmetric grade 2 palatine tonsils and a left-sided posterior oropharyngeal wall bulge. Lateral neck x-ray shows widening of prevertebral soft tissue at the level of nasopharynx and oropharynx.

What is the most likely diagnosis for this patient?

- A. Prevertebral abscess
- B. Retropharyngeal abscess
- C. Adenoid hypertrophy
- D. Vallecular cyst

ANSWER: B

87. A fourth-year public health student asked his teacher about the Universal Numbering System and how to identify permanent maxillary first molar teeth.

What is the number that designates the stated tooth in this case?

- A. 3
- B. 4
- C. 5
- D. 6

ANSWER: A

88. A 45-year-old male visits a hospital for a dental checkup. He gave a clinical history of smoking of one pack of cigarettes per day for the past ten years, and his preference of diet is most of the time food rich in carbohydrate. On physical examination, the health officer recognized he has tooth decay with erosions.

What is the best preventive measure for his condition?

- A. Fill the eroded teeth with amalgam
- B. Use brace to protect from trauma
- C. Proper oral hygiene twice per day
- D. Use oral antibiotics to kill bacteria

ANSWER: C

89. A 29-year-old male patient came to dental clinic with complaints of teeth pain and easy bleeding of three days duration. Associated with this he has a history of fever and bad mouth smell. On examination, he has necrotic gingival ulcers with grayish debris and he has right sub mental lymphadenopathy.

What is the best management for residual condition this patient?

- A. Supra and sub gingival scaling
- B. Scaling of the affected teeth
- C. 2 percent Chlorhexidine mouth wash
- D. Antibacterial cleaning

ANSWER: A

90. A 37-year-old male patient presented to an emergency department after he sustained road traffic accident three hours ago while he was crossing a road. He has a history of bleeding through the mouth after the occasion. On dental evaluation, his lower incisor tooth totally comes out of its socket.

What is the most likely type of dental trauma he sustained?

- A. Extrusion
- B. Intrusion
- C. Avulsion
- D. Fracture of teeth

ANSWER: C

91. A 26-year-old female presented with complaints of three episodes of unexpected intense fear in one week and it reaches a peak within minutes. She has a history of palpitations, sensations of shortness of breath, chest pain, shaking of her hands multiple times in the last one year. Otherwise, she has no history of past psychiatric, medical or substance use problems.

What is the most likely diagnosis for this patient?

- A. Social Phobia
- B. Panic attack
- C. Panic disorder
- D. Agoraphobia

ANSWER: C

92. A 25-year-old male worried about dying by cholera after he watched the news of the cholera outbreak in a neighboring country. He became very careful about ensuring that the food he eats is safe and clean. He started to wash fruits, vegetables, and his hands at least five times before consuming. He spends lots of time in washing and cleaning. He avoids hanging out with his friends.

What is the most likely diagnosis for this case?

- A. Obsessive-compulsive disorder
- B. Illness anxiety disorder
- C. Delusion disorder
- D. Panic disorder

ANSWER: A

93. A 40-year-old single man presented with sleep disturbance, lack of appetite, fatigue, difficulties in making decisions, lack of interest in pleasurable activities, hopelessness and recurring thoughts of suicide for the last two months. No history of medical condition or substance use. What is the most likely diagnosis for this patient?

- A. Major Depressive Disorder
- B. Persistent Depressive Disorder
- C. Disruptive Mood Dysregulation Disorder
- D. Adjustment disorder

ANSWER: A

94. A 29-year-old woman who sustained a road traffic accident two months ago came to a hospital with complaints of sleep disturbance, nightmare, irritability and being easily annoyed. She avoids cars at any cost in her daily life.

What is the most likely diagnosis for this patient?

- A. Acute Stress Disorder
- B. Social Anxiety Disorder
- C. Adjustment Disorders
- D. Posttraumatic Stress Disorder

ANSWER: D

95. A 21-year-old unmarried female has difficulties participating in social activities. As she explained, she has trouble speaking in front of people; even eating food in public places makes her anxious. For this reason, she preferred to stay alone. Otherwise, she has no history of other medical illnesses or substance use.

What is the most likely diagnosis of her condition?

- A. Avoidant personality disorder
- B. Social Anxiety Disorder
- C. Generalized Anxiety Disorder
- D. Agoraphobia

ANSWER: B

96. A 24-year-old male on the third post-surgery day for generalized peritonitis became irritable and started to vomit, diarrhea, and sweating. Physical examination is unremarkable, except tremor of hands and tongue. His attendant reported that he is a social drinker. He is oriented to time, place, and person. No underlying medical condition has been reported.

What is the most likely cause of his current condition?

- A. Alcohol abuse
- B. Alcohol intoxication
- C. Alcohol dependency
- D. Alcohol withdrawal

ANSWER: D

97. A seven-year-old male child, who was hit by his friend with a stone two days back, was brought to an emergency unit with a painful right eye swelling. He has high grade fever and visual impairment. On physical examination, there is proptosis, tender, warm, red eye with periorbital edema of his right eye. CT scan shows opacification posterior to the orbital septum.

What is the most likely diagnosis of this child?

- A. Rhino-orbital mucormycosis
- B. Orbital cellulitis
- C. Preseptal cellulitis
- D. Orbital hematoma

ANSWER: B

98. In a rural area, there is high prevalence of blindness among pediatric age groups. During physical examination of some of the children, more than five hair follicles greater than 0.5 mm in diameter on tarsal conjunctivae, conjunctival scarring with corneal neo-vascularization and granulation tissue formation were found.

What is the most likely risk factor leading to blindness in these children?

- A. Lack of vitamin A supplementation
- B. Lack of topical antibiotic prophylaxis
- C. Lack of appropriate Vaccination
- D. Poor personal and environmental hygiene

ANSWER: D

99. A 40-year-old male patient, who works in leather factory, came to an emergency unit with a complaint of severe eye pain after an unknown chemical spilled into his eyes. On physical examination, there is first-degree burn of the periorbital skin, conjunctival chemosis, and hyperaemia.

What is the best initial treatment option for this patient?

- A. Refer to an ophthalmologist immediately
- B. Give antihistamine medication
- C. Copious irrigation with water and an antihistamine
- D. Copious irrigation with water and topical antibiotics

ANSWER: D

100. A 15-year-old male patient presented to a Hospital with complaints of reddish discoloration of eye associated with severe, sharp pain, blurred vision and photophobia in both his eyes of one week duration. Otherwise, he has no eye discharge or trauma. On physical examination, he has bilateral corneal hyperemia.

What is the most likely cause of the patient's condition?

- A. Conjunctivitis
- B. Keratitis
- C. Foreign body
- D. Allergic conjunctivitis

ANSWER: B

101. A nine-year-old male child was brought to a Hospital with a complaint of a skin lesion associated with broken-off hairs on his scalp of six weeks duration. He was treated by unspecified topical herbal medication, but the lesion did not improve. On physical examination, there are broken-off hairs and hyperkeratosis on his scalp. On KOH solution, long, narrow, septated, and branching hyphae were seen.

What is the most likely diagnosis for this case?

- A. Alopecia areata
- B. Seborrheic dermatitis
- C. Tinea capitis
- D. Lichen simplex chronicus

ANSWER: C

102. A six-year-old female child was brought to a Hospital with a complaint of broken -off hair and scalp lesions of two weeks duration. On physical examination, besides the scalp lesion, she also has slightly yellowish discoloration of her eyes. On laboratory investigation, ALT is increased to 536, AST is increased to 408 and other results were normal.
What is the most appropriate treatment for this child?
A. Griseofulvin 10 to 15 mg/kg/day for 6 to 8 weeks
B. Fluconazole, 6 mg/kg/day × 3 to 6 weeks
C. Terbinafine, 250 mg/day × 2 to 8 weeks
D. Ketoconazole two percent Shampoo 2 times/week
ANSWER: D
103. A six-year-old male child was brought to a Hospital with a complaint of skin lesions around his mouth. On physical examination, there is flesh coloured and centrally umbilicated skin lesion on the left side of his face close to his lips.
What is the most likely diagnosis of this patient?
A. Herpes labialis
B. Varicella zoster
C. Molluscum contagiosum
D. Measles
ANSWER: C
104. Multiple people are coming from a prison in the town with a similar complaint of pruritic skin lesions. Most of them complain severe, intense itching on their hands, axillae and genital areas. On physical examination, most of the prisoners have excoriation marks over the extensor surface of hands, axillae and groin areas.
What is the best prevention strategy to be applied in this case?
A. Providing Benzyl benzoate ten percent lotion
B. Applying Permethrin five percent lotion
C. Personal and environmental hygiene
D. Isolation of the affected prisoners
ANSWER: C
105. A 16-year-old male patient presented to OPD with a complaint of skin lesions on his face, neck and upper back of two months duration. On physical examination, there are diffuse comedos and papulo-pustular lesions mixed with nodules on his face, chest and back. He has no history of any illness or medication intake.
What is the most likely diagnosis of this patient?
A. Folliculitis
B. Acne
C. Tuberous sclerosis
D. Rosacea
ANSWER: B
106. A 27-year-old male patient presented to OPD with a complaint of skin lesions seen on his face and neck after shaving his beard. On physical examination, there are multiple papules and pustules on his face skin extending to the upper neck and jaw.
What is the most likely diagnosis of this patient?
A. Bullous Impetigo
B. Cellulitis
C. Pseudo folliculitis barbae
D. Deep folliculitis
ANSWER: C

107. A pregnant mother came to the health center with a complaint of an increased production of saliva after she became pregnant. The health officer conducted an assessment to recommend appropriate medication for her diagnosed problem.
What is the most likely nutritional problem the pregnant mother faced in this case?
- A. Pica
 - B. Ptyalin
 - C. Constipation
 - D. Vegetarianism
- ANSWER: B
108. A woman came from a Kebele where their common food is maize, and she complained of symptoms such as fatigue, poor appetite, weakness, mild digestive disturbance, anxiety, and irritability. After an appropriate examination and diagnosis, the health officer identified that the health problem is caused due to micronutrient deficiency.
What is the most likely micronutrient deficiency in this case?
- A. Vitamin A
 - B. Vitamin B
 - C. Vitamin C
 - D. Vitamin D
- ANSWER: C
109. A health officer working in a health center uses measurements of physical dimensions and gross composition of the human body at different age levels to identify malnourished mothers among antenatal care visitors of the health center.
What is the most likely nutritional assessment method to be used in this case?
- A. Clinical Method
 - B. Dietary Method
 - C. Biophysical Method
 - D. Anthropometric Method
- ANSWER: D
110. A health officer assigned to a health center conducted nutritional assessment surveillance by measuring the frequency and distribution of nutrition-related diseases using collected and available data from patients in that health center.
What is the most appropriate data collection method at the individual level?
- A. Data from feeding program admissions
 - B. Data from daily dietary intake
 - C. Data from mass screenings
 - D. Data reported from clinics
- ANSWER: B
111. A health officer planned to provide a health education session for mothers which refused to use long-term contraceptives. Before providing the health education session, the health officer prepared on the topic by conducting social assessment to address the real gap in the community.
What is the most likely behavioral model that is used by health officer in this case?
- A. Health Belief Model
 - B. Transtheoretical Model
 - C. Theory of Reasoned Action
 - D. PRECEDE-PROCEED Model
- ANSWER: D

112. A health extension worker who works in that Kebele asked a mother who delivered her second baby at home, why she did not attend delivery in the health centers in a similar manner while delivering the first baby. The mother said "During my first delivery, the health professionals abused me, so I do not want to go to the health center again."
What is the most likely factor that affected the behavior of the mother in this case?
A. Enabling
B. Reinforcing
C. Maintaining
D. Predisposing
ANSWER: D
113. The food and Drug Administration authority put high tax on cigarettes and restricted the sales of cigarettes in pieces by law. Due to unaffordability to buy a box of cigarettes, the number of young smokers decreased dramatically.
What is the most likely factor that affected the behavior of the young in this case?
A. Predisposing
B. Enabling
C. Reinforcing
D. Maintaining
ANSWER: B
114. Professionals from woreda education office conducted an inspection on the nearby school. During the inspection, they found the school location is not free from potential physical and chemical hazards. There are sources of noise and air pollution. Based on the findings, they decided to close the school.
What component of the school health is address in this case?
A. Hygiene
B. Sanitation
C. Safe water supply
D. Health related policy
ANSWER: D
115. A health team conducted inspection in a prison. During the inspection, the prisoners complained about infestation of flea. The health officer recommended that the prisoners should clean their class and expose their beddings to sun or fumigate to control vector infestation.
What is the most likely health problem the prisoners may experience in this case?
A. Scabies
B. Diarrhea
C. Tuberculosis
D. Relapsing fever
ANSWER: D
116. A person came to the health center with complaints of vomiting, headache, nausea and difficulty of breathing. The health officer asked the patient if there were combustible charcoal inside the house, the person responded, he used charcoal to heat his house.
What is the most likely cause of health problem in this case?
A. Radon gas
B. Sulfur dioxide
C. Carbon monoxide
D. Oxides of Nitrogen
ANSWER: C

117. A woman cleaned her compound and separated the wastes into different types. She collected decomposable wastes to use for production of compost, the non- decomposable one to another container for disposal.
What is the most likely waste management activity done by the woman in this case?
A. Disposal
B. Collection
C. Segregation
D. Minimization
ANSWER: C
118. A woman came to a health center with complaints of sweating, seizure, fever and headache. The health officer conducted a physical examination; then sent her for laboratory investigation. The laboratory investigation result shows that she was reactive for malaria.
What is the most likely water associated disease that affected the woman in this case?
A. Water borne
B. Water washed
C. Water based
D. Water related
ANSWER: D
119. A 45-year-old male patient came to a health center with complaints of mild abdominal pain, nausea, and change in appetite, weakness and weight loss. Associated to this, he also saw proglottidis in his faces. When the health officer took a history, the patient mentioned that the problem started after consumption of raw meat with his friends.
What is the most likely food borne disease a man affected in this case?
A. Ascaris
B. Taeniasis
C. Brucellosis
D. Trichinosis
ANSWER: B
120. An epidemic of typhus was reported from the prison. Based on the report, the woreda epidemic investigation team visited the prison. After conducting an investigation, the team reported that an epidemic arose due to vector infestation.
What is the most likely vector that causes the epidemic in this case?
A. Flies
B. Fleas
C. Louse
D. Bedbugs
ANSWER: C
121. A 46-year-old man presented to a health center with weak pulse, dizziness and dehydration. During history taking, a health officer found that a person is a staff at metal industry and had an exposure to high temperature for a relatively long time. The health officer also identified the man had an excessive loss of salt and moisture from his body due to sweating.
What is the most likely occupational hazard that affected the man in this case?
A. Physical hazard
B. Chemical hazard
C. Biological hazard
D. Mechanical hazard
ANSWER: A

122. A woman that gave birth recently came to a health center with complaints of headache, seizure and fever. The health officer conducted an assessment and found the microorganisms that got access to enter the mother uterus during delivery were highly multiplied. To manage the disease, the health officer prescribed appropriate antibiotics. When the woman went back to home, she refused to take medication. As a result, the illness became complicated and finally the woman died.

What is the most likely cause of the maternal death in this case?

- A. Infection
- B. Hemorrhage
- C. Obstructed Labor
- D. Hypertensive disorders

ANSWER: A

123. A couple found in a Kebele started living together after their wedding; the husband paid a lot of cattle for her family to get her as a wife. As a result he always ordered her to serve him; she used a local market trade to support their life. The husband does not want to work; in addition, he always tooks for money to drink and enjoy with his friends.

What is the most likely gender based violence the wife experienced in this case?

- A. Sexual
- B. Physical
- C. Economical
- D. Psychological

ANSWER: C

124. A 19-year-old rural girl came to her relative's hotel, which is located in a nearby town to have a work. During her stay they promised to give her a work and pay her a fair salary. Once upon a time, the owner of the hotel ordered her to have sex with one of the hotel customers. Even if the girl refused, the customer forcefully did it.

What is the most likely sexual and reproductive health problem she experienced in this case?

- A. Early marriage
- B. Commercial sex
- C. Sexual violence
- D. Unwanted pregnancy

ANSWER: C

125. The epidemiologist conducted surveillance using data from weekly report from health center to detect sudden changes in disease occurrence and distribution in order to ensure effective action to be taken to control the disease.

What is the most likely type of surveillance used by the epidemiologist?

- A. Active
- B. Passive
- C. Sentinel
- D. Progressive

ANSWER: B

126. The Ministry of Health applied an integrated disease surveillance system. The system facilitates collaboration between surveillance of focal points at health institution, woreda, regional and national levels. The surveillance system designs to use either weekly or immediate reports.

What is the most likely surveillance type applied in this case?

- A. Disease-specific surveillance
- B. Health facility-based surveillance
- C. Surveillance conducted at laboratories
- D. Designated or representative health facility surveillance

ANSWER: B

127. A Woreda Health office head gives direction to maternal and child health unit to develop an action plan. As part of the action plan, he prepared that "by the end of 2025, 90 percent of eligible children will be vaccinated against six target diseases in the Woreda".
What is the most likely action plan component recommended by the head in this case?
- A. Setting objectives
 - B. Designing the strategy
 - C. Selecting priority problem
 - D. Identifying obstacles and limitations
- ANSWER: A
128. A public health officer observes a person suffering from HIV/AIDS and he feels risk of infection for the community members and he assessed the community's experience on disease prevention. Finally, he gives a recommendation to continue the safe healthy practices.
What is the most appropriate method of health promotion the officer applied in this case?
- A. Develop personal skills
 - B. Reorient health services
 - C. Develop health policy
 - D. Strengthen community action
- ANSWER: D
129. A Woreda administrator wants to know the challenges of maternal health in a Kebele. He planned to identify the challenges by inviting around twelve mothers to tell him and talk to each other on all possible prevalent maternal challenges in the Kebele.
What is the most likely method of health promotion the administrator applies in this case?
- A. Club
 - B. Meeting
 - C. Demonstration
 - D. Group discussion
- ANSWER: D
130. A woman came to a health center to get advice on how she can destroy all forms of micro-organisms, including the most resistant spores or cysts from the water fetched from the river water at her home.
What is the most appropriate water treatment method to be applied in this case?
- A. Boiling
 - B. Filtration
 - C. Chlorination
 - D. Storage before use
- ANSWER: A
131. A person wants advice and recommendation from environmental health expert to construct a liquid waste storage to receive waste matter from kitchen and toilets. The expert visits the home compound and suggests digging hole in the ground compound and to construct using dry masonry to minimize water contamination and cover slab.
What is the most likely liquid waste management method suggested by the expert in this case?
- A. Soak pit
 - B. Cesspool
 - C. Septic tank
 - D. Seepage Pits
- ANSWER: B

132. The medical director of a hospital wants to design and construct a waste management system. The director requests the environmental health expert to give expertise suggestions to design and construct appropriate waste management method to be used to reduce quantity of waste as well as to destroyed the infectiousness of the waste.
What is the most likely waste management method to be used in this case?
- A. Incineration
 - B. Land filing
 - C. Autoclaving
 - D. Compaction
- ANSWER: A
133. According to the Woreda health office annual report, the occurrence of communicable disease in the woreda was high, and the health office decided to reduce specifically diarrhea by promoting personal hygiene.
What is the most likely personal hygiene practice that can be applied in this case?
- A. Face hygiene
 - B. Hand hygiene
 - C. Body hygiene
 - D. Clothe hygiene
- ANSWER: B
134. The mayor of the town organized and deployed team of experts from health bureau, electricity office and fire extinguisher agency to inspect big buildings in the city. The team visited the buildings and gave feedback to all building owners to have and put emergency first aid kit, electricity circuit box and fire extinguishers. The team members also put their contact cell phones to call when they face any emergency disasters.
What is the most likely disaster prevention measure taken by the mayor in this case?
- A. Disaster response
 - B. Disaster recovery
 - C. Disaster preparedness
 - D. Post disaster surveillance
- ANSWER: C
135. A mayor of the city requested the Public Health specialist to design disaster prevention measures in the City.
What is the first most likely step of the disaster prevention measure considered in this case?
- A. Delineation of the areas affected by the disaster
 - B. Identification of the disaster to be monitored
 - C. Monitoring and reporting by local health services
 - D. Rapid, gross detection of cases or presumptive cases
- ANSWER: B
136. A field epidemiologist sets a case definition such as sudden onset of fever greater than 40.5 degree centigrade, neck stiffness, puerperal rash, and laboratory investigation result to identify and manage an outbreak that occurred in the region. Based on the case definition, the epidemiologist sends a specimen and found it positive for the requested laboratory investigation.
What is the most appropriate case definition used as a control measure of an outbreak in this case?
- A. Possible case definition
 - B. Probable case definition
 - C. Suspected case definition
 - D. Confirmed case definition
- ANSWER: D

137. A Hospital medical director planned to decrease childhood morbidity due to vaccine-preventable diseases. To achieve this plan successfully, a medical director assigns a Public health specialist to design an immunization strategy that can be successfully implemented and give the immunization service every day in the Hospital.

What is the most likely immunization strategy used by the manager in this case?

- A. Mopping up immunization
- B. Routine immunization
- C. National immunization days
- D. Outbreak response immunization

ANSWER: B

138. A researcher wants to know the level of mental growth and the status of goiter in an elementary school. The researcher revealed that the children have low mental growth, and a significant number of children also have non-visible goiter.

What is the most likely cause of child morbidity in this case?

- A. Zinc deficiency
- B. Iron deficiency
- C. Iodine deficiency
- D. Calcium deficiency

ANSWER: C

139. A woreda maternal and child health team wants to develop a strategic plan, and the team identifies different cause of maternal mortality. However, due to resource and time shortage, the team agreed to prioritize only on the direct obstetric cause of maternal death.

What is the most likely direct obstetric cause that the team could prioritize in this case?

- A. Control of abortion
- B. Prevention of Anemia
- C. Treatment of Malnutrition
- D. Prevention of HIV/AIDS

ANSWER: A

140. A researcher wants to recommend for the policy makers on the appropriate strategy that helps to reduce maternal morbidity due to malnutrition. To answer the research question, he is interested to measure the current nutritional status of mothers.

What is the appropriate expected measurement that should be used in this case?

- A. Weight-change
- B. Weight-for-age
- C. Weight-for-height
- D. Weight to height ratio

ANSWER: C

141. A Health professional planned to give a School health service for High school students on sexual and reproductive health. He shared a lesson using a key lesson ' Young people with HIV and AIDS need care, support, and treatment with antiretroviral medicines (ARVs) according to protocols'. What is the most likely standard practice of sexual and reproductive health that he planned to implement?

- A. Diagnosis services
- B. Prevention services
- C. Risk reduction services
- D. Treatment and care services

ANSWER: D

142. A woreda administrator who saw a high burden of harmful traditional practices planned to set a strategy to reduce harmful traditional practices. He established teams working within the community and allocated essential resources for the teams. Finally, he implemented successful interventions among the residents inside the community.

What is the most likely strategy implemented to reduce harmful traditional practices in this case?

- A. Clinic-level strategy
- B. Macro level strategy
- C. Community level strategy
- D. Health sector level strategy

ANSWER: C

143. An internist working in a Hospital was planning to implement appropriate foodborne disease control mechanisms. In the implementation process, the internist wants to focus on the control mechanisms to be applied in the Hospital.

What is the most appropriate foodborne disease control mechanism applied by the internist?

- A. Case treatment
- B. Safe water supply
- C. Good personal hygiene
- D. Sanitary disposal of feces

ANSWER: A

144. A regional health bureau conducted a workshop with stakeholders to evaluate the impact of an intervention on reproductive health. A regional health bureau used the percentage of women who visited a health center at least once during their pregnancy as outcome measurement.

What is the most likely reproductive health indicator used in this case?

- A. Postnatal care coverage
- B. Antenatal care coverage
- C. Prevalence of infertility in women
- D. Availability of comprehensive essential obstetric care

ANSWER: B

145. A health promoter gave health education to individuals on how they tend to repeat behaviors that are rewarded.

What is the most appropriate health education principle applied in this case?

- A. Repetition
- B. Practice
- C. Knowledge
- D. Reinforcement

ANSWER: D

146. The municipality of a town decided to displace some of the population living around river basin area forcefully to another location to avoid the risk of disaster.

What is the most likely disaster the community might face in this case?

- A. Volcano
- B. Drought
- C. Flooding
- D. Earth quick

ANSWER: C

147. In district "A" there are 2 hospitals and 3 health centers which are functioning very well. However, the vaccination coverage in this district was reported to below 90 percent in 2000, 60 percent in 2001; and 15 percent in 2003; which could rather have been 100 percent. Based on this information, a researcher wants to investigate the potential barriers related to the vaccination coverage. By doing so, a researcher aspires to spot light new information on the existing body of knowledge, bridges current gaps, and also inform policy formulation.
What is the most appropriate title selection criterion that the researcher considered in this case?
- A. Relevance
 - B. Feasibility
 - C. Ethical acceptability
 - D. Applicability of possible results
- ANSWER: A
148. A wordea health office formed a group of researchers to assess the level of bed net distribution and magnitude of malaria in the worda. The research team tries to consider all kebeles of the worda, and selected households in the selected kebeles for interview and observation.
What is the most appropriate research type to be used for this case?
- A. Survey study
 - B. Correlation study
 - C. Case study
 - D. Intervention study
- ANSWER: A
149. A researcher planned to identify the determinants of pre-term birth among women who gave birth at a midwifery clinic. The researcher recruited 120 women with pre-term babies and 240 women with term babies. The researcher interviewed all the women about their interpregnancy interval, pregnancy and obstetric-related histories, and other potential risk factors.
What is the most appropriate study design used for this case?
- A. Cohort
 - B. Cross-sectional
 - C. Case control
 - D. Case series
- ANSWER: C
150. A clinician wants to conduct a study to determine the prevalence of intestinal parasites among schoolchildren in a selected primary school. To recruit individual students, the clinician plans to select two sections (section "A" and "B") among from the five sections in the school. Then, the clinician collected data from students in both sections "A" and "B".
What is the most likely sampling method a clinician used for this case?
- A. Simple random sampling
 - B. Systematic sampling
 - C. Cluster sampling
 - D. Stratified sampling
- ANSWER: C

151. A trial was conducted to study the effects of medicinal plant extracts on plasmodium species. A pharmacologist then designed to do the study on mice that were infected by the plasmodium species. He administered the plant extracts for five mice, and the other five mice would rather take the standard drug for the treatment of plasmodium species. He controlled the laboratory experiment, administration of the drugs, feeding of the mice, and follow up of the clinical progression of the infection.
What is the most appropriate research approach to be used in this case?
- A. Quazi experimental
 - B. Randomized controlled
 - C. Non randomized
 - D. Uncontrolled experiment
- ANSWER: B
152. A researcher initially determined a sample of 300 patients for the assessment of the knowledge level of cancer patients towards chemotherapy. However, the researcher decided to change the initial sample size by keeping all other factors constant. The researcher only changed the margin of error from 5 percent to 3 percent.
What will be the size of the recalculated sample for this case?
- A. The same as the initial sample
 - B. Larger than the initial sample
 - C. Less than the initial sample
 - D. The initial sample decreased by 3 percent
- ANSWER: B
153. A researcher wants to conduct a study to determine the prevalence of respiratory tract infection among under-five children in a district. He collected the data from mothers through history taking and explained the questions for the mothers when they became confused.
What is the most appropriate data collection tool to be used in this case?
- A. Questionnaires
 - B. Interviews
 - C. Schedules
 - D. Observation
- ANSWER: B
154. A public health officer planned to conduct a study to assess hand-washing practices to prevent cholera in rural district. The officer selected households systematically to conduct the study.
What is the most likely data collection method to be used in this case?
- A. Observation
 - B. In-depth interview
 - C. Focused group discussion
 - D. Face-to-face interview
- ANSWER: A
155. A public health officer wants to study the long-term impacts of a chronic disease management program on patients. The health officer aims to gather comprehensive, individual stories and detailed feedback on the experiences of individual health journeys.
What is the most appropriate data collection method to be used for this case?
- A. Observation
 - B. In-depth interview
 - C. Telephone interview
 - D. Using available information
- ANSWER: B

156. A midwifery professional wants to study the utilization of abortion care services in a health center. The midwifery professional plans to use interviews to collect the data from women of the reproductive age groups who are attending the maternity outpatient department. The midwifery professional then selected the women randomly for the interview and assured them that the information they provided would not be disclosed to anybody.
What is the most appropriate research ethical principle applied in this case?
- A. Confidentiality
 - B. Openness
 - C. Fairness
 - D. Honesty
- ANSWER: A
157. A nurse records the hemoglobin level measurements of eleven patients. Six of the patients have hemoglobin measurements of 11 g/dl each, and the rest of the patients have hemoglobin measurements of 12 g/dl each.
What is the median hemoglobin level of patients for this case?
- A. 11g/dl
 - B. 11.5g/dl
 - C. 12g/dl
 - D. 12.5g/dl
- ANSWER: A
158. A health officer records the total cholesterol measurements of fifteen patients. The total cholesterol measurements are 100 g/dl for each patient.
What is the most likely interquartile range value of these measurements?
- A. 0 g/dl
 - B. 50 g/dl
 - C. 75 g/dl
 - D. 100 g/dl
- ANSWER: A
159. A physician takes the hemoglobin measurement of 100 patients. The measurements spanned from 9 m/dl up to 14 g/dl, and the physician wants to know the crude measure of variation of hemoglobin levels of patients.
What is the best appropriate dispersion measure that fits for this case?
- A. Range
 - B. Interquartile range
 - C. Standard deviation
 - D. Variance
- ANSWER: A
160. A health officer planned to assess the weight, cholesterol level, and blood pressure distribution of type 2 diabetic patients in a chronic follow-up clinic. The health officer recorded weight, cholesterol level and blood pressure from 300 type 2 diabetic patients and tried to present data using an appropriate method of data presentation for the report.
What is the most appropriate data presentation method that fits this case?
- A. Histogram
 - B. Pie chart
 - C. Line graph
 - D. Bar chart
- ANSWER: A

161. In maternity hospitals of a city, there were 50 deaths of women due to pregnancy-related hypertension, 150 deaths during delivery and 55 deaths during post-partum care were reported within a single budget year. There are a total of 15000 live births in that specific year. What is the maternal mortality rate per 100, 000 live births in the hospital?

A. 170
B. 1700
C. 2500
D. 1333

ANSWER: B

162. A university launches a screening program for COVID-19 for new admission students every year. From the previous admissions, the university has 15000 students. In the 2023 academic year, a university received 5000 new students. Screening tests revealed that 300 students had COVID-19.

What is the estimated incidence measure of COVID-19?

A. 2 percent
B. 6 percent
C. 30 percent
D. 33 percent

ANSWER: B

163. A clinician conducted a study to assess the effects of physical exercise on weight reduction among adult patients. The clinician recruited and followed 100 adult patients who regularly do physical exercises, and 100 adult patients who do not do regular exercises for six months. At the end of six month, 75 of the adult patients who do regular physical exercises were having significant weight reduction; while only 10 adult patients who are not doing physical groups reduced weight at the end of six month.

What is the most appropriate calculated measure of association value for this case?

A. 0.1
B. 0.33
C. 0.75
D. 7.5

ANSWER: D

164. The incidence rate of disease A is 40 percent, and its prevalence rate was 15 percent in the period of one year. While, disease B have an incidence rate of 10 percent and its prevalence was 60 percent in one one-year duration.

What is the most likely appropriate explanation for this finding?

A. Diseases A is more sever disease than diseases B
B. There is a long incubation period for disease A than for disease B
C. There is a short incubation period for disease B than for disease A
D. Disease A has slow progression than disease B

ANSWER: A

165. A health center annual report indicates low achievement of organizational objectives against the plan. Hence, the manager investigates the root causes for low performance to incorporate appropriate interventions to improve organizational performance in the next year. What is the most appropriate management principle applied by the manager in this case?
- A. Division of labor
 - B. Learning from experience
 - C. Management by exception
 - D. Management by objectives
- ANSWER: B
166. A medical director visualizes the complex interrelationships in a workplace, understand how the various factors in particular situations fit together and interact with one another and sees the organization as a whole and relationships among its parts in the organization. What is the most likely managerial skill to be applied in this case?
- A. Technical skill
 - B. Diagnostic skill
 - C. Conceptual skill
 - D. Human relations skill
- ANSWER: C
167. A manager designs new strategy to persuade staff of an organization to work willingly and continuously to achieve extraordinary organizational objectives. Due to the new strategy, the performance of staff was improved and the organization became successful. What is the most appropriate managerial role applied in this case?
- A. Leader
 - B. Liaison
 - C. Monitor
 - D. Negotiator
- ANSWER: A
168. A health center receives complaints from clients about the services provided at the health center. The manager considers it a serious issue and discusses with management staff to decide how to deal with the problem appropriately. Then the manager communicates the decision of the management staff to external stakeholders to indicate their areas of responsibility. What is the most appropriate managerial role applied in this case?
- A. Liaison
 - B. Monitor
 - C. Disseminator
 - D. Spokesperson
- ANSWER: D
169. A health center introduced a new health program to respond to community needs. A human resource department of the health center attracts, selects and assigns qualified people to fill all positions in the health center to properly handle activities. What is the most appropriate management function to be used in this case?
- A. Staffing
 - B. Directing
 - C. Organizing
 - D. Planning
- ANSWER: A

170. A manager of a district hospital develops a plan that indicates detailed activities that will be achieved within the physical year. Moreover, he outlined the responsible unit for each task and the required amount of budget for each unit of the hospital.
What is the most appropriate type of plan prepared by the manager in this case?
- A. Strategic plan
 - B. Business plan
 - C. Long-term plan
 - D. Operational plan
- ANSWER: D
171. The head of a health center prepared an organizational plan by adopting different techniques, technologies, and procedures to facilitate the achievement of the health center's goals, objectives and targets.
What is the most appropriate step of planning used to prepare the health center plan in this case?
- A. Situational analysis
 - B. Designing strategies
 - C. Setting objectives and targets
 - D. Identifying potential obstacles and limitations
- ANSWER: B
172. In a district hospital, the planning team reviews annual organizational reports to identify the public health burden imposed by different health and health-related problems as a selection criterion during prioritization in the planning process.
What is the most appropriate prioritization criterion used in this case?
- A. Feasibility
 - B. Magnitude
 - C. Government concern
 - D. Community concern
- ANSWER: B
173. A manager at the district health office organizes a planning team to formulate a five-year strategic plan. The planning team applies SWOT analysis to review internal and external environment factors to achieve organizational objectives. In their review, they identified the availability of international partners' support, well-organized community associations, and strong involvement of communities for administrative actions.
What is the most appropriate SWOT analysis matrix is identified in this scenario?
- A. Threats
 - B. Strengths
 - C. Weaknesses
 - D. Opportunities
- ANSWER: D
174. A newly appointed manager at a health service organization decides to restructure organizational tasks. In the process of organizing, the manager categorizes the whole organizational work into individual jobs and allocates similar jobs into sections, assembles correlated sections into departments, and assigns head to each department to supervise and control the various activities of each department.
What is the most important building block of an organization are considered in this case?
- A. Delegation
 - B. Decentralization
 - C. Division of labour
 - D. Over all coordination to secure unity
- ANSWER: C

175. A manager of an organization assigns specific duties with corresponding authority to a subordinate in an organization and creates the obligation for the subordinate to perform the duties assigned. The manager allows the subordinate to handle tasks with extreme freedom, with minimal periodic checking.

What are the most appropriate characteristics of effective delegation used in this case?

- A. The manager provides balanced responsibility and authority
- B. The manager selects the appropriate person who can do the work
- C. The manager explains to others that some tasks are delegated to 'somebody.'
- D. The manager gives support as needed and follows the progress of the work

ANSWER: A

176. A human resource department report shows frequent entry and exits of organizational employees, and the instability of employees significantly affects organizational performance. The top management undertakes personnel administrative activities such as promotion, compensation, and other benefits to improve organizational success.

What is the most appropriate human resource management activity in this case?

- A. Orientation
- B. Recruitment
- C. Retaining employees
- D. Human resource planning

ANSWER: C

177. An organization systematically measures employees' achievement and compares their achievement with previously established standards to give feedback about the result, to locate the need for additional training and to identify candidates for promotion.

What is the most appropriate human resource management activity applied in this case?

- A. Selection
- B. Promotion
- C. Recruitment
- D. Performance appraisal

ANSWER: D

178. A client visits a health center to get a measles immunization for a child. However, a nurse at the immunization room declines to give the vaccine to the child and tells the family that it is wastage to give a measles vaccination for less than ten children at a time and to come back another day. But, the client reports the situation to the nurse's supervisor, and the supervisor instructs the nurse to vaccinate the child. After a brief discussion, the nurse agrees to give the vaccine to the child to resolve the conflict with the supervisor.

What is the most appropriate conflict management strategy applied in this case?

- A. Avoidance
- B. Competition
- C. Collaboration
- D. Accommodation

ANSWER: D

179. A manager of an organization tries to achieve his own job responsibilities and guide subordinates towards organizational goals. The manager regulates organizational activities in accordance with the vision and mission of his organization.

What is the most appropriate leadership characteristic described in this case?

- A. Focusing longer-term successes
- B. Focusing on results and analysis of failure
- C. Performing functions such as planning and controlling
- D. Commanding subordinates to achieve predetermined organizational objectives.

ANSWER: A

180. A manager of an organization closely supervises and controls subordinates for the handling of organizational tasks. Due to this, employees become irresponsible, show lack of interest, and try to find new jobs.
What is the most likely leadership style applied by the manager in this case?
- A. Autocratic
 - B. Democratic
 - C. Laissez-faire
 - D. Participatory
- ANSWER: A
181. A health service organization determines staff needs through human resources planning. The type and number of personnel needed in the organization currently and in the future is estimated with a human resource audit.
What is the most likely next step that needs to be applied in this case?
- A. Selection
 - B. Orientation
 - C. Recruitment
 - D. Performance appraisal
- ANSWER: C
182. The Ministry of Health of Ethiopia realizes the importance of agriculture, education, transport, and communication offices and invites these organizations during the identification of community health needs to take intervention that has an impact on improving the health of the population.
What is the most appropriate principle of primary health care applied in this case?
- A. Equity
 - B. Decentralization
 - C. Community participation
 - D. Inter-sectoral collaboration
- ANSWER: D
183. A health center gives emphasis on strengthening the involvement of the catchment population to determine their own needs and take appropriate actions, and actively involves in assessment of the situation, problem identification, and priority setting and making decisions, and sharing responsibility in the planning, implementing, monitoring and evaluation of programs.
What is the most likely principle of primary health care emphasized in this case?
- A. Equity
 - B. Decentralization
 - C. Community participation
 - D. Inter-sector collaboration
- ANSWER: C
184. During the health sector development plan IV, the health delivery system of Ethiopia was restructured into three tier system including health post, health centers and district hospitals.
What is the appropriate health tier system that had to be categorized in this case?
- A. Primary level
 - B. Secondary level
 - C. Referral level
 - D. Primarily health care units
- ANSWER: A

185. A Health Science College establishes a group consisting of students from different departments for a community health attachment. Students undergo a series of discussions related to the handling of tasks to make the group successful. After a brief discussion of their differences, the disagreements and the conflicts that arose are addressed, group unity emerged, and members established common goals, norms and ground rules.

What is the most appropriate stage of team development to be applied in this case?

- A. Forming
- B. Norming
- C. Storming
- D. Performing

ANSWER: B

186. A project manager regularly checks the handling of tasks to ensure that outputs are produced and distributed as planned. The project manager reacts immediately if a particular job is not being done. The rapid feedback allows the manager to fine-tune program operations and improve the organizational climate.

What is the most appropriate activity applied by the manager in this case?

- A. Monitoring
- B. Evaluation
- C. Supervision
- D. Controlling

ANSWER: A

187. A 50-year-old female patient came to a primary hospital with complaints of watery diarrhea and dry lips of two days duration. On physical examination and laboratory investigation finding, the patient was diagnosed as gastro enteritis that the health officer managed the patient with respect and empathy for wellbeing of the patient.

What is the most likely moral practice provided as professional in this case?

- A. Norm
- B. Value
- C. Culture
- D. Moral

ANSWER: B

188. The health officer on duty checked the health status of the patient of chronic diabetes and prescribed the monthly medication. Besides, he counseled on life style modification and strictly told the patient that complication can occur if the life style modifications are not performed.

What is the most likely ethical principle applied by the health officer in this case?

- A. Autonomy
- B. Non-maleficence
- C. Truth telling
- D. Beneficence

ANSWER: C

189. A 70-year-old hypertensive male patient has been admitted to the medical ward with a stroke. With one week of treatment, the patient started to improve. Later, after ten days, he started deteriorating and stayed in the hospital for three weeks. Lastly, health care providers and families got in conflicting ideas on whether continuing the treatment is actually helping or hurting the patient, prolonging his suffering.

What is the most likely ethical dilemma reflected in this case?

- A. Refusal of treatment dilemma.
- B. Disclosure truth-telling dilemma.
- C. End of life dilemma.
- D. Disagreement dilemma.

ANSWER: C

190. A health officer working in a primary hospital has seen a patient with cardiac case. The patient has planned to change her place of residence to other region and she needs to take her medical record with her. The health officer does not allow the patient to take her medical record. What is the most likely legal frame that the health officer breached in this case?
- A. Access to medical record.
 - B. Access to get information on medical record.
 - C. Patient's access to own record.
 - D. Access to get right information.
- ANSWER: C
191. A 32-year-old gravida III para II pregnant mother came to a health center with a complaint of pushing down sensation. On assessment, she is term and the cervix was dilated 4cm; otherwise, the membrane was intact. The progress of labor was good and she gave birth after three three-hour stay. While the health officer receiving, the fetus was slip and the scalp of the fetus got swollen. What is the most likely malpractice done by the health officer in this case?
- A. Medical error
 - B. Negligence
 - C. Loss of attention
 - D. Disobedient
- ANSWER: B
192. A 45-year-old male patient, amputated left lower leg below the knee, was discharged a month ago and came for a medico-legal request. The surgeon who performed the amputation procedure was available. The surgeon believed that medical reports must be truthful and secret. What is the most likely medico-legal issue in professional practice that should be focused in this case?
- A. Information should normally be disclosed.
 - B. The surgeon has a duty of care to the patient.
 - C. The surgeon has a duty of confidentiality.
 - D. The surgeon can report patient status easily.
- ANSWER: C
193. A woreda health office head came for supervision to the primary hospital and to give orientation on the development of dignified, respectful care behavior. The head emphasized on a variety of patients in terms of ethnicity, religion, belief, culture, language, age, gender, disability, mental status and social conditions shall be taken in to account when professionals assess, examine, diagnose and plan to manage any health condition. What is the most likely principle of respectful care behavior to be focused in this case?
- A. Communicate with individuals meaningfully.
 - B. Recognize the diversity and uniqueness of individuals.
 - C. Recognize that an individual's surroundings are important.
 - D. Value workplace culture that actively promotes respect.
- ANSWER: B
194. A 29-year-old female patient came to the health center with complaints of shortness of breath and a productive cough of one week duration. The health officer on duty assessed the patient by using history, physical examination and chest x-ray, and she was diagnosed with pneumonia. On interaction with the patient, the health officer was able to understand the patient's perspectives, internal experiences, and feelings without intensive emotional involvement. What is the most likely human-centered care showed by the health officer in this case?
- A. Sympathy
 - B. Empathy
 - C. Kindness
 - D. Humanity
- ANSWER: B

195. A 39-year-old female patient came to the outpatient department of the primary hospital with a complaint of lower abdominal pain of one month duration. The health officer assessed the patient and diagnosed as a Mass. Due to the scope of practice, the health officer needs to consult senior professionals: a surgeon and an internist; then through hot discussion, the patient's diagnosis was found to be an appendicular mass.

What is the most likely principle of the health care team that was seen among the professionals in this case?

- A. Respectful professionalism.
- B. Respectful cooperation.
- C. Respectful collaboration.
- D. Respectful compassionate.

ANSWER: C

196. A health officer working in a health center needs voluntary health services with charity organization in addition to his regular duty. The health officer understands that volunteerism is a service without financial gain and for the common good, and to recognize informal acts of volunteering, taken outside formal organizations. The more contemporary view is that people engage in volunteer work to achieve a positive result for an individual and a community.

What is the most likely principle of volunteerism the health officer understands in this case?

- A. Volunteering benefits the community and the volunteer.
- B. Volunteering is not a substitute for paid work and wage.
- C. Volunteering is always a matter of choice of the individual.
- D. Volunteering respects the rights, dignity and culture of others.

ANSWER: A

197. A 20-year-old female came to a health center with compliant of pain full lesion on the vulva five days duration. She has a history of unsafe sex. The health officer assessed her and from history and physical examination diagnosed as vaginal ulcer and treated with the right regimen for STI and counseled for safe sex. Finally, the health officer on counseling told to the patient, all information regarding the patient is confidential.

What is the most likely communication principle established by the health officer in this case?

- A. Trust
- B. Respect
- C. Dignity
- D. Serve

ANSWER: A

198. An 18-year-old female patient came to primary hospital's maternal and child care unit for antenatal follow up. The midwife on duty assessed the mother by taking her history and physical examination. From the findings of the history and physical examination, the mother was on twenty-four weeks of gestational age, and the midwife prescribed folic acid and counseled on nutrition and personal hygiene. During the communication, the midwife was giving authority to the mother for her own health.

What is the most likely communication skill the midwife showed to the mother in this case?

- A. Mother as active participant.
- B. Interdependence on decision making.
- C. Joint decision making.
- D. Empowerment of the mother in her own health.

ANSWER: D

199. A 50-year-old male patient with cerebral malaria was admitted to medical ward. He was given treatment for malaria and the patient developed hypoglycemia. After two weeks stay in the ward, no response for the treatment was observed; rather, it has been getting worse and worse. The patient communicated the nurse and planned to go against medical advice.

What is the most likely relationship that helps to make decision to keep the wellbeing of the patient?

- A. Mutual problem-solving relationships.
- B. Patient-centered versus physician.
- C. Open relationships on both parties.
- D. Two-way relationships.

ANSWER: B

200. 19-year-old female patient came to a health center with a complaint of a lesion on the vulva with itching sensation of five days duration; she has a history of unsafe sex. A health officer who is her neighborhood assessed her and, from history and physical examination diagnosed her for genital ulcer. The officer treated the patient with the right regimen and counseled on the safer sex behavior. The elder brother of the patient was who is a friend of the health officer, slapped his sister because she is infected with STI.

What was the fault of the health officer on the patient?

- A. Failed to maintain the highest standard of personal conduct and integrity.
- B. Failed to respect patient confidentiality on medical information.
- C. Failed to provide appropriate counseling services to the patient.
- D. Failed to maintain appropriate and effective communication.

ANSWER: B