

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

OPTOMETRY



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority and is essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, leadership and management, ethics, and Health Promotion as outlined in the Information Booklet 2025.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies, including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey—and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Optometry

1. A 75-year-old female patient came to an eye clinic with complaints of a reduction of vision both at distance and near for the last five years.
What is the right approach to take history from this patient?
 A. Increase voice and reduce speaking speed
 B. Put the patient distance from the examiner
 C. Inform guardians to stay out
 D. Use non-verbal communication
 ANSWER: A

2. A 35-year-old patient came to an ophthalmic clinic with a complaint of blurred vision after reading for two hours. On examination, both distance and near visual acuities are normal. The optometrist ordered an accommodation fatigue test to rule out the cause of vision reduction nearby.
What important history must be taken regarding the patient's symptoms?
 A. Manner of onset
 B. Diurnal variation
 C. Precipitating factors
 D. Laterality of the problem
 ANSWER: C

3. A patient came to an ophthalmic clinic with a complaint of blurring of vision with his spectacle prescribed a year back. An optometrist performed refraction and found a 0.50 dioptre increment from the current spectacle.
What is the most important factor that helps to conclude the refractive error increment in this case?
 A. The prescription for the current spectacle
 B. The power of the current spectacle
 C. Duration of spectacle use
 D. Purpose of prescription
 ANSWER: A

4. A 10-year-old child was brought to a paediatric optometry clinic with a complaint of midday outward turning of the right eye in the last six months. The deviation manifests intermittently without causing any visual reduction.
What is the most appropriate technique to measure the total amount of deviation in this case?
 A. Krimsky evaluation
 B. Hirschberg measurement
 C. Alternative prism cover test
 D. Modified Krimsky examination
 ANSWER: C

5. A 20-year-old female student, who has worn spectacles for distance correction, presented to an ophthalmic eye clinic for her annual appointment. The optometrist planned to do an initial measurement before proceeding to the new refraction procedure.
What is the most appropriate initial measurement the optometrist has to do?
 A. Distance between optical centers
 B. Power of current spectacle
 C. Thickness of lenses
 D. Pantoscopic angle
 ANSWER: B

6. A 17-year-old female diabetic patient came to an eye clinic with a complaint of a reduction of vision at near for the last month. Her visual acuity on both eyes was 6/6 and 6/18 at distance and near respectively.
What is the most recommended examination in this case?
A. Accommodation
B. Pupillary
C. Vergence
D. Motility
ANSWER: A
7. A 13-year-old patient has come to an eye clinic for a refractive error check-up. After ascertaining the ocular anomalies, an optometrist planned to perform static retinoscopy. During retinoscopy, the patient is looking in a different direction of gaze.
What is the most appropriate retinoscopy preparation in this case?
A. Prepare a semi-darkened room
B. Avoid working distance lens
C. Give distance attention holding the target
D. Being off-centre from the patient's visual axis
ANSWER: C
8. A 22-year-old patient came to an eye clinic with complaints of intermittent blurring of vision and headache. After taking a physical examination, the optometrist planned to estimate the accommodative response of the patient.
What is the best retinoscopy type that suits the above scenario?
A. Wet
B. Near
C. Static
D. Dynamic
ANSWER: D
9. A 26-year-old male patient came to an eye clinic with a complaint of painless reduction of distance vision. The patient has a far point of 10cm. During retinoscopy, the optometrist observed a dull retinoscopy reflex.
What is the best time-saving retinoscopy mirror effect in this case?
A. Plane
B. Convex
C. Concave
D. Divergent
ANSWER: C
10. A 39-year-old patient came to an eye clinic with a complaint of a reduction of distance vision over the last two years. During physical examination, the vision of patient became worse with a pinhole. Slit lamp examination also showed central corneal opacity. After physical examination, the optometrist planned to perform static retinoscopy to see the retinoscopic reflex.
What is the most appropriate retinoscopy technique to be applied in this case?
A. Using a working distance lens
B. Increasing working distance
C. Adding a high-power lens
D. Being Off-axis
ANSWER: D

11. A 30-year-old patient came to an eye clinic with a complaint of blurring of vision at both distance and near. An optometrist performed static retinoscopy and got the refractive power of $+3.00/-2.00 \times 90$. What is the movement of the retinoscopic reflex seen at a 90-degree axis?

A. Fastest with
 B. Slowest with
 C. Fastest against
 D. Slowest against

ANSWER: A

12. A 25-year-old male patient having hyperopic astigmatism came to a clinic for a refractive error check-up. He also had associated symptoms of bitemporal headache during reading. During static retinoscopy, an optometrist observed retinoscopic reflex having different speeds in the two principal meridians using the minus cylinder technique.

What is the most likely retinoscopic reflex that should be neutralized first in this scenario?

A. Fastest with
 B. Slowest with
 C. Fastest against
 D. Slowest against

ANSWER: B

13. A 33-year-old patient came to eye clinic with a complaint of blurring of vision in the right eye. After physical examination, the optometrist performed retinoscopy and found $+2.50\text{DS}$ on the right eye. The optometrist was performing retinoscopy at 66cm with the expected working distance lens put on.

What is the most appropriate refractive error in this case?

A. $+1.00\text{DS}$
 B. $+2.00\text{DS}$
 C. $+2.50\text{DS}$
 D. $+4.00\text{DS}$

ANSWER: C

14. A 62-year-old patient came to an eye clinic with a complaint of a reduction of distance vision in the left eye. During retinoscopy, the optometrist found that the refractive power of OS is $5.00/-1.00 \times 180$, with VA is CF at 5M. At the end of the subjective sphere check, the optometrist found a refractive power of OS is $-4.00/-1.00 \times 180$ with a VA is 6/60.

What is the most appropriate JCC power to start a cyl-check in this case?

A. $\pm 1.00\text{D}$
 B. $\pm 0.75\text{D}$
 C. $\pm 0.50\text{D}$
 D. $\pm 0.25\text{D}$

ANSWER: A

15. A 47-year-old male patient came to a clinic with a complaint of blurring of near vision on both eyes for one year. During examination, his remaining amplitude of accommodation is found to be 3.00D. The patient's habitual working distance is 33cm.

What is the estimated near add needed in this case?

A. $+2.50\text{DS}$
 B. $+2.00\text{DS}$
 C. $+1.50\text{DS}$
 D. $+0.50\text{DS}$

ANSWER: C

16. A 70-year-old patient came to an optometry clinic with a complaint of blurring of distance and near vision on both eyes for the last three years. At the end of subjective refraction, the optometrist found a refractive power of OD is $+1.00/-1.50 \times 60$ with VA is 6/9, OS is $+0.50\text{DS}$ with VA is 6/6. With this refraction result, the patient can read at near up to N-24.

What should be the starting add lens for near add determination in this case?

- A. $+2.75\text{DS}$
- B. $+2.25\text{DS}$
- C. $+2.00\text{DS}$
- D. $+1.50\text{DS}$

ANSWER: A

17. A 24-year-old patient came to an eye hospital with a complaint of blurring of distance vision on the right eye. During retinoscopy, an optometrist found a refractive power of OD is $+2.00\text{DS}$ with VA is 6/12. During the right eye subjective sphere check using -0.50DS lens, the patient says "the vision is the same with the lens".

What is the refractive error of the patient's right eye immediately after this step?

- A. $+2.50\text{DS}$
- B. $+2.00\text{DS}$
- C. $+1.50\text{DS}$
- D. $+1.00\text{DS}$

ANSWER: B

18. A 47-year-old patient came to an optometry medium clinic with a complaint of a reduction of vision at both distance and near. An optometrist could not see reflex during retinoscopy and planned to perform subjective refraction with stenopaic slit. During refraction with stenopaic slit, the patient reported that "the vision is clear with $+4.00\text{DS}$ when the direction of the slit is at 90 degree and the vision is also clear with $+3.00\text{DS}$ when the direction of the slit is at 180 degree".

What is the refractive power of this patient based on the above scenario?

- A. $+3.00/-2.00 \times 180$
- B. $+3.00/-1.00 \times 90$
- C. $+4.00/-1.00 \times 90$
- D. $+4.00/-1.00 \times 180$

ANSWER: C

19. A 17-year-old patient came to an optometry clinic with a complaint of blurring of vision on the right eye, both at distance and near. At the end of the subjective sphere check, the optometrist found a refractive power of OD is $-1.00/-2.00 \times 30$ with VA is 6/24. During subjective cyl-power check using $\pm 0.5\text{D JCC}$, the patient says "the target becomes clearer and rounder with $+0.50\text{D}$ side of JCC".

What is the refractive power of the patient's right eye just immediately after this step?

- A. $-0.50/-1.00 \times 30$
- B. $-0.75/-1.75 \times 30$
- C. $-1.00/-2.00 \times 30$
- D. $-1.25/-1.50 \times 30$

ANSWER: D

20. A 30-year-old male patient came to an optometry clinic with a complaint of blurred near vision after prolonged work for the last eight months. The patient had no significant refractive error. An optometrist suspects accommodative fatigue.

What is the most appropriate test to be performed to confirm the diagnosis in this case?

- A. Accommodative rock
- B. Dynamic retinoscopy
- C. Binocular indirect ophthalmoscopy
- D. Repeated amplitude of accommodation

ANSWER: D

21. A 27-year-old male patient came to an optometry clinic with a complaint of intermittent blurred near vision after prolonged near work for the last five months. During refraction, he had no significant refractive error. On accommodation assessment, an optometrist planned to assess the monocular accommodation facility first, using a ± 2.00 D flipper.

What is the normal level of monocular accommodation facility in this case?

- A. Within 3 cycles per minute
- B. Within 4 cycles per minute
- C. Within 5 cycles per minute
- D. Within 11 cycles per minute

ANSWER: D

22. A 20-year-old male patient came to an eye clinic with a complaint of intermittent blurring of near vision. After ruling out refractive error, an optometrist planned to measure the amplitude of accommodation by the Duane–Hoffstetter formula for probable amplitude of accommodation.

What is the average amplitude of accommodation in this case?

- A. 15.5D
- B. 14.5D
- C. 13.5D
- D. 12.5D

ANSWER: D

23. A 15-year-old patient came to a clinic with a complaint of blurring of near vision for six months. During refraction, the optometrist found a refractive error of $+6.00$ DS. The patient can read clearly at distances of 40 cm and farther away without spectacle correction.

What is the amplitude of accommodation in this case?

- A. 8.50D
- B. 8.00D
- C. 2.50D
- D. 2.00D

ANSWER: A

24. A 15-year-old hyperopic patient came to an ophthalmic clinic. During dry refraction, an optometrist found $+4.00$ DS. After doing wet refraction, the optometrist found $+6.00$ DS.

What is the most likely symptom the patient has been encountering in this case?

- A. Defective vision with no asthenopia
- B. Asthenopia symptoms with normal vision
- C. Defective vision with asthenopia symptoms
- D. Normal vision with foreign body sensation

ANSWER: C

25. A patient came to the refraction clinic with a complaint of distance vision reduction. The optometrist took all preliminary examinations and started doing retinoscopy. The optometrist performs retinoscopy without a working distance lens at 33 cm and neutralizes it with +2.00 DS.

What is the most likely diagnosis in this case?

- A. Low myopia
- B. Low hyperopia
- C. Moderate myopia
- D. Moderate hyperopia

ANSWER: A

26. A young glaucomatous patient came to an ophthalmic clinic with a complaint of distance visual reduction. The patient has been on pilocarpine eye drops for glaucoma treatment. After performing refraction, the optometrist found a refractive error.

What is the most likely type of refractive error in this case?

- A. Induced myopia
- B. Simple myopia
- C. Nocturnal myopia
- D. Pathological myopia

ANSWER: A

27. A 70-year-old patient had cataract surgery four months ago. The surgical incision was done temporally, and because of this, the horizontal cornea is flattened. During refraction, the optometrist found an induced astigmatism.

What is the most likely type of astigmatism that the patient has in this case?

- A. With-the-rule
- B. Against the rule
- C. Oblique
- D. Irregular

ANSWER: A

28. An ophthalmologist was performing cataract surgery. During the surgery, the ophthalmologist faced zonular disease. As an immediate action, the ophthalmologist put the calculated posterior chamber intraocular lens into the anterior chamber.

What is the most likely refractive error induced by the ophthalmologist in this case?

- A. Myopia
- B. Hyperopia
- C. Astigmatism
- D. Emmetropia

ANSWER: A

29. An adult patient came to an ophthalmic clinic with complaints of blurred vision and eye strain associated with near tasks. During assessment, an optometrist found low positive relative accommodation. The patient fails monocular and binocular accommodative facility with minus lenses. The patient's amplitude of accommodation decreased when measured repeatedly. What is the most likely diagnosis in this case?
- A. Accommodative paralysis
 - B. Accommodative fatigue
 - C. Accommodative infacility
 - D. Accommodative spasm
- ANSWER: B
30. A 44-year-old patient came to an ophthalmic clinic with a complaint of near vision difficulty that lasted for one year. When assessing the patient, the practitioner found that the patient's distance visual acuity is 6/6 and the amplitude of accommodation is normal for that age. But a near add is needed for this patient. What is the most likely diagnosis in this case?
- A. Presbyopia
 - B. Hyperopia
 - C. Accommodative fatigue
 - D. Accommodative insufficiency
- ANSWER: A
31. A 51-year-old patient came to an ophthalmic OPD with a complaint of reduced vision at night. His vision is normal during the daytime. After the best refraction, an optometrist found out that the patient is presbyopic and needs a near add. What is the most likely type of presbyopia that the patient has in this case?
- A. Nocturnal presbyopia
 - B. Functional presbyopia
 - C. Absolute presbyopia
 - D. Premature presbyopia
- ANSWER: A
32. A patient re-visits an ophthalmic clinic complaining of reduced vision at near that lasted for five days. The patient was on an eye drop medication for the last five days, and the visual reduction happened after starting the medication. When the amplitude of accommodation was measured, it was nearly zero. What is the most likely anomaly causing eye drops, the patient has been using?
- A. 1 percent Atropine
 - B. 2 percent Pilocarpine
 - C. 0.3 percent Ciprofloxacin
 - D. 0.1 percent Dexamethasone
- ANSWER: A

33. A four-year-old boy was brought to an eye clinic with Down syndrome. The boy's mother reported that the child has difficulty identifying objects while playing. After thorough examination, the Optometrist found -5.00 DS in both eyes.

For what condition is the boy at greatest risk in this case?

- A. Retinal detachment
- B. Amblyopia
- C. Cataract
- D. Macular hole

ANSWER: B

34. A 36-year-old man came to an eye clinic with a complaint of a reduction of vision at a nearby. His visual acuity was 6/9 and 6/12 at distance and near, respectively, in both the right and left eyes. On cycloplegic refraction, the optometrist found a refractive error of OD: +2.00 DS and OS; + 1.50 DS.

What is the best management option in this case?

- A. Prescribe add
- B. Correct the refractive error
- C. Prescribe vision therapy
- D. Prescribe a bifocal lens

ANSWER: B

35. A 12-year-old boy came to an eye clinic with a complaint of loss of vision in the left eye for two years. On assessment, the patient has a history of surgery following ocular injury in the left eye three years ago. After a detailed examination, the patient is diagnosed with aphakia. With +10.00 DS, the visual acuity improved from CF at 2m to 6/18.

What is the best management option for the patient in this case?

- A. Spectacle correction
- B. Prescribing contact lens
- C. Intraocular lens implantation
- D. Refractive surgery

ANSWER: C

36. A 47-year-old patient came to an eye clinic with a complaint of reduced vision at distance in both eyes for three years. After excluding all other ocular anomalies, an optometrist performed an accurate refraction and found -5.00 DS in the right eye and -2.00 DS in the left eye with a significant visual acuity improvement after subjective refraction.

What test must be performed first to determine the correct spectacle power for this patient?

- A. Binocular refraction
- B. Duochrome
- C. Worth forth dot
- D. Fixation disparity

ANSWER: A

37. A 48-year-old patient came to an eye clinic with a complaint of a reduction of vision at distance in both eyes for four years. After excluding all other ocular anomalies, an optometrist performed refraction and found -3.00 DS in both the right and left eyes with a significant visual acuity improvement after subjective refraction.

What is the most convenient management option for this patient?

- A. Contact lens
- B. Spectacle
- C. Refractive surgery
- D. Intraocular lens implantation

ANSWER: B

38. A 45-year-old man came to an eye clinic with a complaint of a reduction of vision both at distance and near. After excluding other ocular anomalies, the optometrist performed refraction and found OD: +2.00/-1.00x60 and OS: +1.50/-1.75x80 at the end of subjective refraction. The patient's visual acuity is improved from 6/12 to 6/6 at distance, both on right and left eyes, but still 6/12 at near.

What is the correct starting near add for the patient in this case?

- A. +1.00 DS
- B. +1.50 DS
- C. +2.00 DS
- D. +2.50 DS

ANSWER: B

39. A 45-year-old female patient presented with a complaint of a reduction of vision at nearby distances. She has visual acuity of 6/12 and 6/24 at distance and near, respectively both her right and left eyes. After subjective refraction at distance, the prescription reads OD: +1.50 DS with a visual acuity of 6/6 and OS: +1.75 DS with a visual acuity of 6/6.

What is the best management option for the patient in this case?

- A. Correct distance and near
- B. Provide near prescription
- C. Evaluate for accommodation anomaly
- D. Perform the vergence test

ANSWER: A

40. A 29-year-old patient came to an eye clinic with a complaint of blurring of near vision after sustained near work. Both distance and near visual acuities were 6/6. After detailed history and examination, the optometrist performed the amplitude of accommodation test for the fourth time, and that receded from 9.00 D at the first measurement to 4.00 D.

What is the most likely management option for this patient?

- A. Prism
- B. Plus add
- C. Vision therapy
- D. Atropinisation

ANSWER: C

41. A 70-year-old post-bilateral cataract surgery patient came to an eye clinic with a complaint of a reduction of near vision with a distance vision glass.

What is the starting near add power for this patient?

- A. +2.00 DS
- B. +2.50 DS
- C. +2.75 DS
- D. +3.00 DS

ANSWER: D

42. A 45-year-old man came to an eye clinic with complaints of a reduction of vision at distance and near. After excluding other ocular anomalies, the Optometrist performed refraction and found OD: +3.00 DS and OS: +1.00 DS at the end of subjective refraction. His visual acuity improved from 6/18 to 6/6 on the right eye and from 6/9 to 6/6 on the left eye at distance. His estimated add was +1.50 DS.

What is the most likely way to reduce vertical imbalance at the near visual point for this case?

- A. Prescribing a progressive addition lens
- B. Lowering the distance optical center
- C. Select the down curve segment
- D. Prescribe the executive segment

ANSWER: B

43. A 60-year-old patient came to an ophthalmic clinic with a complaint of visual reduction. An optometrist is taking a history.

What is the most relevant family ocular history that the examiner should take?

- A. Optic neuritis
- B. Retinitis
- C. Glaucoma
- D. Cataract

ANSWER: C

44. An optometrist is in an ophthalmic clinic measuring the intraocular pressure by using a Schiottz tonometer. During measuring the IOP of an old patient with 5.5g weight, the Schiottz scale reading needle shows 2.

What is the most appropriate action an optometrist should take in this situation?

- A. Re-measure by replacing the 5.5g weight with a 7.5g weight
- B. Re-measure by replacing the 5.5g weight with a 10g weight
- C. Re-measure by adding 7.5g weight over the 5.5g weight
- D. Re-measure by adding a 10g weight over the 5.5g weight

ANSWER: C

45. A two-month-old infant was brought to an ophthalmic clinic for evaluation of the eye. The optometrist wants to test the visual acuity of the infant.

What is the best visual acuity assessment method to be applied in this case?

- A. Cardiff acuity
- B. Landolt -C chart
- C. Fix and follow
- D. Allen symbols

ANSWER: C

46. A patient came to an ophthalmic OPD for a check-up. The optometrist starts evaluating the ocular adnexa of the patient one-by-one using slit lamp.

For what clinical sign does the optometrist evaluate the patient in this case?

- A. Seeing the lid margin to see if there is a lash
- B. Seeing the punctum, if there is an opening
- C. Seeing the meibomian gland, if there is pouting
- D. Seeing the conjunctiva, if there is a blood vessel

ANSWER: C

47. A patient came to an ophthalmic clinic with complaints of foreign body sensation and tearing. When an optometrist sees the patient's chart, the patient had been treated for blepharitis a year ago. The optometrist wants to evaluate the lid and lash.

What is the most likely technique to be used by the optometrist in this case?

- A. Diffuse illumination
- B. Indirect illumination
- C. Specular illumination
- D. Retro illumination

ANSWER: A

48. An optometrist is evaluating the patient's eye by using slit lamp. The optometrist wants to evaluate the anterior chamber angle by using the Van-Herrick estimation method.

What is the most appropriate illumination technique to be used by the optometrist in this case?

- A. Parallelepiped
- B. Optical
- C. Conical
- D. Specular

ANSWER: B

49. An adult patient came to an ophthalmic OPD with complaints of pain and redness in his eye that lasted for three days. After slit lamp evaluation, the optometrist diagnoses acute anterior uveitis.

What is the most likely finding observed by the optometrist in this case?

- A. Cells and flares in the anterior chamber
- B. Corneal epithelium laceration
- C. Nuclear lens opacity
- D. Misdirected lashes

ANSWER: A

50. A 56-year-old woman came to an optometry clinic with a complaint of inability to see objects to her side. She has had a constant headache that worsened in the last 10 days. Her vision is 6/18 and 6/24 in the right and left eyes, respectively. Her IOP is 39mmHg and 45mmHg in the right and left eyes, respectively.

What is the most likely finding in the optic nerve head of this patient?

- A. Blot haemorrhage could be seen on the disc
- B. The inferior quadrant of the rim may be thicker
- C. The vertical cup-to-disc ratio could be increased
- D. The blood vessels could be at the centre of the disc

ANSWER: C

51. An 18-year-old patient came to an ophthalmic clinic with the complaints of a gradually decreasing night vision and loss of peripheral vision in both eyes since childhood.

What is the most likely finding at the retina of the eyes for this case?

- A. Pigment is dispersed throughout the peripheral retina
- B. Haemorrhages throughout the peripheral retina
- C. Drusen is dispersed throughout the peripheral retina
- D. Narrowing of veins throughout the retina

ANSWER: A

52. A driver complaining of difficulty seeing the red colours of a traffic light came to an optometry clinic. Based on the colour vision assessment, the driver was diagnosed as protanomalous trichromatic.

What is the most likely result from the colour vision assessment in this case?

- A. The patient will see everything visible in the first plate of the Ishara test plate
- B. The patient will report the blue colours of the Ishara test plate as a different colour
- C. The patient will report the green colours of the Ishara test plate as a different colour
- D. The patient will report the red colours of the Ishara test plate as a different colour

ANSWER: D

53. A glaucomatous patient came to an ophthalmic clinic for a regular six-month follow-up. The optometrist wants to check if there is a subtle visual field change within the last six months.

What is the best visual field assessment type preferred by the optometrist in this case?

- A. Automated Static Perimetry
- B. Automated Kinetic Perimetry
- C. Manual Kinetic Perimetry
- D. Manual Tangent Screen

ANSWER: A

54. A 33-year-old man came to the hospital with a complaint of a progressively increasing painless swelling over the upper eyelid for two months. The same swelling had occurred a year before on the lower eyelid, and it was resolved spontaneously. There is a focal swelling of the upper eyelid and obstruction of the meibomian glands.

What is the most likely diagnosis in this case?

- A. Chalazion
- B. Eyelid Neoplasm
- C. Internal Hordeolum
- D. External Hordeolum

ANSWER: A

55. A 45-year-old woman from a rural area came to a hospital with a complaint of foreign body sensation in both eyes. She has a history of lash epilation in both eyes from the upper eyelid. Physical examination showed that there are misdirected lashes rubbing the cornea because the upper eyelid is turned inward.

What is the most likely diagnosis in this case?

- A. Entropion
- B. Ectropion
- C. Epiblepharon
- D. Euryblepharon

ANSWER: A

56. A 30-year-old patient came to an eye clinic with the complaints of burning, foreign body sensation, moderately decreased vision, and excessive tearing. These symptoms are often exacerbated by smoke, wind, low humidity, and prolonged use of the eye. Physical examination showed that an irregular tear meniscus was seen at the inferior eyelid margin.

What is the most likely diagnosis in this case?

- A. Superficial punctate keratopathy
- B. Computer vision syndrome
- C. Dry eye syndrome
- D. Blepharitis

ANSWER: C

57. A 35-year-old woman came to an optometry clinic with complaints of foreign body sensation in the eyes, intermittent lacrimation, stickiness of the eyelids, and mucoid discharge. On examination, there were congested conjunctival blood vessels, follicles, papillary reactions, and scar on the upper palpebral conjunctival and small circular pitted scars around the limbus.

What is the most likely cause of the infection in this case?

- A. *Klebsiella pneumoniae*
- B. *Staphylococcus aureus*
- C. *Escherichia coli*
- D. Chlamydial

ANSWER: D

58. A 50-year-old man came to an optometry clinic with complaints of pain, reduced vision, photophobia, redness, tearing, discharge, and foreign body sensation. He had a history of trauma from a tree branch 15 days before. On slit-lamp examination, he has conjunctival injection, greyish-white stromal infiltrates, corneal ulcer, and anterior chamber reaction.

What is the most likely diagnosis in this case?

- A. *Acanthamoeba* Keratitis
- B. Bacterial Keratitis
- C. Fungal Keratitis
- D. Viral Keratitis

ANSWER: C

59. A 16-year-old boy came to an eye clinic with complaints of gradual blurred vision and whitening of the central eye. On physical examination, the central zone of corneal stroma and epithelial oedema, ring-shaped deep stromal haze, and decreased corneal sensations were observed. The IOP was raised to 26 mmHg.

What is the most likely diagnosis in this case?

- A. Necrotizing stromal keratitis
- B. Neurotrophic ulceration
- C. Disciform keratitis
- D. Epithelial keratitis

ANSWER: C

60. A 56-year-old male patient came to a hospital with complaints of a shadow moving over the peripheral field of vision while sitting, and a shadow on the central vision while he was in a supine position for five days. There was mild RAPD. The underlying choroid was intact.

What is the most likely diagnosis in this case?

- A. Rhegmatogenous Retinal Detachment
- B. Tractional Choroidal Detachment
- C. Exudative Retinal Detachment
- D. Tractional Retinal Detachment

ANSWER: C

61. A 28-year-old male patient came to a hospital with a complaint of gradual reduction of vision at distance and near in both eyes during the night since childhood. On examination, he has vitreous cells, dark pigment in the peripheral retina around the blood vessels, narrowing of arterioles, and waxy optic disc pallor. In gross perimetry ring scotoma was detected.

What is the most likely diagnosis in this case?

- A. Retinoschisis
- B. Retinitis pigmentosa
- C. Pathological myopia
- D. Age-related macular degeneration

ANSWER: B

62. A 25-year-old female patient came to an eye clinic with a complaint of painless, moderately decreased vision in the right eye for a week. She has orbital pain that worsens with eye movement. On examination, she has relative afferent pupillary defect of the right eye, decreased colour vision, and flame-shaped haemorrhages around the disc margin.

What is the most likely diagnosis for this patient?

- A. Optic Atrophy
- B. Optic Neuritis
- C. Papilledema
- D. Retinoblastoma

ANSWER: B

63. A 65-year-old male patient came to an eye clinic with complaints of unilateral severe pain, decreased vision, tearing, and photophobia. His vision is light perception, and the IOP is 45mmHg. On examination, he has conjunctival injection, hyper-mature cataract, and a white material in the anterior chamber and on the anterior lens capsule. The anterior chamber angle was wide open.

What is the most likely diagnosis for this patient?

- A. Phacolytic glaucoma
- B. Lens particle glaucoma
- C. Phacomorphic glaucoma
- D. Phacoanaphylaxis

ANSWER: A

64. A 35-year-old sportsman, who had a boxing game an hour before, came to an eye clinic with a complaint of loss of vision in the right eye after having feast injury. His vision was hand motion. Both of his eyes were healthy, despite mild redness in the right eye.

What is the most likely diagnosis for this patient?

- A. Vitreous detachment
- B. Traumatic cataract
- C. Lens subluxation
- D. Retinal detachment

ANSWER: D

65. A 30-year-old patient, who had a falling injury on his face one day before, came to an eye clinic with a complaint of painless reduction of vision in the right eye. The corneoscleral wall of the globe was intact. There is mild ciliary injection and an intact pupillary reaction.

What is the most likely finding of the affected eye in this scenario?

- A. Corneal tear
- B. Traumatic cataract
- C. Iris prolapse
- D. Vitreous haemorrhage

ANSWER: D

66. A 65-year-old male patient came to an eye clinic with a complaint of painless, gradual reduction of vision in the right eye for five years. He had a history of trauma to the right eye 20 years ago. On examination, he has a clear cornea, clear lens, iris sphincter tear, RAPD, pale excavated disc, uneven iris insertion, and absence of iris processes.

What is the most likely diagnosis for this patient?

- A. Angle recession glaucoma
- B. Primary open angle glaucoma
- C. Endophthalmitis
- D. Panophthalmitis

ANSWER: A

67. A 40-year-old female patient came to an eye clinic with a month-long complaint of eye discharge associated with burning, itching, and foreign-body sensation in both eyes, particularly upon awakening, but improves during the day. Eye examination revealed that there are hard scales, small ulcers, loss of lash on the anterior eyelid margins, and hyperemic bulbar conjunctiva.

What is the first line of treatment in this case?

- A. Dexamethasone 0.1 percent eye drops QID
- B. Chloramphenicol 0.5 percent eye drops QID
- C. Cromolyn sodium 2 percent eye drops TID
- D. Gentamycin 0.3 percent eye drops QID

ANSWER: B

68. A two-year-old child was brought to an eye clinic with a complaint of excessive tearing on one eye since birth. After examinations, all external eye structures looked well and open punctum. There is no eye discharge or ocular inflammation.

What is the definitive treatment for this child?

- A. Massaging of the lacrimal sac by hydrostatic pressure.
- B. Probing of the canalicular system and nasolacrimal duct
- C. Intubation with Silastic tubing of the lacrimal duct
- D. Referring the patient for a surgical procedure

ANSWER: B

69. A 30-year-old man came to a hospital with complaints of redness and watery discharge on both eyes. He has a history of a recent upper respiratory tract infection for the last three days. On examination, his vision was 6/6 in both eyes.

What is the most appropriate treatment option in this case?

- A. Dexamethasone 0.1 percent eye drops QID
- B. Lid hygiene with warm compress
- C. Reassure that it can resolve by itself
- D. Gentamycin 0.3 percent eye drops QID

ANSWER: C

70. A 10-year-old girl came to an eye clinic with a complaint of blurred vision associated with foreign-body sensation, light sensitivity, and pain in the right eye for two weeks. The visual acuity was 6/60 for the right eye, and she had dendritic epithelial ulcers stained with rose Bengal and reduced corneal sensitivity.

What is the most appropriate treatment option in this case?

- A. Acyclovir three percent eye ointment, five times daily
- B. Chloramphenicol one percent eye ointment BID
- C. Erythromycin five percent eye ointment TID
- D. Prednisolone one percent eye ointment QID

ANSWER: A

71. A 60-year-old female patient came to a hospital for an eye check-up. She had type II diabetes mellitus for the last 10 years. On dilated fundus examination, she had microaneurysms, dot-and-blot intraretinal haemorrhages, and hard exudates in both eyes.

What is the most important treatment recommended in this case?

- A. Intensive glycemic control
- B. Anti-VEGF drugs
- C. Corticosteroids
- D. Laser photocoagulation

ANSWER: A

72. A 20-year-old female patient came to an eye clinic with a complaint of a reduction of vision in the left eye associated with ocular pain with eye movement since last week. On examination, the visual acuity was 6/6 for the right eye and counting fingers at two meters for the left eye. An optometrist found relative afferent pupillary defect, poor colour vision, and swollen optic disc in her left eye.

What is the most appropriate management to be done by the optometrist in this case?

- A. Refer to a neuro-ophthalmologist
- B. Prescribe Oral Prednisolone
- C. Offer intravenous steroid therapy
- D. Observation and follow-up

ANSWER: A

73. A 45-year-old male patient came to an eye clinic with a complaint of a gradual reduction of vision in his left eye for the last two years. On eye examination, he had a relative afferent pupillary defect on the left eye. Intraocular pressure, vertical cup to disc ratio, and visual acuity were 19 mmHg and 29mmHg, 0.4 and 0.8, 6/6 and 6/12 for the right and left eye, respectively.

What is the best drug of choice for treatment in this case?

- A. Timolol 0.5 percent eye drops 2 times/day
- B. Betaxolol 0.25 percent eye drops 2 times/day
- C. Brimonidine 0.2 percent eye drops 2 times/day
- D. Dorzolamide 2 percent eye drops 2 times/day

ANSWER: A

74. A 40-year-old male came to a hospital with an injury to his left eye from a car accident within one day. On physical examination, the visual acuity was no light perception; he had severe scleral tear and lacerated cornea with uveal prolapse on the injured eye.

What is the most appropriate management option for the patient?

- A. Enucleation
- B. Evisceration
- C. Oral antibiotics
- D. Exenteration

ANSWER: A

75. A woman came to a hospital with a complaint of foreign body sensation due to an injury on her left eye with her child's finger two hours back. She also has associated photophobia and tearing. On examination, she had hyperaemic conjunctiva and peripheral corneal abrasion stained with fluorescein.

What is the most appropriate management option in this case?

- A. Apply antibiotic eye ointments
- B. Apply steroid eye drops
- C. Bandage contact lens therapy
- D. Apply a potent cycloplegic agent

ANSWER: A

76. A man, who is a football player, came to an eye clinic with a fist injury to his left eye one day ago. He had associated symptoms of blurred vision and pain. On eye examination, there was a collection of blood in the anterior chamber, congested conjunctival vessels, but intact eyeball.

What is the most appropriate primary management in this case?

- A. Bed rest with elevated position and limit physical activities
- B. Exercise and encourage strenuous activity, bending, or heavy lifting
- C. Patch the involved eye at all times to prevent sudden rebreeding
- D. Admission and use of anticoagulant medications

ANSWER: A

77. An optometrist examined a patient in an ophthalmic clinic. During examination, the optometrist suspects glaucoma and wanted to assess the thickness of the cornea with an instrument attached to the slit lamp biomicroscope.

What is the most appropriate instrument to use in this scenario?

- A. Goldmann tonometer
- B. Pachymeter
- C. Hruby lens
- D. Gonio-lens

ANSWER: B

78. An optometrist has conducted a posterior segment evaluation with a direct ophthalmoscope on a patient with age-related macular degeneration. During the process, the optometrist turned the power on well, but the light was not shined into the patient's eye.

Which part of the direct ophthalmoscope is most likely malfunctioning in this case?

- A. The battery
- B. The bulb
- C. The pad
- D. The viewing aperture

ANSWER: B

79. The optometrist wants to use a slit lamp biomicroscope to investigate the vitreous, the retina and other posterior segment structures of the eye.

What is the most appropriate supplementary lens used in this case?

- A. 78D Volk lens
- B. Hruby lens
- C. 20D condensing lens
- D. Gonio-lens

ANSWER: A

80. A 40-year-old male came to an eye clinic for a check-up. He has a family history of glaucoma. In order to examine his visual field, a cyclic grating pattern target has been used. Then, the reliability test was done, and the result showed a normal visual field.

What type of perimetric procedure is most likely to be used in this case?

- A. Kinetic perimetry
- B. Static perimetry
- C. Gross perimetry
- D. Confrontation perimetry

ANSWER: A

81. A retinal sub-specialist put an indirect ophthalmoscope on the head to examine the peripheral retina and tried to adjust the binocularity of the view by targeting own finger. After that, the specialist brought a lens with a white mark around to use along with the ophthalmoscope.

What is the most likely dioptric power of the lens in this case?

- A. +20 D
- B. +60 D
- C. +78 D
- D. +90D

ANSWER: A

82. A 25-year-old male, who has a history of an ocular problem and was treated a month before, came to a hospital. In his medical record chart, there was a picture that showed a color-coded ophthalmoscopic result where the inferior portion of the picture was filled with a blue color.

What is the most likely ocular condition in this case?

- A. Retinal haemorrhage
- B. Retinal opercula
- C. Retinoblastoma
- D. Retinal detachment

ANSWER: D

83. A 55-year-old male patient has been on follow-up in an ophthalmic clinic for an immature cataract. Upon his follow up, the clarity of the crystalline lens was assessed by using the slit lamp biomicroscope in a wide illumination beam and decreased angle between the observing system and illumination bar. As a result, different parts of the crystalline lens became visible.

What type of illumination system is most likely to be used in this case?

- A. Optic section
- B. Parallelepiped
- C. Indirect beam
- D. Conical beam

ANSWER: B

84. Newly admitted optometry students are screened for different visual function tests. One of the tests done is a color vision assessment by using different kinds of plates. The plates have different types of numbers and colors.

What is the most appropriate distance between the plate and the students in this case?

- A. 25 cm
- B. 50 cm
- C. 75 cm
- D. 100 cm

ANSWER: C

85. A seven-year-old child came to an eye clinic with complaints of eye pain and discomfort. On assessment, ocular pain was examined, and there was no positive finding. To rule out the tear secretion abnormality, a 34mm strip was inserted in the lower canthal area, and the size of the wet area was measured to assess the quality of tear secretion.

What is the most appropriate waiting time to finish the procedure in this case?

- A. 3 minutes
- B. 5 minutes
- C. 10 minutes
- D. 15 minutes

ANSWER: B

86. A Jones dye test was performed on a patient suspected with dry eye, as there was no dye appearing on the sterile swab. In the initial phase, there was a significant amount of fluorescein in the fornix. However, the punctum did not allow it to pass.

What is the most appropriate next procedure in this case?

- A. Dilation
- B. Irrigation
- C. Probing
- D. Palpating

ANSWER: A

87. A 25-year-old man came to an eye clinic and got checked for color vision abnormalities using Ishihara color plates. During the test, the person was able to detect the three-color types and discriminated most of the plates; however, he wrongly responded to plates number 2, 5, and 6.

What is the most likely type of color vision defect this patient has?

- A. Protanopia
- B. Protanomaly
- C. Deutanopia
- D. Deutanomaly

ANSWER: B

88. A 45-year-old male glaucoma patient has frequent follow-ups in a glaucoma clinic. Besides the glaucoma, the patient has myopia that has been corrected with an eye glass since childhood. Upon every follow-up visit, one of the procedures undertaken is visual field assessment with frequency doubling technology. Hence, the examiner has instructed the patient to wear his habitual eyeglasses to conduct the procedure.

What is the most likely dioptric power of the patient in this case?

- A. Below 6.00D
- B. Below 9.00D
- C. Below 12.00D
- D. Below 15.00D

ANSWER: A

89. A patient with a head injury has been examined for a visual field defect. The result has shown that the hemifield of both eyes has a defect in a similar direction along the vertical meridian.

What is the most likely visual field defect in this case?

- A. Homonymous hemianopia
- B. Heteronymous hemianopia
- C. Bitemporal hemianopia
- D. Junctional hemianopia

ANSWER: A

90. A 40-year-old patient underwent complete ocular examination during a glaucoma screening campaign. On examination, different sections of the visual field were assessed by the clinicians to detect glaucoma and neurologic field defects.

What is the most common visual field area to be assessed in this case?

- A. Central 15 degree
- B. Central 30 degree
- C. Central 45 degree
- D. Central 60 degree

ANSWER: B

91. A 45-year-old female teacher went to an eye clinic for a check-up. On assessment, among ocular tests, the intraocular pressure had a borderline result.

What is the most likely intraocular pressure of this person?

- A. 16mmHg
- B. 21mmHg
- C. 26mmHg
- D. 31mmHg

ANSWER: B

92. Ocular examination was performed on a client to diagnose glaucoma. The examiner used a special lens with three mirrors in different angles during the slit lamp examination. The result of this test has shown that the client's eyes are in a normal condition.

What is the most likely anatomical structure of the eye observed in this test?

- A. Scleral spur
- B. Trabecular meshwork
- C. Iris root
- D. Schwalbe's line

ANSWER: A

93. Gonioscopic examination has been performed for one of the patients suspected of having glaucoma. After examination, the result has shown that the angle width was grade D and the iris had a convex shape configuration with a lightly pigmented angle structure.

What is the most likely gonioscopic grading method used in this case?

- A. Van Herrick
- B. Schaffer
- C. Spaeth
- D. Scheie

ANSWER: C

94. A female client came to an optical dispensing room with a spectacle prescription. In the middle of the communication, she told to a dispenser that she had an appointment the next day for rashes around her ears that appeared after using metallic ear rings.

What is the most appropriate frame to be ordered for this person?

- A. Magnesium
- B. Aluminium
- C. Titanium
- D. Bronze

ANSWER: C

95. A male client, who uses a long-lasting progressive spectacle, came to an eye clinic. He has changed his spectacles for the third time. A new lens was fitted on his previous metallic old frame with some frame adjustments. Now, he has a distortion in his vision with the new spectacle prescribed.

What is the most likely spectacle change causing visual disturbance in this case?

- A. Vertical depth adjustment
- B. Coating the antireflection on the lens
- C. Vertex distance increment
- D. +0.25D near addition increment

ANSWER: C

96. A male lawyer, who had been using a bifocal spectacle for a long period, came to a hospital, complaining about the difficulty in seeing the dispersed documents from one side of the table to the other. Due to the nature of his work, his near visual field demand is higher. An optometrist decided to prescribe a bifocal with a wider near viewing portion for him.

What is the most appropriate type of bifocal to be prescribed in this case?

- A. Executive segment
- B. Round segment
- C. Flat-top segment
- D. Ribbon segment

ANSWER: A

97. A 55-year-old male patient came to an eye care center complaining about blurring of vision for near objects. He was refracted last month and had a power of OD: +2.00/-1.00x180 and OS: +1.50/-0.75x175 for distance with an addition of +2.50DS.

What is the appropriate near vision prescription for this case?

- A. OD; +2.50/-1.00x180 and OS; +2.50/-0.75X175
- B. OD; +4.50/-1.00x180 and OS; +4.00/-0.75X175
- C. OD; +2.50/-1.00x90 and OS; +2.50/-0.75 X85
- D. OD; +4.50/-1.00x90 and OS; +4.00/-0.75X85

ANSWER: B

98. A 16-year-old female patient came to a dispensing workshop with a prescription of -8.00DS for the right eye and -10.00DS for the left eye.

What is the most suitable frame choice for fitting this prescription?

- A. Metal frames with full rim
- B. Metal frames with small eye size
- C. Plastic frames with rounded edges
- D. Plastic frames with a large eye size

ANSWER: C

99. A 35-year-old man came to a dispensing workshop with a prescription of -4.50/-2.00 x 75 for the right eye and -5.00/-1.50 x 180 for the left eye.

What is the most suitable frame choice for fitting this prescription?

- A. Long vertex distance frames
- B. Wider eye size frames
- C. Rounded edges frames
- D. Half rim frames

ANSWER: C

100. A female patient came to a dispensing workshop and brought a prescription of -1.00DS for both eyes with an anti-reflection coating lens. She had a round face shape.

What is the most suitable frame type likely recommended for her face shape?

- A. Narrow frame with high temple attachment
- B. Wider frame with sharp edge
- C. Round frame with sharp edge
- D. Oval frame with angular edge

ANSWER: A

101. A female patient employed at a bank visits an optical dispensing workshop to have her prescription filled. Her prescription is -1.00 DS in both eyes, and she has chosen lenses with an anti-reflection coating to minimize glare and improve visual clarity, especially for her office work. Upon facial assessment, the patient is noted to have a round face shape, which requires careful selection of frame style to enhance facial balance and aesthetic appeal.

what type of frame is most appropriately recommended for her round face shape?

- A. Narrow frames with high temple attachment
- B. Wider frames with sharp edges
- C. Round frames with sharp edges
- D. Oval frames with angular edges

Answer: A

102. A man, who uses bifocal eyewear which was prescribed from a hospital for the last one month, came with a complaint of difficulty reading with his eyeglasses unless he pushes the eyeglasses upward when reading to see through the bifocal.

What is the most appropriate adjustment to correct this problem?

- A. Decrease the vertex distance
- B. Increase the Pantoscopic tilt
- C. Decrease the temple bend
- D. Bring the nose pads together

ANSWER: D

103. A 50-year-old man, who wears spectacles, came to an optical workshop and complained that the current frame's bottom rim rests on his cheek.

What is the most likely cause of this problem?

- A. The frontal angle is too vertical for the angle of the nose
- B. The crest angle is too narrow compared to the nose
- C. The splay angle is too narrow compared to the nose
- D. The pantoscopic angle is narrow to the face

ANSWER: A

104. A 35-year-old daily labourer, who was prescribed to wear protective eyeglasses, has come to the optical workshop. He had been routinely involved in rough work and physical activities.

What is the most appropriate frame template for this patient?

- A. Skull
- B. Library
- C. Convertible
- D. Riding bow

ANSWER: D

105. A 65-year-old man, who had been wearing spectacles for the past two weeks, came to the optical workshop complaining of a sudden position change of objects when he changed his vision from distance to near. He was using another bifocal lens before this spectacle was comfortable. The lens in his current spectacle was a rounded segment bifocal lens.

What is the most appropriate bifocal lens for this patient?

- A. Flat top segment with 26mm segment width
- B. Flat top segment with 30mm segment width
- C. Flat top segment with 28mm segment width
- D. Flat top segment with 25mm segment width

ANSWER: B

106. A newly prescribed, single vision spectacle user complained about visual discomfort. The optical center of the +2.50DS lens was decentred 2mm to the nasal.

What prismatic effect caused visual discomfort with this lens?

- A. 0.25
- B. 0.5
- C. 2.5
- D. 5.0

ANSWER: B

107. An optometrist prescribed a prism for a patient to correct an eye deviation. Even though it is urgent, an optician could not get a lens with a prism. An optician asked the prescribing optometrist for any option that compensates for the prescription.

What is the most appropriate procedure to advise the optician?

- A. Sphere addition
- B. Lens decentration
- C. Power transposition
- D. Cylinder incorporation

ANSWER: B

108. A progressive lens of a 45-year-old patient is glazed in an optical laboratory. An optician needs to check the removable markings when the prescription is returned from the laboratory.

What is the most appropriate explanation that indicates perfect fitting?

- A. The distance reference point is at the center of the arc
- B. The fitting cross is 4mm above the pupil of the patient
- C. The horizontal dashes are tilted by 15 degrees
- D. The near reference point is at the center of the lens

ANSWER: A

109. A male driver, who is using +1.75DS for both eyes, has come to an eye clinic complaining of difficulty driving at night with the spectacle. And he consulted an optician to recommend for suitable lens. The lens he had been using was white.

What is the most effective lens type the optician should recommend in this case?

- A. Photochromatic
- B. Polarizer
- C. Antireflection coated
- D. Tinted

ANSWER: C

110. A male tourist consulted an optician to recommend suitable spectacles that are used for different purposes. A tourist told the optician that he is an ice skater and he mainly needs the spectacle while ice skating.

What is the most suitable lens an optician should recommend for the tourist?

- A. Mirror-coated
- B. Antireflection coated
- C. Anti-scratch coated
- D. Antifog coated

ANSWER: D

111. A male client who uses +10.00DS spectacle came to an eye clinic complaining of right and left side vision reduction with the spectacle. On assessment, the client reported that sometimes it was difficult to see objects if it is not appear in the central area. An optometrist identified that the problem was due to decreased visual field.

What is the most appropriate management option in this case?

- A. Decrease the Pantoscopic tilt
- B. Decrease the frame diameter
- C. Decrease the back vertex distance
- D. Decrease the space of the nose pads

ANSWER: C

112. A male client came to an eye care center complaining of difficulty blinking with the spectacle with a metal frame. While he is blinking, the eyelash touches his spectacle.

What is the most likely procedure that moves the frame away from the eye?

- A. Spreading the nose pads
- B. Decreasing the effective length of pads
- C. Spreading the temples
- D. Shrinking the bridge

ANSWER: D

113. A spectacle user customer has come to the dispensing room with a complaint of vision distortion after wearing a progressive spectacle for one week. The dispenser was sure that the spectacle was dispensed with the correct powers.

What is the first step the dispenser must take in this case?

- A. Measure the fitting cross height of the patient
- B. Measure the pupillary distance of the patient
- C. Put the progressive markings on the lenses
- D. Adjust the pantoscopic tilt of the patient

ANSWER: C

114. A first-time bifocal user visited an eye care center repeatedly with complaints of discomfort and inability to adapt to the bifocal. The patient did not understand the cause of the discomfort that an eye care professional explained to him.

What is the most likely cause of discomfort in this patient?

- A. Peripheral distortion
- B. Image jump
- C. Prismatic effect
- D. Image magnification

ANSWER: B

115. An eight-year-old male child came to an eye clinic for an ocular examination. His family reported that his eye was deviated and his vision was reduced.

What is the most appropriate history to be asked in this case?

- A. Birth weight
- B. Gestational period
- C. Age of onset
- D. Consciousness

ANSWER: C

116. A five-year-old child was brought to an eye care center. An optometrist wanted to grossly examine the presence of eye deviation, opacities in the visual axis, and isometropia. During the procedure, the optometrist tries to compare the difference between the two eyes. What is the most appropriate test to be applied in this case?
- A. Bruckner
 - B. Hirschberg
 - C. Krimsky
 - D. Amblyoscope
- ANSWER: A
117. A nine-year-old patient was brought to a hospital with a complaint of a reduction in vision. After slit lamp eye examination, an optometrist decided to perform cycloplegic refraction. What is the most appropriate drug used in this case?
- A. Atropine
 - B. Homatropine
 - C. Cyclopentolate
 - D. Tropicamide
- ANSWER: C
118. A 12-year-old child came to an optometry clinic complaining of asthenopic symptoms, especially associated with reading. The distance visual acuity of each eye is 6/9. During subjective refraction, the child accepted +1.00DS, and the visual acuity is 6/6 in each eye. The child further accepted plus lenses up to +2.00DS and still has visual acuity of 6/6. According to the final subjective refraction result. What is the most likely type of hyperopia that the child has?
- A. Latent hyperopia
 - B. Manifested hyperopia
 - C. Facultative hyperopia
 - D. Absolute hyperopia
- ANSWER: B
119. A 20-day-old infant was brought to an ophthalmic clinic by parents with complaints of swollen eyelid, mucopurulent discharge, and chemosis of both eyes that started 10 days ago. During birth, the infant did not get prophylactic tetracycline eye ointment. What is the most likely diagnosis in this case?
- A. Congenital nasolacrimal duct obstruction
 - B. Allergic conjunctivitis
 - C. Ophthalmia neonatorum
 - D. Infective chronic follicular conjunctivitis
- ANSWER: C
120. A two-week-old infant was brought to an eye clinic by parents with complaints of a swollen eyelid and profuse purulent discharge in both eyes, starting from three days after birth. Delivery was through vaginal delivery. The child has chemosis and corneal ulceration. What is the most appropriate management in this case?
- A. Topical antibiotics
 - B. Systemic antibiotics
 - C. Topical antibiotics and tropicamide
 - D. Apply shield
- ANSWER: B

121. A 10-year-old student came to an eye clinic and reported a problem of jumping around on the page while reading a book, trouble following the line, and eyestrain after reading for 50 minutes. On physical examination, ocular health, visual acuities, refractive error, and accommodative function and near point convergence were normal.

What is the most appropriate binocular test to be done in this case?

- A. Cover uncover test
- B. Hirschberg reflex test
- C. Krimsky's reflex test
- D. Fusional vergence test

ANSWER: A

122. A 25-year-old patient came to an eye clinic with a complaint of blurred vision while reading. The visual acuity of the right eye is no light perception, and the left eye has 6/6 vision.

What is the most appropriate method to measure the amplitude of accommodation in this case?

- A. Push-up method
- B. Minus lens method
- C. Using ± 2.00 DS flippers
- D. Monocular estimated method

ANSWER: B

123. An 18-year-old patient came to an eye clinic with complaints of a pulling sensation around the eyes, frontal headache late in the day, diplopia, and eyestrain. The patient has esophorias of 6 prism diopters at near and 2 prism diopters at distance, A C/A ratio of 6:1, and the near point of convergence was closer to the nose. The ocular health and visual acuity were normal.

What is the most likely diagnosis in this case?

- A. Convergence insufficiency
- B. Divergence insufficiency
- C. Convergence excess
- D. Pseudo convergence insufficiency

ANSWER: C

124. A 22-year-old patient came to an eye clinic with a complaint of jump around on the page while reading, trouble following the line, and eyestrain at nearby. The patient prefers reading with one eye. During physical examination, visual acuity is 6/6 in both eyes separately. Ocular health and accommodative function are normal. The refraction result is emmetropic for both eyes. A hesitant and jerking recovery movement at 10cm is noted during the near point of convergence test. The patient also has a slow and jerking recovery movement during the cover test at near.

What is the most likely diagnosis for this patient?

- A. Convergence insufficiency
- B. Decompensated phoria
- C. Fusional vergence dysfunction
- D. Binocular suppression

ANSWER: B

125. A six-year-old child was referred from a paediatric clinic to an eye clinic for ophthalmologic evaluation. The child had 20prism diopter esotropia in the right eye. During the cycloplegic refraction, a refractive error of +4.75DS in the right eye and +3.00DS in the left eye is obtained. The ocular motility and other ocular findings are all quiet.

What is the most appropriate management for this child?

- A. Lateral rectus muscle resection or medial rectus muscle resection
- B. Full cycloplegic refraction prescription and patching the right eye
- C. Penalization of the right eye with one percent atropine sulphate
- D. Orthotic exercise with an evaluation every month

ANSWER: B

126. A 16-year-old patient came to an optometry clinic with complaints of jumping around on the page while reading, trouble following the line, and eye strain after reading for about 50 minutes. During physical examination, ocular health and visual acuity are normal. His near point of convergence is 12cm, the AC/A ratio is 3:1, and he had 10 prism diopter of exophoria at near. The amplitude of accommodation is 11.00 DS.

What is the most appropriate management for this patient?

- A. Vision therapy
- B. Add eyeglasses
- C. Treating latent hyperopia
- D. Prescribe prism

ANSWER: A

127. A 12-year-old patient with ocular-cutaneous albinism came to an eye clinic with a complaint of difficulty copying notes from the blackboard, watching football, and photophobia. His distance visual acuity is 1/60 and 6/60 in the right and left eyes, respectively. With refraction, he has no visual acuity improvement.

What is the most critical factor in assessing for low vision aid selection?

- A. Being a 12-year-old child
- B. The daily activity
- C. Having poor vision
- D. Having ocular-cutaneous albinism

ANSWER: B

128. A 17-year-old male patient came to an eye clinic with a complaint of difficulty copying notes from a blackboard, even from the front desk in the classroom. He had cataract surgery in both eyes 10 years ago. He has visited different clinics to consult about his vision condition and has been told that his vision will not improve further with treatment.

What is the most appropriate next step for this case?

- A. Clinical examination of visual functions
- B. Calculation of required magnifications
- C. Selection of low vision assistive devices
- D. Referral to rehabilitation training

ANSWER: A

129. A 50-year-old patient, who had refractive surgery for myopia three years before, came to an optometry clinic complaining of difficulty driving at night. During physical examination, the patient has 6/6 visual acuity in both eyes. The anterior segment and posterior segments are normal. What is the most appropriate test to be applied next in this case?

- A. Visual field test
- B. Contrast sensitivity
- C. Glare test
- D. Colour test

ANSWER: B

130. A 60-year-old male with AMD came to an eye clinic with a complaint of difficulty of reading medical leaflets and drug expiry dates with his +4.00DS reading glasses. During examination, the near vision with glasses is N16.

What is the most appropriate management for this patient?

- A. Advice to use hand handheld magnifier
- B. Advice to use dome dome-shaped magnifier
- C. Advice to use a spectacle magnifier
- D. Advice to use a stand magnifier

ANSWER: A

131. A 50-year-old male patient with early AMD came to an eye clinic with a complaint of reading difficulty with his reading glasses. The patient is a lawyer. During examination, he reads his reading materials with a +5.00DS lens under good lighting conditions comfortably and fluently. He has no refractive error at distance.

What is the most appropriate magnifier needed for this patient?

- A. Spectacle magnifier with good illumination working room
- B. Hand-held magnifier with good illumination working room
- C. Stand with a well-illuminated working room
- D. Flat-shaped good illumination working room

ANSWER: A

132. A patient who wears contact lenses came to an optometry clinic with complaints of redness, fear of light, and irritation. During examination, mucoid discharge, mild hyperaemic conjunctiva, and cobblestones on tarsal conjunctiva are seen in both eyes.

What is the most appropriate management of this case?

- A. Discontinue contact lenses and give steroids and antibiotics
- B. Give steroids with contact lens wear
- C. Give lubricants with contact lens wear
- D. Discontinue contact lenses and give antibiotic

ANSWER: A

133. A 20-year-old male patient came to an eye clinic with complaints of foreign body sensation, fear of light, and frequent blinking in his left eye for the last five days. On ocular assessment, an optometrist diagnosed a corneal foreign body. However, the frequently blinking eye of the patient restricted the optometrist to view and remove the foreign body.

What is the most appropriate instrument that can help the optometrist in this case?

- A. Clamp
- B. Forceps
- C. Speculum
- D. Hook

ANSWER: C

134. An optometrist planned to clean and decontaminate blood-contaminated surgical equipment that had been used a few days before to make it ready for the upcoming procedures. What is the most appropriate initial step to decontaminate the equipment?
- A. Rinsing
 - B. Drying
 - C. Soaking
 - D. Flaming
- ANSWER: C
135. A five-year-old male child was brought to an eye clinic with a penetrating trauma to his right eye. After a complete ocular examination, he is diagnosed with a corneal tear. An ophthalmologist has scheduled to undertake a corneal tear repair under general anaesthesia. Before surgery, the anaesthetist suggested that the child should fast for eight hours. What is the most likely reason to make the child fast in this case?
- A. To prevent seizure
 - B. To prevent airway closure
 - C. To prevent faecal incontinence
 - D. To prevent urine incontinence
- ANSWER: B
136. An optometrist planned to prepare a patient for retinal detachment surgery. On a pre-operative assessment of the eye, the optometrist observed that the patient had a bacterial conjunctivitis. What is the most likely treatment to be used before surgery in this scenario?
- A. Ciprofloxacin 0.3 percent
 - B. Dexamethasone 0.1 percent
 - C. Flurbiprofen 0.03 percent
 - D. Tetracycline 1 percent
- ANSWER: A
137. A 24-year-old male patient came to an eye clinic with a complaint of painless swelling on the lower lid of the left eye. On assessment, an optometrist diagnosed chalazion and decided to undertake surgery. The optometrist administered a local anaesthetic to make the incision. What is the most appropriate direction of incision in this case?
- A. Horizontal
 - B. Vertical
 - C. Diagonal
 - D. Oblique
- ANSWER: B

138. An optometrist planned to undertake a bilamellar tarsal rotation surgery on a patient with trichomatous trichiasis. Before the surgery, the optometrist placed the patient on a surgical bed and prepared him ready to start the surgery.

What is the most appropriate sitting position for the optometrist to undertake the surgery?

- A. Sitting at the head of the patient
- B. Sitting at the side of the patient
- C. Sitting in front of the patient
- D. Sitting above the patient

ANSWER: A

139. A surgeon conducted enucleation surgery on a patient with phthisis bulbi. During the procedure, the surgeon has removed the conjunctiva and Tenon's capsule.

What is the most appropriate next step in the above scenario?

- A. Cutting the optic nerve
- B. Removing the eyeball
- C. Inserting orbital implant
- D. Separating extraocular muscles

ANSWER: D

140. A surgeon has been conducting a bilamellar tarsal rotation surgery for a trichomatous trichiasis patient. Hence, the surgeon wants to incise the eyelid.

What is the appropriate equipment to be used in this case?

- A. Scissor
- B. Blade
- C. Forceps
- D. Clamp

ANSWER: B

141. Trichomatous trichiasis surgery was conducted for a trichomatous trichiasis patient. But, after the surgery, the operated eyelid manifested a nasal recurrence.

What is the most likely cause of nasal recurrence in this case?

- A. Sutures are too tight
- B. Tarsal bites too low
- C. Incisions are too high
- D. Skin bites too close to lashes

ANSWER: B

142. A surgeon performed a cataract surgery on a patient with a mature senile cataract. After surgery, the postoperative care has been started. However, the operated eye of the patient has developed endophthalmitis.

What is the most likely period to develop endophthalmitis in this case?

- A. Post-operative day 1-3
- B. Post-operative day 4-6
- C. Post-operative day 7-9
- D. Post-operative day 10-12

ANSWER: B

143. A patient came to an eye clinic with a complaint of painless swelling on the right upper lid. On examination, an optometrist diagnosed the patient with a chalazion and decided to conduct surgery. The patient has complained of pain around the right eyelid right after the surgery. What is the most appropriate treatment in this case?
- A. Paracetamol tablet
 - B. Ciprofloxacin tablet
 - C. Augmentin tablet
 - D. Prednisolone tablet
- ANSWER: A
144. A patient who underwent a trichomatous trichiasis surgery came to an eye clinic for post-operative follow-up after six weeks. On assessment, the operated lid has developed a granuloma. What is the most appropriate management in this case?
- A. Excision
 - B. Antibiotics
 - C. Warm compress
 - D. Follow-up
- ANSWER: A
145. An ophthalmologist prepared to conduct a cataract surgery and wanted to use hand scrub before starting the procedure. What is the most appropriate duration of scrubbing in this case?
- A. 1 minute
 - B. 5 minutes
 - C. 10 minutes
 - D. 15 minutes
- ANSWER: B
146. An optometrist assembled reusable plastic equipment in various surgical sets and intends to sterilize them separately. What is the most appropriate type of sterilization in this case?
- A. Steaming
 - B. Soaking
 - C. Autoclaving
 - D. Boiling
- ANSWER: A
147. A patient who has been under topical corticosteroids for severe uveitis requested not to use the medication because of perceived difficulty in hearing following medication use. The health professional accepted the complaint and agreed to stop prescribing the medication, knowing the possible hearing side effects of the medication. What is the most likely ethical principle applied in this case?
- A. Justice
 - B. Autonomy
 - C. Beneficence
 - D. Nonmaleficence
- ANSWER: C

148. A 76-year-old patient with long-term Diabetic Mellitus was admitted and scheduled for eye surgery. The patient's child is with him to take care of him. However, the consent was not taken for the surgery because the patient was unable to comprehend due to the serious illness and was using medication. But the procedure was done successfully, and vision was preserved.

What ethical principle is most likely violated in this case?

- A. Paternalism
- B. Autonomy
- C. Beneficence
- D. Nonmaleficence

ANSWER: B

149. A 35-year-old patient visited an eye clinic for frequent headaches associated with vomiting. On examination, the optometrist diagnosed the case as a serious disorder and informed the patient that only one oral medication is available for treatment. However, the patient refuses to take the medication due to anticipated gastric irritation. The optometrist immediately accepts the patient's refusal and appoints the patient for another time.

What is the most likely ethical principle applied by the optometrist in this case?

- A. Veracity
- B. Fidelity
- C. Beneficence
- D. Autonomy

ANSWER: D

150. An elective cataract surgery was scheduled, and informed consent was taken after appropriate discussion with the patient. Unfortunately, formaldehyde was injected instead of lidocaine for anesthetizing the eye during the cataract surgery. The professionals were accused in front of the law of the malpractice.

What is the most likely ethical principle violated that leads to a lawsuit in this case?

- A. Paternalism
- B. Autonomy
- C. Beneficence
- D. Nonmaleficence

ANSWER: D

151. In a national eye bank, thousands of clients are waiting for a lens transplant from donors. Today, one of the well-known artists registered and is included in the waiting list to receive the donation. On the same day, the staff contacted this popular artist and did cross-matching for transferring the eye lens. However, one senior optometrist resists the procedure and said, "We have to be ethical".

What ethical principles best support the senior optometrist's resistance?

- A. Autonomy
- B. Beneficence
- C. Paternalism
- D. Justice

ANSWER: D

152. A 28-year-old male patient came to an emergency room with an eye injury due to a car accident. The health care worker explains the extent of the problem and informs that he needs to have surgery. The health care worker assesses the patient's capacity to reason and make an autonomous decision based on discussions about the surgery. What is the most appropriate component of informed consent that the health care worker tried to address?
- A. Competence
 - B. Disclosure
 - C. Comprehension
 - D. Voluntary choice
- ANSWER: A
153. A 63-year-old male illiterate patient was having preoperative counselling. The patient was rushed to sign consent without enough understanding of the nature of the procedure, since he needed the procedure to be done so quickly. As a result, the procedure was initiated immediately. What is the most likely type of consent explained in this case?
- A. Consent by word
 - B. Consent by action
 - C. Informed consent
 - D. Uninformed consent
- ANSWER: D
154. A junior optometrist is always blamed for his poor patient handling and compassionate care. The optometrist was emotionally stable, had less concern for patients' feelings, and was less reactive to the feelings of patients. As a result, focused capacity-building training was organized for the professionals to improve the level of compassionate care. What is the most likely compassionate character to be addressed in the capacity building?
- A. Empathy
 - B. Sensitive
 - C. Sympathetic
 - D. Distress tolerance
- ANSWER: A
155. A senior optometrist usually gives adequate time for patients to express their feelings, takes consent for every procedure, discusses patients' information with hospital staff, and uses a screen to keep privacy of the patient. However, the optometrist was accused of unethical behaviour that he showed during patient management in the hospital. What is the most likely reason for the health professional's accusation in this case?
- A. Time allocation issue
 - B. Consent utilization issue
 - C. Confidentiality issue
 - D. Privacy issue
- ANSWER: C

156. An optometrist is taken as a role model in the hospital for his patient handling and compassionate care. He talks openly to his patients, and he can listen and appropriately recognize them. However, he is emotionally stable and less reactive to the feelings of patients.

Which compassionate trait is most likely lacking in this optometrist's behavior?

- A. Empathetic
- B. Motivation
- C. Sensitive
- D. Sympathetic

ANSWER: C

157. An optometrist was doing a comprehensive assessment on follow-up patients. Between the assessments, the professional identified incorrect statements about previous visit documentation and started to correct the information.

Which documentation practice applies the correct method in this case?

- A. Line out incorrect data with a single line
- B. Use white fluid and write the correct one over it
- C. Use an eraser to write the replacement with the correct information
- D. Use a separate sheet and transcribe the correct information

ANSWER: A

158. An optometrist is deployed to a rural district of a country with a population of 25,000. The country does not have any eye care service provision centers. So, the optometrist plans to initiate the service for the community.

What is the most appropriate method of delivering the service?

- A. Establishing an eye care hospital
- B. Establishing an eye care sub-center
- C. Establishing a primary eye care unit
- D. Establishing a community eye care unit

ANSWER: C

159. An eye care team conducts a comprehensive ocular assessment on secondary school students. A visual acuity examination shows that a significant number of students have reduced visual acuity.

What was the most likely way of screening for refractive error in this case?

- A. Near Pinhole vision
- B. Distance Pinhole vision
- C. Near Stenopic slit vision
- D. Distance Stenopic slit vision

ANSWER: B

160. A patient came to an outreach camp with a complaint of a reduction of vision in both eyes lasting for 10 years. On physical examination, the patient's unaided vision is 6/36 and 6/60 in the right and left eye, respectively. The optometrist wants to classify the patient's visual impairment category based on the International Classification of Diseases 11(ICD11).

What is the most appropriate classification category of the patient's condition in this case?

- A. Mild visual impairment
- B. Severe visual impairment
- C. Very severe visual impairment
- D. Moderate visual impairment

ANSWER: D

161. An eye care team provides a comprehensive eye care service in a rural district. After an ophthalmic investigation, it was found that the community had a 10 percent prevalence of trachomatous inflammation among school-age children.
What is the most appropriate way of management approach to be employed in this case?
- A. Mass topical treatment
 - B. Family systemic treatment
 - C. Individual topical treatment
 - D. Individual systemic antibiotic
- ANSWER: A
162. An optometrist is involved in an eye campaign. In the campaign, patients who had an operable cataract underwent cataract surgery. Then, the optometrist planned to conduct post-cataract surgery refraction.
What is the most appropriate time to perform post-operative refraction in this case?
- A. One week
 - B. Three weeks
 - C. Six weeks
 - D. Ten weeks
- ANSWER: C
163. A group of eye care professionals is screening school-age children for xerophthalmia. During ophthalmic evaluation, a significant number of students had a foamy plaque on the temporal aspect of the bulbar conjunctiva.
What is the most likely clinical sign of the students in this case?
- A. Conjunctival xerosis
 - B. Corneal xerosis
 - C. Keratomalecia
 - D. Bitot's spot
- ANSWER: D
164. A 21-year-old patient came to a primary eye care clinic with a reduction in vision since childhood. After careful ophthalmic investigation, an optometrist prescribed a pair of spectacles.
What is the most likely level of prevention applied by the optometrist in this case?
- A. Primary prevention
 - B. Secondary prevention
 - C. Tertiary prevention
 - D. Primordial prevention
- ANSWER: B
165. An eye care team is providing health education in the diabetic clinic about proper follow-up and early screening for cataract.
What is the most likely level of prevention applied by the eye care team in this case?
- A. Primary prevention
 - B. Secondary prevention
 - C. Tertiary prevention
 - D. Primordial prevention
- ANSWER: A

166. A patient came to a primary eye care unit with a complaint of ocular pain of two weeks duration. During ophthalmic investigation, the patient has unaided vision of 6/9 and 6/6 on the right and left eye, respectively, and conjunctival redness on both eyes. What is the most appropriate intervention for this patient?
- A. Advise to wash eyes regularly
 - B. Give a topical antibiotic
 - C. Give allergy medications
 - D. Refer urgently
- ANSWER: D
167. An optometrist who is engaged in an outreach eye care service identified a 30-year-old female patient with severe ocular discomfort. After an ophthalmic examination, she has a large number of eyelashes which has turned in and touch the cornea. What is the most appropriate management for this patient?
- A. Antibiotic therapy
 - B. Eyelid surgery
 - C. Trimming
 - D. Epilation
- ANSWER: B
168. An optometrist is engaged in a trachoma elimination program in a rural district of a country. The program has a series of activities, including health education about trachoma prevention strategies. What is the most appropriate topic in this case?
- A. Environmental hygiene
 - B. Fortification of food
 - C. Immunization against infection
 - D. Trauma and anoxia at birth
- ANSWER: A
169. A pregnant woman came to an outreach camp from a severely drought-affected area with a complaint of foreign body sensation. After an ophthalmic examination, conjunctival and corneal xerosis were found. What is the most likely management in this case?
- A. Daily dose of 20,000 IU of vitamin A orally for 2 weeks
 - B. Daily dose of 10,000 IU of vitamin A orally for 2 weeks
 - C. Daily dose of 10,000 IU of vitamin A orally for 1 week
 - D. Daily dose of 20,000 IU of vitamin A orally for 1 week
- ANSWER: B
170. An eye care team conducted eye health screening among primary school students in a rural part of the country. During ophthalmic investigation, some students presented with keratomalacia. What is the most likely stage of xerophthalmia in this case?
- A. X2A
 - B. X2B
 - C. X3A
 - D. X3B
- ANSWER: D

171. An optometrist conducted on-the-job refreshment training to general health care professionals regarding the recommendation of screening of children for an eye disease.

What is the highest-priority ocular condition that should be addressed?

- A. Vitamin A deficiency
- B. Refractive error
- C. Nystagmus
- D. Amblyopia

ANSWER: C

172. An optometrist who is involved in an eye campaign encountered a patient with a gradual, painless reduction of vision. After careful assessment of visual acuity, the patient has a right eye vision of 6/12 and a left eye vision of 6/18. So, an optometrist wanted to assess the function of the optic nerve to rule out glaucoma as a cause of visual reduction.

What is the most appropriate method of screening in this case?

- A. Colour vision
- B. Ocular motility
- C. Pupillary function
- D. Contrast sensitivity

ANSWER: C

173. Health extension workers in a community are involved in delivering health education about personal hygiene, the causes of trachoma, and prevention methods for trachoma.

What is the most appropriate practice of the health extension workers in this case?

- A. Preventive
- B. Curative
- C. Rehabilitative
- D. Promotive

ANSWER: D

174. An eye care professional who is newly employed as a community eye health professional in a health center aims to reduce the burden of visual impairment and blindness in the district.

What is the most appropriate description of the eye care professional's action in this case?

- A. Focus on individual health
- B. Foundation on clinical skills
- C. Apply a survey as a diagnostic tool
- D. Focus on a medical intervention

ANSWER: C

175. An optometrist works on comprehensive eye care services and delivers monthly health talks about medication adherence to glaucoma patients at a glaucoma clinic.

What is the most appropriate method of delivering the health talks in this case?

- A. Insists on a formal atmosphere
- B. The speaker behaves like a school teacher
- C. A two-way communication
- D. Subject matter theoretical

ANSWER: C

176. An eye care team conducts a trachoma screening among primary school children in a rural part of the country. A team identifies a student with poor facial cleanliness and trichomatous follicles in the palpebral conjunctiva.

What is the most appropriate method of management in this case?

- A. Tetracycline eye ointment once daily for 6 weeks
- B. Tetracycline eye ointment twice daily for 4 weeks
- C. Tetracycline eye ointment twice daily for 6 weeks
- D. Tetracycline eye ointment once daily for 4 weeks

ANSWER: C

177. A researcher conducted a study to determine the prevalence of trachoma and its association with sharing a towel among pregnant women who had antenatal care follow-up at public hospitals at the same point in time.

What is the most appropriate study design to be used in this case?

- A. Cohort
- B. Case-control
- C. Experimental
- D. Cross-sectional

ANSWER: D

178. Ten freshman optometry students were recorded by the registrar of a university. The record showed that the ages of the students are 17, 20, 21, 18, 19, 20, 22, 23, 24, and 21.

What is the range of the students' ages in this case?

- A. 7
- B. 6
- C. 5
- D. 4

ANSWER: A

179. A researcher conducted a study to determine the new occurrence of trachoma among 300 elementary school students from August 1st to the end of October 2021. During his follow-up, about 13, 15, and 10 children developed trachoma in August, September, and October, respectively.

What is the appropriate measure of trachoma disease occurrence in this case?

- A. Point prevalence
- B. Incidence density
- C. Period prevalence
- D. Cumulative incidence

ANSWER: D

180. A researcher conducted a study to determine the prevalence of conjunctivitis among reproductive age women who visited public hospitals in a district in the year 2022. The data were collected through history taking from the women in person, and additional explanation was provided to them when they got confused with the questions.

What is the most likely data collection method used in this case?

- A. Observation
- B. Mailed questionnaire
- C. Face-to-face interview
- D. Focus group discussions

ANSWER: C

181. An annual report of a region shows that there were 50 trachoma patients in the region. Among them, 10 became blind in the same year.

What is the blinding rate of trachoma per 100 population in this case?

- A. 10
- B. 20
- C. 40
- D. 50

ANSWER: B

182. The seven optometry students' research method course exams score out of 10 were 4, 7, 3, 5, 6, 8, and 6. The central exam score location of all students was required and calculated to determine how many students varied in their exam scores.

What is the median of the students' exam results in this case?

- A. 5
- B. 6
- C. 7
- D. 8

ANSWER: B

183. An optometrist conducted a study to evaluate the effect of pilocarpine on the reduction of intraocular pressure of glaucoma patients in a town in the year 2023. The optometrist grouped patients randomly into two groups: one group received pilocarpine, but the other did not.

What is the most likely study design used in this case?

- A. Cohort
- B. Case-control
- C. Experimental
- D. Cross-sectional

ANSWER: C

184. A quality assurance director of a hospital wanted to evaluate patients' satisfaction with the services provided in the optometry department. The sample size was determined to be 50 patients who visited the optometry department during the data collection period. Then, the data collector interviewed all of them, consequently, until he collected 50 data.

What is the most likely sampling technique used in this case?

- A. Quota
- B. Snowball
- C. Purposeful
- D. Convenience

ANSWER: D

185. An optometrist wishes to research to determine the prevalence of iatrogenic eye trauma experienced by patients who visited the optometry department. The minimum sample size computed to conduct this research was 424. However, 55 patients refused to participate in the study, and they did not provide data at all.

What was the response rate observed in this case?

- A. 94 percent
- B. 87 percent
- C. 84 percent
- D. 97 percent

ANSWER: B

186. An optometry department head planned to evaluate lecturers' teaching methods at the optometry department. Then, he designed highly structured questionnaires accordingly. Finally, the technical assistants provided the questionnaires simultaneously for all optometry students in the class, and the students filled them out and returned it back to them.

What is the most likely data collection method used in this case?

- A. Mailed questionnaire
- B. Face-to-face interview
- C. Focus group discussions
- D. Self-administered questionnaire

ANSWER: D

187. An optometrist wanted to develop a research hypothesis to be investigated by exploring problems experienced by patients who visited the optometry department. For this case, from the total patients who were available in the optometry department at the time of data collection, he invited 5 patients from the outpatient department and 2 patients from the inpatient department. The invited patients argued about the problems they faced, and finally, they reported to him the agreed-upon points.

What is the most likely data collection tool used by the optometrist in this case?

- A. Self-administered
- B. Mailed questionnaire
- C. Face-to-face interview
- D. Focus group discussions

ANSWER: D

188. An optometrist wanted to determine the prevalence of glaucoma among adult communities after he enrolled the required sample size through a systematic random sampling technique. After the data was collected and analysed, the result showed a high prevalence of glaucoma existed among adult communities, and the optometrist made a link for identified patients to hospitals to get care and treatment.

Which research ethics principle was applied by the researcher in this case?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-Maleficence

ANSWER: C

189. An optometrist was one of the dedicated committees that were involved in an outbreak investigation. After all the necessary workup was performed, the outbreak of trachoma was diagnosed. Then, the optometrist was requested to provide training on trachoma for all optometrists working at a hospital.

What is the best training area for this case?

- A. Primary prevention of trachoma
- B. Treatment outcome of trachoma
- C. Laboratory investigation of trachoma
- D. Diagnosis and treatment of trachoma

ANSWER: D

190. An optometrist promoted on television about giving free cataract and glaucoma diagnoses to the communities living in a region.
Which communication component do the communities living in the region represent?
A. Source
B. Channel
C. Message
D. Audience
ANSWER: D
191. Training was provided for 10 optometrists who came from different hospitals for six days. The trainers oriented the trainees to have active participation during their training stay.
What is the training phase that is explained in this case?
A. Preparation
B. Training
C. Follow-up
D. Post-training
ANSWER: B
192. A newly established general hospital is going to organize staff members for each unit. The hospital developed a job description, specification, and qualifications needed. The hospital posted internal and external vacancies in the newspaper and received a pool of applicants.
What is the most appropriate next step in human resource planning for this case?
A. Recruitment
B. Selection
C. Job analysis
D. Retention
ANSWER: B
193. A health care system leader has identified that the employees are working on unrelated activities that do not help the organization to reach its goal. The employees were busy focusing on issues that do not contribute to the organization's success.
What is the most likely measure that the health care system leader should take in this case?
A. Sharing priority objectives
B. Changing leadership style
C. Motivating team member
D. Delegating authority
ANSWER: A
194. An optometry staff member accused a supervisor of denying sick leave. The manager called the supervisor to explore the conflict. The supervisor stated that the reason for denying the sick leave was due to repeated requests more than three times within a month. After hearing both sides, the manager was less concerned about the conflict and left the conflict unsolved.
What is the most likely conflict management strategy the manager applied in this case?
A. Avoidance
B. Smoothing
C. Compromise
D. Collaboration
ANSWER: A

195. A hospital manager and a supervisor of the ophthalmic unit were having a rough discussion. The manager heard that the newly hired employees were unfamiliar with the working environment, and the new employees are disorganized due to they were directly engaged to work by the supervisor. What is the most likely human resource planning step the supervisor missed in this case?
- A. Orientation
 - B. Knowledge test
 - C. Reference checking
 - D. Performance evaluation
- ANSWER: A
196. A charity organization, which is working on elderly people's eye care, conducted a workshop and developed a document incorporating the overall goal of the organization, objectives, values, mission, and vision statement for the coming five years. What is the most appropriate planning document that the team developed in this case?
- A. Strategic plan
 - B. Synergic plan
 - C. Continuing plan
 - D. Operational plan
- ANSWER: A
197. A chief executive director of a primary hospital is awarded by staff members for being a good leader. The subordinates mentioned good communication and motivation as qualities of the director. The subordinates also acknowledged that the director provides clear instructions about procedures and methods in a manner that the employees follow willingly and with enthusiasm. What is the most likely managerial function that describes the director's role in this case?
- A. Directing
 - B. Planning
 - C. Organizing
 - D. Coordination
- ANSWER: A
198. A busy manager has been working on a deadline project which should be submitted on the due date. The manager is also expected to submit a report, which is not yet finalized. At the same time, the manager's supervisor requested for procurement plan as soon as possible, which is not ready. What is the most likely management principle that helps the manager in this case?
- A. Functional authority
 - B. Communication
 - C. Dissemination
 - D. Delegation
- ANSWER: D

199. A project working on better eye care for all has designed a specific objective as “train ophthalmic professionals on community level prevention of trachoma by the end of the third year of the project cycle.”

What is the most likely objective-setting criterion that is missed in this case?

- A. Specific
- B. Realistic
- C. Measurable
- D. Time-bound

ANSWER: C

200. A project manager working on quality care for glaucoma patients has been tracking the progress of each planned activity with the cost incurred and the timeline for carrying out each activity. The manager observed that there is an over-budget utilization for one of the activities in the plan. The manager also checked that two activities have not yet started, which should be accomplished before the end of the current quarter.

Which project management phase is the manager most likely executing in this case?

- A. Project plan
- B. Project goal
- C. Project evaluation
- D. Project monitoring

ANSWER: D