

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

NURSING



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority and is essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

The EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and to the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modelled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, leadership and management, ethics, and Health Promotion as outlined in the Information Booklet 2024.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and gaps in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging with this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey—and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights to the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions—both public and private—whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Nursing

1. A 16-year-old boy presents to a primary hospital with complaints of fever, headache, and malaise. He said that he usually wakes up at midnight due to apnea and he has difficulty of swallowing. Upon examination, the boy has a bad breath odor, a red pharynx with yellow exudates around the adenoids, mouth breathing, and nasal speech. The nurse in charge suspected tonsillitis.

What is the most likely clinical manifestation to suspect tonsillitis in this case?

- A. Midnight apnea and fever
- B. Red pharynx, yellow exudes, and fever
- C. Headache, difficulty of swallowing and malaise
- D. Bad breath odor and difficulty of swallowing

ANSWER: B

2. A 38-year-old female came to a hospital with complaints of cough, yellowish sputum, fever, and chest pain for two days. The symptoms worsen with breathing. Upon examination, the patient looks severely ill, uses of respiratory muscle, and crackle is noted on auscultation. Her temperature is 39.5 degrees centigrade; pulse rate is 112 beats/min.; respiratory rate is 32 breaths/min. A physician diagnosed the patient with pneumococcal pneumonia.

What is the most likely typical manifestation to set the diagnosis in this case?

- A. Cough with yellowish sputum and crackles with fever
- B. Severely ill appearance and pulse rate of 112 beats/minute
- C. Chest pain worse with breathing and tachycardia
- D. Respiratory rate 32 breath/min and use of respiratory muscle for breathing

ANSWER: A

3. A 32-year-old female came to a hospital with complaints of cough with sputum and chest pain of five days duration. Upon examination, the patient looks severely ill, in respiratory distress, and the trachea is slightly deviated to the right. Absent breath sound on auscultation and dull sound on percussion are noted. Examination findings show a temperature of 38.5 degrees centigrade; pulse rate of 109 beats/min; and respiratory rate of 32 breaths/min. The physician diagnosed the patient with pleural effusion.

What is the most likely manifestation consistent with the diagnosis of pleural effusion?

- A. Cough with sputum and chest pain
- B. Use of respiratory muscle on breathing and fever
- C. Absence of breath sound and dull not on percussion
- D. Respiratory rate of 32 breaths/ min and pulse rate of 109 beat/minute

ANSWER: C

4. A 28-year-old male patient, who attended a wedding ceremony the day before and consumed lots of alcohol, presented to a hospital with a complaint of voice loss. Upon physical examination, he has a harsh cough, and sore and dry throat.

What is the most likely diagnosis in this case?

- A. Tonsillitis
- B. Pharyngitis
- C. Laryngitis
- D. Epiglottitis

ANSWER: C

5. A 38-year-old female patient came to a hospital with complaints of fever, cough with rusty purulent sputum, and chest pain of two days duration. On assessment, the patient looks severely ill and uses accessory muscle while breathing, and crackles noted on auscultation of left lower lobe. Her temperature is 39.5 degree centigrade, pulse rate is 120beats/min, and respiratory rate is 28/min.

What is the most likely diagnosis of this patient?

- A. Acute bronchitis
- B. Pneumonia
- C. Tuberculosis
- D. Pleurisy

ANSWER: B

6. A 28-year-old patient presents to a hospital with a diagnosis of Laryngitis. He also has a harsh cough, sore throat, and dry throat. He had a concert a day before he developed the current problem.

What is the most likely priority nursing diagnosis in this case?

- A. Altered body image
- B. Ineffective airway clearance
- C. Ineffective role performance
- D. Impaired verbal communication

ANSWER: D

7. A 56-year-old male presents to a hospital with complaints of cough and loss of appetite of four weeks duration. He also reports weight loss, night sweating, and bloody sputum. On auscultation, crackles were noted. His temperature is 38 degrees centigrade and his respiratory rate is 28 breaths/min. Laboratory investigation shows increased lymphocyte count and ESR. Chest X-ray shows apical and posterior lung infiltrates. The physician suspects pulmonary tuberculosis.

What is the most likely priority nursing diagnosis in this case?

- A. Activity intolerance
- B. Ineffective breathing pattern
- C. Ineffective airway clearance
- D. Imbalanced nutrition less than the body requirement

ANSWER: C

8. A 32-year-old male came to a hospital with complaints of cough and chest pain of five days duration. He was diagnosed with lung cancer two months ago and underwent thoracic surgery. On assessment, the patient uses accessory muscles while breathing. The trachea is slightly deviated to the left, breath sounds absent, and dull on percussion of the right lung.

What is the most appropriate respiratory care modality recommended in this case?

- A. Chest drainage
- B. Postural Drainage
- C. Thoracentesis
- D. Tracheotomy

ANSWER: A

9. A 23-year-old male patient came to a hospital with complaints of sudden onset of chest pain and fever for two days. He also reported the chest pain aggravated by deep breathing and coughing. He has no history of admission, and other chronic diseases. On assessment, he has shortness of breath and use of accessory muscles in respiration. Vital signs showed respiratory rate of 36 breaths per minute, pulse rate of 113 beats per minute, temperature of 39.1 degrees centigrade, and oxygen saturation of 87 percent.

What is the most likely medical diagnosis of the patient in this case?

- A. Aspiration pneumonia
- B. Hospital-acquired pneumonia
- C. Community-acquired pneumonia
- D. Pneumonia in immune-compromised patient.

ANSWER: C

10. A 45-year-old male client came to a hospital with complaints of food regurgitation, chest pain, and heartburn. His vital signs are stable. X-ray study shows esophageal dilation above the narrowing at the gastroesophageal junction. The patient is diagnosed with Achalasia.

What is the most likely additional manifestation required to diagnose this case?

- A. Chewing difficulty
- B. Swallowing difficulty
- C. Breathing difficulty
- D. Speaking difficulty

ANSWER: B

11. A 55-year-old male client came to a hospital with a complaint of mid-epigastric burning pain that radiated to the back for two weeks duration. He has a history of heartburn and constipation. On abdominal assessment, epigastric tenderness and abdominal distension are noted. The patient is diagnosed with severe gastric ulcer.

What is the most likely additional manifestation required to diagnose this case?

- A. Hematochezia
- B. Hematemesis
- C. Pain occurs 2-3 hours after a meal
- D. Hyper secretion of stomach acid

ANSWER: B

12. A 50-year-old male client with a history of PUD comes to a hospital with a complaint of sudden abdominal pain radiating to the right shoulder. Upon examination, it was found that a rigid board-like abdomen, BP of 70/40mmHg, PR of 120 beats/minute, RR of 28 breaths/minute.

What is the most likely medical diagnosis in this case?

- A. Acute gastritis
- B. Perforated PUD
- C. Gastric outlet obstruction
- D. Zollinger-Ellison syndrome

ANSWER: B

13. A 41-year-old male client came to an emergency OPD with a complaint of diffuse abdominal pain of two days duration. Upon physical examination, abdominal rigidity, and absent bowel sounds were noted with the following vital signs; BP of 80/60mmHg, PR of 110 beats/minute, RR of 26 breaths/minute, and temperature 38.5 degrees centigrade.

What is the most appropriate medical diagnosis in this case?

- A. Gastritis
- B. Peritonitis
- C. Diverticulitis
- D. Appendicitis

ANSWER: B

14. A 55-year-old client came to an emergency OPD with complaints of watery diarrhea and vomiting of one-day duration. Physical examination shows a dry oral mucosa, sunken eyes, blood pressure of 60/40 mmHg, pulse rate of 120 beats/minute, respiratory rate of 28 breaths/minute, and temperature of 37.5 degrees centigrade.

What is the most appropriate priority nursing intervention in this case?

- A. Intravenous fluid administration
- B. Patient education on hygiene
- C. Nasogastric tube feeding
- D. Oral fluid administration

ANSWER: A

15. A 43-year-old male patient is recovering from a cholecystectomy surgery. After the patient recovered from anesthesia, a nurse working in the unit conducted a complete assessment and noticed his vital signs are smooth and stable.

What is the most likely appropriate position for the patient?

- A. Prone position
- B. Supine position
- C. Low fowler's position
- D. Trendelenberg position

ANSWER: C

16. A 51-year-old female patient is admitted to a surgical ward with complaints of severe abdominal pain, back pain, nausea, and vomiting. She reported that her pain is aggravated by meals and not relieved by anti-acid. Laboratory investigation shows elevated serum amylase and lipase levels.

What is the priority nursing intervention in this case?

- A. Provide potent intravenous analgesic
- B. Insert NGT to remove stomach content
- C. Administer cleansing enema
- D. Provide oral mild anti-pain

ANSWER: A

17. A 23-year-old farmer came to a hospital with complaints of localized itching, erythema, swelling, and pruritic vesicles over his bare parts of the body. In history, the patient declared that he touched a green leaf known as poison ivy; and then, the itching became intense in areas that have contact with the poison ivy. Upon physical examination, a nurse noted a red rash, itching, and localized swelling over the bare part of the body.

What is the most likely diagnosis in this case?

- A. Allergic dermatitis
- B. Irritant dermatitis
- C. Atopic dermatitis.
- D. Exfoliative Dermatitis

ANSWER: A

18. A 35-year-old male patient was admitted to a surgical ward with second-degree flame burn on his entire anterior trunk, anterior and posterior right arm, and anterior part of his left leg.

What is the most likely f total body surface area (TBSA) burnt in this case?

- A. 18 percent
- B. 27 percent
- C. 36 percent
- D. 45 percent

ANSWER: C

19. A 70-year-old male stroke patient was brought to a health facility by his neighbors. On admission, the patient was diagnosed with bedsore. Upon physical examination, a wide open ulcer of the epidermis and papilla of the dermis with a red-pink bed without slough is observed at the back (around the buttocks) of the patient.

What is the most likely stage of bedsore in this case?

- A. STAGE 3
- B. STAGE 4
- C. STAGE 2
- D. STAGE 1

ANSWER: C

20. A 62-year-old man with HIV AIDS came to a hospital with a complaint of lesions around his right shoulder that do not pass the midline of the body. On assessment, lesions have blister and tingling sensation. He experiences severe pain over the entire dermatome supplied by the affected nerve root. The patient avoids movement and does not want clothing or bed linen to touch the affected shoulder.

What is the most appropriate priority management for the patient?

- A. Administer antibiotics
- B. Administer Acyclovir
- C. Administer Tramadol
- D. Administer calamine lotion

ANSWER: C

21. A 62-year-old female patient came to an emergency OPD with an open wound on her left leg, 30 minutes after she was bitten by a dog.

What is the most appropriate priority wound management for this patient?

- A. Suturing
- B. Wound debridement
- C. Wound dressing
- D. Wound cleansing

ANSWER: D

22. A 45-year-old male patient came to a hospital with a major burn injury of 35 percent total surface area. He was admitted to burn unit for 14 days. A plastic surgeon performed dermal replacement surgery. The patient complains of pain over the grafted area. Upon physical examination, the patient was stable and clear red spotting from the donor receptor site, otherwise, he is good.

What is the most appropriate nursing intervention in this case?

- A. Put the grafted area open.
- B. Institute additional pain management.
- C. Change dressing from the donor and receptor site in the next post-operative day.
- D. Encourage the patient to engage active exercise in the next post-operative day.

ANSWER: B

23. An 85-year-old male patient comes to a hospital with a complaint of generalized itching after caustic substance flow on his leg and hand. Upon examination, a nurse identified soreness, discomfort, itching, redness, swelling, and vesiculation on his hand and lower leg.

What is the most likely prioritized nursing diagnosis in this case?

- A. Acute pain.
- B. Disturbed sleep.
- C. Risk for infection.
- D. Risk for impaired skin integrity related.

ANSWER: A

24. A 25-year-old female is admitted to a hospital after having a road traffic accident. She had traction for two weeks. The physician decided removal of the traction weight and planned for closed and full leg cast. An assigned nurse is assessing the patient's fitness to the cast.

What is the most likely priority of nursing assessment in this case?

- A. Neurovascular status
- B. Mobility and exercise
- C. Muscle contracture
- D. Pain and comfort

ANSWER: A

25. A 45-year-old female patient has a radial fracture after falling injury. After the full arm cast was applied, the patient complained of mild pain and burning sensation under the applied cast area.

What is the most appropriate intervention is contraindicated in this case?

- A. Elevate the site
- B. Clean edge of cast with saline
- C. Administer prescribed anti-pain
- D. Remove other constrictive materials

ANSWER: B

26. A 45-year-old female patient, who is managed with the cast, complained of unrelieved pain in the cast covered area. Upon examination, a nurse noticed impaired tissue perfusion in the affected area. What is the most likely priority intervention needed in this case?

- A. Remove the cast
- B. Give anti-pain
- C. Report to physician
- D. Warmth the area

ANSWER: C

27. A 67-year-old male with diabetes came to a hospital with complaints of swollen, puffy, painful, and warm knee joints. He also reports some level of fever and chills. After several investigations, he is diagnosed with septic arthritis.

What is the most appropriate nursing intervention in this case?

- A. Aspirate accumulated fluid
- B. Keep jogging exercise
- C. Immobilize and rest
- D. Prepare for surgery

ANSWER: C

28. A 55-year-old male driver has been admitted to a hospital for multiple bone fractures following a traffic accident. He is now to be discharged with some internal fixation plate for the fractured ribs which had caused a flail chest. The surgical incision wound almost healed and there is no need for dressing. He asked a nurse how long the metal stays inside his body.

What is the most appropriate response of the nurse in this case?

- A. Life long
- B. Until bone heals
- C. Until next visit
- D. Three months

ANSWER: A

29. A 65-year-old female patient was admitted to a hospital with a full arm cast for the fractured radial bone following a falling injury. Three days after the cast has been applied, she is complaining of pain and paresthesia that is increasing in its severity. A nurse assessed the patient and suspected neurovascular impairment due to pressure from the cast.

What additional information is most likely needed to support the diagnosis in this case?

- A. Lice on cast
- B. Stained cast
- C. Bluish fingers
- D. Reddish nail beds

ANSWER: C

30. A patient with a closed arm fracture has stayed on traction for about two weeks. The bone healing is progressing well. A nurse prepares the patient for discharge from the hospital with a full arm cast. The nurse plans to teach the patient what he may face while he stays his home.

What should be the most appropriate priority education the nurse gives to this patient?

- A. Fat embolism
- B. Infection
- C. Compartment syndrome
- D. Muscle atrophy

ANSWER: D

31. A 68-year-old known diabetic and hypertensive female patient has above-knee amputation following complications of peripheral vascular disease. Despite the surgical procedure being performed several days before, she still complains of pain or unusual sensations, numbness, tingling, muscle cramps, and as if the extremity presents, crushed, cramped, or twisted in an abnormal position at the severed part. A nurse developed a nursing diagnosis of "Disturbed sensory perception".

What is the most appropriate nursing intervention in this case?

- A. Keeping the patient active
- B. Keep the room quiet
- C. Consult psychiatrist
- D. Remove elastic bandage

ANSWER: A

32. A male nurse working in a medical ward had a small superficial cut on his hand while preparing bandages a day before. He is now on duty in the ward and preparing to give medications to the admitted patients. As per the guideline of the hospital, it is a must to comply with standard precautions.

What should the most likely a nurse do according to the standard precautions?

- A. Cover the cut with occlusive and waterproof dressing
- B. Ask the hospital administration to replace him
- C. Take sick leave for a month so that the wound can completely heal
- D. Ask colleagues to cover your tasks and stay away from patients

ANSWER: A

33. A 25-year-old male client came to an emergency unit with a complaint of bleeding from knife stab injury at his left lower abdomen. He reported that the robbers attacked him looking for his mobile phone. While a nurse examines, the patient's shirt is fairly soaked with blood. Fresh blood is still oozing from the injury site while the patient lifts his hand from it.

What is the most likely kind of wound is it in terms of cleanliness?

- A. Clean
- B. Clean-contaminated
- C. Contaminated
- D. Dirty and infected

ANSWER: C

34. A nurse, who has the responsibility to control the traffic in the operation room, wants a patient's family to stay in an area near to operation room where surgery is ongoing.

What is the most likely area the nurse would allow the patient's family to stay with their usual cloth?

- A. Unrestricted area
- B. Semi restricted area
- C. Restricted area
- D. Operating area

ANSWER: A

35. A 25-year-old female patient is admitted to a medical ward with a diagnosis of chronic glomerulonephritis. On assessment, she has flank pain, low urine output, and bilateral crackles. Laboratory investigation revealed a urine protein +3, elevated blood urea nitrogen, and creatinine.

What is the most appropriate assessment finding best indicates a positive prognosis after intervention?

- A. Urine protein of +1
- B. Decreased BUN level
- C. Reduced flank pain
- D. Blood pressure of 140/85mmHg

ANSWER: A

36. A 23-year-old male patient with a history of seven episodes of vomiting within four hours came to a hospital with a complaint of sudden onset of vomiting and diarrhea. Assessment findings reveal a respiratory rate of 31 breaths/minute, blood pressure of 90/50mmHg, and pulse rate of 131 beats/minute. What is the most likely finding that indicates impaired tissue perfusion?

- A. Seven episodes of vomiting
- B. Pulse rate 131beats/minute
- C. Blood pressure 90/50mmHg
- D. Respiratory rate 31breath /minute

ANSWER: C

37. A 34-year-old known hypertensive patient came to a renal clinic with complaints of a very small quantity of urine, nausea, and fatigue for the last three days. Laboratory investigation shows an elevated BUN and creatinine level, and a decreased urinary sodium concentration.

What is the most appropriate information to be included in the nurse's teaching plan?

- A. Increase protein intake
- B. Encourage monosaturated fat
- C. Encourage polysaturated fats
- D. Encourage a diet in simple carbohydrates

ANSWER: D

38. A 17-year-old female patient came to a renal clinic with complaints of malaise, fatigue, and generalized body swelling that started from the face. On assessment, she had pitting edema on the sacrum, ankles, and hands. Urine analysis showed many WBC and urinary casts. Then, the patient was diagnosed with nephrotic syndrome.

What is the most likely additional finding required to confirm the diagnosis in this case?

- A. Hyperlipidemia
- B. Marked Proteinuria
- C. Grade III pitting edema
- D. Increased blood urea nitrogen

ANSWER: B

39. A 65-year-old male patient with a history of anorexia, nausea, and vomiting came to a hospital with a complaint of easy fatigability. . Assessment has shown generalized bilateral edema, engorged neck veins, a blood pressure of 165/100mmHg, and a 24-hour urine output of 375ml. Laboratory investigation showed an elevated BUN and creatinine level.

What is the most likely priority nursing diagnosis for this case?

- A. Imbalanced nutrition
- B. Fluid volume excess
- C. Activity intolerance
- D. Disturbed body image

ANSWER: B

40. An 18-year-old female client with a recent history of upper respiratory infection came to a renal clinic with complaints of facial swelling and flank pain. She reported that she has not taken any medications for the infection. Physical examination showed a blood pressure of 165/98 mmHg, and bilateral pitting edema of the feet. The physician diagnosed glomerulonephritis.

What is the most appropriate additional confirmatory investigation in this diagnosis?

- A. Urine protein
- B. WBC count
- C. Hematocrit level
- D. Hemoglobin concentration

ANSWER: A

41. A 23-year-old male patient comes to a hospital with a complaint of severe chest pain of two hours duration. Assessment findings reveal crackles, rapid, irregular, and thready pulse, ventricular gallop, a mean arterial pressure of 54 mmHg, and urine output of 20 ml/hr.

What is the most likely assessment finding best indicates Cardiogenic shock?

- A. Ventricular gallop
- B. Urine output less than 20 ml/hour
- C. Mean arterial pressure of 54 mm Hg
- D. Rapid, irregular, thready pulse

ANSWER: A

42. A 67-year-old known hypertensive patient reports that he has suffocation, chest discomfort at rest, easily fatigability, anorexia, and frequent nighttime urination. Physical examination findings show a pulse rate of 115 beats/minute and crackles on the left lobe of the lung.

What is the most appropriate assessment finding most likely indicates left sided heart failure?

- A. Easily fatigability
- B. Crackles on the left lobe
- C. Pulse rate of 115beats/minute
- D. Frequent nighttime urination

ANSWER: B

43. A 65-year-old known chronic heart failure patient is admitted to a hospital with complaints of shortness of breathing worsened by activity, easily fatigability, anorexia, and nausea. Upon assessment, blood pressure is 155/95 mmHg, pulse rate is 131 breaths/minute, and respiratory rate is 28 breaths/minute.

What is the most likely probable risk factor for the increased pulse rate?

- A. Increased blood pressure
- B. Decreased cardiac output
- C. Increased fluid in the lungs
- D. Increased irritation of the lung mucosa

ANSWER: B

44. A 65-year-old male patient is admitted to an emergency department after sustaining a falling down injury from a seven-story building. Physical examination reveals patient has bilateral leg swelling with flaccid paralysis, a pulse rate of 55 beats/minute, and a blood pressure of 80/50 mmHg. Diagnostic investigation shows an injury involving the thoracic vertebra.

What is the most likely cause for the presenting symptoms in this case?

- A. Immobility
- B. Spinal cord injury
- C. Age of the patient
- D. Bilateral edema

ANSWER: B

45. A 45-year-old male patient is admitted to a hospital with a diagnosis of left-side heart failure. On history, he has mild dyspnea and difficulty in performing activities of daily living. Upon physical examination, there is bilateral jugular vein distention, diastolic murmur, delayed capillary refill time, and decreased peripheral pulse.

What is the most likely priority nursing diagnosis in this case?

- A. Activity intolerance
- B. Impaired tissue perfusion
- C. Impaired gas exchange
- D. Ineffective breathing pattern

ANSWER: B

46. A client has complaints of fatigue and lower extremity numbness. Pertinent assessment findings include beefy red tongue, slight jaundice, and a positive Schilling test.

What is the most appropriate medical management for this case?

- A. Long-term RBC transfusion
- B. Adhere to a strict vegetarian diet
- C. Vitamin B12 supplementation
- D. Peripheral blood stem cell transplant

ANSWER: C

47. A 65-year-old male patient is admitted to a medical ward. On 12:00 pm progress assessment, he opens his eyes to verbal communication. He tries to localize the pinch site and he knows where he is and the name of the nurse in charge of his care.

What is the most likely Glasgow coma scale score in this case?

- A. 12
- B. 13
- C. 14
- D. 15

ANSWER: A

48. A 32-year-old male patient presents to a hospital with complaints of headache, fever, and stiff neck. From diagnostic evaluation, a lumbar puncture is performed. The nurse is going to reposition the patient just after the procedure.

What is the most likely recommended patient position after the procedure?

- A. Prone
- B. Lithotomy
- C. Side-lying
- D. High fowlers

ANSWER: A

49. A 45-year-old male patient was brought to a hospital with complaints of fever and headache of two days duration. Upon assessment, the patient has nuchal rigidity, a positive Kernig sign, and Brudzinski's sign. The nurse in charge has a diagnosis of risk for impaired tissue perfusion related to increased intracranial pressure.

What is the most likely nursing intervention for the stated diagnosis?

- A. Provide proper positioning
- B. Provide nutritional support
- C. Monitor fluid intake and output
- D. Implement isolation techniques

ANSWER: C

50. A 46-year-old female patient diagnosed with cancer is admitted for 3rd cycle of chemotherapy. Her numeric pain intensity scale is 9.

What is the most likely management in this case?

- A. Aspirin
- B. Morphine
- C. Acetaminophen
- D. Indomethacin

ANSWER: B

51. A 58-year-old male patient is admitted to a medical ward with a diagnosis of ischemic stroke. Upon assessment, the patient has right-side motor weakness, confusion, difficulty swallowing, wide pulse pressure, and erythema on the back.

What is the most likely priority nursing diagnosis in this case?

- A. Activity intolerance
- B. Impaired skin integrity
- C. Ineffective cerebral perfusion
- D. Imbalanced nutrition less than the body requirement

ANSWER: C

52. A 35-year-old known psychiatric female patient came to a hospital with complaints of fatigue, hair loss, brittle, nails, and dry skin for the last six months. She also reported that she had an intentional weight gain and cold intolerance. A physician diagnosed severe hypothyroidism.

What is the most likely additional finding required to diagnose this case?

- A. Sleep apnea
- B. Deafness
- C. Slow speech
- D. Amenorrhea

ANSWER: A

53. An assessment finding of a 72-year-old male known diabetic patient revealed a drop in systolic blood pressure, weak and rapid pulse rate, and blood glucose level of 312mg/dl.

What is the most likely indicator to assess the severity of metabolic acidosis?

- A. Plasma osmolarity
- B. Urine osmolarity
- C. Urine ketone level value
- D. Serum level of bicarbonate

ANSWER: D

54. A 34-year-old female patient is to undergo a sub-total thyroidectomy. A physician has prescribed iodine solution to be taken for pre-operatively.

What is the most likely indication for the intervention?

- A. To minimize post-operative blood loss
- B. To decrease the signs of hyperthyroidism
- C. To reduce the heart rate and blood pressure
- D. To decrease the serum concentration of calcium

ANSWER: A

55. A 45-year-old female client arrived at a hospital with complaints of numbness, tingling, and cramps in the upper and lower extremities. Upon physical examination, a nurse notes that she has positive Trousseau's sign and Chvostek's sign.

What is the most likely information should the nurse include in the patient's teaching plan?

- A. To take a diet high in calcium and low in phosphorus.
- B. To take a diet low in calcium and low in phosphorus.
- C. To take a diet low in calcium and high in phosphorus.
- D. To take a diet high in calcium and high in phosphorus.

ANSWER: A

56. A 45-year-old female comes to an endocrine clinic with complaints of hot intolerance, irritability, and nervousness accompanied by difficulty in breathing. Upon assessment, bilateral smooth anterior neck swelling, bulging eyes, and startled facial expression were noted.

What is the most appropriate initial medical management in this case?

- A. Radioisotopes
- B. Propranolol
- C. Methimazole
- D. Levothyroxine

ANSWER: C

57. A 45-year-old known diabetic and hypertensive patient was admitted to an emergency department. Physical examination shows signs of dehydration and hypovolemia. Laboratory investigation also shows a blood glucose level of 345mg/dl, bicarbonate level of 10.5mEq/L, and serum pH of 6.7.

What is the most appropriate fluid to be used by the nurse in the initial phase of therapy?

- A. 0.9 percent normal saline
- B. 0.45 percent normal saline
- C. Dextrose 5 percent in water
- D. Dextrose 50 percent in water

ANSWER: B

58. A 28-year-old male client came to a hospital for an eye examination to register himself to a driving license school. A nurse is in charge of assessing the visual acuity of the man by Snellen's chart. The visual acuity score is 20/70 on Snellen's chart.

What is the most likely interpretation of the finding in this case?

- A. This patient can be labeled as legally blind
- B. This patient has visual impairment of category 3
- C. The patient saw the letter at 20 feet; a normal person can see it at 70 feet
- D. A normal eye can see at a distance of 20 feet; the patient can see it at 70 feet

ANSWER: C

59. A 12-year-old male student presented to a hospital with complaints of eye pain and tearing of the eyes for two weeks. Upon assessment, inward growth of eyelashes was seen. The physician diagnosed trachoma.

What is the most likely stage of the disease in this case?

- A. Trachoma intense
- B. Trachoma scarring
- C. Trachoma trichiasis
- D. Trachoma folliculitis

ANSWER: C

60. A 65-year-old known diabetic patient came to a hospital with complaints of pain in and around eyes, and blurred vision. He has decreased visual acuity and visual field. He has also nausea and vomiting. The tonometer reading is 28mmHg.

What is the most appropriate priority nursing diagnosis in this case?

- A. Acute pain
- B. Impaired vision
- C. Fluid volume deficit
- D. Disturbed sensory perception

ANSWER: A

61. A 25-year-old male patient came to an eye care center with a complaint of a foreign body in the eye. A nurse planned to perform eye irrigation.

What is the most appropriate solution to irrigate the eye in this case?

- A. Warm tap water
- B. Isotonic solution
- C. Hypotonic solution
- D. Hypertonic solutions

ANSWER: B

62. A 37-year-old known hypertensive patient came to an eye care center with complaints of blurred and distorted vision. He cannot resist bright light. He does not have pain. Upon assessment, the pupil is milky white.

What is the most likely diagnosis in this case?

- A. Glaucoma
- B. Cataract
- C. Keratoconus
- D. Trachoma

ANSWER: B

63. A long-distance truck driver is found to be HIV positive. After post-test counseling, a nurse working in the unit wants to identify his disease status.

What is the most likely indicator of an advanced HIV disease in this case?

- A. CD4 cell count greater than 500cells/mm³
- B. CD4 cell count greater than 200cells/mm³
- C. CD4 cell count greater than 200cells/mm³
- D. WHO stage 1 and 2

ANSWER: C

64. A 25-year-old female client is admitted to a medical ward with a diagnosis of stage 3 HIV/AIDS. On assessment, she has a history of nausea and vomiting, diarrhea, dysphagia, and odynophagia. Upon physical examination, there is dry oral mucosa, BP is 50/40 mmHg, PR is 110 beats/minute, RR is 24 breaths/minute, and temperature is 36.8 degrees centigrade.

What is the most likely nursing diagnosis?

- A. Imbalanced nutrition: less than body requirements
- B. Altered thermoregulation
- C. Fluid volume deficit
- D. Knowledge deficit

ANSWER: C

65. A 20-year-old female client who is diagnosed with rheumatic fever is administered procaine penicillin via the intramuscular route. After two minutes, the client manifests severe dyspnea, cyanosis, and hypotension.

What is the most likely diagnosis in this case?

- A. Dermatitis Medicamentosa
- B. Anaphylactic reaction
- C. Angioneurotic edema
- D. Allergic rhinitis

ANSWER: B

66. A 25-year-old HIV-positive client came to an emergency OPD with complaints of sudden onset of productive cough, chest pain, fever, and shortness of breath. Laboratory investigations reveal a negative gene X-pert result and a right lower lobe consolidation on the X-ray.

What is the most likely opportunistic infection in this case?

- A. Pulmonary TB
- B. Bacterial pneumonia
- C. Pneumocystis pneumonia
- D. Extrapulmonary TB

ANSWER: B

67. A 44-year-old female client, who has been on ART for the last two years, has a significant weight reduction as compared to previous measurements. A nurse working in the unit wants to assess her adherence to ART regimens.

What is the most appropriate monitoring mechanism in this case?

- A. Pharmacy refill record monitoring
- B. Monitoring by self-report
- C. Viral load monitoring
- D. Monitoring by pill count

ANSWER: C

68. A 25-year-old woman is admitted to an emergency ward with the diagnosis of anaphylaxis. Upon assessment, she has a blood pressure of 60/50 mmHg, pulse rate of 120 beats/minute, and a respiratory rate of 28 breaths/minute.

What is the most likely response to this treatment?

- A. Increased blood pressure
- B. Increased respiration rate
- C. Decreased urine output
- D. Increased pulse rate

ANSWER: A

69. A 30-year-old male client came to an emergency OPD with complaints of headache, weakness, joint pain, chills, and fever. He has a history of travel to a low land area, and he has intermittent fever. Physical examination showed blood pressure of 70/50mmHg, pulse rate of 100 beats/minute, respiratory rate of 20 breaths/minute and temperature of 39.5 degrees centigrade.

What is the most likely initial management in this case?

- A. Imaging studies
- B. Administering Anti pain
- C. Administering antipyretic
- D. Laboratory Investigation

ANSWER: C

70. A 25-year-old female client is admitted to an emergency unit with a diagnosis of cholera. She has a history of watery diarrhea and vomiting for the last 12 hours. Assessment findings show sick-looking, dry oral mucosa and sunken eyes. Her vital signs showed a blood pressure of 50/40mmHg, pulse rate of 110 beats/minute, respiratory rate of 25 breaths/minute and temperature of 37.8 degrees centigrade.

What is the most likely priority nursing intervention needed for this client?

- A. Fluid administration
- B. Blood transfusion
- C. Administer antibiotic
- D. Administer anti helminths

ANSWER: A

71. An 18-year-old male client comes to an adult OPD with complaints of fever, headache, and vomiting for two days duration. His fever and headache are increasing continuously. He has abdominal pain and diarrhea. Upon physical examination, his vital sign are stable. Laboratory investigation results indicate Hgb of 14gm/dl, reactive widal test for H & O antibody.

What is the most likely diagnosis in this case?

- A. Malaria
- B. Typhus
- C. Typhoid fever
- D. Relapsing fever

ANSWER: C

72. An adolescent male client came to a hospital with complaints of sudden onset fever, headache, shaking, chills, and sweating. On laboratory investigation, spirochetes are detected from the blood sample.

What is the most likely diagnosis of this case?

- A. Typhoid fever
- B. Typhus fever
- C. Relapsing fever
- D. Malaria

ANSWER: C

73. A 28-year-old woman is admitted to a medical ward with severe malaria. A nurse working in the medical ward assessed her for hypoglycemia, anemia, and convulsion. Laboratory results showed Hgb of 4gm/dl and random blood sugar of 70mg/100ml. The nurse administered quinine and two units of blood as per prescription.

What is the most appropriate indicator of prognosis in this case?

- A. Hemoglobin level
- B. Body fluid status
- C. Blood glucose level
- D. Body temperature level

ANSWER: A

74. A four-year-old boy, who was diagnosed with malnutrition, was admitted to a pediatric ward with a complaint of respiratory distress. Physical examination shows swollen and pale skin, and decreased subcutaneous tissue in the legs, arms, buttocks, and face. The abdomen is distended and tender on palpation. A vital sign shows a respiratory rate of 56/minute, a heart rate of 122 beats per minute, and an axillary temperature of 36.8 degree centigrade.

What is the most likely additional finding to be used to diagnose malnutrition in this case?

- A. MUCA 12.5cm
- B. WFH greater than or equal to -2z score
- C. Tachypnea
- D. Tachycardia

ANSWER: C

75. A three-year-old child boy is admitted to a pediatric ward due to complicated severe acute malnutrition for one week. He is on F-75 therapeutic milk and taking it through a nasogastric tube. He has abdominal distension and weight gain of more than 12g/kg/ day.

What is the most appropriate treatment phase for this patient?

- A. Phase I
- B. Transition Phase
- C. Phase II
- D. Outpatient

ANSWER: A

76. A two-year-old child girl is brought to a pediatric outpatient clinic with her mother due to concern about poor growth and development compared with the same child's age. Her weight and height fall below the normal range. A nurse gave health education about good feeding practices and appointed the child after two weeks.

What is the most likely expected outcome that indicates good progress on the appointment date?

- A. Weight gain
- B. Increasing height
- C. Increasing chest circumference
- D. MUAC 11.4

ANSWER: A

77. A four-year-old girl is admitted to a pediatric ward with severe acute malnutrition. She has visible wasting. A nurse gave an appropriate therapeutic feeding, antibiotic, and ReSomal solution based on a physician's order. On the next day, her vital signs showed a temperature of 38 degree centigrade, a respiratory of 38 breaths/minute, and saturation of 90 percent.

What is the most likely typical sign that indicates complicated SAM in this case?

- A. MUAC 11.2cm
- B. Edema (3+) of both feet
- C. WFH less than -3Z-score
- D. WFH less than -3Z-score

ANSWER: C

78. A five-year-old male child was admitted to a pediatric ward after he was diagnosed with severe acute malnutrition. He is conscious, has a sunken eye ball, and manifested with the signs of wasting. His vital signs showed a temperature of 37.8 degree centigrade, oxygen saturation of 84 percent, a respiratory rate of 58 breaths per minute, and a pulse rate of 122 per minute.

What is the most likely priority nursing care should be given by the nurse in this case?

- A. Antibiotic
- B. Oxygen
- C. Anti-pain
- D. Intravenous fluid

ANSWER: B

79. A seven-year-old male neonate whose birth weight is 2300 grams and born at 37th week of gestation is brought to an under five unit by his mother with a complaint of difficulty of feeding. The vital signs show a body temperature of 36.6⁰C, respiratory rate of 68 breaths per minute, and 82 percent oxygen saturation. The neonate is admitted to the neonatal intensive care unit (NICU) with a diagnosis of neonatal sepsis.

What is the most likely finding that made the neonate to be admitted to the NICU in this case?

- A. Low birth weight
- B. Hypothermia
- C. Hyperthermia
- D. Respiratory distress

ANSWER: D

80. A one-hour-old newborn male child whose gestation is 35 weeks with a birth weight of 2300 grams is transferred to post-natal ward. A nurse who is assigned to newborn child care evaluated the infant. He is sucking well, active, and temperature is in the normal range on the evaluation.

What is the most likely diagnosis in this case?

- A. Extremely preterm and very low birth weight
- B. Very preterm and low birth weight
- C. Preterm and low birth weight
- D. Preterm and normal birth weight

ANSWER: C

81. A one-day-old newborn child with cleft lip and palate is admitted to a neonatal ward. The baby has breastfeeding difficulty and vomiting following every breast milk feeding. Otherwise, all vital signs are in the normal range.

What is the most appropriate nursing diagnosis in this case?

- A. Infection related to malformation
- B. Effective Infant Feeding Pattern related to the defect
- C. Ineffective Infant Feeding Pattern related to the defect
- D. Aspiration related to tongue and palate deformities

ANSWER: C

82. A five-day-old infant is brought to a hospital with protrusion of the spinal cord and meninges through a defect in the spinal cord. A nurse conducted an assessment and diagnosed it as myelomeningocele. A nurse covered the defect with a sterile saline-soaked dressing, and she gave the antibiotic based on the order and plans to identify complications.

What is the most likely complication in this in this case?

- A. Meningitis
- B. Hyperthermia
- C. Feeding difficulty
- D. Neonatal jaundice

ANSWER: A

83. A 38-week-baby of gestational age was born from the primigravida mother at a hospital. The baby did not cry immediately after birth and breathing poorly with poor muscle tone. A nurse assists in keeping warm, dry, stimulate, and clears secretions from the mouth and the nose. The newborn still has not started to breathe. Thus, the nurse wanted to provide ventilation.

What are the most likely primary indicators to be evaluated during resuscitation in this case?

- A. Respiration, heart rate, and color
- B. Heart rate and oxygen saturation
- C. Respiration, temperature, and color
- D. Temperature, heart rate, and saturation

ANSWER: A

84. A 15-month-old male baby is brought to an EPI unit with his mother for the second dose of measles vaccine (MCV2). He took the first dose of the measles vaccine five months ago. A nurse is preparing the vaccine dose and the right site.

What is the most appropriate dose, site, and route of the MCV2 for this child respectively?

- A. 0.5ml, Left upper arm and Subcutaneous route
- B. 0.5 ml, Right deltoid, and subcutaneous route
- C. 0.5 ml, Right deltoid, and intramuscular route
- D. 0.5 ml, Left anterolateral thigh and intramuscular route

ANSWER: B

85. A four-day-old infant was transferred from a maternity ward to an immunization clinic for vaccination. A nurse identified that there was no vaccination given for this infant previously and she gave him BCG and Oral polio zero vaccines. She scheduled the infant for PENTA, PCV, ROTA, and OPV.

What is the most appropriate age of the infant for next vaccine in this case?

- A. After 4 weeks of BCG given
- B. At 4 weeks of postnatal age
- C. At 6 weeks of postnatal age
- D. At 8 weeks postnatal age

ANSWER: C

86. A 50-day-old male infant was brought to a pediatric outpatient clinic by his mother due to irritability and high-grade fever, diarrhea, and intermittent vomiting. His body temperature is 38 degrees centigrade and others are in normal range. The mother tells a nurse that the infant received his second dose of vaccines two days ago.

What is the most likely vaccine that causes the above adverse effects?

- A. OPV
- B. Penta vaccine
- C. Rota vaccine
- D. PCV

ANSWER: B

87. A nurse works in a district health center where polio, BCG, measles, DPT+hepB, PCV, and TT vaccines are found in the store. She put the heat-sensitive vaccines in the coldest area in the refrigerator. What is the most likely heat-sensitive vaccine to be stored in the coldest area of the refrigerator?

- A. Polio vaccine
- B. Polymerase conjugated vaccine
- C. TT vaccine
- D. DPT+hepB vaccine

ANSWER: A

88. A 15-month-old child was brought to a health center for vaccination. A nurse identified that the child has already completed all vaccines as per the schedule.

What is the most likely vaccine that the child is eligible for right now?

- A. OPV
- B. Rotavirus vaccine
- C. DPT-Hib-HepB 3
- D. Measles containing vaccine(MCV2)

ANSWER: D

89. A three-year-old male child was brought to a health center by his mother with complaints of cough and shortness of breath for two days duration. He has a loss of appetite and increased body temperature. The vital signs indicated a body temperature of 38.5 degrees centigrade and a respiratory rate of 44 breaths per minute while the oxygen saturation is 87 percent. Severe chest in drawing and grunting were noted on the examination.

What would be the most likely typical sound report on auscultation in this case?

- A. Wheezing
- B. Crackles
- C. Stridor
- D. Bronchial

ANSWER: B

90. A seven-year-old male child known RVI patient was on ART for the last six months. Currently, he is admitted to a hospital with complicated pulmonary TB. His family reported that the child interrupted ART drugs for two months after he started Anti-TB at a health center. He is undernourished on examination. After three days of treatment at the hospital, the child is discharged to follow treatment at home.

What is the most appropriate counsel to be offered before discharging the patient?

- A. Balanced nutrition
- B. Adherence to ART and anti-TB drugs
- C. TB & HIV transmission
- D. Schooling

ANSWER: B

91. While a medical ward nurse was caring for a 12-year-old girl with pulmonary tuberculosis, the father informed the nurse that there is a similar case in their neighbor. The father is worrying about his children because he feels that they could acquire TB and he asks the nurse the way he can decrease the chance of acquiring this disease.

What is the most appropriate precautionary method the nurse should plan to teach?

- A. Airborne
- B. Droplet
- C. Contact
- D. Hygiene

ANSWER: A

92. A 12-year-old child was admitted to a surgical ward for post-operative care of tonsillectomy for five days. While the child was taking soup, the child suddenly developed choking secondary food particles inhaled to the airway and the nurse was called for help.

What is the most appropriate first action to be done to manage the child?

- A. Suck the airway
- B. Put on oxygen
- C. Encourage to cough
- D. Apply Heimlich maneuver

ANSWER: C

93. A pediatric emergency head nurse identified the knowledge and skill gap of newly recruited nurses working in an emergency unit. The nurse planned to give short-term training on the principles and techniques of cardiopulmonary resuscitation. He recommended to nurses that severely ill children should be rapidly evaluated in a deliberate sequence.

What is the most appropriate sequence the nurses should follow in the principle of CPR?

- A. Circulation-Drug-Airway-Breathing
- B. Breathing-Airway-Drug-Circulation
- C. Airway-Breathing-Circulation-Drug
- D. Drug-Airway-Breathing-Circulation

ANSWER: C

94. A seven-year-old child presented to a hospital at midnight with complaints of dry cough and shortness of breath associated with chest tightness. Cyanosis, nasal flaring, and wheezing are noted while a body temperature of 37.8 degrees centigrade, respiration rate of 41 breaths per minute, and 87 percent oxygen saturation are recorded. The nurse diagnosed the child with severe pneumonia and immediately put the child on oxygen therapy.

What is the most likely indication for immediate oxygenation in the above case?

- A. Wheezing
- B. Peripheral cyanosis
- C. Hypoxia
- D. Chest tightness

ANSWER: C

95. A six-year-old boy was brought to a pediatric OPD with a complaint of extreme fatigue. His parents reported that this child can't cope with playing with his friends. He also has shortness of breath. His vital signs showed a body temperature of 36.5 degrees centigrade, pulse rate of 142 beats per minute, and his blood pressure was elevated between 90th and 95th percentiles. During examination, he has bilateral pitting edema on his feet and ankle. Auscultation shows diffused crackles. He is diagnosed with congestive heart failure.

What is the most likely additional typical feature that supports the diagnosis in this child?

- A. Hematuria
- B. High-grade fever
- C. Unable to walk up the stairs
- D. Weight loss

ANSWER: C

96. A 14-year-old patient presented to a pediatric outpatient department with a complaint of shortness of breath. Upon examination, severely swollen joints and subcutaneous nodules without edema were identified. The patient diagnosed with arthritis and responded quickly after the administration of oral aspirin. He is also positive for maneuver milkmaid's grip and confirming Sydenham chorea. Laboratory result showed antistreptolysin O antibody titer positive.

What is the most likely expected long-term complication if the case is poorly managed?

- A. Acute glomerulonephritis
- B. Congestive Heart failure
- C. Pulmonary tuberculosis
- D. Bacterial skin infections

ANSWER: B

97. A 13-year-old male child is presented to a health facility with complaints of history of anorexia, fatigue, and malaise accompanied predominantly by severe bone pain in the lower extremities that awake the child at night. Upon examination, pallor, purpuric and petechial skin lesions, lymphadenopathy, and splenomegaly were noted. The child is suspected to have leukemia.

What is the most likely typical clinical feature missing from the above report?

- A. Epistaxis
- B. Low grade fever
- C. High level of platelets
- D. High level of white blood cells

ANSWER: A

98. A 12-year-old child with a history of difficulty of swallowing, high body temperature, and exudative patches in the upper roof of the soft palate is admitted to a pediatric ward due to high-grade fever and arthralgia. Upon examination, pea-sized subcutaneous lesions and main joints were edematous. What is the most likely risk factor for this child's problem?

- A. Chronic ear infection
- B. Recurrent pneumonia
- C. Recurrent Tonsillopharyngitis
- D. Recurrent lower respiratory tract infections

ANSWER: C

99. A five-year-old child with a history of recurrent upper respiratory tract infections gets tired with a small-scale exercise. On examination, a respiratory rate of 36 breaths/minute, pulse rate of 120 beats/minute and murmuring on auscultation were noted. The child also squats while playing and is suspected to have congenital heart disease.

What could be the most likely confirmatory clinical feature to reach at this diagnosis?

- A. Higher pulse rate
- B. Exercise intolerance
- C. Squatting during play
- D. Recurrent respiratory infection

ANSWER: C

100. A six-year-old boy presented to a health center with a complaint of acute abdominal pain of 10 hours duration that was gradually associated with anorexia, nausea, and vomiting. The child did not pass stool after he experienced the acute pain. He is lethargic. His body temperature is 38°C and his respiratory rate is 45 breaths per minute. A nurse suspected appendicitis based on the location of the pain by superficial palpation.

What is the most likely location of the pain in the above case?

- A. Left upper abdominal quadrant
- B. Right upper abdominal quadrant
- C. Left lower abdominal quadrant
- D. Right lower abdominal quadrant

ANSWER: D

101. A four-year-old boy was brought to a health center with a complaint of watery diarrhea of 36 hours duration with episodes of two to three times per day. A mother also reported that the child has intermittent abdominal cramps. A nurse identified that the child is not immunized for all vaccine-preventable diseases. His body temperature is 37.6 degree centigrade, respiratory rate count is 38 breaths per minute, and pulse rate is 120 beats per minute are recorded. The nurse diagnosed the child problem as acute gastroenteritis.

What is the most appropriate nursing intervention in this case?

- A. Antibiotics and ORS
- B. Ant protozoa and ORS
- C. Zinc and ORS
- D. Anti-parasite and ORS

ANSWER: C

102. A 12-year-old male child presented to a health center with acute and severe abdominal pain of 36 hours duration which began centrally and later shifted to the right iliac fossa. He was referred urgently to a hospital. A nurse recognized that the child had appendicitis. Upon examination, rebound tenderness, abdominal wall rigidity, paucity of body motion, and absent bowel sounds were noted. The laboratory studies also reveal a peripheral WBC count greater than 15,000 cells/mm³.

What is the most likely complication identified in this case?

- A. Peritonitis
- B. Perforation
- C. Shock
- D. Sepsis

ANSWER: A

103. A seven-year-old child was brought to a pediatric emergency with colic, intermittent abdominal pain, and crying while drawing the legs to the abdomen. A physician diagnosed as moderate intestinal obstruction secondary to intussusceptions. The physician preferred hydrostatic reduction to surgery and ordered the nurse to perform it. The nurse put the child on NPO status, and inserted a nasogastric tube that is connected to low suction to reduce intussuscepted intestine to monitor patient progress. If reduction is failed, the patient will be planned for surgery.

What is the most likely expected outcome for spontaneous resolution of the obstruction?

- A. Reduced in the colicky pain
- B. Passage of normal stool
- C. Decreased abdominal distention
- D. Aspiration of clear gastric content

ANSWER: B

104. A three-week-old newborn with a history of frequent vomiting was admitted to a hospital. The infant started to have forceful vomiting following breastfeeding three days ago. Along breastfeeding, the infant started intravenous fluid as preoperative care. A nurse anticipated some accidental problems associated with vomiting and counseled the mother to hold the baby in a proper position during vomiting.

What is the most likely nursing diagnosis in this case?

- A. Risk for aspiration
- B. Risk for fluid volume deficit
- C. Risk for nutritional deficit
- D. Risk for blood volume deficit

ANSWER: A

105. A 12-year-old female child was admitted to a hospital with impaired urinary illumination secondary to renal problems. A nurse took a detail history of the child. Upon physical examination, urinary catheterization was decided for the child. A nurse rules out as if there is no contraindication for catheter insertion.

What is the most appropriate position the nurse should select?

- A. Supine position with knees bent
- B. High flower's position with knees extended
- C. Prone position with knees bent
- D. Knee-chest position with knees extended

ANSWER: A

106. A four-year-old male child known nephrotic syndrome was admitted to a pediatric ward for the last one week. The child has generalized body edema and most of the time he is confined to a bed. A physician suspected steroid dependency nephrotic syndrome on close follow-up. Urine analysis reveals proteinuria (2+) and little red blood cell.

What is the most likely priority nursing intervention needed in this case?

- A. Observe the skin for signs of redness and cyanosis.
- B. Change position at least every two hours
- C. Encourage the child to perform exercise
- D. Restrict not to take high protein diet

ANSWER: B

107. A three-year-old male child was admitted to a pediatric ward with the diagnosis of nephrotic syndrome. The child developed generalized body edema for which the child stays in a bed most of the time. A mother reported that the urine output is decreased significantly over the last 24 hours. A nurse noted that there is a significant weight gain over the last two days.

What is the most appropriate decision to be taken by the nurse in this case?

- A. Increase intake to increase output
- B. Fluid restriction and enhance voiding
- C. Measure intake and output every 3 to 4 days
- D. Increase intake with the increasing weight

ANSWER: B

108. A seven-year-old boy was admitted to an emergency unit with a complaint of abdominal pain. The abdominal pain was accompanied by nausea and vomiting. He has a shallow and rapid breathing. He also has sunken eyes and a low capillary refill. A mother reported that he has frequent urination day and night. Based on the assessment findings, he is diabetic.

What another additional cardinal feature is most likely reported in this case?

- A. Weight gain
- B. Hypocalcemia
- C. Hematuria
- D. Glucosuria

ANSWER: D

109. A 14-year-old known diabetes Mellitus patient was admitted to a pediatric emergency unit with severe acquired pneumonia. Laboratory results revealed that blood glucose of 250mg/dl, ketone of 3+, decreased PH-value and elevated bicarbonate.

What is the most likely laboratory result that suggests severe diabetic ketoacidosis in this case?

- A. PH 7.20
- B. CO₂ level 13mEq/L
- C. PH 5mEq/L
- D. CO₂ level 26meq/L

ANSWER: C

110. A 12-year-old boy was brought to a health center for follow-up. A parent asks for advice on whether to let him play football the next day or not. A nurse sends laboratory requests to identify his level of RBS and ketone. The laboratory result of RBS is 250mg/dl, and ketone +.

What is the most appropriate nursing intervention in this case?

- A. Advice on postponing exercise
- B. Adjusting insulin amount prior to the exercise
- C. Advice on giving snacks between the game
- D. Change injection site to less active muscles.

ANSWER: A

111. A 12-year-old known diabetic mellitus child was admitted to a pediatric emergency unit with a complaint of abdominal pain, frequent urination, and thirst. A nurse takes the sample for further investigation. The laboratory investigations provide evidence of diabetic Ketoacidosis. After appropriate treatment is given, the glucose drops below 150 mg/dL.

What is the most likely appropriate nursing intervention in this case?

- A. Change to 10 percent DNS
- B. Increase fluid infusion rate
- C. Low insulin infusion rates
- D. Give a snack between the injections

ANSWER: C

112. A six-month-old child was brought to a pediatric emergency clinic with complaints of vomiting, fever, and generalized weakness. Upon vital signs assessment: the child's body temperature is 38-degrees centigrade, respiratory rate is 50 breaths per minute and oxygen saturation is 75 percent. A nurse identified that the infant had a history of seizures.

What is the most likely clinical manifestation of seizure in this case?

- A. Grunting & High-grade fever
- B. Generalized weakness
- C. Respiratory distress
- D. Jerking extremities

ANSWER: D

113. A 10-year-old unconscious boy with a history of high-grade fever, headache, and nuchal rigidity is admitted to a pediatric emergency unit of a hospital. A pediatric nurse examined the child for the Brudzinski sign that indicated involuntary flexion of the knees and hips after passive flexion of the neck while in the supine position. A nurse was ordered to take 10ml of blood for culture examination. She took 10ml of blood from one site of injection.

What is the most likely error the nurse made while collecting the sample in this case?

- A. Collecting high-volume blood for culture
- B. Using the same site for taking blood
- C. Taking blood for culture examination
- D. Not considering the contraindications

ANSWER: B

114. A two-year-old child was admitted to a pediatric ward with uncontrolled seizure disorder. She has repeated seizures lasting longer than 30 minutes. He also has a history of intermittent vomiting. Upon examination, cyanotic and grunting were noted. The body temperature is 37 degrees centigrade and 44 breathings per minute were recorded.

What is the most appropriate nursing diagnosis in this case?

- A. Fluid volume deficit
- B. Risk for fluid over load
- C. Risk for decreased cardiac output related
- D. Ineffective breathing pattern

ANSWER: D

115. A seven-year-old girl presented to a hospital with complaints of fever and acute pain in right lower leg for two days duration. The child has a poor appetite and irritability. The mother also reported that the child has intermittent chills. Her recorded axillary body temperature is 38.5 degree centigrade and other vital signs are stable. A nurse suspected septic arthritis.

What is the most likely typical feature used to confirm the above diagnosis?

- A. Refusal to move the affected limb
- B. Refusal to bear weight on leg
- C. Local tenderness
- D. Swollen and tender joint

ANSWER: D

116. A 10-year-old boy was admitted to a hospital for surgery secondary to chronic Osteomyelitis on the left lower leg. There is a purulent discharge from the affected site and the child's mobility is limited due to fear of pain. The child is febrile associated with the inflammatory process of the disease. The child undergoes surgery and starts antibiotics.

What are the most likely outcome criteria indicating improvement after two days of management in this case?

- A. Remaining as febrile
- B. Mobilizing the affected limb
- C. Severe pain of the affected limb
- D. Pus containing discharge

ANSWER: B

117. A 10-year-old male child was admitted to a hospital with chronic Osteomyelitis on the left tibia bone for which he was on treatment in the last three months. There was continued purulent discharge from the affected site for the last three weeks, which was started to be limited to 1-2 times per day since the fourth week of its duration. The affected limb becomes very thin and mild loss of function is observed. What are the most appropriate set of nursing intervention needed in this case?

- A. Encourage exercise, low calcium diet, administer antibiotics
- B. Daily wound care, high protein diet, administer antibiotics
- C. Wound care every other day, skin assessments once a day, encourage exercise
- D. Skin assessments once a day, wound care every other day, high calcium diet

ANSWER: B

118. A five-year-old girl was brought to a hospital with a complaint of severe acute joint pain. The pain limited the child's range of motion, overlying skin warmth; redness and fever are observed from the affected site. The child has history of trauma to the affected joint. A nurse listed two differential diagnoses of the child as 'Osteomyelitis and septic arthritis'. Then, the nurse confirmed the case as 'septic arthritis' after getting additional assessment findings.

What would be the most likely additional finding that supports the diagnosis in this case?

- A. Increased systematic white blood cell
- B. Turbid, purulent, cloudy synovial fluid
- C. Restriction of movement of the affected limb
- D. Swelling and tenderness of the affected site

ANSWER: B

119. A two-year-old child was brought to a pediatric outpatient by child's mother due to the abnormal skin manifestations. The onset of the lesion is vesicular which rapidly turned into a crusted plaque at multiple places on his face. The child is diagnosed with impetigo.

What is the most likely feature of lesion that supports the diagnosis in this case?

- A. Pruritic vesicles
- B. Honey-colored
- C. Papules with burrows
- D. Erythematous papules

ANSWER: B

120. A two-year-old male child was brought to a health center with complaints of erythematous and pruritic papules with linear burrows that initially started between his wrist and interdigital spaces. The eggs of the parasite are seen when the burrowed hand lesion is seen under the light microscopy.

What is the most likely diagnosis in this case?

- A. Scabies
- B. Cellulitis
- C. Eczema
- D. Impetigo

ANSWER: A

121. A four-year-old male child visited a health center with his mother with a complaint of night itchy skin rash. Physical examination shows erythematous, pruritic papules with linear burrows on his wrist and inter-digital spaces. The nurse identified it as scabies.

What is the most appropriate management in this case?

- A. Treating pruritus with anti-histamine
- B. Applying permethrin 5 percent to the entire body
- C. Treating close contact
- D. Advice washing clothes in hot water

ANSWER: B

122. A four-year-old boy, who used topical corticosteroids with moisturizers to treat dermatitis (eczema), was appointed to visit a hospital after a week. A nurse advised the mother on proper utilization of medication.

What is the most appropriate expected outcome indicating good progress in this case?

- A. Absence of lesions
- B. Degrees of swelling
- C. Severity of itching
- D. Decrease body temperature

ANSWER: C

123. A 12-year-old boy was admitted to a pediatric ward with complaints of cola-colored urination, chills, and mild swelling on his face which is typically observed during the morning time. Two weeks ago, he had a history of skin infection for which he did not receive adequate therapy. His body temperature is 37.9 degrees centigrade. Laboratory results showed positive antistreptolysin 'O' titer. What is the most likely complication the patient would develop in this case?

- A. Acute rheumatic fever
- B. Acute post-streptococcal infection
- C. Septic arthritis
- D. Urinary tract infection

ANSWER: B

124. A one-week-old male newborn was referred from a health center to a hospital with a complaint of severe hypothermia. The baby was admitted to NICU for the essential therapies. The gestational age of the child and the birth weight are 35 weeks and 1800 grams respectively. He is not feeding well. Upon the examination, he moved only when stimulated. His body temperature is 35.3 degrees centigrade and the respiration rate count is 62 breaths per minute.

What is the most appropriate disease classification of the neonate in this case?

- A. Critical illness
- B. Very severe disease
- C. Pneumonia
- D. Local bacterial infection

ANSWER: B

125. A two-month-old infant female child was brought to a health center by her mother due to shortness of breath and oral thrush for two days duration. The birth weight of the child was 3000 grams. Her baby has no history of danger signs. Her body temperature is 37 degrees centigrade and her respiration rate is 61 breaths per minute. Laboratory results revealed a positive HIV antibody test for her child. Her mother is on antiretroviral therapy.

What is the most appropriate HIV classification of the infant in this case?

- A. HIV infected
- B. HIV exposed
- C. HIV status unknown
- D. HIV infection unlikely

ANSWER: B

126. A mother brought her seven-day-old female newborn to a pediatric under-five unit with a complaint that her baby is unable to suck breast for the last 16 hours. He has no danger signs. Physical examination showed red, swelling, and pus discharging from the umbilicus. Her body temperature is 37.9 degrees centigrade, and her respiratory rate is 65 breaths per minute.

What is the most appropriate treatment for the neonate according to IMNCI?

- A. Give amoxicillin for five days
- B. Apply Chlorohexidine on the cord
- C. Advice mother on home care treatment
- D. Refer immediately after first dose of antibiotics

ANSWER: D

127. A two-year-old female child was brought to a pediatric outpatient department by her mother with a complaint of cough, which lasted for more than two weeks. Her mother reported that her baby stopped breast feeding three months ago. Laboratory results revealed the mother is positive for HIV meanwhile her baby's HIV antibody test is negative.

What is the most appropriate HIV status classification for the child in this case?

- A. HIV infected
- B. HIV exposed
- C. HIV infection unlikely
- D. HIV infection unlikely

ANSWER: C

128. A father brought his four-year-old male child to a health center with a complaint of shortness of breath for three days duration. Auscultation of the lungs shows diffuse wheezing during expiration. His respiratory rate is 39 breaths per minute with the visible chest in drawing. He has no history of TB contact. He has no other severe classification.

What is the most appropriate treatment for this child?

- A. Refer urgently to hospital
- B. Give amoxicillin for three days
- C. Soothe the throat with a safe remedy
- D. Give an inhaled bronchodilator for 5 days

ANSWER: D

129. A 19-year-old Primigravida woman visited a health center for ANC follow-up. She had a history of nausea, fatigue, loss of appetite, and amenorrhea since November 02, 2016 E.C. After conducting a comprehensive prenatal assessment, the nurse calculated the expected date of delivery (EDD) for the woman.

What would be the most likely calculated EDD for the woman?

- A. July 05/2016 E.C
- B. August 09/2016 E.C
- C. September 07, 2017 E.C
- D. October 02, 2017 E.C

ANSWER: B

130. A 25-year-old Primipara woman with a gestational age of 13 weeks referred from a health center to a hospital. She has a history of severe vomiting and difficulty of performing day-to-day activities. Physical examination showed a blood pressure of 90/55 mmHg, pulse rate of 110 beats per minute, respiratory rate of 20 breaths per minute, body temperature of 36.5 degrees centigrade, and decreased body weight by 2 kg. Laboratory investigation shows ketone +3 and RBS 90mg/dl.

What is the most likely priority nursing diagnosis in this case?

- A. Activity Intolerance
- B. Imbalanced Nutrition
- C. Deficient, Fluid Volume
- D. Imbalanced body temperature

ANSWER: C

131. A 30-year-old Multigravida woman with a gestational age of 36 weeks came to a health center with complaints of severe frontal headache, epigastric pain, and blurring of vision for the last one week. Physical examination shows BP of 160/120mmHg, PR of 85 beats per minute, Temperature of 37.2 degree centigrade, and RR of 14 breaths per minute. Depending on clinical features, a nurse suspects severe preeclampsia.

What is the most appropriate laboratory test required to make these diagnoses?

- A. Complete blood count
- B. Protein dipstick test
- C. Renal function test
- D. Liver function test

ANSWER: B

132. A 24-year-old primipara woman with a gestational age of 26 weeks came to a health center for her third ANC follow-up. During the abdominal examination, a nurse gently grasped the lower portion of the abdomen just above the symphysis pubis with his thumb and fingers to determine whether it was soft or firm.

What is the most likely Leopold maneuvering the nurse performed in this case?

- A. 1st Leopold Maneuver
- B. 2nd Leopold Maneuver
- C. 3rd Leopold Maneuver
- D. 4th Leopold Maneuver

ANSWER: C

133. A 26-year-old pregnant woman with a gestational age of 32 weeks came to a hospital with complaints of severe headache and epigastric pain for two days. Upon physical examination, a blood pressure of 150/115mmHg and respiratory rate of 16 breaths minute was detected. Laboratory investigation reveals a WBC count of 9×10^3 and proteinuria of +3.

What is the most appropriate management option for this woman?

- A. Expectant management
- B. Prepare the woman for CS
- C. Immediate delivery of the baby by SVD
- D. Management with magnesium sulphate

ANSWER: D

134. A 24-year-old HIV-positive woman came to a hospital with a complaint of pushing down pain for 12 hours. Upon physical examination, a nurse identified that the presentation is breech with six cm cervical dilation. The fetal heart rate is 140 beats/minute and the membrane is intact. After consulting an obstetrician, the woman was transferred to the operation room for Cesarean delivery.

What is the most likely indication for cesarean delivery?

- A. Age of the woman
- B. Fetal heartbeat
- C. Fetal presentation
- D. HIV infection status

ANSWER: C

135. A pregnant woman at her 39th week of gestation came to a labor unit with pushing down pain for three hours. She has no history of ANC and was never tested for HIV infection. Physical examination shows that the cervix is four cm dilated and the fetal heart rate is 140 beats per minute. A nurse conducted an HIV antibody test and the result became positive.

What is the most appropriate intervention for the mother?

- A. Initiate ART drug at early postnatal
- B. Provide co-trimoxazole prophylaxis for woman
- C. Provide INH prophylaxis to prevent TB infection
- D. Prepare for infant HIV testing immediately after delivery

ANSWER: A

136. A pregnant woman at her 41th week of gestation came to a health center with complaints of pushing down pain and perspiration for 10 hours. A nurse identified that full dilated cervix, vertex presentation, four uterine contractions in 10 minutes that lasted for 55 seconds, and a distended perineum by the head of the fetus. The nurse performed an episiotomy to assist the labor.

What is the most likely indication for the episiotomy in this case?

- A. Gestational age
- B. Cervical dilation
- C. Uterine contraction
- D. Distended perineum

ANSWER: D

137. A 34-year-old multipara woman gave spontaneous vaginal delivery at a primary hospital. The woman was discharged after 24 hours as everything was normal. On 4th day of her postnatal, she presents with generalized malaise, headache, nausea, vomiting, chills, severe pain, and tender swelling in one quadrant of the breast. Upon examination, a temperature of 38.5 degrees centigrade, presence of swelling in one breast and overlying skin is red, hot and tender were identified.

What is the most likely priority nursing diagnosis in this case?

- A. Impaired Home Maintenance
- B. Risk for Activity Intolerance
- C. Ineffective breastfeeding
- D. Self-Care Deficit

ANSWER: C

138. A 28-year-old multipara woman comes to a hospital with excessive vaginal bleeding after she gave birth at home three hours ago. The placenta is not delivered and the cord is visible outside the vagina. The vital signs show BP of 90/60mmHg, PR of 110 beats/minute, and RR of 22 Breaths/minute. A nurse makes a diagnosis of Postpartum Hemorrhage.

What is the most appropriate outcome criterion for the prioritized nursing diagnosis?

- A. Pain will be relived
- B. Placenta will be removed
- C. Vital signs will be stabilized
- D. Bleeding will be controlled

ANSWER: C

139. A 34-year-old known HIV-positive pregnant woman gave spontaneous vaginal delivery in the labor ward of a health center. A nurse had completed fetal evaluation, and the APGAR score was 7/10 and 10/10 at the 1st & 5th minute respectively.

What is the most appropriate next priority nursing intervention needed for HIV exposed infant?

- A. Provide HIV testing
- B. Provide cotrimoxazole
- C. Provide ART prophylaxis
- D. Provide vaccination as protocol

ANSWER: C

140. A nurse is providing essential newborn care for a baby who cried strongly at birth. After drying the baby with a dry towel, the nurse assessed the breathing and colors of the baby. Since breathing was normal, the nurse performed cord care.

What is the most appropriate next logical newborn care in this case?

- A. Apply chlorohexidine on the cord
- B. Apply tetracycline eye ointment
- C. Initiate breastfeeding
- D. Give vitamin K

ANSWER: C

141. A couple visited a health center for family planning services. Both of them are known HIV-positive and have been taking a recommended ART regimen containing protease inhibitors for the last one year. Following counseling on family planning options, both of them agreed to use Progestin-only pills (POPs).

What is the most likely additional contraceptive the couple needs to use in this case?

- A. Jadelle
- B. Female Condom
- C. Injectable contraceptives
- D. Intrauterine Contraceptive Device

ANSWER: B

142. A 22-year-old girl and her partner came to a health center with a complaint of sexual discomfort due to the IUCD string. The girl had inserted IUCD three months ago, and the nurse checked that IUCD was in place.

What is the most likely nursing intervention in this case?

- A. Remove and replace with a new method
- B. Cut the string to a level of the cervix then counsel
- C. Counsel that this is the natural course of the method
- D. Reassure that it's a sign of feeling appropriate insertion

ANSWER: B

143. A 20-year-old new COC user came to a health center with complaints of nausea and vomiting. A nurse reassures the client by saying "Many women use COC experience as a side effect and also it is not harmful since it is normal after the first few months of use."

What would be the most likely additional side effect the client experienced?

- A. Migraine Headache
- B. Suspected pregnancy
- C. Irregular vaginal bleeding
- D. Severe lower abdominal pain

ANSWER: C

144. A 23-year-old student came to a hospital with excessive vaginal bleeding for six days. She also has frequent period even twice per month. The girl is irritable and feels moody during her period. Her vital signs show a blood pressure of 100/65mmHg, pulse rate of 90 beats/minute, and normal body temperature.

What is the most likely intervention for this woman?

- A. Uterine curettage
- B. Routine perineal care
- C. Counsel on safe sex practice
- D. Hormonal therapy with COC

ANSWER: D

145. A 22-year-old primipara with gestational age of nine week came to gynecology OPD with complaints of vaginal bleeding followed by uterine contraction. A nurse suspected of inevitable abortion. What is the most likely typical clinical manifestation that suggested the diagnosis in this case?

- A. Dilatation of cervix
- B. Passage of blood and some tissue
- C. Passes all of the contents of the uterus
- D. Retention of a failed intrauterine pregnancy

ANSWER: A

146. A 21-year-old lady came to a health center with a complaint of excessive vaginal bleeding during the past three menstrual cycles. The woman is unable to identify her menstrual period in most of her cycle. She bleeds for an average of 10 days and the cycle is irregular.

What is the most likely pattern of menstruation in this case?

- A. Metrorrhagia
- B. Menorrhagia
- C. Olymenorrhea
- D. Menometrorrhagia

ANSWER: D

147. A 19-year-old unmarried girl came to a hospital with a complaint of abdominal bloating for three days. The girl is irritable and feels mood swings during her period. She feels restlessness and sleeplessness during few days before her menstruation. She said that she is in a state of inability to manage her condition.

What is the most likely nursing diagnosis in this case?

- A. Post-Trauma Syndrome
- B. Compromised Family Coping
- C. Ineffective Individual Coping
- D. Readiness for Enhanced Family Coping

ANSWER: C

148. An 11-year-old male child is brought to a psychiatric clinic with his mother. During the assessment, a health professional identified that the child faced difficulty of complying with parent's directions, problems with concentration, difficulties of following instructions in school, and the child has emotionally labile, explosive, lacks focus and is irritable.

What is the most likely diagnosis of this child?

- A. Attention Deficit/Hyperactive disorder
- B. Autistic Spectrum disorder
- C. Intellectual disability
- D. Learning Disorder

ANSWER: A

149. A known schizophreniform male patient came to a psychiatric clinic with an improved symptom of excessive fear of being harmed by his family. As a result, he isn't willing to take any foods that were prepared by his family. Because of this behavior, his social interaction has been significantly impaired.

What is the most likely type of delusional symptoms presented in this case?

- A. Delusion of Erotomania
- B. Delusion of Grandiosity
- C. Delusion of Persecutory
- D. Delusion of Jealousy

ANSWER: C

150. A 30-year-old single woman came to a psychiatric clinic with her older brother. The woman has urged to go out from home, talk alone, and assault people verbally and physically. During the interview, she is highly agitated, and non-cooperative for the past two months. She was saying that there is a person who is whispering in her ear to hurt the people around.

What is the most likely diagnosis in this case?

- A. Schizophrenia Disorder
- B. Schizoaffective Disorder
- C. Brief Psychotic Disorder
- D. Schizophreniform Disorder

ANSWER: D

151. A 30-year-old woman came to a psychiatric clinic with complaints of feeling depressed and inability to do daily activities. The woman has significant impairment on interpersonal interaction, is unable to sleep, and thought of self-killing for the last one month. A physician diagnosed her with Major Depression Disorder.

What is the most likely priority nursing care in this case?

- A. Working on sleep problem
- B. Working on Suicidal behaviors
- C. Working to alleviate a depressed mood
- D. working on interpersonal impairment

ANSWER: B

152. An 18-year-old male patient arrived at a psychiatry emergency unit with his relatives. The patient has an overt and aggressive behavior towards his relative as well as the health workers. He also tries to break objects and has self-destructive behavior. The nurse made a nursing diagnosis of "Risk for violence of self-directed and other-directed behavior related to catatonic excitement".

What is the most likely nursing intervention in this case?

- A. Observe the client's behavior after sometimes
- B. Provide frequent assistance with self-care needs
- C. Provide information for the family about the illness
- D. Maintain a low level of stimuli in the client's environment

ANSWER: D

153. After a month experiencing a Bajaj accident, a 20-year-old single female who was born into socio-economically deprived family patient came to a psychiatric clinic with symptoms of recurrent, involuntary, and intrusive distressing memories of the accident, and blaming herself for the accident. She has an inability to express her feeling, especially in conditions that make everyone happy.

What is the most appropriate stage of the traumatic event that predisposes to poor prognosis in this case?

- A. Pretraumatic factors
- B. Peritraumatic factors
- C. Posttraumatic factors
- D. Psychosocial factors

ANSWER: A

154. A 30-year-old female patient came to a psychiatry clinic with complaints of flashbacks, intrusive recollections, and nightmares for the last one month. The patient also has psychological numbness which is related to an event and de-realization. A physician diagnosed her with Acute Stress Disorder.

What is the most likely nursing diagnosis in this case?

- A. Stress overload
- B. Ineffective Coping
- C. Dysfunctional Grieving
- D. Post-trauma Syndrome

ANSWER: D

155. A 16-year-old woman who was raped by an unknown person came to a psychiatric clinic with her mother. She has a significant distressing feeling about the situation. As she explained to a nurse, she has a flashback related to the event and fear of discussing it with others.

What is the most likely prior intervention in this case?

- A. Assess for self-destructive ideas and behavior of a patient
- B. Encourage the client to talk about the trauma at her own pace
- C. Assess the impact of the trauma on the client's ability to resume regular activities
- D. Respect client's wishes regarding interaction with individuals of opposite sex

ANSWER: D

156. A 22-year-old woman who was raped by one of her family members was admitted to a hospital ward. On nursing assessment, the nurse found that the client has the feeling of being abandonment by others, impaired cognition fostering a negative view of self, and low self-esteem.

What is the most likely initial nursing intervention in this case?

- A. Accept the client and spend time with her
- B. Allow client to participate in goal setting regarding own care
- C. Reassure the client that she is not alone when feeling inadequate
- D. Help client identify areas of life situation that are under own control

ANSWER: A

157. A 22-year-old male who lives with his family came to a clinic for a psychiatric evaluation. Upon history, he used Cannabis for feeling relaxation on a daily base with a constant amount. However, today, he had smoked a lump of cannabis despite his friend's warning.

What is the most likely symptom the patient would currently experience?

- A. Tremors, insomnia, and seizures
- B. Inflated self-esteem and heightening
- C. Increased heart rate and blood pressure
- D. Depersonalization and Derealization

ANSWER: D

158. An 18-year-old male patient is brought to a rehabilitation center by his parent. The patient used Chat frequently; as a result, his interaction with the family is severely impaired. However, the patient states that he has no addiction of chat and strongly believes that he can stop anytime without therapy. He said that he only used Chat just for social pleasure.

What is the most likely priority nursing diagnosis in this case?

- A. Ineffective Denial
- B. Ineffective Coping
- C. Ineffective attitude
- D. Ineffective relationship

ANSWER: A

159. After conducting a thoroughly assessment, a nurse set an outcome criterion for a 21-year-old female student patient. The nurse stated the outcome criterion as "The patient will be able to verbalize adaptive coping mechanisms to use, instead of alcohol use for stress reliving in two months".

What is the most likely nursing intervention in this case?

- A. Give her positive reinforcement for her ability to delay gratification
- B. Explore options available to assist her with stress-related behaviors
- C. Acknowledge the association between personal problems and substance use
- D. Set limits on manipulative behavior and take measures when violated

ANSWER: D

160. A 65-year-old mother was brought to a clinic for the management of Neurocognitive disorder (NCD). The mother has faced significant difficulties in engaging with familiar activities and familiar environments. She is unable to use kitchen and feeding utensils properly. She has also difficulty of recognizing familiar faces.

What is the most likely type of NCD stage in this case?

- A. Mild type
- B. Moderate type
- C. Major type
- D. Unspecified type

ANSWER: C

161. A 92-year-old patient was admitted to a Geriatrics care center for three weeks. During the nursing round, a team found that he has difficulty of accurately interpreting the environment in which he was admitted, recognizes people around him and can only remembered things which were happened remotely. Based on their findings, the team set a nursing diagnosis as he has a disturbed thought process related to cerebral degeneration.

What is the most appropriate intermediate nursing plan in this case?

- A. The client will be interpreting the environment accurately and maintain orientation
- B. The client will accomplish activities of daily living to the best of his or her ability
- C. The client will not experience injury to the best of his ability not to harm
- D. The client will discuss positive aspects of self and life.

ANSWER: A

162. After taking a detailed assessment of a psychiatric patient, a nurse set a nursing diagnosis of "Risk for suicide related to depressed mood secondary awareness in the decline of physical capability". Taking the patient's current self-harm thought, the nurse set an outcome as the patient will not experience self-injury after getting an initial management.

What is the most likely nursing intervention in this case?

- A. Arrange furniture in the client's room
- B. Discourage rumination of delusional thinking
- C. Frequently orient the client to time, place, and person
- D. Keep explanations simple and use face-to-face interaction

ANSWER: A

163. A primary hospital manager had organized an annual staff meeting. At the meeting, the manager guides and encourages all unit heads to present their achieved results, challenges faced, and measures taken during the budget year. In addition, at the end of the meeting, staff members with outstanding achievements were recognized.

What is the most likely function of management performed by the manager in this case?

- A. Planning
- B. Controlling
- C. Staffing
- D. Leading

ANSWER: D

164. A regional health office organized a training on leadership and management for its experts working in different departments. The training is intended to establish logical, sequential, and special relations between activities, and functional and structural relations among people working in the health office.

What is the most likely management principle stressed in this training?

- A. Convergence of work
- B. Management by exception
- C. Shortest decision pathway
- D. Function determines structure

ANSWER: A

165. The Ethiopian Ministry of Health has introduced a new plan aimed at reducing maternal mortality which will be implemented throughout the country for the coming ten years.

What is the most likely type of plan expressed in this case?

- A. Contingency
- B. Strategic
- C. Tactical
- D. Operational

ANSWER: B

166. A manager of a health center has developed a SWOT analysis matrix and disseminated it to all department heads and staff for their reflection. The compiled analysis result revealed the presence of shortage of supply, poor allocation of finance, and shortage of trained personnel in all departments of the facility.

What is the most likely SWOT component revealed by this analysis result?

- A. Threat
- B. Strength
- C. Opportunity
- D. Weakness

ANSWER: D

167. A zonal health office has experience in purchasing and distributing required equipment like cotton, laboratory chemicals, paper, and syringes for the sectors under its supervision biannually. What is the most likely category of materials explained in this situation?

- A. Basic
- B. Expendable
- C. Compulsory
- D. Non-expendable

ANSWER: B

168. A head of a health office carried out an annual evaluation on the achievement status of planned activities by each of the staff members. The aim of the evaluation is to give feedback, recognize staff members who made outstanding achievements, locate the need for additional training, and identify candidates for promotion.

What is the most likely maintenance function of group behavior that focused on in this evaluation?

- A. Training and development
- B. Performance appraisal
- C. Coaching and controlling
- D. Separation and transfer

ANSWER: B

169. A department head has assigned supervisors for different units of the department. As a principle, one supervisor is supposed to receive a report only from a single unit head to prevent conflicting demands, conflicting priorities, and no-win situations between the units and the assigned supervisors.

What is the most likely principle of organizational concept applied in this case?

- A. Authority
- B. Decentralization
- C. Work specialization
- D. Unity of Command

ANSWER: D

170. A health facility manager held one-on-one discussion sessions with staff members to introduce the need to change less effective old behaviors, structures, and processes of the health facility. A potential advantage of the planned change is to increase the productivity of the facility.

What is the most likely change stage occurring in this situation?

- A. Unfreezing
- B. Movement
- C. Changing
- D. Refreezing

ANSWER: A

171. A head of a health center decided new perspective on activities related to personnel deployment, purchases, and specific work assignments.

What type of decision is mostly explained in this scenario?

- A. An End
- B. Operational
- C. Unstructured
- D. Non-programmable

ANSWER: B

172. A private clinic manager wants to identify all possible causes of service quality problems and its contributing factors in terms of people, process, equipment, environment, and management.

What is the most appropriate quality improvement tool utilized in this scenario?

- A. Run chart
- B. Flow chart
- C. Pareto chart
- D. Fishbone diagram

ANSWER: D

173. An estimated ANC coverage in a district was 86 percent. Despite this high coverage of ANC, only 56 percent of pregnant women gave birth in health institutions and 44 percent of them gave birth at home. Women who give birth at home are more prone to maternal and perinatal morbidity and mortality.

What is the most likely research problem that could be identified in this case?

- A. Identify contributing factors for high ANC coverage
- B. Identify contributing factors for health institution delivery
- C. Identify associated factors for maternal morbidity and mortality
- D. Identify contributing factors for home delivery

ANSWER: D

174. A researcher planned to assess the burden of bronchial asthma as well as its contributing factors among adults in a district after reviewing the prior community health problems. The data collection was done at a point in time.

What would be the most likely research objective of the study?

- A. To identify associated factors of bronchial asthma.
- B. To determine the incidence of bronchial asthma.
- C. To identify the treatment option of bronchial asthma
- D. To identify the diagnostic modality of bronchial asthma

ANSWER: A

175. A nurse wants to study the association between oral contraceptive use and breast cancer among women in reproductive age groups. Then, the nurse selected those women with breast cancer and an equal number of women without breast cancer and asked about their prior use of oral contraceptives.

What is the most likely study design used by the nurse in this case?

- A. Cohort
- B. Ecological
- C. Case-control
- D. Cross-sectional

ANSWER: C

176. A researcher studied the prevalence of HIV infection among rural and urban communities in a district. During the process of sampling, the researcher selected 300 samples equally from urban and rural communities.

What is the most likely sampling technique used by the researcher in this case?

- A. Quota sampling
- B. Purposive sampling
- C. Snowball sampling
- D. Convenience sampling

ANSWER: A

177. A nurse wants to assess measles vaccine coverage among under-five children in a district. The nurse decided to use an assumption of 95 percent confidence level, and a 5 percent margin of error to compute the number of children using a single population proportion formula.

What is the most likely missed criterion to determine sample size?

- A. Variance
- B. Proportion
- C. Standard deviation
- D. The mean of the sample

ANSWER: B

178. A researcher assigned a nurse to collect data on community-based health insurance utilization levels of the community in rural districts of a region. The nurse interviewed only people who were volunteers and agreed to take part in the study during the data collection process.

What is the most likely type of error committed by the nurse in this case?

- A. Recall bias
- B. Selection bias
- C. Information bias
- D. Observation bias

ANSWER: B

179. A researcher wants to explore the families' perception of their daughters' early relationship with male school students, which the researcher is not sure about the possible answer of the family.

What is the most likely data collection tool the researcher needs to prepare?

- A. Checklist
- B. Low flexible type of questions
- C. Closed-ended type of questions
- D. Open-ended type of questions

ANSWER: D

180. A 40-year-old patient came to a hospital emergency OPD with a complaint of cervical bone fracture secondary to a vehicle accident for 4 hours duration. After all the necessary assessment and investigation, a physician ordered analgesics until the CT scan result is accessible. Unfortunately, a nurse gives the wrong medication to the victim.

What is the most likely professional malpractice committed by the nurse?

- A. Burnout
- B. Negligence
- C. Disappointment
- D. Work overload

ANSWER: B

181. A 25-year-old male and a 20-year-old female came to a hospital with complaints of forearm fracture and shortness of breath respectively. A nurse tries to assess and sort out the patients' problems accordingly without any discrimination.

Which ethical principle is the most appropriately used to encourage the nurse to manage these cases?

- A. Fidelity
- B. Beneficence
- C. Nonmaleficence
- D. Respect for autonomy

ANSWER: A

182. A 60-year-old male patient who was admitted to a medical ward has just been diagnosed with diabetes mellitus. After the diagnosis, a physician ordered insulin until the serum sugar level was maintained. Before discharging, a nurse plans to teach how to inject himself and the sites to be injected. What is the most likely therapeutic communication applied by the nurse in this case?

- A. Simulation
- B. Demonstration
- C. Group instruction
- D. One-on-one discussion

ANSWER: B

183. A 60-year-old male patient presented to a surgical ward of a hospital with a diagnosis of large bowel obstruction scheduled for laparotomy surgery before a week. A surgical nurse was assigned as a scrub nurse at the operation room. The nurse believed that dignified care is very important and valuable which is creating an opportunity to develop self-esteem, self-worth, and modesty in every activity performed in the operation room.

What is the most likely dignified care component identified by the nurse in this case?

- A. Respect
- B. Autonomy
- C. Empowerment
- D. Communication

ANSWER: C

184. A 70-year-old female patient scheduled to have surgery due to her colorectal cancer, which was diagnosed two weeks ago. A surgical nurse has taken informed consent and introduced herself to take time to put the patient at ease and appreciate the patient's circumstances with respect and kindness.

What is the most likely determinant of health care provider exercised by the nurse in this case?

- A. Attitude
- B. Behavior
- C. Autonomy
- D. Communication

ANSWER: B

185. A medical nurse has been working in a medical ward for the last five years. As a nurse professional, he has a good understanding of legal boundaries, good communication skills, ethical principles, and values. He focused on demanding the best care for patients while delivering nursing care.

What is the most likely principle of professionalism identified by the medical nurse?

- A. Altruism
- B. Humanism
- C. Excellence
- D. Accountability

ANSWER: C

186. A 34-year-old pregnant woman came to a district hospital's obstetric ward for delivery. A duty nurse left the woman alone while she was in labor. In addition to this, there is a failure of a nurse to monitor a woman and to attend the labor when needed.

What is the most likely disrespectful character experienced by the duty nurse in this case?

- A. Non-consented care
- B. Abandonment of care
- C. Detention in facilities
- D. Non-confidential care

ANSWER: B

187. A 70-year-old male and 11-year-old female patient came to a chronic clinic for follow-up. Both are known as Diabetic patients. However, there is insulin shortage in the hospital, and currently, there is only one of vial insulin available in the hospital. A nurse should be able to pare the options down to a few realistic alternative actions and other relevant issues need to be examined while attempting to weigh the alternative action.

What is the most appropriate step used by the nurse to solve an ethical dilemma?

- A. Consider the choices of possible courses of action
- B. Decide on differences opinion action
- C. Collect, analyze, and interpret the data of each course of action
- D. Analyze the advantages and disadvantages of each course of action

ANSWER: D

188. A 40-year-old woman was admitted to a surgical ward after performed cholyolithiasis surgery since three days ago. A surgeon ordered wound care two times a day with aseptic techniques. A duty nurse is performing the wound care with normal saline 0.9 percent by wearing examination gloves. However, she has a poor wound healing process.

What is the most appropriate activity needed to be breached by the duty nurse in this case?

- A. Negligence
- B. Malpractice
- C. Responsibility
- D. Accountability

ANSWER: B

189. A 40-year-old woman was admitted to a surgical ward after performing large bowel obstruction surgery two days ago. A surgeon ordered Ceftriaxone 1gram intravenously two times a day. A duty nurse changed a drug to Cloxacillin and administered 500 milligrams two times a day.

What is the most likely liability experienced by the duty nurse?

- A. Negligence
- B. Malpractice.
- C. Accountability
- D. Responsibility

ANSWER: A

190. A 23-month-old child was brought by his mother to the under-5 OPD with complaints of loss of appetite and growth problems. A nurse in charge suspects the child's problem is mild acute malnutrition. What is the most appropriate nutritional assessment method to be used to diagnose this child?

- A. MUAC
- B. Weight for age
- C. Height for age
- D. Recent weight loss

ANSWER: A

191. A researcher in a regional health bureau wants to identify institutions that serve the population subgroups of interest. During their review, they want to obtain data regarding a disease within specified period.

What is the most likely type of surveillance preferable for this case?

- A. Active surveillance
- B. Sentinel surveillance
- C. Passive surveillance
- D. Mixed surveillance

ANSWER: B

192. A 23-year-old woman was admitted with labor pain to a delivery room of a health center. During the vaginal examination, a nurse on duty found that there was partial excision of labia majora together with total excision of the clitoris and vaginal inner lip.

What is the most likely type of female genital mutilation observed in this case?

- A. Type I
- B. Type II
- C. Type III
- D. Type IV

ANSWER: C

193. A two-year-old child came with his mother to the under-five OPD with complaints of growth problems and frequent illness. During the physical examination, a nurse found that there was visible severe wasting and an observable break in the surface of the transparent layer that covered the pupil and iris of both eyes. Weight for height indicates 70 percent.

What is the most likely vitamin-A deficiency examined in this case?

- A. Bitot's spots
- B. Corneal scar
- C. Corneal clouding
- D. Corneal ulceration

ANSWER: D

194. A nurse working in a prison clinic identified an increased in the number of patients' who showed similar sign and symptoms. The clinical manifestations shown by the prisoners were prolonged fever, headache, anorexia, chills, malaise, abdominal pain, and diarrhea. During history taking, the nurse found that the prisoners developed this after consuming foods that were prepared within the prison's cafeteria. What is the most likely disease manifested in these prisoners?

- A. Shigellosis
- B. Salmonellosis
- C. Typhoid fever
- D. Food Poisoning

ANSWER: C

195. A nurse is assigned as a malaria officer in a district. The nurse designed a malaria prevention and control mechanism. This mechanism includes the discharging of stagnation of water that serves as a mosquitoes breeding site to limit the breeding of mosquitoes.

What is the most likely appropriate vector control measure applied by the nurse in this case?

- A. Use of chemical
- B. Basic sanitation
- C. Physical measure
- D. Biological method

ANSWER: B

196. A nurse who works in a school clinic always faces cases from the students like skin infections, abdominal discomfort, eye infections, gastroenteritis, and tonsillitis. Hence, the nurse plans to control those cases by applying environmental health services to the school.

What is the most likely environmental health service need to be applied by the nurse in this case?

- A. Nutrition
- B. Physical exercise
- C. Mental health service
- D. Personal hygiene and sanitation

ANSWER: D

197. A student nurse is working as a team member of community outreach service in a high school which is found in the catchment area. During the health needs assessment, the student nurse and director of the high school reported that most of the students engaged in unprotected sexual intercourse and consequently some of the girls quit their education.

What is the most likely health problem faced by high school girls in this case?

- A. Behavioral
- B. Mental health
- C. Socioeconomic
- D. Communicable diseases

ANSWER: A

198. A nurse who is working in a primary hospital attends a health education session in the patient waiting area. During the session, the nurse selects one of the top ten diseases in the hospital and explains the ideas using medical words frequently. At the end of the session, the nurse asked the audience about any unclear issues, but the audience refused to respond.

What is most likely principle of health education missed by the nurse in this case?

- A. The health educator has to make himself acceptable
- B. It is necessary to discuss freely on health problems
- C. The educational process has to take into account the learning process
- D. Health educator has to adjust his talk and action to suit the group

ANSWER: D

199. A 40-year-old woman came to a health center with complaints of headache and blurring of vision. An adult OPD staff, who was on duty and very busy shouted at her. Even one health care provider stated "We don't want to waste our time to the complaint of headache please come back if it worsens and when we are less busy". The woman does not turn back to the health center despite her illness being worsened.

What is the most likely predisposing factor that affects the woman's behavior?

- A. Value
- B. Belief
- C. Norms
- D. Attitude

ANSWER: D

200. A health center in collaboration with a woreda health office developed a poster that shows the benefits of health facility delivery for the mother and her newborn baby. The prepared poster was posted in front of the MCH clinic. However, most of the clients at the MCH clinic area walked and passed the poster without paying attention to it.

What is the most likely stage of communication is failed in this case?

- A. Stage 1
- B. Stage 2
- C. Stage 3
- D. Stage 4

ANSWER: B