

Prepare for Exit Exam! Join our Telegram Channel for all Department
@ExitExamPrep

NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:03

Question 4

Not yet
answeredMarked out of
1.00Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Prepare for Exit Exam! Join our Telegram Channel for all Department
@ExitExamPrep

Question 5

Not yet
answeredMarked out of
1.00Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Prepare for Exit Exam! Join our Telegram Channel for all Department
@ExitExamPrep

Next page

Lenovo

Ac63

Time left 1:14:22

Question 81

Not yet
answered

Marked out of
1.00

Flag question

A clinical nurse gave health education to a community about prevention mechanisms of infectious disease. Then, all the community members practiced the prevention approaches of infectious diseases each day. The health status of the community was maintained. Finally, the community members gave a credit for the primary prevention. What is the most appropriate communication stage for this scenario?

- ☐ a. Attracting the audience of attention
- ☐ b. Understanding the message
- ☐ c. Reaching the intended audience
- ☐ d. Improving to health

Question 82

Not yet
answered

Marked out of
1.00

Flag question

A 28-year-old woman came to a gynecology OPD with complaints of fever and vaginal discharge of one week duration. She had offensive vaginal discharge and high grade fever. Physical examination revealed the presence of lower abdominal tenderness, cervical motion tenderness, adnexal tenderness and pelvic mass. Laboratory finding suggested leukocytosis of $11,500/\text{mm}^3$. What is the most likely diagnosis in this case?

- ☐ a. Torsion of ovarian cyst
- ☐ b. Vulvovestibulitis
- ☐ c. Ectopic pregnancy

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:43:44

Question 36

Not yet
answeredMarked out of
1.00

Flag question

A biostatistics instructor wanted to assess the daily performance of nursing students and has given quizzes on different days. The result is distributed in descending order 10, 9, 8.5, 8, 8, 7.5, 7, 7, 6, 6, 5.5, 5.5, 4, 4, 4, 3, 3 and 2. What is the mean of the students' result in this case?

- ☐ a. 6
- ☐ b. 8
- ☐ c. 5.5
- ☐ d. 7

Question 37

Not yet
answeredMarked out of
1.00

Flag question

A 28-month-old young male child was brought to an emergency unit due to difficulty of breathing and coughing that happened suddenly. Examination revealed wheezing, asymmetric breath sounds and decreased air entry. What is the most likely location of foreign body in this case?

- ☐ a. Trachea
- ☐ b. Bronchus
- ☐ c. Lung
- ☐ d. Nasal area

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 3
A nurse was exposed to a gush of amniotic fluid that splashed to the conjunctiva and oral mucosa while attending delivery. Since there was no additional assistance at the moment, the nurse continued the management for half an hour. Upon assessment of the source case, the patient was found to be HIV positive without clinical symptoms with CD4 cell count of 800 cells/mm³. Later on, the detail of the incident together with the laboratory finding was sent to the post exposure local person (PELP). What is the most appropriate immediate action the PELP should take?

- ☐ a. Start ART treatment as soon as possible
- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question 4
A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5
A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep



NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:03

Question 4

Not yet
answeredMarked out of
1.00Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5

Not yet
answeredMarked out of
1.00Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Next page

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

☐ c. 10+3 weeks

☐ d. 16+4 weeks

Time left 2:21:46

Question 8

Not yet answered

Marked out of 1.00

Flag question

A 67-year-old female patient came to a hospital with major burn injury with total surface area of 35 percent. The weight of the patient was 60kg. A nurse administered 4200ml of fluid for the first eight hours and 4200ml in the second 16 hours. After 16 hours, the nurse assessed the overall status of the patient. Upon assessment, her blood pressure was 120/80mmHg, pulse rate was 90/min and urine output was 56ml/hr. On laboratory investigation, hemoglobin level of 9mg/dl, serum sodium level of 138mg/dl and serum potassium level of 4.1mg/dl were noted. What is the most appropriate next nursing intervention for this patient?

- ☐ a. Assess breath sounds and respiratory rate
- ☐ b. Elevate head of patient's bed and elevate burned extremities
- ☐ c. Decrease the rate of intravenous fluid administration
- ☐ d. Monitor arterial blood gas values and pulse oxymetry reading

Question 9

Not yet answered

Marked out of 1.00

Flag question

A 40-year-old known patient with diabetes came to an emergency department with vomiting, abdominal pain and muscle cramps. Laboratory findings show random blood sugar of 400mg/dL, serum and urine ketone bodies, and elevated blood urea nitrogen, creatinine and hemoglobin. What is the most appropriate initial nursing intervention in this case?

- ☐ a. Administer anti-pain
- ☐ b. Give oral hypoglycemic agents
- ☐ c. Hydrate the patient
- ☐ d. Give insulin

Question 10

Not yet answered

A primigravida woman at 32-weeks of gestational age came to a clinic for a routine antenatal care follow-up. On examination, the fetus is in a longitudinal lie, and the upper uterine segment is occupied by a soft irregular mass. The nurse decided to determine the fetal presentation in the next assessment. What Leopold maneuver should the nurse perform?

Lenovo

Le63

☐ c. 10+3 weeks

☐ d. 16+4 weeks

Time left 2:21:46

Question 8

Not yet answered

Marked out of 1.00

Flag question

A 67-year-old female patient came to a hospital with major burn injury with total surface area of 35 percent. The weight of the patient was 60kg. A nurse administered 4200ml of fluid for the first eight hours and 4200ml in the second 16 hours. After 16 hours, the nurse assessed the overall status of the patient. Upon assessment, her blood pressure was 120/80mmHg, pulse rate was 90/min and urine output was 56ml/hr. On laboratory investigation, hemoglobin level of 9mg/dl, serum sodium level of 138mg/dl and serum potassium level of 4.1mg/dl were noted. What is the most appropriate next nursing intervention for this patient?

- ☐ a. Assess breath sounds and respiratory rate
- ☐ b. Elevate head of patient's bed and elevate burned extremities
- ☐ c. Decrease the rate of intravenous fluid administration
- ☐ d. Monitor arterial blood gas values and pulse oxymetry reading

Question 9

Not yet answered

Marked out of 1.00

Flag question

A 40-year-old known patient with diabetes came to an emergency department with vomiting, abdominal pain and muscle cramps. Laboratory findings show random blood sugar of 400mg/dL, serum and urine ketone bodies, and elevated blood urea nitrogen, creatinine and hemoglobin. What is the most appropriate initial nursing intervention in this case?

- ☐ a. Administer anti-pain
- ☐ b. Give oral hypoglycemic agents
- ☐ c. Hydrate the patient
- ☐ d. Give insulin

Question 10

Not yet answered

A primigravida woman at 32-weeks of gestational age came to a clinic for a routine antenatal care follow-up. On examination, the fetus is in a longitudinal lie, and the upper uterine segment is occupied by a soft irregular mass. The nurse decided to determine the fetal presentation in the next assessment. What Leopold maneuver should the nurse perform?

Lenovo

Le63

- ☐ c. GIVE NVP syrup prophylaxis
- ☐ d. Give Co-trimoxazole preventive therapy

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question 20

Not yet
answered

Marked out of
1.00

Flag question

A three-year-old child who had a history of ingesting uncooked vegetables was brought to a health center complaining watery stool accompanied with flatus, bulky stool, abdominal cramps and weight loss for a week. On laboratory test, ova of parasite are seen. What is the most likely etiologic agent in this case?

- ☐ a. Salmonella
- ☐ b. Campylobacter jejuni
- ☐ c. Shigella
- ☐ d. Giardia lamblia

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Previous page

Next page



- ☐ c. GIVE NVP syrup prophylaxis
- ☐ d. Give Co-trimoxazole preventive therapy

Question 20

Not yet
answered

Marked out of
1.00

Flag question

A three-year-old child who had a history of ingesting uncooked vegetables was brought to a health center complaining watery stool accompanied with flatus, bulky stool, abdominal cramps and weight loss for a week. On laboratory test, ova of parasite are seen. What is the most likely etiologic agent in this case?

- ☐ a. Salmonella
- ☐ b. Campylobacter jejuni
- ☐ c. Shigella
- ☐ d. Giardia lamblia

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Previous page

Next page



Question 21Not yet
answeredMarked out of
1.00

Flag question

Time left 1:48:55

A 28-year-old multiparous woman visited a hospital for contraception service. She was on her 14 days of postpartum period and not breastfeeding. She had no history of chronic disease. What is the most appropriate contraceptive method for the client?

- ☐ a. Progestin only injectables
- ☐ b. Combined oral contraceptive
- ☐ c. Female sterilization
- ☐ d. Combined vaginal ring

Question 22Not yet
answeredMarked out of
1.00

Flag question

An epidemiologist wanted to follow male diabetic patients aged 40 to 60 years in a population for one year and report the rate of death of diabetics men among midyear population in the age of 40 to 60 years. What is the most appropriate measure of mortality for this scenario?

- ☐ a. Sex-specific death rate
- ☐ b. Age specific death rate
- ☐ c. Cause specific death rate
- ☐ d. Proportionate mortality ratio

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 21

Not yet
answered

Marked out of
1.00

Flag question

Time left 1:48:55

A 28-year-old multiparous woman visited a hospital for contraception service. She was on her 14 days of postpartum period and not breastfeeding. She had no history of chronic disease. What is the most appropriate contraceptive method for the client?

- ☐ a. Progestin only injectables
- ☐ b. Combined oral contraceptive
- ☐ c. Female sterilization
- ☐ d. Combined vaginal ring

Question 22

Not yet
answered

Marked out of
1.00

Flag question

An epidemiologist wanted to follow male diabetic patients aged 40 to 60 years in a population for one year and report the rate of death of diabetics men among midyear population in the age of 40 to 60 years. What is the most appropriate measure of mortality for this scenario?

- ☐ a. Sex-specific death rate
- ☐ b. Age specific death rate
- ☐ c. Cause specific death rate
- ☐ d. Proportionate mortality ratio

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 23

Not yet
answeredMarked out of
1.00

Flag question

A 20-year-old woman has been admitted to a gynecology ward with complaints of fever and abdominal pain for two day. She looks sick and anxious, has chills and rigors. She had gone through illegal abortion three days back. On examination, her vital sign shows temperature of 38.6 degree centigrade, pulse rate of 92/min, and respiratory rate of 24/min. Pelvic examination revealed offensive, and purulent vaginal discharge with tender uterus. What is the most priority nursing diagnosis in this case?

- ☐ a. Risk for infection related to dilated cervix
- ☐ b. Acute pain related to uterine cramping
- ☐ c. Anxiety related to medical interventions
- ☐ d. Risk for deficient fluid volume related to bleeding

Question 24

Not yet
answeredMarked out of
1.00

Flag question

During curriculum development, an instructor wanted to assess the availability of skill lab equipment, reference and procedure books in order to deliver the program effectively. What is the most likely component of curriculum stated in this case?

- ☐ a. Quality assurance mechanism
- ☐ b. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

97 98 99 100

Finish attempt



Question 23

Not yet
answeredMarked out of
1.00

Flag question

A 20-year-old woman has been admitted to a gynecology ward with complaints of fever and abdominal pain for two day. She looks sick and anxious, has chills and rigors. She had gone through illegal abortion three days back. On examination, her vital sign shows temperature of 38.6 degree centigrade, pulse rate of 92/min, and respiratory rate of 24/min. Pelvic examination revealed offensive, and purulent vaginal discharge with tender uterus. What is the most priority nursing diagnosis in this case?

- ☐ a. Risk for infection related to dilated cervix
- ☐ b. Acute pain related to uterine cramping
- ☐ c. Anxiety related to medical interventions
- ☐ d. Risk for deficient fluid volume related to bleeding

Question 24

Not yet
answeredMarked out of
1.00

Flag question

During curriculum development, an instructor wanted to assess the availability of skill lab equipment, reference and procedure books in order to deliver the program effectively. What is the most likely component of curriculum stated in this case?

- ☐ a. Quality assurance mechanism
- ☐ b. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

97 98 99 100

Finish attempt

- ☐ b. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

Question 25

Not yet
answered

Marked out of
1.00

Flag question

Researchers conducted a research on the prevalence and associated factors of perinatal asphyxia among newborns admitted to intensive care unit (ICU). What is the most appropriate specific objective for this research problem?

- ☐ a. To determine the prevalence of perinatal asphyxia among newborn admitted to ICU
- ☐ b. To understand the prevalence of perinatal asphyxia among newborns admitted to ICU
- ☐ c. To study factors associated with perinatal asphyxia among newborns admitted to ICU
- ☐ d. To appreciate factors associated with perinatal asphyxia among newborns admitted to ICU

Previous page

Next page



- ☐ b. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

Question 25

Not yet
answered

Marked out of
1.00

Flag question

Researchers conducted a research on the prevalence and associated factors of perinatal asphyxia among newborns admitted to intensive care unit (ICU). What is the most appropriate specific objective for this research problem?

- ☐ a. To determine the prevalence of perinatal asphyxia among newborn admitted to ICU
- ☐ b. To understand the prevalence of perinatal asphyxia among newborns admitted to ICU
- ☐ c. To study factors associated with perinatal asphyxia among newborns admitted to ICU
- ☐ d. To appreciate factors associated with perinatal asphyxia among newborns admitted to ICU

Previous page

Next page



- ☐ b. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

Question 25

Not yet
answered

Marked out of
1.00

Flag question

Researchers conducted a research on the prevalence and associated factors of perinatal asphyxia among newborns admitted to intensive care unit (ICU). What is the most appropriate specific objective for this research problem?

- ☐ a. To determine the prevalence of perinatal asphyxia among newborn admitted to ICU
- ☐ b. To understand the prevalence of perinatal asphyxia among newborns admitted to ICU
- ☐ c. To study factors associated with perinatal asphyxia among newborns admitted to ICU
- ☐ d. To appreciate factors associated with perinatal asphyxia among newborns admitted to ICU

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Previous page

Next page



- ☐ a. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question **25**

Not yet
answered

Marked out of
1.00

Flag question

Researchers conducted a research on the prevalence and associated factors of perinatal asphyxia among newborns admitted to intensive care unit (ICU). What is the most appropriate specific objective for this research problem?

- ☐ a. To determine the prevalence of perinatal asphyxia among newborn admitted to ICU
- ☐ b. To understand the prevalence of perinatal asphyxia among newborns admitted to ICU
- ☐ c. To study factors associated with perinatal asphyxia among newborns admitted to ICU
- ☐ d. To appreciate factors associated with perinatal asphyxia among newborns admitted to ICU

Previous page

Next page



- ☐ a. Learning resources
- ☐ c. Student evaluation measures
- ☐ d. Teaching strategies

Question 25

Not yet
answered

Marked out of
1.00

 Flag question

Researchers conducted a research on the prevalence and associated factors of perinatal asphyxia among newborns admitted to intensive care unit (ICU). What is the most appropriate specific objective for this research problem?

- ☐ a. To determine the prevalence of perinatal asphyxia among newborn admitted to ICU
- ☐ b. To understand the prevalence of perinatal asphyxia among newborns admitted to ICU
- ☐ c. To study factors associated with perinatal asphyxia among newborns admitted to ICU
- ☐ d. To appreciate factors associated with perinatal asphyxia among newborns admitted to ICU

[Previous page](#)

[Next page](#)



Time left 1:46:36

Question 26

Not yet
answered

Marked out of
1.00

Flag question

A postmenopausal woman was admitted to a hospital with a diagnosis of endometrial cancer. She had irregular postmenopausal bleeding, watery offensive discharge, colicky abdominal pain, and pallor. She experienced fear, feeling of guilty, anxiety and denial. What is the most appropriate nursing intervention for the patient?

- ☐ a. Providing emotional therapy
- ☐ b. Providing antibiotics
- ☐ c. Providing radiation therapy
- ☐ d. Providing chemotherapy

Question 27

Not yet
answered

Marked out of
1.00

Flag question

A health manager planned to reduce the burden of communicable diseases in a community, and proposed that health education is the most crucial step to prevent the disease. In order to achieve the goal, the manager assigned skilled man power, and allocated budget and different materials. What characteristic of good planning is missing in this scenario?

- ☐ a. Mobilizing resources needed for the task
- ☐ b. Identifying clear picture of the task to be accomplished
- ☐ c. Community participation
- ☐ d. Setting clear objective

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:46:36

Question 26

Not yet
answered

Marked out of
1.00

Flag question

A postmenopausal woman was admitted to a hospital with a diagnosis of endometrial cancer. She had irregular postmenopausal bleeding, watery offensive discharge, colicky abdominal pain, and pallor. She experienced fear, feeling of guilty, anxiety and denial. What is the most appropriate nursing intervention for the patient?

- ☐ a. Providing emotional therapy
- ☐ b. Providing antibiotics
- ☐ c. Providing radiation therapy
- ☐ d. Providing chemotherapy

Question 27

Not yet
answered

Marked out of
1.00

Flag question

A health manager planned to reduce the burden of communicable diseases in a community, and proposed that health education is the most crucial step to prevent the disease. In order to achieve the goal, the manager assigned skilled man power, and allocated budget and different materials. What characteristic of good planning is missing in this scenario?

- ☐ a. Mobilizing resources needed for the task
- ☐ b. Identifying clear picture of the task to be accomplished
- ☐ c. Community participation
- ☐ d. Setting clear objective

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:46:36

Question 26

Not yet
answered

Marked out of
1.00

Flag question

A postmenopausal woman was admitted to a hospital with a diagnosis of endometrial cancer. She had irregular postmenopausal bleeding, watery offensive discharge, colicky abdominal pain, and pallor. She experienced fear, feeling of guilty, anxiety and denial. What is the most appropriate nursing intervention for the patient?

- ☐ a. Providing emotional therapy
- ☐ b. Providing antibiotics
- ☐ c. Providing radiation therapy
- ☐ d. Providing chemotherapy

Question 27

Not yet
answered

Marked out of
1.00

Flag question

A health manager planned to reduce the burden of communicable diseases in a community, and proposed that health education is the most crucial step to prevent the disease. In order to achieve the goal, the manager assigned skilled man power, and allocated budget and different materials. What characteristic of good planning is missing in this scenario?

- ☐ a. Mobilizing resources needed for the task
- ☐ b. Identifying clear picture of the task to be accomplished
- ☐ c. Community participation
- ☐ d. Setting clear objective

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:46:36

Question 26

Not yet
answered

Marked out of
1.00

Flag question

A postmenopausal woman was admitted to a hospital with a diagnosis of endometrial cancer. She had irregular postmenopausal bleeding, watery offensive discharge, colicky abdominal pain, and pallor. She experienced fear, feeling of guilty, anxiety and denial. What is the most appropriate nursing intervention for the patient?

- ☐ a. Providing emotional therapy
- ☐ b. Providing antibiotics
- ☐ c. Providing radiation therapy
- ☐ d. Providing chemotherapy

Question 27

Not yet
answered

Marked out of
1.00

Flag question

A health manager planned to reduce the burden of communicable diseases in a community, and proposed that health education is the most crucial step to prevent the disease. In order to achieve the goal, the manager assigned skilled man power, and allocated budget and different materials. What characteristic of good planning is missing in this scenario?

- ☐ a. Mobilizing resources needed for the task
- ☐ b. Identifying clear picture of the task to be accomplished
- ☐ c. Community participation
- ☐ d. Setting clear objective

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:46:36

Question 26

Not yet
answered

Marked out of
1.00

Flag question

A postmenopausal woman was admitted to a hospital with a diagnosis of endometrial cancer. She had irregular postmenopausal bleeding, watery offensive discharge, colicky abdominal pain, and pallor. She experienced fear, feeling of guilty, anxiety and denial. What is the most appropriate nursing intervention for the patient?

- ☐ a. Providing emotional therapy
- ☐ b. Providing antibiotics
- ☐ c. Providing radiation therapy
- ☐ d. Providing chemotherapy

Question 27

Not yet
answered

Marked out of
1.00

Flag question

A health manager planned to reduce the burden of communicable diseases in a community, and proposed that health education is the most crucial step to prevent the disease. In order to achieve the goal, the manager assigned skilled man power, and allocated budget and different materials. What characteristic of good planning is missing in this scenario?

- ☐ a. Mobilizing resources needed for the task
- ☐ b. Identifying clear picture of the task to be accomplished
- ☐ c. Community participation
- ☐ d. Setting clear objective

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti culex

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti culex

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti culex

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti culex

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti culex

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 28

Not yet
answeredMarked out of
1.00

Flag question

A 30-year-old man came to a health center with a complaint of chills and abdominal pain of 1-week duration. The detail history showed that the patient had fever, headache and diarrhea for the last one week. On physical examination, there was enlargement of spleen. In addition the blood film result was positive for hemo-parasite. What is the most likely vector for disease transmission in the above case?

Time left 1:46:15

- ☐ a. Biomphalaria
- ☐ b. Female anopheles mosquito
- ☐ c. Borrelia recurrentis
- ☐ d. Wuchereria bancrofti

Question 29

Not yet
answeredMarked out of
1.00

Flag question

A newborn infant was brought to a clinic with complaints of swollen and red eye, pus discharging from both eyes and persistent crying of three days duration. The mother had multiple sexual partners. She also had hypertension, DM, and has been treated for malaria during pregnancy. A nurse admitted the newborn in neonatal ICU for further treatment. What is the most likely cause for the newborn problems?

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Mellitus

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 30

Not yet
answered

Marked out of
1.00

Flag question

Investigators informed study participants before starting data collection saying: "This data will not be disseminated to third party for any purpose without your voluntariness, and the principal investigator will keep the data in a computer with protected password. In addition there are no any identification parameters like name and others from the questionnaire." What is the most appropriate principle of research ethics applied in this scenario?

- ☐ a. Beneficence
- ☐ b. Respect for communities
- ☐ c. Autonomy
- ☐ d. Justice

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 30

Not yet
answered

Marked out of
1.00

Flag question

Investigators informed study participants before starting data collection saying: "This data will not be disseminated to third party for any purpose without your voluntariness, and the principal investigator will keep the data in a computer with protected password. In addition there are no any identification parameters like name and others from the questionnaire." What is the most appropriate principle of research ethics applied in this scenario?

- ☐ a. Beneficence
- ☐ b. Respect for communities
- ☐ c. Autonomy
- ☐ d. Justice

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 30

Not yet
answered

Marked out of
1.00

Flag question

Investigators informed study participants before starting data collection saying: "This data will not be disseminated to third party for any purpose without your voluntariness, and the principal investigator will keep the data in a computer with protected password. In addition there are no any identification parameters like name and others from the questionnaire." What is the most appropriate principle of research ethics applied in this scenario?

- ☐ a. Beneficence
- ☐ b. Respect for communities
- ☐ c. Autonomy
- ☐ d. Justice

- ☐ a. Maternal malaria
- ☐ b. Maternal sexually transmitted disease
- ☐ c. Maternal hypertension disease
- ☐ d. Maternal diabetes Miletus

Question 30

Not yet
answered

Marked out of
1.00

Flag question

Investigators informed study participants before starting data collection saying: "This data will not be disseminated to third party for any purpose without your voluntariness, and the principal investigator will keep the data in a computer with protected password. In addition there are no any identification parameters like name and others from the questionnaire." What is the most appropriate principle of research ethics applied in this scenario?

- ☐ a. Beneficence
- ☐ b. Respect for communities
- ☐ c. Autonomy
- ☐ d. Justice

Time left 1:43:44

Question 36

Not yet
answeredMarked out of
1.00

Flag question

A biostatistics instructor wanted to assess the daily performance of nursing students and has given quizzes on different days. The result is distributed in descending order 10, 9, 8.5, 8, 8, 7.5, 7, 7, 6, 6, 5.5, 5.5, 4, 4, 4, 3, 3 and 2. What is the mean of the students' result in this case?

- ☐ a. 6
- ☐ b. 8
- ☐ c. 5.5
- ☐ d. 7

Question 37

Not yet
answeredMarked out of
1.00

Flag question

A 28-month-old young male child was brought to an emergency unit due to difficulty of breathing and coughing that happened suddenly. Examination revealed wheezing, asymmetric breath sounds and decreased air entry. What is the most likely location of foreign body in this case?

- ☐ a. Trachea
- ☐ b. Bronchus
- ☐ c. Lung
- ☐ d. Nasal area

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

○ d. Solid foods should be delayed until age 9 months

Time left 1:54:03

97 98 99 100

Finish attempt

Question 18

Not yet
answered

Marked out of
1.00

Flag question

A child who was relatively well three hours ago was brought to an emergency unit due to sudden loss of consciousness. The child has no history of foreign body aspiration. Physical examination revealed absent breathing and cold extremities. The child is suspected to have cardiac arrest. What is the classic sign to confirm the diagnosis?

- ☐ a. Increased heart rate
- ☐ b. Absent pulse rate
- ☐ c. Increased blood pressure
- ☐ d. Decreased urine output

Question 19

Not yet
answered

Marked out of
1.00

Flag question

A 12-hour baby born at home from HIV positive mother was brought to a hospital. HIV status of the mother was confirmed 12 hours after birth. The baby was on exclusive breastfeeding. All vital signs were in the normal range. What is the most priority nursing intervention in this case?

- ☐ a. Start NVP + 3TC prophylaxis
- ☐ b. Start NVP + AZT prophylaxis
- ☐ c. GIVE NVP syrup prophylaxis
- ☐ d. Give Co-trimoxazole preventive therapy

Question 16

Not yet
answered

Marked out of
1.00

Flag question

A medical director of a hospital assessed the overall activities of the organization and found out different problems that hindered the achievement of the set goals. The director ranked the listed problems according to their criticality and frequency. What is the most likely next step of planning process in this scenario?

- ☐ a. Selecting objectives and targets
- ☐ b. Situational analysis
- ☐ c. Identifying potential obstacles
- ☐ d. Designing strategies

Question 17

Not yet
answered

Marked out of
1.00

Flag question

A mother brought her three-month-old infant to a pediatric OPD to ask when to start solid foods for her baby. Assessment result shows all primitive reflexes are intact and anthropometric measurements are in normal range. What is the most appropriate advice to be given in this case?

- ☐ a. Start pureed solids at this age if fed via the bottle
- ☐ b. Infants do not require solid food until 12 months of age
- ☐ c. Good time to start solid foods is at age 6 months
- ☐ d. Solid foods should be delayed until age 9 months

Time left 1:54:36

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 16

Not yet
answered

Marked out of
1.00

Flag question

A medical director of a hospital assessed the overall activities of the organization and found out different problems that hindered the achievement of the set goals. The director ranked the listed problems according to their criticality and frequency. What is the most likely next step of planning process in this scenario?

- ☐ a. Selecting objectives and targets
- ☐ b. Situational analysis
- ☐ c. Identifying potential obstacles
- ☐ d. Designing strategies

Question 17

Not yet
answered

Marked out of
1.00

Flag question

A mother brought her three-month-old infant to a pediatric OPD to ask when to start solid foods for her baby. Assessment result shows all primitive reflexes are intact and anthropometric measurements are in normal range. What is the most appropriate advice to be given in this case?

- ☐ a. Start pureed solids at this age if fed via the bottle
- ☐ b. Infants do not require solid food until 12 months of age
- ☐ c. Good time to start solid foods is at age 6 months
- ☐ d. Solid foods should be delayed until age 9 months

Time left 1:54:36

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 14

Not yet
answeredMarked out of
1.00

Flag question

A 20-year-old female patient came to an emergency outpatient unit with severe burn to the perineal area. The emergency area was full of other patients and there was no enough space to deliver care as per the patient's need. The nurse planned to give care and informed the patient. However, the patient was not willing to accept the care. What is the most likely reason for refusal of the planned care?

- ☐ a. The patient preferred to get care from physician other than nurse
- ☐ b. The patient feared that the pain become worst during care
- ☐ c. The patient felt that her case does not need the planned care
- ☐ d. The patient felt that the caring environment was not convenient

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question 15

Not yet
answeredMarked out of
1.00

Flag question

A head of a health center wanted to employ two nurses to deliver holistic care services for the populations in the catchment areas. There were a number of applicants who applied for the job. The head saw the applicants' profiles, gave exam, conducted pre-employment interview, and identified potential applicant who best suited for the job. What is the most likely human resource management activities applied in this scenario?

- ☐ a. Selection
- ☐ b. Orientation
- ☐ c. Planning
- ☐ d. Recruitment

☐ c. Cervical eversion

☐ d. Cervical ectopy

Question 13

Not yet
answered

Marked out of
1.00

Flag question

A community health team assessed the health status of a community and identified lack of latrine and inappropriate waste disposal system in the community. Moreover, the team listed activities to construct latrine and prepared sessions to teach the community on how they would handle wastes. At the end, the activities were distributed to each team members and the necessary resources to accomplish the activities were allocated. What is the most likely component of action plan lacking in this scenario?

- ☐ a. Responsible person
- ☐ b. Time to do the activities
- ☐ c. Required materials
- ☐ d. Tasks to be implemented

Question 14

Not yet
answered

Marked out of
1.00

A 20-year-old female patient came to an emergency outpatient unit with severe burn to the perineal area. The emergency area was full of other patients and there was no enough space to deliver care as per the patient's need. The nurse planned to give care and informed the patient. However, the patient was not willing to accept the care. What is the most likely reason for refusal of the planned care?

89	90	91	92	93	94
97	98	99	100		

Finish attempt



Time left 1:59:19

Question 11

Not yet
answered

Marked out of
1.00

Flag question

A 26-year-old young man came to an outpatient department for consultation stating that he is unable to stop use of substance, and he has developed headache, aggressiveness, and generalized tonic-clonic seizures while he does not consume. What is the most appropriate diagnosis in this case?

- ☐ a. Caffeine dependency
- ☐ b. Amphetamine dependency
- ☐ c. Alcohol dependence
- ☐ d. Cannabis dependence

Question 12

Not yet
answered

Marked out of
1.00

Flag question

A 43-year-old known HIV positive multigravida woman came to a hospital with complaints of vaginal discharge and bleeding. She has postcoital bleeding and foul smelling yellowish vaginal discharge. On speculum examination, there is an exophytic cervical mass that bleeds on contact. What is the most likely diagnosis in this case?

- ☐ a. Cervical cancer
- ☐ b. Cervical polyp
- ☐ c. Cervical eversion
- ☐ d. Cervical ectopy

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

- ☐ c. 384
- ☐ d. 600

Question 10

Not yet
answered

Marked out of
1.00

 Flag question

A nursing association evaluated all nursing curricula every five years for the purpose of accreditation and in order to decide the standard of education system. What is the most appropriate type of curriculum evaluation stated in this case?

- ☐ a. Formative evaluation
- ☐ b. Summative evaluation
- ☐ c. Placement evaluation
- ☐ d. Diagnostic evaluation

[Previous page](#)

[Next page](#)



Question 8

Not yet
answeredMarked out of
1.00

Flag question

A patient informed a nurse that he chews khat every day to calm his stress and to stop his craving. He has been chewing khat repeatedly, and he develops headache, nervousness, and restlessness if he does not get khat. For what condition is the patient at greatest risk?

- ☐ a. Neurological dependency
- ☐ b. Physical dependency
- ☐ c. Psychological dependency
- ☐ d. Substance abuse

Question 9

Not yet
answeredMarked out of
1.00

Flag question

A nurse wanted to conduct a research on the prevalence and associated factors of hospital acquired infection among admitted patients. A 95 percent confidence interval is desired to estimate the proportion. Prevalence rate of previous similar study was 40 percent with margin error of 4 percent. What is the most appropriate sample size for this scenario?

- ☐ a. 294
- ☐ b. 576
- ☐ c. 384
- ☐ d. 600

97 98 99 100

Finish attempt...

Question 6

Not yet
answered

Marked out of
1.00

Flag question

101: A 34-year-old woman, who has been treated for breast cancer, came to a health center to receive contraception. She has a history of deep vein thrombosis. What is the most appropriate contraceptive method?

- ☐ a. Copper-bearing intrauterine device
- ☐ b. Combined vaginal ring
- ☐ c. Progestin-only injectables
- ☐ d. Combined oral contraceptive pills

Question 7

Not yet
answered

Marked out of
1.00

Flag question

A head of an operational theater room of a hospital received and recorded all materials which are purchased from different shops in the country and abroad. He labeled all the equipment based on their group and put them in to the appropriate place. What is the most likely material management principle applied in this scenario?

- ☐ a. Controlling
- ☐ b. Storing
- ☐ c. Issuing
- ☐ d. Ordering

9	10	11	12
17	18	19	20
25	26	27	28
33	34	35	36
41	42	43	44
49	50	51	52
57	58	59	60
65	66	67	68
73	74	75	76
81	82	83	84
89	90	91	92
97	98	99	



Question 5

Not yet
answered

Marked out of
1.00

Flag question

- ☐ b. 0.5 ml
- ☐ c. 5 ml
- ☐ d. 1 ml

A newborn infant, who was delivered from a mother with a history of vaginal discharge, was brought to a hospital with complaints of fever and poor breast feeding of one day duration. The mother had high grade fever before and during labor and was put on IV medications. Physical examination shows body temperature of 38 degree centigrade, respiratory rate of 66/min, and stamped umbilical cord. Maintenance fluid of 10 percent glucose and antibiotics were started after admission. What is the most likely risk factor for this newborn?

- ☐ a. Maternal infection
- ☐ b. Complicated umbilical stump
- ☐ c. Unhygienic hospital environment
- ☐ d. Complication of central IV lines

Next page

Question 3

Not yet
answered

Marked out of
1.00

Flag question

Time left 2:13:05

A 20-day-old newborn who was delivered at home was brought to a clinic after developing severe red-swollen and pussy discharge from eyes of two days duration. The mother had a history of sexually transmitted infections. What is the best management of this case?

- ☐ a. Topical corticosteroid application
- ☐ b. Give epinephrine
- ☐ c. Washing eyes
- ☐ d. Give Ceftriaxone IM

Question 4

Not yet
answered

Marked out of
1.00

Flag question

An immunization clinic nurse administered vaccine using 16-mm needle through anterolateral thigh muscle. The administered vaccine was important to protect a child from development of fast breathing, chest in drawing and cough. What is the appropriate vaccine dose in this case?

- ☐ a. 0.1 ml
- ☐ b. 0.5 ml
- ☐ c. 5 ml
- ☐ d. 1 ml

Question 5

Not yet
answered

A newborn infant, who was delivered from a mother with a history of vaginal discharge, was brought to a hospital with complaints of fever and poor breast feeding of one day duration. The mother had



Question 3

Not yet
answered

Marked out of
1.00

Flag question

Time left 2:13:05

A 20-day-old newborn who was delivered at home was brought to a clinic after developing severe red-swollen and pussy discharge from eyes of two days duration. The mother had a history of sexually transmitted infections. What is the best management of this case?

- ☐ a. Topical corticosteroid application
- ☐ b. Give epinephrine
- ☐ c. Washing eyes
- ☐ d. Give Ceftriaxone IM

Question 4

Not yet
answered

Marked out of
1.00

Flag question

An immunization clinic nurse administered vaccine using 16-mm needle through anterolateral thigh muscle. The administered vaccine was important to protect a child from development of fast breathing, chest in drawing and cough. What is the appropriate vaccine dose in this case?

- ☐ a. 0.1 ml
- ☐ b. 0.5 ml
- ☐ c. 5 ml
- ☐ d. 1 ml

Question 5

Not yet
answered

A newborn infant, who was delivered from a mother with a history of vaginal discharge, was brought to a hospital with complaints of fever and poor breast feeding of one day duration. The mother had



Question 1

Not yet
answered

Marked out of
1.00

Flag question

A person who is working in an organization has a problem of making an effective communication and understanding the feelings of other co-workers. The co-workers in the organization have strong disagreement with this person because he undermines and insults them verbally and uses unnecessary name-calling. What is the most likely type of violence in this case?

- ☐ a. Emotional abuse
- ☐ b. Sexual abuse
- ☐ c. Economic abuse
- ☐ d. Physical abuse

Question 2

Not yet
answered

Marked out of
1.00

Flag question

A 20-day-old neonate is admitted to NICU due to vomiting and poor breast feeding of one day duration. The newborn has convulsion with a body temperature of 35 degree centigrade, heart rate of 120/min, and breathing rate of 50/min. The nurse has provided 4ml/kg bolus and maintenance fluid of 150ml/kg 10 percent glucose for 24-hours. What is the most likely sign that helped the nurse to decide the treatment?

- ☐ a. Convulsion
- ☐ b. Hypothermia
- ☐ c. Tachypnea
- ☐ d. Tachycardia

Time left 2:14:52

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 1

Not yet
answered

Marked out of
1.00

Flag question

A person who is working in an organization has a problem of making an effective communication and understanding the feelings of other co-workers. The co-workers in the organization have strong disagreement with this person because he undermines and insults them verbally and uses unnecessary name-calling. What is the most likely type of violence in this case?

- ☐ a. Emotional abuse
- ☐ b. Sexual abuse
- ☐ c. Economic abuse
- ☐ d. Physical abuse

Question 2

Not yet
answered

Marked out of
1.00

Flag question

A 20-day-old neonate is admitted to NICU due to vomiting and poor breast feeding of one day duration. The newborn has convulsion with a body temperature of 35 degree centigrade, heart rate of 120/min, and breathing rate of 50/min. The nurse has provided 4ml/kg bolus and maintenance fluid of 150ml/kg 10 percent glucose for 24-hours. What is the most likely sign that helped the nurse to decide the treatment?

- ☐ a. Convulsion
- ☐ b. Hypothermia
- ☐ c. Tachypnea
- ☐ d. Tachycardia

Time left 2:14:52

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 3
A nurse was exposed to a gush of amniotic fluid that splashed to the conjunctiva and oral mucosa while delivering a newborn. Since there was no additional assistance at the moment, the nurse continued the management for half an hour. Upon assessment of the source case, the patient was found to be HIV positive without clinical symptoms with CD4 cell count of 800 cells/mm³. Later on, the detail of the incident together with the laboratory finding was sent to the post-exposure local person (PEEP). What is the most appropriate immediate action the PEEP should take?

- ☐ a. Start ART treatment as soon as possible
- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Question 4
A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5
A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

Lenovo

Lc63

Question 3
A nurse was exposed to a gush of amniotic fluid that splashed to the conjunctiva and oral mucosa while delivering a newborn. Since there was no additional assistance at the moment, the nurse continued the management for half an hour. Upon assessment of the source case, the patient was found to be HIV positive without clinical symptoms with CD4 cell count of 800 cells/mm³. Later on, the detail of the incident together with the laboratory finding was sent to the post-exposure local person (PEEP). What is the most appropriate immediate action the PEEP should take?

- ☐ a. Start ART treatment as soon as possible
- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Question 4
A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5
A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

Lenovo

Lc63

NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:03

Question 4

Not yet
answeredMarked out of
1.00Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5

Not yet
answeredMarked out of
1.00Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Next page

Lenovo

Ac63

NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:03

Question 4

Not yet
answeredMarked out of
1.00Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5

Not yet
answeredMarked out of
1.00Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Next page

Lenovo

Ac63

Prepare for Exit Exam! Join our Telegram Channel for all Department
@ExitExamPrep

NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:02

Question 4

Not yet
answered
Marked out of
1.00

Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5

Not yet
answered
Marked out of

Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Next page

Lenovo

Ac63

NLE IB

- ☐ b. Start PEP containing two basic ART drugs
- ☐ c. Manage as low risk exposure
- ☐ d. Start PEP containing three basic ART drugs

Time left 2:25:02

Question 4

Not yet
answered
Marked out of
1.00

Flag
question

A 22-year-old primigravida woman has been admitted to labor ward after she presented with a complaint of pushing down pain for two days. During initial assessment, she has term sized uterus, 3 uterine contractions lasting for 30 seconds, descent of 3/5, cervical dilatation of 4cm and intact membrane. Four hours after initial examination, she has 4 uterine contractions lasting for 60 seconds, descent of 2/5, cervical dilatation of 6cm and ruptured membranes. What signs are most suggestive of the unsatisfactory progress of labor in this scenario?

- ☐ a. Cervical dilatation and descent
- ☐ b. Uterine contraction and cervical dilatation
- ☐ c. Cervical dilatation and rupture of membranes
- ☐ d. Uterine contraction and descent

Question 5

Not yet
answered
Marked out of
1.00

Flag
question

A 62-year-old unconscious male patient was brought to an emergency ward. He has sunken eye and dry lip. His blood pressure is 90/60mmHg, respiratory rate is 25/min, pulse rate is 90/min and temperature is 36.7 degree centigrade. His Glasgow Coma Scale is 6. A nurse administered intravenous fluids, but inappropriate fluid distribution is noted. What is the most appropriate initial nursing intervention in this scenario?

- ☐ a. Nutritional support
- ☐ b. Proper positioning
- ☐ c. Vasoactive medications
- ☐ d. Fluid replacement

Next page

Lenovo

Ac63

Time left 2:24:05

A 15-year-old female patient came to a hospital with complaints of dysuria, itching and burning pain during urination of five days duration. On assessment, a nurse identified that the patient has knowledge deficit about perineal cleaning. Then, the nurse gave her norfloxacin medication prescription and nursing education on how to clean the perineal area. What is most likely nursing intervention that addresses the patient's knowledge deficit?

- ☐ a. Wiping perineal area by water from back to front after defecation
- ☐ b. Wiping perineal area by warm sitz bath two times a day for 20 minutes
- ☐ c. Wiping perineal area by water from front to back after defecation
- ☐ d. Wiping perineal area by soap bath two times a day for 20 minutes

A 19-year-old primigravida presented to antenatal care unit to start routine follow-up on 17/05/2014 E.C. Her LMP was on 14/01/2014 E.C. She reported that she does not feel her fetal movement since amenorrhea. Her urine HCG test was positive and ultrasound revealed that fetal cardiac activity is positive. What is the most likely gestational age in this case?

- ☐ a. 17+3 weeks
- ☐ b. 17+4 weeks
- ☐ c. 16+3 weeks
- ☐ d. 16+4 weeks

A 67-year-old female patient came to a hospital with major burn injury with total surface area of 35 percent. The weight of the patient was 60kg. A nurse administered 4200ml of fluid for the first eight hours and 4200ml in the second 16 hours. After

Quiz n

1	2
9	10
17	18
25	26
33	34
41	42
49	50
57	58
65	66
73	74
81	82
89	90
97	98

Finish attempt

Lenovo

Be63

NLE IB

question

- ☐ b. Wiping perineal area by warm sitz bath two times a day for 20 minutes
- ☐ c. Wiping perineal area by water from front to back after defecation
- ☐ d. Wiping perineal area by soap bath two times a day for 20 minutes

Time left 2:23:48

Question 7

Not yet answered

Marked out of 1.00

Flag question

A 19-year-old primigravida presented to antenatal care unit to start routine follow-up on 17/05/2014 E.C. Her LNMP was on 14/01/2014 E.C. She reported that she does not feel her fetal movement since amenorrhea. Her urine HCG test was positive and ultrasound revealed that fetal cardiac activity is positive. What is the most likely gestational age in this case?

- ☐ a. 17+3 weeks
- ☐ b. 17+4 weeks
- ☐ c. 16+3 weeks
- ☐ d. 16+4 weeks

Question 8

Not yet answered

Marked out of 1.00

Flag question

A 67-year-old female patient came to a hospital with major burn injury with total surface area of 35 percent. The weight of the patient was 60kg. A nurse administered 4200ml of fluid for the first eight hours and 4200ml in the second 16 hours. After 16 hours, the nurse assessed the overall status of the patient. Upon assessment, her blood pressure was 120/80mmHg, pulse rate was 90/min and urine output was 56ml/hr. On laboratory investigation, hemoglobin level of 9mg/dl, serum sodium level of 138mg/dl and serum potassium level of 4.1mg/dl were noted. What is the most appropriate next nursing intervention for this patient?

- ☐ a. Assess breath sounds and respiratory rate
- ☐ b. Elevate head of patient's bed and elevate burned extremities
- ☐ c. Decrease the rate of intravenous fluid administration
- ☐ d. Monitor arterial blood gas values and pulse oxymetry reading

Question 9

A 40-year-old known patient with diabetes came to an emergency department with vomiting, abdominal pain and muscle

Lenovo

Le63

NLE 1B

question

- ☐ a. Assess breath sounds and respiratory rate
- ☐ b. Elevate head of patient's bed and elevate burned extremities
- ☐ c. Decrease the rate of intravenous fluid administration
- ☐ d. Monitor arterial blood gas values and pulse oximetry reading

Time left 2:19:34

Question 9

Not yet answered
Marked out of 00

Flag question

A 40-year-old known patient with diabetes came to an emergency department with vomiting, abdominal pain and muscle cramps. Laboratory findings show random blood sugar of 400mg/dL, serum and urine ketone bodies, and elevated blood urea nitrogen, creatinine and hemoglobin. What is the most appropriate initial nursing intervention in this case?

- ☐ a. Administer anti-pain
- ☐ b. Give oral hypoglycemic agents
- ☐ c. Hydrate the patient
- ☐ d. Give insulin

Question 10

Not yet answered
Marked out of 00

Flag question

A primigravida woman at 32 weeks of gestational age came to a clinic for a routine antenatal care follow-up. On examination, the fetus is in a longitudinal lie, and the upper uterine segment is occupied by a soft irregular mass. The nurse decided to determine the fetal presentation in the next assessment. What Leopold maneuver should the nurse perform?

- ☐ a. Deep pelvic palpation
- ☐ b. Lateral palpation
- ☐ c. Pawlick's grip
- ☐ d. Fundal palpation

Previous page

Next page

Lenovo

Ke63

- urse. What is the most likely ethical principle the nurse breached in this case?
- ☐ a. Beneficence
 - ☐ b. Veracity
 - ☐ c. Fidelity
 - ☐ d. Non-maleficence

Question 100

Not yet
answered

Marked out of
1.00

Flag question

A 35-year-old man has been brought by his wife to a psychiatry department with symptoms of sleep disturbance and refusal to eat for the last seven months. The patient stated that his wife wants to kill him and he sees that his wife may put something on his food. He also hears that his friends were calling his name and insulting him. His family and the community have good support. His symptoms have become worsen when he gets into chewing chat and drinking alcohol. What is the most likely cause of the patient's condition in this scenario?

- ☐ a. Genetic
- ☐ b. Substance abuse
- ☐ c. Psychological problem
- ☐ d. Social problem

Previous page

Finish attempt ...

NLE IB

Question 93

Not yet
answered

Marked out of
1.00

Flag question

A 20-year-old female patient came to a health center for abortion on the sixth week of pregnancy. The nurse identified that the reason of abortion was conflict with her boyfriend. On further assessment, she had sign of anemia, hopelessness, and suicidal ideation. What is the best mechanism in which the nurse can support the patient?

Time left 1:12:03

Finish attempt

- ☐ a. Refer to hospital for abortion where there is blood transfusion
- ☐ b. Try to call her family, discuss with them to help her
- ☐ c. Take informed consent, conduct abortion and provide folic acid
- ☐ d. Provide counseling and give her iron sulphate

Question 94

Not yet
answered

Marked out of
1.00

Flag question

A medical ward team leader is following whether or not the nursing care services are being delivered according to the standard performance using checklist. Furthermore, he measures and evaluates each activities based on the standard indicators and gives feedback to each staff members. What is the most likely management function practiced by the leader?

- ☐ a. Controlling
- ☐ b. Directing
- ☐ c. Organizing
- ☐ d. Planning

Question 95

A graduate nurse took out DPT and TT vaccine from the refrigerator. Before administering those

Marked out of
1.00

Flag question

decision, though it was the right of the patient to accept or reject the decision. Finally, the patient was convinced to accept the amputation. What is the most likely ethical principle promoted by the nurse?

- ☐ a. Autonomy
- ☐ b. Non-maleficence
- ☐ c. Veracity
- ☐ d. Beneficence

Question 85

Not yet
answered

Marked out of
1.00

Flag question

A fourth year nursing student wanted to assess the nutritional status of a mother and measured her weight which was 56kg and height 1.65 meter in order to determine body mass index of the mother. What is the most likely nutritional assessment method applied in this case?

- ☐ a. Biophysical method
- ☐ b. Clinical method
- ☐ c. Anthropometric method
- ☐ d. Dietary method

Marked out of
1.00

Flag question

decision, though it was the right of the patient to accept or reject the decision. Finally, the patient was convinced to accept the amputation. What is the most likely ethical principle promoted by the nurse?

- ☐ a. Autonomy
- ☐ b. Non-maleficence
- ☐ c. Veracity
- ☐ d. Beneficence

Question 85

Not yet
answered

Marked out of
1.00

Flag question

A fourth year nursing student wanted to assess the nutritional status of a mother and measured her weight which was 56kg and height 1.65 meter in order to determine body mass index of the mother. What is the most likely nutritional assessment method applied in this case?

- ☐ a. Biophysical method
- ☐ b. Clinical method
- ☐ c. Anthropometric method
- ☐ d. Dietary method

Time left 1:14:22

Question 81

Not yet
answered

Marked out of
1.00

Flag question

A clinical nurse gave health education to a community about prevention mechanisms of infectious disease. Then, all the community members practiced the prevention approaches of infectious diseases each day. The health status of the community was maintained. Finally, the community members gave a credit for the primary prevention. What is the most appropriate communication stage for this scenario?

- ☐ a. Attracting the audience of attention
- ☐ b. Understanding the message
- ☐ c. Reaching the intended audience
- ☐ d. Improving to health

Question 82

Not yet
answered

Marked out of
1.00

Flag question

A 28-year-old woman came to a gynecology OPD with complaints of fever and vaginal discharge of one week duration. She had offensive vaginal discharge and high grade fever. Physical examination revealed the presence of lower abdominal tenderness, cervical motion tenderness, adnexal tenderness and pelvic mass. Laboratory finding suggested leukocytosis of $11,500/\text{mm}^3$. What is the most likely diagnosis in this case?

- ☐ a. Torsion of ovarian cyst
- ☐ b. Vulvovestibulitis
- ☐ c. Ectopic pregnancy

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:14:22

Question 81

Not yet
answered

Marked out of
1.00

Flag question

A clinical nurse gave health education to a community about prevention mechanisms of infectious disease. Then, all the community members practiced the prevention approaches of infectious diseases each day. The health status of the community was maintained. Finally, the community members gave a credit for the primary prevention. What is the most appropriate communication stage for this scenario?

- ☐ a. Attracting the audience of attention
- ☐ b. Understanding the message
- ☐ c. Reaching the intended audience
- ☐ d. Improving to health

Question 82

Not yet
answered

Marked out of
1.00

Flag question

A 28-year-old woman came to a gynecology OPD with complaints of fever and vaginal discharge of one week duration. She had offensive vaginal discharge and high grade fever. Physical examination revealed the presence of lower abdominal tenderness, cervical motion tenderness, adnexal tenderness and pelvic mass. Laboratory finding suggested leukocytosis of $11,500/\text{mm}^3$. What is the most likely diagnosis in this case?

- ☐ a. Torsion of ovarian cyst
- ☐ b. Vulvovestibulitis
- ☐ c. Ectopic pregnancy

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

NLE IB

Question 73

Not yet
answeredMarked out of
1.00

Flag question

A 40-year-old woman came to a psychiatric clinic with drowsiness, cardiac arrhythmia, weight gain and orthostatic hypotension after she took Amitriptyline 100mg PO per day. She cannot perform her daily activities due to the problem. What is the most appropriate management option in this case?

Time left 1:28:09

Finish attempt...

- ☐ a. Shift to another tricyclic group
- ☐ b. Decrease Amitriptyline dose
- ☐ c. Increase Amitriptyline dose
- ☐ d. Wait for one month

Question 74

Not yet
answeredMarked out of
1.00

Flag question

A 45-year-old male patient was brought to a hospital by his brother with complete leg fracture and profuse bleeding secondary to car accident. The surgeons decided that the patient's leg should be amputated at below the knee. The surgeons were busy by other cases and informed the nurse to prepare the patient immediately for operation. What is the most appropriate initial step in the preparation of the patient in this case?

- ☐ a. Take informed consent from his brother
- ☐ b. Clean a surgical site with iodine solution
- ☐ c. Give preoperative fluids and antibiotics
- ☐ d. Assess the patient's ability for informed consent

Question 75

Time left 1:28:33

Question 71

Not yet
answeredMarked out of
1.00

Flag question

A child was brought to a health center with a complaint of tender swelling behind the ear. Body temperature showed 38 degree centigrade. The child had a history of recurrent upper respiratory infection. Then, the child was treated accordingly. What is the most appropriate treatment for this child?

- ☐ a. Amoxicillin PO
- ☐ b. Ampicillin and Chloramphenicol IV/IM
- ☐ c. Gentamycin IV/IM
- ☐ d. Erythromycin and Amoxicillin IV/IM

Question 72

Not yet
answeredMarked out of
1.00

Flag question

A three-year-old child who had been on ART for three months was brought to a pediatric OPD due to weight loss. The child has no history of diarrhea, cough and body swelling. Physical examination showed that MUAC is 11.6cm, weight for height is between -2 and -3 Z-score and there is no bilateral pitting edema. The child was treated for malnutrition for more than a month though there was no response yet. What is the most likely clinical stage of the child in this case?

- ☐ a. Stage III
- ☐ b. Stage IV
- ☐ c. Stage II
- ☐ d. Stage I

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:28:33

Question 71

Not yet
answeredMarked out of
1.00

Flag question

A child was brought to a health center with a complaint of tender swelling behind the ear. Body temperature showed 38 degree centigrade. The child had a history of recurrent upper respiratory infection. Then, the child was treated accordingly. What is the most appropriate treatment for this child?

- ☐ a. Amoxicillin PO
- ☐ b. Ampicillin and Chloramphenicol IV/IM
- ☐ c. Gentamycin IV/IM
- ☐ d. Erythromycin and Amoxicillin IV/IM

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question 72

Not yet
answeredMarked out of
1.00

Flag question

A three-year-old child who had been on ART for three months was brought to a pediatric OPD due to weight loss. The child has no history of diarrhea, cough and body swelling. Physical examination showed that MUAC is 11.6cm, weight for height is between -2 and -3 Z-score and there is no bilateral pitting edema. The child was treated for malnutrition for more than a month though there was no response yet. What is the most likely clinical stage of the child in this case?

- ☐ a. Stage III
- ☐ b. Stage IV
- ☐ c. Stage II
- ☐ d. Stage I

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

answered

Marked out of
1.00

Flag question

evaluate the success of the course by taking oral examination, written test and practical exam." What is the most appropriate component of course syllabus stated in this scenario?

- ☐ a. Course description
- ☐ b. Teaching method
- ☐ c. Course schedule
- ☐ d. Assessment method

Question 70

Not yet
answered

Marked out of
1.00

Flag question

A 20-year-old woman came to a clinic for family planning service. She never used any family planning method before. The nurse in charge greeted the client and started to provide counseling on family planning methods. But the woman was fearful to communicate her issue for the nurse. What is the most appropriate counseling principle the nurse should follow?

- ☐ a. Provide information on the a clinic preference on family planning method
- ☐ b. Explore the client's satisfaction with the method
- ☐ c. Try to involve another nurse for discussion
- ☐ d. Engage in two way communication with the client

Previous page

Next page

Question 68

Not yet
answeredMarked out of
1.00

Flag question

A head of a health center prioritized different health care services in order to decide the amount of materials and professionals needed for each department's activities. Then after, he engaged in assigning professionals, setting schedules, giving out materials, and preparing budgets according to their activities. What is the most likely role of a manager applied in this scenario?

- ☐ a. Resource allocator
- ☐ b. Disturbance handler
- ☐ c. Negotiator
- ☐ d. Entrepreneur

Time left 1:30:34

Finish attempt

Page Summary

Search page...

Developer Console

Question 69

Not yet
answeredMarked out of
1.00

Flag question

A instructor was writing from the syllabus during the first day of a class "Based on the course objectives, all students will evaluate the knowledge, attitude and skills; those attending the course will evaluate the success of the course by taking oral examination, written test and practical exam." What is the most appropriate component of course syllabus stated in this scenario?

- ☐ a. Course description
- ☐ b. Teaching method
- ☐ c. Course schedule
- ☐ d. Assessment method

Question 68

Not yet
answeredMarked out of
1.00

Flag question

A head of a health center prioritized different health care services in order to decide the amount of materials and professionals needed for each department's activities. Then after, he engaged in assigning professionals, setting schedules, giving out materials, and preparing budgets according to their activities. What is the most likely role of a manager applied in this scenario?

- ☐ a. Resource allocator
- ☐ b. Disturbance handler
- ☐ c. Negotiator
- ☐ d. Entrepreneur

Time left 1:30:34

Finish attempt

Page Summary

Search page...

Developer Console

Question 69

Not yet
answeredMarked out of
1.00

Flag question

A instructor was writing from the syllabus during the first day of a class "Based on the course objectives, all students will evaluate the knowledge, attitude and skills; those attending the course will evaluate the success of the course by taking oral examination, written test and practical exam." What is the most appropriate component of course syllabus stated in this scenario?

- ☐ a. Course description
- ☐ b. Teaching method
- ☐ c. Course schedule
- ☐ d. Assessment method

answered

Marked out of
1.00

Flag question

applied in this scenario?

- ☐ a. Clinical simulation
- ☐ b. Demonstration
- ☐ c. Case study
- ☐ d. Discussion

Question 60

Not yet
answered

Marked out of
1.00

Flag question

A nurse working at medical ward had five patients. One of the patients was on end stage of life. After many supports, the doctors confirmed that the patient will not survive and the nurse also confirmed that. Then, the nurse withholds basic care for the patient and focused only on the other patients. Meanwhile, the head nurse saw the patient, gave warning to the nurse and ordered to continue the necessary cares until death is confirmed. What is the most likely reason for the order by the head nurse in this case?

- ☐ a. The nurse feared the family complaint that will happen after death
- ☐ b. The patient should get dignified care irrespective of the condition
- ☐ c. It is not human authority to predict death of the patient
- ☐ d. The nurse suspected the physicians evidence

Previous page

Next page

ENG

☐ d. Provide Amoxicillin for five days

Time left 1:32:34

97 98 99 100

Finish attempt...

Question 58

Not yet
answered

Marked out of
1.00

Flag question

A three-day-old male neonate was brought by his father after his mother died due to maternal complication. A father needs health education about formula feeding. What is the most appropriate health education message for the father?

- ☐ a. Sterilizing bottles and nipples for 30 seconds with tap water
- ☐ b. Dilute with extra water to extend the amount of formula
- ☐ c. Use bottled water without fluoride for mixing infant formula
- ☐ d. Always follow manufacturers instruction about preparation

Question 59

Not yet
answered

Marked out of
1.00

Flag question

An instructor showed students about the procedure of female catheterization. After the discussion, all students tried to do the procedure based on the steps. What is the most likely teaching method applied in this scenario?

- ☐ a. Clinical simulation
- ☐ b. Demonstration
- ☐ c. Case study
- ☐ d. Discussion

Question 60

Time left 1:32:58

Question 56

Not yet
answeredMarked out of
1.00

Flag question

After attending a wedding ceremony, many people showed signs of sudden onset of vomiting, watery diarrhea, fever and abdominal cramp within a short period of time. The people who experienced those signs were taken to a nearby hospital to get emergency medical treatment and the diagnosis of all the people was food intoxication. What is the most likely type of outbreak in this scenario?

- ☐ a. Point common source epidemic
- ☐ b. Propagative/progressive epidemic
- ☐ c. Intermittent common source
- ☐ d. Continuous common source epidemic

Question 57

Not yet
answeredMarked out of
1.00

Flag question

A two-year-old child was brought to a health center with complaints of cough and fever of three days duration. On examination, oxygen saturation is 95 percent with a respiratory rate of 38/min. What is the most likely management for this child?

- ☐ a. Refer the child urgently to the hospital after giving first line drugs
- ☐ b. Advise the mother to give fluid remedies
- ☐ c. Provide bronchodilators and send home with appointment
- ☐ d. Provide Amoxicillin for five days

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:32:58

Question 56

Not yet
answeredMarked out of
1.00

Flag question

After attending a wedding ceremony, many people showed signs of sudden onset of vomiting, watery diarrhea, fever and abdominal cramp within a short period of time. The people who experienced those signs were taken to a nearby hospital to get emergency medical treatment and the diagnosis of all the people was food intoxication. What is the most likely type of outbreak in this scenario?

- ☐ a. Point common source epidemic
- ☐ b. Propagative/progressive epidemic
- ☐ c. Intermittent common source
- ☐ d. Continuous common source epidemic

Question 57

Not yet
answeredMarked out of
1.00

Flag question

A two-year-old child was brought to a health center with complaints of cough and fever of three days duration. On examination, oxygen saturation is 95 percent with a respiratory rate of 38/min. What is the most likely management for this child?

- ☐ a. Refer the child urgently to the hospital after giving first line drugs
- ☐ b. Advise the mother to give fluid remedies
- ☐ c. Provide bronchodilators and send home with appointment
- ☐ d. Provide Amoxicillin for five days

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Flag question

- ☐ a. Start cognitive therapy
- ☐ b. Refer to specialist
- ☐ c. Start medication therapy
- ☐ d. Start speech therapy

Question 55

Not yet
answered

Marked out of
1.00

Flag question

A three-year-old girl was admitted to a pediatric ward with a diagnosis of severe acute malnutrition. The child came after three episodes of watery diarrhea. She had dry mouth, sunken eye ball, eagerness to drink and skin lying in folds. What is the most likely complication in this case?

- ☐ a. Dehydration
- ☐ b. Anemia
- ☐ c. Hypoglycemia
- ☐ d. Hypothermia

Previous page

Next page



Flag question

- ☐ a. Start cognitive therapy
- ☐ b. Refer to specialist
- ☐ c. Start medication therapy
- ☐ d. Start speech therapy

Question 55

Not yet
answered

Marked out of
1.00

Flag question

A three-year-old girl was admitted to a pediatric ward with a diagnosis of severe acute malnutrition. The child came after three episodes of watery diarrhea. She had dry mouth, sunken eye ball, eagerness to drink and skin lying in folds. What is the most likely complication in this case?

- ☐ a. Dehydration
- ☐ b. Anemia
- ☐ c. Hypoglycemia
- ☐ d. Hypothermia

Previous page

Next page



Flag question

- ☐ a. Give adequate cow's milk
- ☐ b. Advice to take fat or cholesterol
- ☐ c. Juice intake of 120 to 180mlper day
- ☐ d. Advice to provide fortified cereals

Question 35

Not yet
answered

Marked out of
1.00

Flag question

A child had been on ART for one year. During ART initiation, the child was on WHO Stage-II. However, the child developed pulmonary TB, PJP and esophageal candidiasis. Due to this the ART provider suspected treatment failure. The CD4 count was 400cells/mm³, and viral load level was unknown. What is the most likely treatment failure in this case?

- ☐ a. Clinical failure
- ☐ b. Immunological failure
- ☐ c. Virological failure
- ☐ d. ART failure

Previous page

Next page



Flag question

- ☐ a. Give adequate cow's milk
- ☐ b. Advice to take fat or cholesterol
- ☐ c. Juice intake of 120 to 180mlper day
- ☐ d. Advice to provide fortified cereals

Question 35

Not yet
answered

Marked out of
1.00

Flag question

A child had been on ART for one year. During ART initiation, the child was on WHO Stage-II. However, the child developed pulmonary TB, PJP and esophageal candidiasis. Due to this the ART provider suspected treatment failure. The CD4 count was 400cells/mm³, and viral load level was unknown. What is the most likely treatment failure in this case?

- ☐ a. Clinical failure
- ☐ b. Immunological failure
- ☐ c. Virological failure
- ☐ d. ART failure

Previous page

Next page



NLE IB

Question 33

Not yet answered

Marked out of 1.00

Flag question

A 27-year-old man, who was diagnosed with generalized anxiety disorder, presented to a clinic with a complaint of extreme worry about his education and his family. What is the best psychological intervention in this case?

- ☐ a. Cognitive behavioral therapy
- ☐ b. Insight oriented therapy
- ☐ c. Group therapy
- ☐ d. Supportive therapy

Time left 1:44:32

Finish attempt...

Question 34

Not yet answered

Marked out of 1.00

Flag question

A seven-month-old girl whose mother has died last month was brought by her father with a complaint of easily fatigability of 15 days duration. She is on supplementary diet. Physical examination finding shows palmar pallor and pale conjunctivitis. Otherwise, her vital signs and growth assessment findings are in normal range. What most appropriate measure that has to be taken in this case?

- ☐ a. Give adequate cow's milk
- ☐ b. Advice to take fat or cholesterol
- ☐ c. Juice intake of 120 to 180ml per day
- ☐ d. Advice to provide fortified cereals

Question 35

A child had been on ART for one year. During ART initiation, the child was on WHO Stage-II.

ENG

2:00 PM

10/04/2024



NLE IB

Question 33

Not yet
answered

Marked out of
1.00

Flag question

A 27-year-old man, who was diagnosed with generalized anxiety disorder, presented to a clinic with a complaint of extreme worry about his education and his family. What is the best psychological intervention in this case?

- ☐ a. Cognitive behavioral therapy
- ☐ b. Insight oriented therapy
- ☐ c. Group therapy
- ☐ d. Supportive therapy

Time left 1:44:32

Finish attempt...

Question 34

Not yet
answered

Marked out of
1.00

Flag question

A seven-month-old girl whose mother has died last month was brought by her father with a complaint of easily fatigability of 15 days duration. She is on supplementary diet. Physical examination finding shows palmar pallor and pale conjunctivitis. Otherwise, her vital signs and growth assessment findings are in normal range. What most appropriate measure that has to be taken in this case?

- ☐ a. Give adequate cow's milk
- ☐ b. Advice to take fat or cholesterol
- ☐ c. Juice intake of 120 to 180ml per day
- ☐ d. Advice to provide fortified cereals

Question 35

A child had been on ART for one year. During ART initiation, the child was on WHO Stage-II.

ENG

2:00 PM



Time left 1:45:01

Question 31

Not yet
answered

Marked out of
1.00

Flag question

During a school health program, a health extension worker assessed the school children and encountered a nine-year-old child who had a complaint of abdominal pain, nausea, dyspnea and passing of worm in the feces. The health extension worker took the child to the health center to perform stool examination for identifying the cause of the child's problem. What is the most likely diagnosis of the child in this scenario?

- ☐ a. Entrobiasis
- ☐ b. Trichuriasis
- ☐ c. Ascariasis
- ☐ d. Hook worm

Question 32

Not yet
answered

Marked out of
1.00

Flag question

A four-year-old male child was brought to an emergency outpatient department with severe bleeding after he sustained car accident. A nurse on duty diagnosed severe anemia that needed immediate management. However, a physician was not around. What is the most appropriate next step the nurse should take in the management of the child?

- ☐ a. Wait until the physician comes back
- ☐ b. Take informed consent from the family
- ☐ c. Consult nurses to transfuse or not
- ☐ d. Start blood transfusion directly

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Time left 1:45:01

Question 31

Not yet
answered

Marked out of
1.00

Flag question

During a school health program, a health extension worker assessed the school children and encountered a nine-year-old child who had a complaint of abdominal pain, nausea, dyspnea and passing of worm in the feces. The health extension worker took the child to the health center to perform stool examination for identifying the cause of the child's problem. What is the most likely diagnosis of the child in this scenario?

- ☐ a. Entrobiasis
- ☐ b. Trichuriasis
- ☐ c. Ascariasis
- ☐ d. Hook worm

Question 32

Not yet
answered

Marked out of
1.00

Flag question

A four-year-old male child was brought to an emergency outpatient department with severe bleeding after he sustained car accident. A nurse on duty diagnosed severe anemia that needed immediate management. However, a physician was not around. What is the most appropriate next step the nurse should take in the management of the child?

- ☐ a. Wait until the physician comes back
- ☐ b. Take informed consent from the family
- ☐ c. Consult nurses to transfuse or not
- ☐ d. Start blood transfusion directly

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 46Not yet
answeredMarked out of
1.00

Flag question

A nurse working in surgical ward always tolerates the crying of patients due to severe pain during wound care. The nurse also advises other nurses not to insult patients; tells them that the patients are crying because of the pain. And the nurse tries to help the other nurses to understand and feel the patients' emotion. What component of dignified care best describes the nurse's behavior?

- ☐ a. Empathy
- ☐ b. Sympathy
- ☐ c. Caring
- ☐ d. Motivation

Question 47Not yet
answeredMarked out of
1.00

Flag question

A nurse made an assessment for a patient admitted to a health center with chronic kidney disease, and found symptoms of mild distress, and significant impairment in social functioning. However, the patient had better self-care and coping skill. The nurse diagnosed the patient with adjustment disorder secondary to medical condition. What is the most appropriate nursing intervention in this case?

- ☐ a. Establish social support system
- ☐ b. Teach deep breathing and relaxation techniques
- ☐ c. Use supportive touch if client responds well
- ☐ d. Discuss self-harm thoughts

Time left 1:39:44

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				



ENG

2:40 PM

10/14/2024



Question 46Not yet
answeredMarked out of
1.00

Flag question

A nurse working in surgical ward always tolerates the crying of patients due to severe pain during wound care. The nurse also advises other nurses not to insult patients; tells them that the patients are crying because of the pain. And the nurse tries to help the other nurses to understand and feel the patients' emotion. What component of dignified care best describes the nurse's behavior?

- ☐ a. Empathy
- ☐ b. Sympathy
- ☐ c. Caring
- ☐ d. Motivation

Question 47Not yet
answeredMarked out of
1.00

Flag question

A nurse made an assessment for a patient admitted to a health center with chronic kidney disease, and found symptoms of mild distress, and significant impairment in social functioning. However, the patient had better self-care and coping skill. The nurse diagnosed the patient with adjustment disorder secondary to medical condition. What is the most appropriate nursing intervention in this case?

- ☐ a. Establish social support system
- ☐ b. Teach deep breathing and relaxation techniques
- ☐ c. Use supportive touch if client responds well
- ☐ d. Discuss self-harm thoughts

Time left 1:39:44

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Question 41

Not yet
answered

Marked out of
1.00

Flag question

A four-day-old male neonate was born via SVD at GA of 35 weeks with birth weight of 2.3kg. He is on the second day of admission after he had yellowish discoloration. On physical examination, all primitive reflexes and other physical examination findings are normal. What is the most likely classification of the neonate according to IMNCI protocol?

- ☐ a. Term and normal birth weight
- ☐ b. Term and low birth weight
- ☐ c. Pre term and low birth weight
- ☐ d. Preterm and normal birth weight

Question 42

Not yet
answered

Marked out of
1.00

Flag question

A 15-year-old girl was admitted to a health center with severe weight loss due to anorexia nervosa. The child denies that she is underweight. She is hospitalized to initiate treatment because her weight has dropped to 39.5kg. A nurse planned to improve the child's nutritional status, setting a goal weight of 50 kg. What is the most appropriate nursing management for the child?

- ☐ a. Weigh monthly at the same time wearing the same type of clothes
- ☐ b. Test urine for glucose and regularly evaluate her skin turgor
- ☐ c. Supervise intake by observing her during and after meals
- ☐ d. Encourage the child from pushing herself beyond her physical limits

Time left 1:40:53

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				



ENG

2:38 PM

10/10/2024



Question 43

Not yet
answered

Marked out of
1.00

Flag question

- ☐ d. Encourage the child from pushing herself beyond her physical limits

Time left 1:40:29

97 98 99 100

Finish attempt

A child came to an emergency clinic due to motor cycle accident. The child was irritable with pain on the right leg. Then, the child was admitted and screened for soft tissue injuries via X-ray. The child's family was nervous even though there was no severe bleeding. What is the priority nursing diagnosis in this case?

- ☐ a. Risk of x-ray side effect
- ☐ b. Pain related to swelling
- ☐ c. Family disturbance related to accident
- ☐ d. Risk of compartment related to cast application

Question 44

Not yet
answered

Marked out of
1.00

Flag question

A nurse wanted to give health education to a community about COVID-19 prevention protocols. The education includes that minimizing exposures is a very important way to prevent the disease. What is the most appropriate level of prevention in this case?

- ☐ a. Secondary prevention
- ☐ b. Primordial prevention
- ☐ c. Tertiary prevention
- ☐ d. Primary prevention



Flag question

- ☐ a. Secondary prevention
- ☐ b. Primordial prevention
- ☐ c. Tertiary prevention
- ☐ d. Primary prevention

Question 45

Not yet
answered

Marked out of
1.00

Flag question

A nurse received new refrigerator to keep vaccines. He received 100 different types of vaccines from regional health office and he arranged them in the new refrigerator. To maintain vaccines potency, the nurse should adjust refrigerator temperature. What is the recommended temperature range in this scenario?

- ☐ a. -8 degree centigrade and +2 degree centigrade
- ☐ b. -2 degree centigrade and +8 degree centigrade
- ☐ c. -8 degree centigrade and -2 degree centigrade
- ☐ d. +2 degree centigrade and +8 degree centigrade

Previous page

Next page



Time left 1:30:57

Question 66

Not yet
answeredMarked out of
1.00

Flag question

A new team has been created to manage acute febrile illness in a hospital. The team consists of 5 nurses, 2 health officers, 5 physicians and 2 cleaners. The individuals in the team have refused to provide quality services for patients as per the standard. What is the most likely action of the individuals in this case?

- ☐ a. Initiating
- ☐ b. Following
- ☐ c. Opposing
- ☐ d. Observing

Question 67

Not yet
answeredMarked out of
1.00

Flag question

A 56-year-old male patient came to a hospital with urinary tract obstruction. A physician decided to perform cystotomy and communicated with a nurse to take informed consent. After the nurse explained in detail about the disease and treatment alternatives for the patient, the nurse saw the sign of acceptance from the patient's facial expression. What is the best next step the nurse should do in this case?

- ☐ a. Provide the patient pre-operative care
- ☐ b. Take the signature of the patient
- ☐ c. Inform the physician the patient's acceptance
- ☐ d. Give the patient a chance to talk to identify his need

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

1.00
Flag question

- ☐ a. Treat for Hypomagnesaemia
- ☐ b. Treat for hypocalcaemia
- ☐ c. Treat for hypoglycemia
- ☐ d. Initiate anti convulsants

Question 65

Not yet
answered

Marked out of
1.00

Flag question

A 35-year-old man came to a psychiatric clinic with sleep disturbance for the last nine months. He believes that his family wants to kill him by poisoned food because he is famous. He has weight loss as a result of refusing to eat and drink. He is happy most of the day; he is extravagant, dances alone and has sleep disturbance. He has self-care deficiency and social isolation. On examination, his blood pressure is 80/40mmHg, and pulse rate is 100/min. What is the priority nursing intervention for this scenario?

- ☐ a. Reduce environmental stimuli
- ☐ b. Provide physical activities
- ☐ c. Supplement diet with vitamins and minerals
- ☐ d. Administer tranquilizing medication as ordered

Previous page

Next page



☐ d. Co-trimoxazole

Time left 1:31:32

97 98 99 100

Finish attempt

Question 63

Not yet
answered

Marked out of
1.00

Flag question

A director wanted to improve service quality of a hospital at the end of the budget year. To achieve the plan, he fulfills different resources and helps the staff to get ready for promotion. What is the most likely factor promoting staff motivation in this scenario?

- ☐ a. Environment
- ☐ b. Responsibility
- ☐ c. Advancement
- ☐ d. Achievement

Question 64

Not yet
answered

Marked out of
1.00

Flag question

A two-hour-old newborn developed a brief and strange movement. On evaluation, the airway was clear, and there was normal breathing pattern and adequate circulation. Smacking and breast feeding like lip movement and eye deviation were noted frequently. What is the most appropriate primary nursing intervention in this case?

- ☐ a. Treat for Hypomagnesaemia
- ☐ b. Treat for hypocalcaemia
- ☐ c. Treat for hypoglycemia
- ☐ d. Initiate anti convulsants

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Question 61

Not yet
answered

Marked out of
1.00

Flag question

A two-year-old child was presented to a clinic with runny nose and barking cough. Physical examination revealed hoarseness, respiratory rate of 38/min, and body temperature of 36.8 degree centigrade. Otherwise, the child has clear chest sounds. What is the most likely etiologic agent for this case?

- ☐ a. Para influenza virus type
- ☐ b. Corona Virus
- ☐ c. Rhino virus
- ☐ d. Respiratory syncytial virus

Question 62

Not yet
answered

Marked out of
1.00

Flag question

A seven-year-old child was brought to a clinic with fever and pain on urination of one day duration. Examination revealed flank pain. There were no microscopic urine bacteria. Then, a nurse discharged the patient with PO medication and appointed for after seven days. What is the most likely medication given by the nurse?

- ☐ a. Gentamycin
- ☐ b. Amoxicillin
- ☐ c. Ampicillin
- ☐ d. Co-trimoxazole

Time left 1:31:46

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

answered

Marked out of
1.00

Flag question

...edly diminished interest in everything, significant weight loss, sleep disturbance, and hearing sounds that tell her to kill herself. She told to the nurse that she does not want to live anymore. What is the most likely priority nursing diagnosis in this case?

- ☐ a. Powerlessness
- ☐ b. Sleep disturbance
- ☐ c. Hopelessness
- ☐ d. Risk for suicide

Question 80

Not yet
answered

Marked out of
1.00

Flag question

A seven-year-old child suddenly started coughing and had a difficulty of breathing while eating. The mother immediately called for help and the neighbor nurse arrived soon. What is the most appropriate nursing intervention for this child?

- ☐ a. Tilt position to open the airway
- ☐ b. Give five times of Heimlich maneuver
- ☐ c. Give five chest thrusts
- ☐ d. Give five back blows

Previous page

Next page



Time left 1:27:29

Question 76

Not yet
answeredMarked out of
1.00

Flag question

A two-year-old child, who has a travel history to the lowland areas of Ethiopia, was brought to a clinic with a complaint of high grade fever. Blood smear test is not available. What is the most likely diagnosis in this case?

- ☐ a. Malaria
- ☐ b. Measles
- ☐ c. Pneumonia
- ☐ d. Fever

Question 77

Not yet
answeredMarked out of
1.00

Flag question

A seven-year-old child with a previous history of trauma to the right leg was brought to a pediatric OPD due to painful right leg swelling associated with high grade fever and malaise. On examination, erythematous and tender swelling with indistinct lateral margins was appreciated. What is the most likely skin disorder in this scenario?

- ☐ a. Eczema
- ☐ b. Cellulites
- ☐ c. Impetigo
- ☐ d. Boil

1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100				

Flag question

What is the most appropriate initial step in the preparation of the patient in this case?

- ☐ a. Take informed consent from his brother
- ☐ b. Clean a surgical site with iodine solution
- ☐ c. Give preoperative fluids and antibiotics
- ☐ d. Assess the patient's ability for informed consent

Question 75

Not yet
answered

Marked out of
1.00

Flag question

A married woman who has five children living in a definite community makes great effort to fulfill basic needs of the family. The woman manages the family by selling vegetables at the market far from home. The husband does not support the woman; instead, he always comes late at night, and also slaps and kicks her. What is the most likely type of violence in this case?

- ☐ a. Economic abuse
- ☐ b. Sexual abuse
- ☐ c. Psychological abuse
- ☐ d. Physical abuse

Previous page

Next page





Milki health science tutorial center



+251910690594, +251911058885


Tajaajiloota nuti kenninu



Barnoota dabalataa kennuu
Exit Exam'f/ COCf isin qopheessuu.
Leenjiwwan gaggabaaboo kennuu fi K.K.F.



Teessoo: Magaalaa Naqamtee, Gamoo OSSA, Darbii 3ffaa



Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Nursing (Comprehensive)

Back

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Time left 4:11:49

Question 100

Not yet
answered

Marked out of
1.00

Flag question

A nurse observed medication administration error made by his friend working in medical ward. Due to this, the patient developed minor complication and treated. What is the first thing the nurse must do?

- ☐ a. Follow secretly and supervise
- ☐ b. Report to the nurse director
- ☐ c. Report to the legal judiciary body
- ☐ d. Confront his friend

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Previous page

Finish attempt ...

Time left 4:12:00

98
out of
question

A woman, who has been using IUCD for six months, came to a clinic with complaints of lower abdominal pain and pain during sexual intercourse. On vaginal examination, uterine and cervical tenderness were identified. What is the priority nursing diagnosis for this woman?

- ☐ a. Risk for injury
- ☐ b. Risk for trauma
- ☐ c. Risk for bleeding
- ☐ d. Risk for infection

Previous page

Next page

64
73
82
91
100

hp

Time left 4:12:07

on 97

ed

d out of

question

A nurse researcher studied a hypothesis which read as "Frequent consumption of fruits and vegetables is protective for the occurrence of stomach cancer". He recruited 198 subjects recently diagnosed with stomach cancer and 396 subjects free of stomach cancer. All subjects were asked about their fruit and vegetable consumption during the past one year. What is the most likely study design that the researcher used?

- ☐ a. Ecological
- ☐ b. Case-control
- ☐ c. Cross-sectional
- ☐ d. Cohort

previous page

Next page

rsing (Comprehensive)

Time left 4:12:15

A nurse was making physical examination for a postnatal mother. The mother was restless, irritable, had pressured speech, and unusual behavior. She had also obsessional and delusional thinking. What is the most likely diagnosis for this woman?

- ☐ a. Postpartum depression
- ☐ b. Postpartum Psychosis
- ☐ c. Postpartum schizophrenia
- ☐ d. Postpartum blue

Next page

Nursing (Comprehensive)

Time left 4:12:22

Question 94

Let
red
ed out of

ing question

A 26-year-old female patient was admitted to a medical ward with a diagnosis of DKA. The patient selected a treatment option that is best for her after being informed about all treatment options. After five days of treatment, the patient progressed well and was discharged. However, since she has no money to pay, she is kept in the ward till the hospital manager will come and decide. What is the most likely encountered disrespect to the patient?

- ☐ a. Non consented care
- ☐ b. Detention in facility
- ☐ c. Abandonment of care
- ☐ d. No dignified care

Previous page

Next page

5
64
73
82

Nursing (Comprehensive)

Time left 4:12:18

Question 95

yet
ered
ed out of

lag question

A four-year-old child was presented to the health center with diarrhea of five days duration. On physical examination, the skin pinch goes back within two seconds, and the child was very eager to drink water and had WFH z-score greater than -1. What is the most appropriate management of this child according to IMNCI?

- ☐ a. Give ORS solution in each loss
- ☐ b. Give RL solution intravascular over three hours
- ☐ c. Give ReSoMAL solution in each loss
- ☐ d. Give ORS over 4-hours period

previous page

Next page



Nursing (Comprehensive)

Time left 4:12:27

Question 93

Not yet
answeredMarked out of
100

Flag question

A 26-year-old Gravid II Para I pregnant woman came to ANC clinic with complaints of headache, dyspnea, and fatigue for three days. Her pulse rate was 78/min, Blood pressure was 100/60 mmHg and fetal heart rate was 140/min. On physical examination, she had pale conjunctiva and 32 weeks gestational age pregnancy. Her hemoglobin concentration was 10g/dl. What is the most likely indicative characteristic for the woman's problem?

- ☐ a. Dyspnea
- ☐ b. Hgb 10 g/dl
- ☐ c. Headache
- ☐ d. Pale conjunctiva

Previous page

Next page

Nursing (Comprehensive)

Time left 4:12:30

Question 92

yet
wered
ked out of
Flag question

An infant was referred from private clinic to a health center with the diagnosis of severe anemia. On physical examination, the nurse found that infant's MUAC was less than 11.5cm, WFL less than -3Z score, bilateral pitting edema and RBS less than 40mg/dl. What is the most appropriate management for this infant?

- ☐ a. Admit the infant on OTP program
- ☐ b. Give oral amoxicillin for seven days
- ☐ c. Give folic acid supplement for anemia
- ☐ d. Give 1st dose of antibiotic and refer

Previous page

Next page

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Answer saved

Marked out of 1.00

 Flag question

A four-year-old child was brought to a health center with a complaint of cough of three days duration. On assessment, the nurse observed that the child had inspiratory stridor and barking cough. What is the most likely diagnosis of this patient?

- ☐ a. Croup
- ☒ b. Pneumonia
- ☐ c. Tonsilo-pharyngitis
- ☐ d. Pulmonary TB

[Clear my choice](#)[Next question](#)**Exam Navigation**

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Question 2

Not yet answered

Marked out of 1.00

Flag question

A 24-year-old male patient was admitted to an orthopedic inpatient unit with a diagnosis of Osteomyelitis. The nurse has taught the patient about nutritional needs used to maintain important amino-acids to facilitate the healing process. What is the most likely anticipated complication prevented with the above intervention?

- ☐ a. Protein deficiency
- ☐ b. Carbohydrate deficiency
- ☐ c. Vitamin deficiency
- ☐ d. Mineral deficiency

[Previous question](#)[Next question](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Marked out of 1.00

 Flag question

A 43-year-old patient came to a hospital with complaint of fever, fatigue, chest pain and night sweat for two weeks. On physical examination, the patient has tachycardia, productive cough and uses accessory muscles for breathing. What is the priority nursing diagnosis for this patient?

- ☐ a. Fluid volume deficit
- ☐ b. Ineffective airway clearance
- ☐ c. Fatigue
- ☐ d. Hyperthermia

[Previous question](#)[Next question](#)

Exam Navigation

[Flag question](#)

A 36-year-old male patient came to an emergency department with complaints of abdominal pain, persistent vomiting and abdominal distension of six hours duration. On physical examination, the patient had high pitched bowel sounds and visible peristaltic waves, tachycardia and his urine output was decreased. What is the priority nursing intervention for this patient?

- ☐ a. Maintain NG tube as prescribed to decompress bowel
- ☐ b. Measure and record all intake and output
- ☐ c. Place the patient in Fowler's position to minimize pain
- ☐ d. Administer IV fluids and electrolyte as prescribed

[Previous question](#)[Next question](#)

**Question 6**

Not yet answered


Marked out of 1.00

Flag question

A 44-year-old female patient with complaint of itching and rashes came to the dermatology clinic. On physical examination, line of scratching is visible on the skin. The patient reported that she has no information on how to care herself. What is the priority nursing diagnosis this patient?

- ☐ a. Disturbed body image related to visible rash
- ☐ b. Impaired skin integrity related to scratching
- ☐ c. Anxiety related to fear of death
- ☐ d. Deficient knowledge about disease process

[Previous question](#)[Next question](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone


Marked out of 1.00

Flag question

A four-year-old child was brought to pediatric OPD for follow up care. The child was on supplementary feeding in the community based management of acute malnutrition. On examination, child's weight-for-height was 65 percent and no other co-morbidity. What would be the interpretation of the anthropometric measurement of this child?

- ☐ a. Mild acute malnutrition
- ☐ b. Moderate acute malnutrition
- ☐ c. Severe acute malnutrition
- ☐ d. Severe chronic malnutrition

[Previous question](#)[Next question](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Question 7

Not yet answered

Marked out of 1.00

Flag question

A 48-year-old woman came to a medical OPD with complaints loss of appetite, fatigue and progressive weight gain. She also stated that she feels cold even at warm environment and has hoarseness of voice. What is the most likely medical diagnosis of this woman?

- ☐ a. Graves' disease
- ☐ b. Thyroid storm
- ☐ c. Hypothyroidism
- ☐ d. Myxedema

[Previous question](#)[Next question](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Marked out of 1.00

Flag question

A 45-year-old female patient with the diagnosis of acute renal failure was admitted to the medical ward. She had edema, difficulty of breathing and urinate 250 mL of urine in the past 24 hours. Her serum Creatinine level was 1.9 mg/dL (Normal range is of creatinine is 0.6–1.2 mg/dL). What is the pathophysiological stage of the patient's renal failure?

- ☐ a. Oliguric Phase
- ☐ b. Diuretic Phase
- ☐ c. Recovery Phase
- ☐ d. Initiation Phase

[Previous question](#)[Next question](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Flag question

A nurse was caring and preparing nursing care plan for patient with an end stage renal disease. A physician prescribed oral Furosemide (Lasix) to be added to the therapy. What is the most likely expected serious risk of the above drug?

- ☐ a. Hyponatremia
- ☐ b. Hypokalemia
- ☐ c. Hypernatremia
- ☐ d. Hyperkalemia

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

Previous question

Next question

Exam Navigation

1	2	3	4	5	6	7	8
---	---	---	---	---	---	---	---

9	10	11	12	13	14	15	16
---	----	----	----	----	----	----	----

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Flag question

A nurse prepared a patient for surgery by inserting NG tube. After inserting the tube, the nurse confirmed whether the tube is in the stomach. What is the most appropriate sign used for confirmation?

- ☐ a. There is no air bubble from the tip of NG tube in a cup of water
- ☐ b. The color of litmus paper is not changed after GI aspiration was applied
- ☐ c. There is central cyanosis and shortness of breath
- ☐ d. There is no gurgle sound auscultated over abdomen

[Previous question](#)[Next question](#)**Exam Navigation**

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Flag question

A 25-year-old male patient was admitted to an ICU and put on mechanical ventilator following car accident. On assessment; his eyes are not open to any stimuli, produces incomprehensible sounds and has abnormal flexion of motor response. What is the level of consciousness of this patient using Glasgow coma scale?

- ☐ a. Stupors
- ☐ b. Somnolent
- ☐ c. Comatose
- ☐ d. Semiconscious

[Previous question](#)[Finish exam ...](#)

Exam Navigation

1

2

3

4

5

6

7

8

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

Flag question

A 25-year-old male patient was admitted to an ICU and put on mechanical ventilator following car accident. On assessment; his eyes are not open to any stimuli, produces incomprehensible sounds and has abnormal flexion of motor response. What is the level of consciousness of this patient using Glasgow coma scale?

- ☐ a. Stupors
- ☐ b. Somnolent
- ☐ c. Comatose
- ☐ d. Semiconscious

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

[Previous question](#)[Finish exam ...](#)

Exam Navigation

1

2

3

4

5

6

7

8

Time left 0:12:40

A 30-year-old pregnant woman came to a health center to attend her first ANC visit. On examination, the nurse identified that woman's amniotic fluid index to be 24, fetal weight is 10 percent greater than the gestational age, fetal heart rate is 110/min and maternal blood pressure is 90/60mmHg. She was diagnosed for amniotic fluid disorder. What is the best characteristic that led to this diagnosis?

- ☐ a. Increased Fetal weight
- ☐ b. Fetal heart beat 110/min
- ☒ c. Amniotic fluid index 24
- ☐ d. Blood pressure 90/60mmHg

[Clear my choice](#)[Next page](#)

Lenovo

Nursing (Comprehensive)

[Back](#)**Question 59**

Answer saved

Marked out of
1.00[Flag question](#)

Time left 0:1

A 43-year-old male patient comes to a hospital with complaint of penetrated skin of two days duration. He also complains hematuria and dysuria. On laboratory, there are cercariae ova in urine. Most of the time the patient has swimming habit from their local river. What is the most appropriate management for this case?

- ☐ a. Clearing of vegetation in water bodies
- ☐ b. Administration of Diclofenac
- ☒ c. Administration of Praziquantel
- ☐ d. Avoid swimming in water bodies

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo

27

aved

out of


question

A 27-year-old patient comes to a hospital with the complaints of acute onset profuse and purulent urethral discharge, urgency, itching sensation of the urethra and burning sensation during micturition. The patient have also painless and regular ulcer on genitalia. What is the most likely diagnosis of this patient?

- ☐ a. Herpes genitalia
- ☒ b. Gonorrhea
- ☐ c. Chancroid
- ☐ d. Syphilis

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo



MoEEE

MoE - Exit Exam

Nursing (Comprehensive)

Back

Time left 0:00

Question 74

Answer saved

Marked out of 1.00

Remove flag

A 23-year-old 34 weeks pregnant mother came to a hospital for her third ANC visit. On assessment, her blood pressure is 150/90mmHg measured two times six hours apart and her pulse rate 75/min. abdominal assessment shows cephalic presentation and positive fetal heart beat. Laboratory investigation revealed 2+ proteins in urine, negative for glucose and ketone in the urine. She was diagnosed with mild preeclampsia. What is the most appropriate management for this mother?

- ☒ a. Follow the mother as an outpatient twice a week
- ☐ b. Give anticonvulsants and antihypertensive before discharge
- ☐ c. Admit the mother and administer diuretics
- ☐ d. Admit the mother and initiate magnesium sulfate

Clear my choice

Previous page

Next page

Lenovo

MoEEE

Nursing (Comprehensive)

Time left 0:01:00

Question 1
Not saved
Marked out of 5
Flag question

A 35-year-old male patient who is on TB treatment came to a TB clinic after completing his fifth month of treatment. The nurse working in TB unit sent him to laboratory for AFB test to evaluate his treatment condition. Unfortunately, the laboratory result indicates positive sputum smear for acid fast bacilli (AFB) test. What is the most likely treatment category of this patient?

- ☐ a. Previously treated
- ☐ b. Treatment after loss of follow-up
- ☐ c. Treatment after failure
- ☒ d. Relapse

Clear my choice

Next page

Lenovo

Exam Center:

Assosa Univeristy

Department:

Nursing (Comprehensive)

[Back](#)

Time left 0:23

Question 45

Answer saved

Marked out of 1.00


[Remove flag](#)

A 60-year-old female patient was admitted to a medical ward with a diagnosis of chronic renal failure. Laboratory result shows serum potassium level of 7mEq/L (normal value 3.5-5.0). On assessment she has ECG changes (tall-tented T waves and widened QRS). The health care provider diagnosed hyperkalemia. What early signs and symptoms could support the diagnosis?

- ☐ a. Blood pressure and fluid retention
- ☐ b. RBC count, hemoglobin and hematocrit level
- ☒ c. Muscle weakness and diarrhea
- ☐ d. Fever, chest pain and additive breath sound

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo



Nursing (Comprehensive)

Back

Question 42

Answer saved

Marked out of 1.00

Remove flag

Time left 0:25:08

A 32 weeks gestational age pregnant woman is admitted in obstetric ward for pregnancy-induced hypertension. During admission, the woman had blurred vision, blood pressure 160/110mmHg, protein urea 0.3gm and oliguria. The woman was conscious and she had no any seizure. She also took the recommended intervention to manage her problem. What is the most appropriate outcome criterion to evaluate the woman's prognosis?

- ☐ a. Urine output 600ml/24hrs
 - ☐ b. Absence of blurred vision
 - ☐ c. Proteinuria 1+
 - ☒ d. Blood pressure 140/90mmHg
- Clear my choice

Previous page

Next page

Que	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
-----	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

Lenovo



Nursing (Comprehensive)

Back

Time left 0:20:50

Question 49

Answer saved

Marked out of
1.00

Remove flag

A group of people came to a hospital with increasing fever, dyspnea, stridor, hypoxia, and hypotension after eating raw meat prepared for the cultural celebration. Blood culture is positive for *Bacillus anthracis* bacteria. What is the most appropriate drug management for these patients?

- ☒ a. Chloramphenicol
- ☐ b. Penicillin
- ☐ c. Ceftriaxone
- ☐ d. Gentamicin

Clear my choice

Previous page

Next page

Lenovo



MoEEE

Ministry of Education, Science and Technology

Exam Center:

Assosa Unveristy

Department:

Nursing (Comprehensive)

[Back](#)

Time left 0:30:16

Question 40

Answer saved

Marked out of 1.00


[Remove flag](#)

A physician decided to discharge a 26-year-old mother who gave normal birth after six hours of follow-up. The head nurse requested the nurse in charge to facilitate the process as per the Ethiopian National Maternal Discharge Protocol. What is the most likely initial nursing consideration to be taken?

- ☒ a. Documenting Discharge checklist
- ☐ b. Complete a discharge summary
- ☐ c. Communicating the planned discharge
- ☐ d. Insuring clinician criteria for discharge

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo



Exam Center:

Assosa Unveristy

Department:

Nursing (Comprehensive)

Back

Question 32

Answer saved

Marked out of 1.00

Remove flag

Time left 0:17:46

A 67-year-old patient is admitted at a hospital for the diagnosis of COPD. A nurse has developed a nursing diagnosis of ineffective airway clearance and planed chest physiotherapy. While the nurse explains how to provide the procedure using percussion, vibration and postural drainage, the patient asked for further clarification about the purpose of the procedure. What should be the most appropriate response of the nurse?

- ☐ a. It helps to stimulate blood circulation
- ☒ b. It helps to remove secretion from the lung
- ☐ c. It helps to feel relax
- ☐ d. It helps to prevent blood clotting

Clear my choice

Lenovo

MoEEE

Time left 0:28:43

Question 41

er saved
ed out of
remove flag

A 33-year-old patient came to a hospital with complaints of peripheral edema and coca-colored urine. The patient reported a history of frequent attack of tonsillitis and skin infections. On laboratory investigation, the patient's Glomerular Filtration Rate reads 100ml/minute/1.73 m² (normal GFR is 125mL/min/1.73 m²). What is the most likely fluid and electrolyte balance change expected from the patient?

- ☐ a. Decreased urine output
- ☐ b. Hypernatremia
- ☒ c. Hypokalemia
- ☐ d. Marked hypertension

Clear my choice

Previous page

Next page

Lenovo



MoEEE

MoE - Exit Exam

Nursing (Comprehensive)

Back

Question 28

Answer saved

Marked out of 1.00

Remove flag

A 26-year-old patient came to the ENT clinic with the diagnosis of chronic otitis media. The patient has purulent ear discharge, tenderness around the external ear, and complained reduced hearing. What is the priority nursing diagnosis for this patient?

- ☐ a. Impaired tissue integrity
- ☐ b. Pain
- ☒ c. Body image disturbance
- ☐ d. Risk for infection

Clear my choice

Time left 0:11:27

Previous page

Next page

Lenovo

MoEEE

Time left 0:28:43

Question 41

er saved
ed out of
remove flag

A 33-year-old patient came to a hospital with complaints of peripheral edema and coca-colored urine. The patient reported a history of frequent attack of tonsillitis and skin infections. On laboratory investigation, the patient's Glomerular Filtration Rate reads 100ml/minute/1.73 m² (normal GFR is 125mL/min/1.73 m²). What is the most likely fluid and electrolyte balance change expected from the patient?

- ☐ a. Decreased urine output
- ☐ b. Hypernatremia
- ☒ c. Hypokalemia
- ☐ d. Marked hypertension

Clear my choice

Previous page

Next page

Lenovo



MoEEE

MoE - Exit Exam

Nursing (Comprehensive)

Back

Question 28

Answer saved

Marked out of 1.00

Remove flag

A 26-year-old patient came to the ENT clinic with the diagnosis of chronic otitis media. The patient has purulent ear discharge, tenderness around the external ear, and complained reduced hearing. What is the priority nursing diagnosis for this patient?

- ☐ a. Impaired tissue integrity
- ☐ b. Pain
- ☒ c. Body image disturbance
- ☐ d. Risk for infection

Clear my choice

Time left 0:11:27

Previous page

Next page

Lenovo



MoEEE

MoEE - Exit Exam

Nursing (Comprehensive)

[Back](#)

Time left 0:27:33

Question 42

Answer saved

Marked out of 1.00

Remove flag

A 32 weeks gestational age pregnant woman is admitted in obstetric ward for pregnancy-induced hypertension. During admission, the woman had blurred vision, blood pressure 160/110mmHg, protein urea 0.3gm and oliguria. The woman was conscious and she had no any seizure. She also took the recommended intervention to manage her problem. What is the most appropriate outcome criterion to evaluate the woman's prognosis?

- ☒ a. Urine output 600ml/24hrs
- ☐ b. Absence of blurred vision
- ☐ c. Proteinuria 1+
- ☐ d. Blood pressure 140/90mmHg

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo



MoEEE

MoE - Exit Exam

Exam Center:

Assosa Unveristy

Department:

Nursing (Comprehensive)

Back

Question 27

Answer saved

Marked out of 1.00

Remove flag

A 35-year-old female patient was admitted to a gynecologic inpatient unit with the diagnosis of fistula. The gynecologist planned to do surgical intervention and requested the newly assigned nurses to prepare appropriate surgical equipments for this procedure. What is the most likely category of equipment that should be prepared for this procedure?

- ☐ a. Probing and dilating
- ☐ b. Exposing and retracting
- ☒ c. Grasping and clamping
- ☐ d. Cutting and dissecting

Previous page

Next page

Lenovo





MoEEE

MoE - Exit Exam

Exam Center:

Assosa Unveristy

Department:

Nursing (Comprehensive)

Back

Question 27

Answer saved

Marked out of
1.00

Remove flag

A 35-year-old female patient was admitted to a gynecologic inpatient unit with the diagnosis of fistula. The gynecologist planned to do surgical intervention and requested the newly assigned nurses to prepare appropriate surgical equipments for this procedure. What is the most likely category of equipment that should be prepared for this procedure?

- ☐ a. Probing and dilating
- ☐ b. Exposing and retracting
- ☒ c. Grasping and clamping
- ☐ d. Cutting and dissecting

Previous page

Next page

Time left 0:

Lenovo



MoEEE

MoE - Exit Exam

Nursing (Comprehensive)

Back

Time left 0:26:08

Question 43

Answer saved

Marked out of 1.00

Flag question

During a seminar presentation, a graduating student presents about the mode of transmission of leprosy. But students were confused about the main method of transmission and asked the presenter for clarification. What is the best answer to be explained by the presenter?

- ☐ a. Through respiratory secretion
- ☐ b. Through mother to child trans placental
- ☐ c. Through sexual contact
- ☒ d. Through broken skin and drainage

Clear my choice

Previous page

Next page

Lenovo



Nursing (Comprehensive)

[Back](#)

Question 28

Answer saved

Marked out of 1.00

[Remove flag](#)

A 26-year-old patient came to the ENT clinic with the diagnosis of chronic otitis media. The patient has purulent ear discharge, tenderness around the external ear, and complained reduced hearing. What is the priority nursing diagnosis for this patient?

- ☐ a. Impaired tissue integrity
- ☐ b. Pain
- ☒ c. Body image disturbance
- ☐ d. Risk for infection

[Clear my choice](#)[Previous page](#)

Lenovo



MoEEE

MoE - Exit Exam

Nursing (Comprehensive)

Back

Time left 0:20:

Question 15

Answer saved

Marked out of 1.00

Remove flag

A 55-year-old woman is scheduled to have a surgery on her lower abdomen for an inguinal hernia. The surgeon told that anesthetizing her lower body part is enough to perform the procedure safely. The Anesthetist decides to anesthetize the whole body part just below to the anesthesia injection site. What kind of anesthesia is most appropriate in this case?

- ☐ a. Nerve blocks
- ☐ b. Spinal and epidural blocks
- ☐ c. Local anesthesia
- ☒ d. General anesthesia

Clear my choice

Previous page

Next page

Lenovo

Nursing (Comprehensive)

Time left: 02:34

Question 12

Answer saved

Marked out of 1.00

Remove flag

A 36-year-old female patient comes to a health institution with complaint of skin rash on her genital area for two months. She has hard symmetrical chancre but do not itch and the serological test (VDRL) is positive. The chancre was painless and disappeared within one month. Her husband tells her that he had similar case for the last six months. What is the best clinical feature to diagnose this case?

- ☐ a. Positive serological test
- ☒ b. Hard symmetrical chancre
- ☐ c. Non itchy chancre
- ☐ d. Chancre on genital area

Clear my choice

Previous page

Next page

Lenovo



Nursing (Comprehensive)

[Back](#)**Question 99**Not yet
answeredMarked out of
1.00

Flag question

A 72-year-old patient came to a medical unit with complaints of sudden onset of visual hemi-paresis, confusion, trouble speaking, and impaired mobility. Assessment findings showed history of poorly controlled hypertension. The physician of the patient stroke and put the patient on oxygen. What is the initial treatment option for this patient?

- ☐ a. Anti-hypertensive
- ☐ b. Thrombolytic
- ☐ c. Anti-platelet
- ☐ d. Analgesics treatment

[Next](#)

Lenovo

Nursing (Comprehensive)

[Back](#)**Question 14**

Answer saved

Marked out of
1.00[Remove flag](#)

Time left

A male patient with closed arm fracture has stayed on traction for about two weeks. The bone healing is progressing well. The nurse prepares the patient for discharge from hospital with full arm cast. During discharge, the nurse wanted to give advice to the patient about the problem he may face during his home stay. What is the most likely problem that the nurse should focus on giving patient education?

- ☐ a. Fat embolism
- ☐ b. Compartment syndrome
- ☒ c. Infection
- ☐ d. Muscle atrophy

[Clear my choice](#)

Lenovo





Nursing (Comprehensive)

[Back](#)

Time left: 1

Question 13

Answer saved

Marked out of 1.00

[Remove flag](#)

A 45-year-old male patient with persistent severe asthma is transferred from emergency department to medical ward for admission. A nurse responsible for the care received the patient and provided him with inpatient treatment bed. What is the most appropriate position for maintaining the patient?

- ☐ a. Semi fowler
- ☐ b. Supine
- ☒ c. Left lateral
- ☐ d. Prone

[Clear my choice](#)[Previous page](#)[Next page](#)

Lenovo

Nursing (Comprehensive)

Back

Time left 3:34:34

Question 81

Not yet
answered

Marked out of
1.00

Flag question

A nurse was giving care for pregnant women at ANC clinic. The MUAC measurement result for a 26 week normotensive mother is 21 cm, for a 30 weeks HIV positive mother 21 cm, for a 38 week RH negative mother 21cm and for 16 week primigravida mother 21cm. Which mother mostly requires nutritional counseling?

- ☐ a. Primigravida mother
- ☐ b. HIV positive mother
- ☐ c. RH negative mother
- ☐ d. Normotensive mother

Previous page


Next page

Quiz n

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
82	83
91	92



Type here to search



Lenovo

71°F Sun



Milki health science tutorial center



+251910690594, +251911058885

Tajaajiloota nuti kenninu



Barnoota dabalataa kennuu
Exit Exam'f/ COCf isin qopheessuu.
Leenjiwwan gaggabaaboo kennuu fi K.K.F.



Teessoo: Magaalaa Naqamtee, Gamoo OSSA, Darbii 3ffaa

Time left 0:46:34

An investigator conducted formal evaluation for a screening program and found out that survival among screened individuals appears longer than that for non-screened individuals. What type of screening bias did she introduce?

- ☐ a. Lead-time bias
- ☐ b. Recall bias
- ☐ c. Length-biased sampling
- ☐ d. Volunteer bias

Next question

question

Time left 0:45:00

A manager of health office said, "In resource poor countries like us one does not go in search of a cure for ring worm while leaving leprosy unattended." In above quotation, which step of the planning is emphasized?

- ☐ a. Problem identification
- ☐ b. Situational analysis
- ☐ c. Problem prioritization
- ☐ d. Objective setting

Next question

Question 81

Not yet

answered

Marked out of
1.00

Flag question

A nurse was giving care for pregnant women at ANC clinic. The MUAC measurement result for a 26 week normotensive mother is 21 cm, for a 30 weeks HIV positive mother 21 cm, for a 38 week RH negative mother 21cm and for 16 week primigravida mother 21cm. Which mother mostly requires nutritional counseling?

- ☐ a. Primigravida mother
- ☐ b. HIV positive mother
- ☐ c. RH negative mother
- ☐ d. Normotensive mother

A 26-year-old Gravid II Para I pregnant woman came to ANC clinic with complaints of headache, dyspnea, and fatigue for three days. Her pulse rate was 78/min, Blood pressure was 100/60 mmHg and fetal heart rate was 140/min. On physical examination, she had pale conjunctiva and 32 weeks gestational age pregnancy. Her hemoglobin concentration was 10g/dL. What is the most likely indicative characteristic for the woman's problem?

- ☐ a. Headache
- ☐ b. Pale conjunctiva
- ☐ c. Dyspnea
- ☐ d. Hgb 10 g/dl

Question 43

Not yet
answered

Marked out of
1.00

Flag question

Time left 3:52:36

A nurse observed medication administration error made by his friend working in medical ward. Due to this, the patient developed minor complication and treated. What is the first thing the nurse must do?

- ☐ a. Confront his friend
- ☐ b. Follow secretly and supervise
- ☐ c. Report to the nurse director
- ☐ d. Report to the legal judiciary body

Previous page

Next page

Time left 3:32:22

Question 88

Not yet
answeredMarked out of
1.00

Flag question

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☐ a. Jaundice
- ☐ b. No jaundice
- ☐ c. Severe jaundice
- ☐ d. Uncomplicated jaundice

Previous page

Next page

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
	75

Bạch Hà ãlën
today at 5:12 PM



688 of 693

...sing (Comprehensive)

25

... of

question

... page

Time left: 1:00:00

A 42-year-old male patient was going to have amputation due to gangrene. If he is amputated, he will lose his leg, which will be transferred to his brother according to cultural practices in the area. If the procedure is not done, the patient will die. The nurse in charge convinced the patient about the procedure. What principle is most likely applied in the case?


- ☐ a. Compassion
- ☐ b. Paternalism
- ☐ c. Veracity
- ☐ d. Proportionality

Next page

type here to search



Lenovo



Question 43

Not yet
answered

Marked out of
1.00

Flag question

Time left 3:52:36

A nurse observed medication administration error made by his friend working in medical ward. Due to this, the patient developed minor complication and treated. What is the first thing the nurse must do?

- ☐ a. Confront his friend
- ☐ b. Follow secretly and supervise
- ☐ c. Report to the nurse director
- ☐ d. Report to the legal judiciary body

Previous page

Next page

48

A nurse working in ART clinic, heard about HIV positive patient who is in love with his neighbor during counseling on his status. The nurse counseled him to stop the relationship. The patient offended and shouted; "I want to infect her and many more so shut-up your mouth". What is the appropriate action to be taken by the nurse?

- ☐ a. Accuse him for endangering the health of the public
- ☐ b. Further communication and counseling with the patient
- ☐ c. Tell his HIV status to the emergency nurse
- ☐ d. Tell his status to management team



Exam Center:

Assosa Univeristy

Department:

Nursing (Comprehensive)

[Back](#)

Question 6

Answer saved

Marked out of 1.00

[Remove flag](#)

A 27-year-old patient comes to a hospital with the complaints of acute onset profuse and purulent urethral discharge, urgent itching sensation of the urethra and burning sensation during micturition. The patient have also painless and regular ulcer of genitalia. What is the most likely diagnosis of this patient?

- ☒ a. Syphilis
- ☐ b. Herpes genitalia
- ☐ c. Chancroid
- ☐ d. Gonorrhea

[Clear my choice](#)

Time left

[NCOMPREHENSIVE055: Nursing \(Comprehensive\) Part 1 Exit E...](#)[Nursing \(Comprehensive\) Part 1 Exit Exam Sene 2016 \(page 6...](#)[Next page](#)

Lenovo

Nursing (Comprehensive)

Back

Question 18

Not yet
answered

Marked out of
1.00

Flag question

A nurse researcher studied a hypothesis which read as "Frequent consumption of fruits and vegetables is protective for the occurrence of stomach cancer". He recruited 198 subjects recently diagnosed with stomach cancer and 396 subjects free of stomach cancer. All subjects were asked about their fruit and vegetable consumption during the past one year. What is the most likely study design that the researcher used?

- ☐ a. Ecological
- ☐ b. Case-control
- ☐ c. Cohort
- ☐ d. Cross-sectional

Time left 4:19:52

Quiz navig

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48
55	56	57
64	65	66
73	74	75

Previous page

Next page

Type here to search



68°F Mostly sunny

Lenovo

Nursing (Comprehensive)

Back

Question 56

Not yet
answered

Marked out of
1.00

Flag question

Time left 3:45:49

A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment kebeles. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues. What is the most likely problem prioritization criterion addressed by the head?

- ☐ a. Magnitude
- ☐ b. Sustainability
- ☐ c. Community concern
- ☐ d. Feasibility

Previous page

Next page



Type here to search



Nursing (Comprehensive)

Back

Question 43

Not yet
answered

Marked out of
1.00

Flag question

A nurse observed medication administration error made by his friend working in medical ward. Due to this, the patient developed minor complication and treated. What is the first thing the nurse must do?

- ☐ a. Confront his friend
- ☐ b. Follow secretly and supervise
- ☐ c. Report to the nurse director
- ☐ d. Report to the legal judiciary body

Previous page

Time left 3:52:36

Next page

Quiz na

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
82	83
91	92



Type here to search



Lenovo



Question 92

Not yet
answered

Marked out of
1.00

Flag question

A neonate, who was diagnosed with neonatal sepsis, was admitted in a hospital. The nurse in charge examined the baby and the findings showed central cyanosis, chest in drawing, tachycardia and tachypnea. What is the most likely nursing diagnosis of this case?

- ☐ a. Ineffective breathing pattern
- ☐ b. Ineffective airway clearance
- ☐ c. Ineffective tissue perfusion
- ☐ d. High risk for hypoventilation

Previous page

Next page

Question 56

Not yet
answeredMarked out of
1.00

Flag question

A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment kebeles. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues. What is the most likely problem prioritization criterion addressed by the head?

- ☐ a. Magnitude
- ☐ b. Sustainability
- ☐ c. Community concern
- ☐ d. Feasibility



Matching (comprehensive)

[Back](#)

Question 47

Not yet
answeredMarked out of
1.00

Flag question

Time left 3:50:44

A 16-year-old boy with brief psychotic disorder came to a clinic with contractions of muscles that result in abnormal movement, tongue protrusion and difficulty of breathing. Symptoms developed after taking two doses of prescribed haloperidol 5 mg orally. What is the best initial treatment for this boy?

- ☐ a. Reduce dose of antipsychotic
- ☐ b. Give 10 mg diazepam intravenously
- ☐ c. Discontinue the drug and refer
- ☐ d. Switch to another antipsychotic


[Previous page](#)[Next page](#)

1	2
3	4
5	6
7	8
9	10
11	12
13	14
15	16
17	18
19	20
21	22
23	24
25	26
27	28
29	30
31	32
33	34
35	36
37	38
39	40
41	42
43	44
45	46
47	48
49	50
51	52
53	54
55	56
57	58
59	60
61	62
63	64
65	66
67	68
69	70
71	72
73	74
75	76
77	78
79	80
81	82
83	84
85	86
87	88
89	90
91	92
93	94
95	96
97	98
99	100

Finish attempt



Type here to search



A 26-year-old Gravid II Para I pregnant woman came to ANC clinic with complaints of headache, dyspnea, and fatigue for three days. Her pulse rate was 78/min, Blood pressure was 100/60 mmHg and fetal heart rate was 140/min. On physical examination, she had pale conjunctiva and 32 weeks gestational age pregnancy. Her hemoglobin concentration was 10g/dL. What is the most likely indicative characteristic for the woman's problem?

- ☐ a. Headache
- ☐ b. Pale conjunctiva
- ☐ c. Dyspnea
- ☐ d. Hgb 10 g/dl

Question 81

Not yet
answeredMarked out of
1.00

Flag question

A nurse was giving care for pregnant women at ANC clinic. The MUAC measurement result for a 26 week normotensive mother is 21 cm, for a 30 weeks HIV positive mother 21 cm, for a 38 week RH negative mother 21cm and for 16 week primigravida mother 21cm. Which mother mostly requires nutritional counseling?

- ☐ a. Primigravida mother
- ☐ b. HIV positive mother
- ☐ c. RH negative mother
- ☐ d. Normotensive mother

48

A nurse working in ART clinic, heard about HIV positive patient who is in love with his neighbor during counseling on his status. The nurse counseled him to stop the relationship. The patient offended and shouted; "I want to infect her and many more so shut-up your mouth". What is the appropriate action to be taken by the nurse?

- ☐ a. Accuse him for endangering the health of the public
- ☐ b. Further communication and counseling with the patient
- ☐ c. Tell his HIV status to the emergency nurse
- ☐ d. Tell his status to management team

48

A nurse working in ART clinic, heard about HIV positive patient who is in love with his neighbor during counseling on his status. The nurse counseled him to stop the relationship. The patient offended and shouted; "I want to infect her and many more so shut-up your mouth". What is the appropriate action to be taken by the nurse?

- ☐ a. Accuse him for endangering the health of the public
- ☐ b. Further communication and counseling with the patient
- ☐ c. Tell his HIV status to the emergency nurse
- ☐ d. Tell his status to management team

48

A nurse working in ART clinic, heard about HIV positive patient who is in love with his neighbor during counseling on his status. The nurse counseled him to stop the relationship. The patient offended and shouted; "I want to infect her and many more so shut-up your mouth". What is the appropriate action to be taken by the nurse?

- ☐ a. Accuse him for endangering the health of the public
- ☐ b. Further communication and counseling with the patient
- ☐ c. Tell his HIV status to the emergency nurse
- ☐ d. Tell his status to management team

Exam Center:

Debre Tabor University

Department:

g (Comprehensive)

Time left 0:45:47

A researcher sets a level of significance 90% to compute confidence interval for mean age of women at first child birth. What would happen to the width of confidence interval if the level of confidence increases to 99%?

- ☐ a. Unchanged
- ☐ b. Decreased
- ☐ c. Increased
- ☐ d. Precise

Next question

Previous question

Ex

1	
10	
19	
28	
37	3
46	47
55	56
64	65
73	74

Time left 0:45:00

A manager of health office said, "In resource poor countries like us one does not go in search of a cure for ring worm while leaving leprosy unattended." In above quotation, which step of the planning is emphasized?

- ☐ a. Problem identification
- ☐ b. Situational analysis
- ☐ c. Problem prioritization
- ☐ d. Objective setting

Next question

Question 56

Not yet
answered

Marked out of
1.00

Flag question

A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment kebeles. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues. What is the most likely problem prioritization criterion addressed by the head?

- ☐ a. Magnitude
- ☐ b. Sustainability
- ☐ c. Community concern
- ☐ d. Feasibility

Question 56

Not yet
answered

Marked out of
1.00

Flag question

A head of health center discussed the bottlenecks of health care insurance service utilization with community leaders who came from the catchment kebeles. During discussion, the community leaders urged the health center head to address frequent hospital supply stock-out issues. What is the most likely problem prioritization criterion addressed by the head?

- ☐ a. Magnitude
- ☐ b. Sustainability
- ☐ c. Community concern
- ☐ d. Feasibility

Time left 3:32:22

Question 88

Not yet
answeredMarked out of
1.00

Flag question

A two-day-old male neonate was admitted to NICU for thermal care. On physical examination, the newborn has pink palms and soles, and yellow discoloration in the face and eyes. Based on IMNCI classification, what is the most likely diagnosis of the neonate?

- ☐ a. Jaundice
- ☐ b. No jaundice
- ☐ c. Severe jaundice
- ☐ d. Uncomplicated jaundice

Previous page

Next page

1	2
10	11
19	20
28	29
37	38
46	47
55	56
64	65
73	74
	75



Question 59

Not yet answered

Marked out of 1.00

 [Flag question](#)

A nurse who is working as a ward head in a hospital developed annual plan with hospital staff teams. However, they failed to achieve the plan because of inability of mobilizing the human and material resources from the hospital. What is the most likely management function that causes low achievement of this ward?

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

- ☐ a. Planning
- ☐ b. Controlling
- ☒ c. Organizing
- ☐ d. Directing

[Clear my choice](#)

[Previous page](#)

[Next page](#)

Quiz navigation



**Question 60**

Time left 3:19:59

Not yet answered

Marked out of 1.00

 [Flag question](#)

A 30-year-old female patient was diagnosed with ovarian cancer and the oncologist decided the need for radiation therapy. The nurse in charge explained the risk of infertility secondary to radiation therapy. She refused radiation therapy and at the same time she had strong desire to have children. The procedure was done to save her life; she was cured of ovarian cancer and develop infertility What ethical principle most likely forced the health care team's action?

- ☐ a. Autonomy
- ☒ b. Beneficence
- ☐ c. Justice
- ☐ d. Nonmaleficence

[Clear my choice](#)

**EXAM ZONE**
MASTER EVERY EXAM

ExamZone

 Flag question

A 25-year-old male patient was admitted to an ICU and put on mechanical ventilator following car accident. On assessment; his eyes are not open to any stimuli, produces incomprehensible sounds and has abnormal flexion of motor response. What is the level of consciousness of this patient using Glasgow coma scale?

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

- ☐ a. Stupors
- ☐ b. Somnolent
- ☐ c. Comatose
- ☐ d. Semiconscious

Prepare for Exit Exam! Join our Telegram Channel for all Department @ExitExamPrep

[Previous question](#)[Finish exam ...](#)

Exam Navigation

1	2	3	4	5	6	7	8
---	---	---	---	---	---	---	---