

ETHIOPIAN HEALTH PROFESSIONALS LICENSING EXAMINATION(EHPLE) PRACTICE TEST BOOKLET 2025

DENTAL MEDICINE



ጤና ሚኒስቴር - ኢትዮጵያ
MINISTRY OF HEALTH-ETHIOPIA

የዜጎች ጤና ለሃገር ብልጽግና!
HEALTHIER CITIZENS FOR PROSPEROUS NATION!



Message from the State Minister, Ministry of Health – Ethiopia



Mrs. Frehiwot Abebe

Improving the quality of healthcare remains a critical global priority, essential for achieving sustainable development and universal health coverage. One of the most effective ways to ensure high standards in healthcare delivery is through the consistent assessment of professional competencies. In line with this commitment, the Ministry of Health of Ethiopia launched the Ethiopian Health Professionals Licensing Examination (EHPLE) in 2019 for undergraduates in seven core health disciplines. Since its inception, the program has expanded to encompass 17 health professions, reflecting our continued effort to build a competent and responsive health workforce.

EHPLE was established with the primary aim of identifying and certifying health professionals who possess the minimum competencies required to provide safe and effective care. This ensures that all practicing professionals are equipped to meet the expectations of the communities they serve. The exam is coordinated by the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (LEO), which consists of four specialized desks. The LEO plays a vital role in system strengthening, enabling the exam to be conducted with increased efficiency, organization, and quality assurance.

It is important to recognize that the EHPLE is not a conventional academic or recruitment examination. Rather, it is a rigorous, competency-based assessment designed to evaluate the practical skills and critical thinking abilities that are essential for professional practice. This *Practice Items Booklet* is part of our broader strategy to support examinees and educators in preparing for the licensing process. It offers a realistic overview of the exam structure, promotes a deeper understanding of its objectives, and fosters transparency and sustainability in implementation.

The development of this booklet has been made possible through the collaborative efforts of experts from higher education institutions, the Addis Ababa University Testing Center, and professional associations. Their insights and dedication have been instrumental in shaping this resource. I extend my heartfelt thanks to all contributors and the Ministry of Health team for their tireless work and steadfast commitment to excellence in healthcare regulation.

We believe this booklet will serve as a valuable tool for candidates preparing for the examination and for educators guiding them toward success. Together, we are building a stronger, more competent health workforce for Ethiopia.

Introduction EHPLE

Welcome to the 2025 Edition of the Ethiopian Health Professionals Licensing Examination Practice Test Booklet.

This resource is designed to support candidates in their preparation for the national licensing examination, ensuring they are well-equipped with the knowledge and clinical reasoning skills required to deliver competent and ethical healthcare across Ethiopia.

In Ethiopia, the licensing examination serves as a pivotal tool in evaluating the competency of health professionals who have completed their academic training and are ready to transition into independent practice. The exam not only ensures public safety by verifying professional readiness but also upholds national and international standards in healthcare service delivery. As the healthcare system continues to expand and evolve, a strong, well-regulated professional workforce becomes even more essential.

This practice booklet contains 200 multiple-choice questions modeled after the structure, content, and format of the actual licensing examination. The questions span across core clinical domains essential for entry-level practice. This includes areas such as Patient care, leadership and management, ethics, and Health Promotion, as outlined in the 2024 Information Booklet.

The booklet is intended to:

- Familiarize candidates with the structure and cognitive level of the actual licensing examination.
- Offer opportunities for self-assessment and identification of strengths and areas for improvement in knowledge.
- Encourage critical thinking and clinical decision-making in scenarios reflective of real-world healthcare practice in Ethiopia.
- Reduce exam-related anxiety by providing realistic practice experiences.

Administered under the auspices of the Ministry of Health and relevant regulatory bodies, the Ethiopian Health Professionals Licensing Examination is grounded in evidence-based frameworks and aligned with both national needs and global best practices. It has been structured to assess a range of competencies, including medical knowledge, communication, professionalism, and safe clinical practice.

This practice test is not a predictive tool, and performance on it does not guarantee similar outcomes on the actual exam. However, it provides an invaluable opportunity for familiarization with question styles, pacing strategies, and content review. We strongly recommend that candidates simulate real testing conditions while using this resource, including timing their responses and reviewing rationale for both correct and incorrect answers.

The goal of licensing examinations is not only to regulate entry into the profession but also to inspire excellence and lifelong learning. By engaging in this practice test, you are taking a proactive step toward professional development, accountability, and a commitment to serving your patients with competence and compassion.

We wish you success on your licensing journey and beyond.

Acknowledgements

This *Practice Test Items Booklet for the Ethiopian Health Professionals Licensing Examination* is the result of a collaborative effort among dedicated professionals committed to advancing the quality of health sciences education in Ethiopia. The Ministry of Health extends its sincere appreciation to all those who contributed their time, expertise, and insights into the development of this essential resource.

Special recognition is given to the item developers and reviewers from various higher education institutions, both public and private, whose academic rigor and practical experience were instrumental in shaping the content and ensuring its relevance to real-world healthcare practice. Their contributions have helped create a high-quality, standardized practice tool that reflects national competencies and global benchmarks.

We are especially grateful to the team at the Addis Ababa University Testing Center, whose technical support and academic guidance were critical throughout the process. Their leadership in assessment design and educational evaluation has significantly strengthened the quality of this booklet.

We also thank the committed staff of the Ministry of Health, particularly those within the Health and Health-Related Institutions and Professionals' Regulatory Lead Executive Office (HHRIPR LEOs), whose coordination, oversight, and unwavering dedication ensured the successful completion of this project.

Together, these collective efforts reflect our shared vision of building a competent, ethical, and resilient health workforce for Ethiopia.

Dental Medicine

1. A 40-year-old edentulous patient presented to dental setup with a complaint of poor jaw relation compared to her previous denture. Extraoral examination shows a symmetric face and an intact temporomandibular joint. On intraoral examination, the residual ridge is thick enough to accept a new denture. However, the patient is curious about the size of the teeth.

What is the most appropriate facial feature to be considered for this patient?

- A. Facial form
- B. Facial profile
- C. Peri-oral features
- D. Lower facial height

ANSWER: A

2. A 50-year-old male patient with a complete denture came to the dental clinic with poor retention and excessive tissue irritation. He had a history of sialadenectomy a year ago after getting diagnosed with sialolithiasis. Intraoral examination shows reddish discoloration around the margin of the denture with associated dry mouth, and the remaining saliva is thick, ropy, which alters the seat of the denture.

What is the classification of saliva based on viscosity?

- A. Class 1
- B. Class 2
- C. Class3
- D. Class4

ANSWER: C

3. A 50-year-old male patient visited a dental clinic with a chief complaint of complete tooth loss in the lower jaw. He has been diabetic for the past 10 years and is on medication for it. During clinical examination, he was found to have an edentulous maxilla with a residual ridge of acceptable width and height. The dentist planned a complete denture, loaded the tray with alginate, and removed it from the patient's mouth after setting.

What is the most appropriate next step in impression taking?

- A. Border molding
- B. Pouring the cast
- C. Checking for the defects
- D. Trimming the impression

ANSWER: C

4. A 25-year-old male patient came to dental OPD after sustaining maxillofacial trauma. On clinical examination, all upper front teeth are missing, and the remaining maxillary teeth are fractured with pulp exposure. He has pain secondary to exposed pulp chambers. On radiographic examination, all upper front teeth are avulsed, and there is a crown-root fracture.

What is the most appropriate treatment plan for this patient?

- A. Analgesics followed by an implant-supported denture
- B. Endodontic therapy followed by an overdenture
- C. Extraction followed by treatment denture
- D. Splinting followed by an immediate denture

ANSWER: B

5. A Patient who has a complete denture came to a dentist with a complaint of a clicking sound while eating or drinking. The dentist corrected premature contact and overextended margins a week ago during his checkup on the second visit. Otherwise, there is no drooling at the corners of the mouth, neuromuscular, and temporomandibular joint problems.

What further evaluation is needed for the final treatment plan of this patient?

- A. Vertical dimensions at occlusion
- B. Vertical dimension at rest
- C. Facial profile
- D. Facial form

ANSWER: A

6. A 65-year-old male patient came to dental OPD with a complaint of pain in the upper jaw. He came by carrying the denture in his pocket because the pain gets aggravated when he inserts the denture. Before the occurrence of these problems, he had been wearing the denture day and night, not adhering to the precautions. He has been a well-known diabetic patient for the past 10 years. On clinical examination, there is a glandular type of erythematous lesion on denture-bearing areas of the hard palate associated with angular cheilitis.

What is the most likely initial management for this patient?

- A. Reline the denture
- B. Change the denture
- C. Treating diabetes
- D. Treat the infection

ANSWER: D

7. A 50-year-old female patient came to a dental OPD with a complaint of a burning sensation on the supporting structures of the denture and tongue. The pain starts in the morning and aggravates during the day, and with hot drinks. Clinically, there are no overt clinical signs and symptoms.

What is the most effective management for this case?

- A. Surgery
- B. Analgesics
- C. Antibiotics
- D. Counseling

ANSWER: D

8. A 40-year-old male patient came to dental OPD with a complaint of mobility of previous artificial teeth. On clinical examination, he has a removable partial denture, which is mobile because of loss of supporting tissue on the retromolar region. The doctor planned to do relining of the denture over mobile tissue on the crest of the ridge.

What is the most appropriate impression material for this patient?

- A. Alginate
- B. Zinc oxide eugenol
- C. Polysulfide silicone
- D. Mouth temperature wax

ANSWER: B

9. A 55-year-old male patient came with a complaint of missing teeth on the lower back jaw. On clinical examination, he has missing teeth numbers 34, 35, and 36, but all remaining teeth in his oral cavity are there with good periodontal support.

What is the classification of the partially edentulous arch of this patient?

- A. Class I
- B. Class II
- C. Class III
- D. Class IV

ANSWER: C

10. A 32-year-old female patient visited a dental clinic with multiple missed teeth on the lower jaw. She has a dental history of extraction due to caries and failed root canal therapy. On clinical examination, she has missing teeth numbers 46, 47, 48, 36, and 34.

What is the classification of the partially edentulous arch of this patient?

- A. Class I modification I
- B. Class II modification I
- C. Class III modification III
- D. Class IV modification IV

ANSWER: B

11. A 35-year-old female patient came to a dental clinic with a complaint of multiple missing teeth on the upper jaw. She has no medical condition that contraindicates prosthodontic procedures. On clinical examination, teeth numbers 16 up to 26 are missing, but the remaining four upper posterior teeth are periodontally intact with no mobility. The residual ridge is broad and flat.

What is the most appropriate treatment plan for this patient?

- A. Removable partial denture
- B. Conventional fixed partial denture
- C. Resin-bonded fixed partial denture
- D. Implant-supported fixed partial denture

ANSWER: D

12. A 57-year-old male patient came to the dental OPD with a chief complaint of multiple missed teeth on the lower jaw. On intraoral examination, tooth numbers 38 and 48 have an acceptable inclination, but tooth numbers 37 and 47 are buccally tipped; otherwise, they are periodontally intact. The doctor wanted to construct a removable partial denture.

What is the most appropriate treatment plan for this patient?

- A. Crown
- B. Splinting
- C. Orthodontics
- D. Enameloplasty

ANSWER: A

13. A 25-year-old male patient came to the dental clinic to replace a missing tooth. There was a fractured root remnant, which was extracted a year ago on tooth number 11, and the site is completely healed. Clinical examinations show a missing tooth number 11 from the oral cavity. On intraoral periapical X-ray, there is no root remnant. The doctor planned fixed prosthodontic therapy, but the patient didn't want the preparation of the anterior tooth as an abutment. The dentist used a permanent first molar as an abutment to replace the missing 11.

What is the most appropriate management of this patient?

- A. Modified fixed removable partial denture
- B. Spring cantilever fixed partial denture
- C. Metal-ceramic fixed partial denture
- D. Acrylic fixed partial denture

ANSWER: B

14. A 25-year-old female patient came to a clinic with a history of sharp pain when she takes cold drinks and food, but it subsides briefly. On intraoral examination, there is a cavity on the upper first premolar; no discoloration or mobility is detected, and percussion is negative. Intraoral periapical radiograph shows that lamina dura is normal, but there is radiolucency involving enamel and dentin.

What is the most likely diagnosis in this case?

- A. Acute pulpitis
- B. Necrotized pulp
- C. Reversible pulpitis
- D. Irreversible pulpitis

ANSWER: C

15. A 48-year-old male patient came to a clinic with a history of a pink tooth. He had been a box player for 20 years. The radiographic result revealed radiolucent enlargement of the pulp canal.

What is the most likely diagnosis in this case?

- A. Pulp stone
- B. Pulp necrosis
- C. Pulp degeneration
- D. Internal resorption

ANSWER: D

16. A 40-year-old male patient came to a clinic with a complaint of swelling, which has been slightly enlarged over time, on the upper right lateral incisor. Clinically, the tooth is discolored, and there is palatal expansion. On intraoral radiographs, a round-shaped radiolucency, outlined by a narrow radiopaque margin at the apex of the tooth, is detected.

What is the most likely diagnosis in this case?

- A. Radicular cyst
- B. Dentigerous cyst
- C. Periapical granuloma
- D. Acute alveolar abscess

ANSWER: A

17. A 23-year-old female patient came to a clinic with a complaint of slight discomfort when food enters the cavity on 25 teeth. After she brushes her teeth, the discomfort subsides. On IOPA, the cavity doesn't involve the pulp.

What is the most appropriate treatment plan for this patient?

- A. Pulpotomy
- B. Restoration
- C. Pulpectomy
- D. Root canal treatment

ANSWER: B

18. A 33-year-old male patient came to a clinic with a history of swelling and discoloration on 11 and 12th for more than 3 years. He said that he had a previous trauma 6 years ago. On clinical examinations, percussion is negative, and there is palatal expansion. On IOPA, at the apex of 11 & 12 ovoid-shaped radiolucency, outlined by a narrow radiopaque margin, is revealed.

What is the most appropriate initial step in the management of this patient?

- A. Pulpotomy
- B. Obturation
- C. Apicectomy
- D. Root canal treatment

ANSWER: D

19. A 45-year-old male patient came to a clinic with a complaint of a discolored tooth in the upper front area for a year's duration. On clinical examination, there is swelling in the apical region of tooth number 21. An intern is planning to perform root canal treatment.

What is the most appropriate chair position for this case?

- A. 12 o'clock
- B. 11 o'clock
- C. 9 o'clock
- D. 7 o'clock

ANSWER: B

20. A 23-year-old patient came to a clinic with a complaint of tooth pain, which is spontaneous and continuous, in the upper left quadrant. The pain is aggravated on mastication and on lying down. On clinical examination, there is a class II cavity on 26, and percussion is positive. On IOPA, it reveals that there is periapical radiolucency. The doctor has planned to extirpate the pulp.

What is the most appropriate instrument to be used in this case?

- A. File
- B. Reamer
- C. spreader
- D. Barbed broach

ANSWER: D

21. A 24-year-old female patient came to a clinic with a complaint of a toothache in the lower left quadrant of four weeks duration. On clinical examination, there is a class II cavity on 34, and percussion is positive. On radiograph, it reveals periapical radiolucency. The doctor asked the assistant to arrange blue and white color files to start the root canal treatment.

What is the most appropriate name for the files?

- A. 15mm & 30mm
- B. 20mm & 35mm
- C. 25mm & 60mm
- D. 45mm & 50mm

ANSWER: B

22. A 10-year-old male patient came to a clinic with a complaint of a tooth fracture on tooth 11 of one day duration. On clinical and radiographic examination, it was determined that it involved the pulp with an open apex and needed root canal treatment.

What is the most appropriate root canal sealer in this case?

- A. AH 26
- B. Endofill
- C. Calcium hydroxide
- D. Zinc oxide eugenol

ANSWER: C

23. A 40-year-old male patient came to a clinic with a complaint of tooth pain on tooth 36 of one month duration. On clinical and radiographic examination, it was determined that the cavity involved the pulp with periapical radiolucency and needs root canal treatment.

What is the appropriate step in this patient's care?

- A. Cleaning and shaping, Access opening, Obturation
- B. Access opening, Obturation, Cleaning, and shaping
- C. Obturation, Cleaning, shaping, and Access opening
- D. Access opening, Cleaning and shaping, and Obturation

ANSWER: D

24. A 29-year-old male patient came to the OPD with a complaint of tooth pain on tooth 26 of six months duration. On clinical and radiographic examination, it was determined that there is periapical radiolucency and needs root canal treatment. The doctor is trying to locate the mesiobuccal canal orifice.

What is the most likely location of the canal in this patient?

- A. Distally
- B. Palatally
- C. Under the mesiobuccal cusp
- D. Under the disto buccal cusp

ANSWER: C

25. A 32-year-old female patient came to the OPD with a complaint of tooth pain on tooth 46 after root canal treatment. On radiographic examination, there is still periapical radiolucency, and the gutta-percha is short of the apical stop. The tip of the gutta-percha is a little away from the center of the canal, and there is no perforation. Condensation of the obturation is good.

What is the most likely cause in this case?

- A. Ledge formation
- B. Inadequate irrigation
- C. Inadequate recapitulation
- D. Incomplete instrumentation

ANSWER: A

26. A 12-year-old male patient has come to OPD with a complaint of a tooth fracture from a falling accident on his 11th tooth, of one day duration. It was diagnosed as Ellis's class III.

What is the most appropriate way to check the vitality of the pulp?

- A. Thermal
- B. Palpation
- C. Inspection
- D. Percussion

ANSWER: A

27. A 21-year-old male patient came to OPD with a complaint of a fall onto his tooth 11 of one day duration. Clinically, the tooth is displayed away from its long axis. What is the most likely appropriate assessment?
- A. Subluxation
 - B. Lateral luxation
 - C. Intrusive luxation
 - D. Extrusive luxation
- ANSWER: B
28. A nine-year-old male patient came to OPD with a complaint of a tooth that is completely displaced out of its socket of tooth 21, from a fall accident of one and a half hours duration. The tooth was re-implanted after local anesthesia was given. What is the most appropriate initial step in this patient's care?
- A. Soak it in citric acid for fifteen minutes
 - B. Soak it in five percent doxycycline for five minutes
 - C. Soak it in one percent stannous fluoride for thirty minutes
 - D. Soak it in five percent sodium hypochlorite for ten minutes
- ANSWER: B
29. A 43-year-old female patient came to a dental clinic with a complaint of tooth sensitivity on the maxillary first molar. The patient had a history of composite restoration on the causative teeth for six years. Clinically, the tooth is negative to percussion and palpation. On exploration, there is a catching point around the restoration. The dentist takes to rule out dental caries. IOPAR shows radiolucency under the radiopaque area. What is the most likely type of caries in this case?
- A. Root
 - B. Primary
 - C. Residual
 - D. Recurrent
- ANSWER: D
30. A 20-year-old female patient came to the dental clinic for the cleaning of her teeth. Clinical examination reveals detectable cavities in enamel due to dental caries. What is the most likely class of the lesion?
- A. D1
 - B. D2
 - C. D3
 - D. D4
- ANSWER: B
31. A 52-year-old female patient has come to a dental OPD complaining of pain in the upper first molar while biting. Clinically, there is a discoloration of 16 but no sign of a cavity. The dentist wants to rule out a class II cavity. What is the most appropriate examination method to be used in this case?
- A. Palpation
 - B. Percussion
 - C. Exploration
 - D. Auscultation
- ANSWER: C

32. A 23-year-old patient has come to a dental OPD complaining of food stuck between her right mandibular first and second molars for the past two months. Clinically, there is no cavitated tooth that is visualized. The dentist has planned further investigation into the proximal area.

What is the most appropriate type of radiograph to be used in this case?

- A. Intraoral
- B. Periapical
- C. Conventional
- D. Bitewing radiographs

ANSWER: D

33. A 41-year-old male patient has come to the dental clinic for a checkup. Clinically, the patient has poor oral hygiene and has a class II cavity on 47 with food debris inside the cavity. Radiographically, there is radiolucency that involves the enamel.

What is the most appropriate initial step in the management of this case?

- A. Liner application
- B. Removal of debris
- C. Permanent restoration
- D. Local anesthesia administration

ANSWER: B

34. A 29-year-old female patient came to the dental clinic for tooth sensitivity in a cold environment. Clinically, the patient had a class III malocclusion with edge-to-edge bite, and she also had a large class III cavity, which induced tooth sensitivity. The dentist plans to fill the cavity with composite.

What is the most effective type of composite to be used in this case?

- A. Macrofilled
- B. Microfilled
- C. Conventional
- D. Small particle

ANSWER: D

35. A 23-year-old female patient has come to dental OPD with a toothache duration of five days. Clinically, there is a class I cavity on 27 with percussion and palpation negative. Radiographically, there is radiolucency, which involves enamel only. The dentist planned to clean and fill the cavity without further cavity preparation.

What is the most effective restorative material of choice for this patient?

- A. GIC
- B. Amalgam
- C. Composite
- D. Cast metal

ANSWER: A

36. A 35-year-old female patient came to dental OPD with a complaint of a toothache in the upper left quadrant of two days duration. Clinically, there is a class I cavity. The dentist plans to prepare a cavity for restoration. What is the third step of the preparation in this case?

- A. Convenience form
- B. Primary resistance form
- C. Primary retention form
- D. Outline form and initial depth

ANSWER: C

37. A 29-year-old male patient has come to a dental clinic for tooth restoration. Clinically, the patient has a class I cavity on 27. Caries affects the outer layer of dentin only. The dentist plans to restore the tooth with an amalgam restoration.

What is the most appropriate minimal occlusal thickness of restoration in this case?

- A. 1mm
 - B. 1.5mm
 - C. 2mm
 - D. 2.5mm
- ANSWER: B

38. A 23-year-old male patient has come to a dental clinic for tooth restoration. Clinically, the patient has a class I cavity on 27. Clinically, only the outer layer of dentin is affected. The dentist plans to restore the tooth with cast metal.

What is the most appropriate step in increasing retention in this case?

- A. Elastic compression of dentin
- B. Giving a slight undercut in dentin near the pulpal wall
- C. Making an occlusal dovetail to prevent the tilting of the restoration
- D. Give a reverse bevel in class I, class II, and MOD preparations to prevent tipping movements.

ANSWER: C

39. A 35-year-old female patient with diabetes has come to the dental OPD for a checkup. Due to her medical condition, she has a fear of developing dental caries. The patient brushes her teeth twice a day, and she has good oral hygiene. Clinically, there is occlusal groove darkness on the molars.

What is the most appropriate initial step in this patient's care?

- A. Systemic fluoride
- B. Antibiotic treatment
- C. Pit and fissure sealant
- D. Oral hygiene instruction

ANSWER: C

40. A nine-year-old male patient has come to dental OPD for an appointment for tooth restoration. Clinically, there is a moderate class I cavity on 84. The dentist plans to fill the cavity with GIC.

What is the most appropriate bonding mechanism in this case?

- A. Physical
- B. Chemical
- C. Mechanical
- D. Micromechanical

ANSWER: B

41. A 35-year-old patient has come to dental OPD for tooth restoration. Clinically, there is a class I cavity on 36. The dentist has restored the tooth with an amalgam restoration.

What is the most appropriate time to polish the restoration?

- A. 3 hours
- B. 6 hours
- C. 12 hours
- D. 24 hours

ANSWER: D

42. A 20-year-old female patient came to dental clinic for the restoration of her molar tooth. A dentist prepares a cavity for amalgam restoration. He puts an initial depth of the cavity of 0.2 to 0.8 mm into the dentin. What is the most likely step of preparation in this case?

A. Retention
B. Resistance
C. Convenience
D. Removal of remaining caries

ANSWER: B

43. A 20-year-old male patient came to the dental clinic with a complaint of moderate pain during cold weather, which subsided after removal of the initiator of pain. Clinical examination revealed vital pulp and class I deep caries on 16. Restoration is done using calcium hydroxide and temporary restorations.

What is the most likely next appointment for this patient in weeks?

A. Two to Four
B. Four to Six
C. Six to Eight
D. Eight to Ten

ANSWER: C

44. A 30-year-old female patient came to the dental clinic with a complaint of discoloration of the restoration on tooth 47, which was restored 10 years ago. She has no history of pain after the restoration was done. She did not like the color of the previous restoration. The restoration is intact, and the tooth is vital.

What was the most likely previous restoration in this case?

A. Amalgam
B. Composite
C. Cast metal
D. Gold filling

ANSWER: A

45. A 35-year-old male patient has come to a dental clinic complaining of tooth sensitivity. Clinically, there is chipped enamel on the incisal and occlusal area.

What is the most likely diagnosis in this case?

A. Erosion
B. Attrition
C. Abrasion
D. Dentin hypoplasia

ANSWER: B

46. A 45-year-old male patient has come to the dental OPD for a second appointment. His first chief complaint was tooth discoloration on the right upper tooth. Previously, there was a history of trauma 10 years before, which resulted in pulpal necrosis. In his previous appointment, root canal treatment was done.

What is the most appropriate bleaching technique in this case?

A. Home
B. In office
C. Laser-assisted
D. Closed chamber

ANSWER: A

47. A patient came to the dental clinic for treatment. He directly went to the record office, and the officer registered his data, such as name, address, age, sex, and occupation. After the registration, the patient is directed to a physician for evaluation and management.

What is the next immediate component of history to be asked in this patient care?

- A. Social History
- B. Family History
- C. Personal History
- D. Chief complaint History

ANSWER: D

48. A patient came to the dental department with a complaint of mobile swelling of the left lower jaw region for three months. He has no history of trauma. On examination, the swelling was a soft tissue mass originating from the submandibular salivary gland. Imaging was required to diagnose the case.

What is the most appropriate initial imaging modality in this patient care?

- A. Ultrasound
- B. CT scan
- C. OPG
- D. MRI

ANSWER: A

49. A 45-year-old male patient came to the dental clinic with complaints of a burning sensation and ulceration of the oral cavity for two weeks. The ulceration comes and disappears by itself on a regular basis. He had no history of smoking and physical as well as chemical burns of his mouth. He has no history of the same problem in his family. On physical examination, there are round, oval, painful ulcers with erythematous borders. Vitamin B12 and folic acid were prescribed. After some time, the patient came back with improvement.

What is the most likely diagnosis of the oral ulcer in this patient?

- A. Kaposi's sarcoma
- B. Aphthous ulcer
- C. Erythroplakia
- D. Leukoplakia

ANSWER: B

50. A 28-year-old pregnant woman came to the hospital complaining of severe dental pain localized to the left side of the upper jaw that had persisted for the last three days. There was slight swelling for which she had taken medication. She is in the second trimester of pregnancy and has regular follow-ups without any complications. On physical examination, all vital signs are within the normal range, and she had a grossly decayed root stump of the mandibular third molar, which is not impacted. After the investigation, it was decided to remove the offending tooth since it is highly infected.

What is the best position of the patient on the dental chair for the procedure?

- A. Ninety-degree sitting position
- B. Lateral decubitus position
- C. Proper standing position
- D. Routine Supine position

ANSWER: B

51. A 35-year-old male patient came to the hospital complaining of discomfort and pain in his mouth, which is aggravated by hot and cold drinks. On medical history, the patient reported ulceration of the genital organ and a painless ulcer of the tongue and lip a year ago. He visited a physician for his general health problem. Physical examination showed intraoral ulceration with a raised, firm border on the tongue and palate, and Saddle nose.

What is the most likely diagnosis?

- A. Primary syphilis
- B. Tertiary syphilis
- C. Secondary syphilis
- D. Congenital Syphilis

ANSWER: B

52. A 21-year-old female patient came to hospital complaining of oral ulceration of five days duration. She had a history of profuse purulent urethral discharge with frequent micturition, dysuria, fever, and headache. She had nasal stiffness, epistaxis, and hoarseness. On intraoral examination, there are multiple, painful and round, elevated gray white eroded spots, without pseudo-membrane, and gingivitis was observed.

What is the most likely diagnosis?

- A. Gonococcal stomatitis
- B. Secondary Syphilis
- C. Tertiary syphilis
- D. Herpes simplex

ANSWER: A

53. A patient came to the dental set up with a complaint of difficulty chewing and swallowing for ten days. He also complains of headache, abdominal pain, nausea, and vomiting. He had a sudden onset sore throat, and he could not chew any food. On examination, he had redness of the tonsil, oral mucosa, and pharynx, palatal petechiae, yellow tonsillar exudates, and cervical lymphadenopathy. The patient was diagnosed with Streptococcal Tonsillitis and Pharyngitis. But he is allergic to the penicillin group.

What will be the next possible treatment?

- A. Cephalosporin
- B. Erythromycin
- C. Gentamycin
- D. Cloxacillin

ANSWER: B

54. A patient came to the dental clinic with a complaint of sharp pain on the left side of the face for two months. The pain is a sharp, shooting type and aggravated by light touch. He had a history of oral vesicles that rapidly evolved into small superficial ulcers that coalesced to give rise to larger irregularly outlined ulcers, which were diagnosed as shingles three months ago.

What is the most likely diagnosis of facial pain in this case?

- A. Orofacial pain
- B. Postherpetic neuralgia
- C. Infectious mononucleosis
- D. Ramsay Hunt syndrome pain

ANSWER: B

55. A 42-year-old male patient came to the dental clinic with a complaint of pain and discomfort. He had a medical history of HIV infection for which he is on ART. On intraoral examination, the dentist visualized an erythematous plaque intermingled with a white patch on the buccal mucosa.

What is the most likely diagnosis?

- A. Pseudomembranous candidiasis
- B. Candida-associated lesions
- C. Erythematous candidiasis
- D. Hyperplastic candidiasis

ANSWER: A

56. A 55-year-old male patient came to dental clinic with a complaint of pain and discomfort due to an oral sore for the last two months. Examination revealed ulceration of the palate and gingival with mobility of teeth. OPG radiograph showed bone loss around the upper teeth. There is macular diffuse pigmentation of the maxillary gingiva. The patient had pigmentation in the malar region for a couple of months, which was diagnosed as cutaneous melanoma.

What is the most likely diagnosis?

- A. Leukoplakia
- B. Erythroplakia
- C. Haemangioma
- D. Mucosal melanoma

ANSWER: D

57. A patient came to the dental clinic with a complaint of severe pain in his mouth, which lasted for three days. The pain is initiated when cold and hot drinks are consumed. On history taking, the patient points to the left side of the upper jaw but cannot specify a tooth. On examination, the dentist observed a carious tooth on the same side as the pain.

What is the most likely diagnosis?

- A. Pulpal pain
- B. Neuralgic pain
- C. Periodontal pain
- D. Paranasal sinus-related pain

ANSWER: A

58. A patient came to the dental clinic with a complaint of a headache for two weeks. The pain is a dull aching type distributed bilaterally around the Peri-orbital region. The headache is also associated with nasal obstruction, nasal purulence, and postnasal drip, alteration in the sense of smell, cough, fever, fatigue, lethargy, halitosis, pharyngitis, and otalgia.

What is the most likely diagnosis?

- A. Medication overuse headache
- B. Headache associated with TMD
- C. Headache associated with sinusitis
- D. Headache associated with Ischemic stroke

ANSWER: C

59. A 60-year-old male patient came to dental clinic with a complaint of hoarseness of voice and difficulty swallowing for a long time. He has superficial blisters and chronic ulceration of the skin and mucosa. On examination, he also had shallow intraoral ulcers with margins of the ulcer showing evidence of tissue tags on buccal mucosa, palate, and gingiva.

The ulcers extended to the pharynx. What is the most likely diagnosis?

- A. Pemphigus vegetans
- B. Pemphigus vulgaris
- C. Pemphigus foliaceus
- D. Pemphigus erythematosus

ANSWER: B

60. A 26-year-old male patient came to dental clinic with a complaint of ulceration in his mouth for a long time. He had no other associated problem. On intraoral examination, he has a white line on the buccal mucosa bilaterally at the level of the occlusal plane of adjacent teeth. The line is more pronounced adjacent to the posterior teeth.

What is the most likely diagnosis?

- A. Linea Alba
- B. Thermal burn
- C. Atypical granuloma
- D. chronic cheek chewing

ANSWER: A

61. A six-year-old male patient was brought to a dental clinic with a complaint of toothache and swelling of two weeks duration. On clinical examination, there are grossly decayed upper anterior teeth and swelling with a sinus tract. The child is extremely uncooperative, fearful, and anxious. The dentist decided on the extraction of 51, 52, 61, and 62.

What is the best approach for the management of this patient?

- A. Sedation
- B. Local anesthesia
- C. Topical anesthesia
- D. General anesthesia

ANSWER: D

62. A nine-year-old girl came to dental clinic for a checkup. On examination, a dentist found peg-shaped both maxillary lateral incisors. OPG detected no other abnormality.

What is the most appropriate initial step in the management of this patient?

- A. Extraction and replacement
- B. Orthodontic treatment
- C. Root canal treatment
- D. Composite filling

ANSWER: D

63. A three-year-old male child was brought to the dental clinic with a complaint of tooth decay. Examinations showed caries-affected teeth on maxillary incisors, canines, and second primary molars.

What is the most likely diagnosis for this case?

- A. Pit and fissure caries
- B. Rampant caries
- C. Nursing care
- D. Arrested caries

ANSWER: C

64. A six-year-old girl came to the dental clinic with a complaint of a toothache on the right upper jaw for a week, which is aggravated on taking cold and hot foods. The dentist felt irregularity during probing the interproximal area of 54 and 55, negative on horizontal and vertical percussion.

What is the best next step in the diagnosis of this case?

- A. OPG x-ray
- B. IOPA x-ray
- C. Bitewing x-ray
- D. Occlusal x-ray

ANSWER: C

65. An 11-year-old girl visited a dentist for treatment of a toothache on the right lower jaw of four days duration. She complains of severe pain in biting. Clinical examination showed class II caries on 46, tender on percussion, no mobility, and swelling. IOPA x-ray shows interrupted lamina dura around the apex of the tooth.

What is the most likely diagnosis for this case?

- A. Acute apical periodontitis
- B. Acute alveolar abscess
- C. Condensing osteitis
- D. Reversible pulpitis

ANSWER: A

66. A three-year-old girl was brought to the pediatric dental clinic for a dental checkup. Examination showed crown discoloration on 51 with deep class III dental caries, positive on vertical percussion, negative on electric pulp test. IOPA x-ray showed no evidence of pathological conditions. She is very cooperative.

What is the most effective management of this case?

- A. Apexification
- B. Pulp Capping
- C. Pulpectomy
- D. Extraction

ANSWER: C

67. A five-year-old female patient came to the dental clinic with a complaint of a toothache on the left upper jaw of one week duration, which is aggravated by cold temperature and during food impaction. Dental examination showed a class I cavity on 65, negative on percussion. IOPA x-ray showed a cavity very close to the pulp horn. A dentist planned pulp therapy, and during cavity preparation, a very small pulp horn was exposed, for which bleeding stopped with simple cotton pellet pressure.

What is the best next step in the management of this case?

- A. Dressing with calcium hydroxide
- B. Extraction of the affected tooth
- C. Apply the form cresol
- D. Direct GIC filling

ANSWER: A

68. An 11-year-old female patient was presented to the dental emergency after three hours of fighting. On clinical examination, there is bleeding from the upper anterior and a shortened permanent right maxillary lateral incisor. Dental x-ray showed a displaced tooth 12 apically, discontinuous periodontal ligament, no crown, and root fracture.

What is the most likely diagnosis?

- A. Avulsion
- B. Intrusion
- C. Extrusion
- D. Concussion

ANSWER: B

69. An eight-year-old female patient was brought to a dental clinic five minutes after falling down an accident on the school playground. The examination showed complete displacement of the maxillary tooth on tooth 11. The tooth is in her vestibule, and the alveolar socket is not collapsed. She is conscious and can open her mouth.

What is the best initial step in the management of this case?

- A. Root canal treatment
- B. Socket debridement
- C. Antibiotic coverage
- D. Re-implantation

ANSWER: D

70. A two-and-a-half-year-old child was brought to a dental clinic after falling injury of one day duration. Dental examination showed intruded both maxillary central incisors were intruded. Lateral x-ray revealed apical displacement of central incisors toward the labial direction.

What is the most effective management of this case?

- A. Extraction
- B. Orthodontic traction
- C. Spontaneous reposition
- D. Repositioning and splinting

ANSWER: C

71. A three-year-old child came to the dental clinic for a checkup. He was diagnosed with Down syndrome and on follow-up.

What is the most likely dental and oral problem the child will be diagnosed with?

- A. Macrodonia
- B. Microglossia
- C. Delayed eruption
- D. Supernumerary teeth

ANSWER: C

72. A seven-year-old boy was brought to a dental OPD with a complaint of a tooth cavity on the left lower jaw. The child has autism spectrum disorder on psychiatric follow-up.

What is the most appropriate initial step in the management of this patient?

- A. Cavity preparation
- B. Fluoride application
- C. Behavior modification
- D. Anesthesia administration

ANSWER: C

73. A 32-year-old male patient came to a dental clinic with a complaint of spontaneous bleeding of the gingiva. Examination revealed gross calculus deposition, loss of stippling, and grade II mobility of lower anterior teeth. The doctor has planned to measure pocket depth.

What amount of probing force is well tolerated by the patient during the probing process?

- A. 0.25N
- B. 0.5N
- C. 0.75N
- D. 1N

ANSWER: C

74. A 47-year-old male patient has been diagnosed with necrotizing ulcerative gingivitis. Stepwise treatment is planned.

What is the goal of initial therapy in the treatment plan for this case?

- A. Restoration of bone and attachment integrity
- B. Restoration of the esthetic condition of the gingiva
- C. Reduction of microbial load and removal of necrotic tissue
- D. Correction of systemic conditions that contribute to the infection

ANSWER: C

75. A 17-year-old patient came to the dentist with a periodontal abscess for the third time. On examination, the dentist noticed severely inflamed gingival and deep periodontal pockets. Additionally, radiographic examination confirmed horizontal bone loss in the molar region. The dentist suspects systemic disease.

What is the most likely suspected systemic disease in this case?

- A. Leukemia
- B. Diabetes mellitus
- C. Hodgkin's lymphoma
- D. Acquired immune deficiency syndrome

ANSWER: B

76. A three-year-old girl was brought to the dental OPD with a complaint of a loose tooth. On examination, she has multiple loose teeth, with missing teeth 84, 85, 64, and 75. She also has hyperkeratotic skin on her sole and palm.

What is the most likely diagnosis for this patient?

- A. Dawn's syndrome
- B. Chediak-Higashi syndrome
- C. Papillon Lefevre syndrome
- D. Leukocyte adhesion deficiency

ANSWER: C

77. A 30-year-old male patient presented to the dental clinic with enlargement of the gingival for six months. He underwent a renal transplant eight months ago. On examination, non-erythematous and painless enlargement of the gingiva all over the labial surfaces of both arches was found.

What is the most likely cause of the enlargement?

- A. Nutritional deficiency
- B. Drug-induced
- C. Neoplastic
- D. Idiopathic

ANSWER: B

78. A 55-year-old male patient with a history of coronary artery disease who is taking nifedipine, presented to the dental OPD with a complaint of gum overgrowth. On clinical examination, he has moderate plaque and calculus on the lower anterior teeth with moderate crowding. He also has bead-like enlargement of the facial and lingual gingiva. In social history, the patient smoked one pack of cigarettes per day for five years.

What is the most likely condition that caused the gingival enlargement?

- A. Malocclusion
- B. Poor oral hygiene
- C. Cigarette smoking
- D. Calcium channel blocker

ANSWER: D

79. A 33-year-old patient came for regular follow-up after full and successful treatment of generalized chronic periodontitis. Radiographic evaluation showed a distinct breakdown of the alveolar bone of the lower arch.

What should be the next plan for this patient?

- A. Bone graft
- B. Gingivoplasty
- C. General health review
- D. Scaling and root planning

ANSWER: C

80. A 25-year-old male patient came to a dental clinic with the complaint of dull and localized pain in the lower right quadrant. On the intraoral examination, a 6mm periodontal pocket and positive vitality test were recorded. Radiographic examination showed periodontal angular bone loss and radiolucency.

What is the most likely diagnosis in this case?

- A. Pulpal Abscess
- B. Gingival Abscess
- C. Periodontal abscess
- D. Pericoronal Abscess

ANSWER: C

81. A 35-year-old male patient presented with a complaint of severe pain and swelling on the right lower back and jaw of six days' duration. On examination, there were heavily restored 44.

Which clinical feature of the tooth makes the diagnosis periodontal abscess?

- A. Loss of vitality
- B. Absence of bone loss
- C. Tenderness to vertical percussion
- D. Radiolucency on the lateral side

ANSWER: D

82. A 55-year-old patient came to the dental OPD with a complaint of intermittent exudation from the lower first premolar. He explained that the tooth had no pain, but the exudation bothered him. On examination, deep vertical pockets and deep furcation defects were found.

What is the most appropriate management in this case?

- A. Splinting
- B. Antibiotics
- C. Surgical treatment
- D. Scaling and root planning

ANSWER: C

83. A 40-year-old male patient presented to the dental OPD with a complaint of intense gum pain of one day duration, which aggravates on mastication. On examination, he has diffuse swelling of the gingiva, which is sensitive to palpation. He has an 8mm pocket measurement on tooth 44. Extra-orally, he has lymphadenopathy and elevated temperature.

What is the most appropriate first step for the management of this patient?

- A. Removal of tooth 44
- B. Systemic antibiotics
- C. Incision and drainage
- D. Drainage through the access cavity

ANSWER: B

84. A 43-year-old male patient presented to the dental OPD with a complaint of gum swelling. On examination, he has fibrous enlargement of the gingiva with shallow pocket depth.

What is the best treatment for this patient?

- A. Gingivectomy
- B. Gingivoplasty
- C. Gingival curettage
- D. Coronally advanced flap

ANSWER: A

85. A five-year-old girl presented to the dental OPD with a complaint of soreness of the mouth of two days duration, which interferes with eating. Her temperature was 38.1 degrees centigrade. On examination, she has diffused, edematous, erythematous, and enlargement of gingiva, which tends to bleed.

What is the most appropriate treatment option for this patient?

- A. Scaling and root planning
- B. Amoxicillin 125 mg/5ml PO, TID for seven days
- C. Acyclovir 15mg/kg five times a day for seven days
- D. Three percent Hydrogen peroxide PO, BID for seven days

ANSWER: C

86. A 34-year-old male patient is visiting his periodontist for a regular checkup. He was diagnosed with a primary periodontal secondary endodontic lesion. During this visit, he is going to have root canal treatment. What is the most appropriate phase of treatment the patient is currently on?

- A. Surgical phase
- B. Etiothropic phase
- C. Restorative phase
- D. Maintenance phase

ANSWER: B

87. A 45-year-old male patient came to a dental clinic with a complaint of swelling on the right posterior lower jaw and difficulty swallowing of five days duration. On intraoral examination, tooth 46 has a cavity and has had pain for a long time. Incision and drainage were done after admission. 48 hours after the procedure, the patient shows no apparent change.

What is the most appropriate follow-up modality for this case?

- A. ESR
- B. CBC
- C. CT scan
- D. Drainage output

ANSWER: C

88. A 19-year-old patient came to the hospital with a chief complaint of a fall accident of one hour duration. On examination, there was bleeding, mobility of teeth 11 & 21, and gum laceration.

What is the most likely imaging modality to rule out root fracture in this case?

- A. PA view of anterior maxilla
- B. Intraoral periapical view
- C. Panoramic view
- D. PNS view

ANSWER: B

89. A 53-year-old female patient came to a dental clinic for a planned tooth extraction of the right maxillary molar tooth due to chronic apical periodontitis. She had a history of myocardial infarction and was on anticoagulation therapy. After local anesthesia administration, the extraction was done.

What is the most important history to ask this patient?

- A. Duration of myocardial infarction
- B. Previous history of extraction
- C. Previous history of allergy
- D. Type of medication taken

ANSWER: A

90. A 28-year-old female patient came to a dental clinic with a severe toothache in the upper right jaw of four days duration. She has no history of chronic diseases. On examination, there was an un-restorable carious tooth, tooth 17, and percussion was positive. She was on holiday, fasting, and not willing to take food before the planned dental procedure.

What is the most likely expected complication in this patient if the procedure is performed?

- A. Orthostatic hypotension
- B. Loss of consciousness
- C. Hypoglycemia
- D. Syncope

ANSWER: D

91. A 72-year-old patient had a severe toothache overnight, for which the patient went to a dental clinic. Clinical examination revealed there was a severe caries on the mandibular last molar. Extraction of the tooth was agreed upon, but immediately after administration of lidocaine, the patient developed stridor with cardiac dysrhythmia and hypotension.

What is the initial step in this patient's management?

- A. Observe in the office for a few minutes
- B. Prescribe the oral antihistamine
- C. Place the patient in a sitting position
- D. Initiate basic life support

ANSWER: D

92. A 32-year-old patient came to the dental department after she developed a toothache in the right maxillary area. She was diagnosed with irreversible pulpitis of the first molar, and extraction of the tooth was planned under greater palatine nerve block. During anesthesia injection, the patient started vomiting and developed hypoxia.

What is the initial step in this patient's care?

- A. Place the patient on the right side in a horizontal position
- B. Administer theophylline 250 mg IV slowly
- C. Monitor vital signs for 30 minutes
- D. Observe in the office for one hour

ANSWER: A

93. A 32-year-old woman in the second trimester of pregnancy came to the maxillofacial surgery unit with complaints of pain and swelling of the left lower posterior jaw of two months duration. On examination, there was pus discharge and a necrotic wound on the left angle of the mandible, both intraoral and extraoral. The surgeon reached on diagnosis of chronic osteomyelitis. Surgical debridement and irrigation of necrotic tissues were planned under general anesthesia.

What is the solution for hand scrubbing in this patient care setting?

- A. Iodophors
- B. Chlorhexidine
- C. Hexachlorophene
- D. Sterile saline solution

ANSWER: B

94. A 40-year-old female patient presented to a dental clinic with a chief complaint of a toothache of one week duration. She had a history of intermittent coughing for the past month and bleeding of the gums for the past two years. On examination, vital signs show blood pressure of 120/80mmHg, heart rate of 76/min, respiratory rate of 16/min, and temperature of 36.9 degree centigrade recorded. Intraoral, there are grossly carious teeth with 2/3 crown removed and percussion positive, and good oral hygiene. The surgeon decided to extract the tooth under local anesthesia.

What is the most appropriate preoperative investigation in this patient's care?

- A. CBC
- B. RBS
- C. Chest X-ray
- D. Coagulation profile

ANSWER: D

95. A 20-year-old girl came to an emergency OPD after stone injury to her lower lip two hours ago. On examination, there was ecchymosis over the mucosa and in subcutaneous tissue on the lower lip, and the lower anterior teeth were tender on percussion.

What is the most effective management for this patient?

- A. Cleaning with surgical soap and antiseptic
- B. Application of ice or pressure dressing
- C. Suturing and systemic antibiotics
- D. Debridement and gentle dressing

ANSWER: B

96. An 8-year-old child was brought to a dental clinic after sustaining a dog bite injury on her face of two hours. On clinical examination, there was a crushed type of wound on the cheek with abrasion of the upper lip. After thorough irrigation of the wound with normal saline and tissue debridement, the patient was sent home with oral medication.

What is the most likely first-line drug in this case?

- A. Metronidazole
- B. Doxycycline
- C. Amoxicillin
- D. Augmentin

ANSWER: D

97. A 32-year-old male patient came to the hospital with bleeding from the face for three hours secondary to a knife injury. On examination, there is laceration, bleeding, edema, and erythema of the wound over the left buccal region. No debris and signs of infection around the wound.

What is the most appropriate wound healing for this patient?

- A. Tertiary
- B. Primary
- C. Secondary
- D. Delayed primary

ANSWER: B

98. A 34-year-old female patient came to the maxillofacial OPD with painful swelling on the upper anterior jaw of three days duration. On examination, there was discoloration of 11 & 21 teeth. There was swelling over the muco-buccal vestibule of 11, 12, 21, & 22 teeth. Intraoral periapical radiograph showed a periapical radiolucency with a diagnosis of radicular cyst. A trapezoidal flap was planned with a vertical releasing incision mesial to 12 and 22 for enucleation.

What is the most likely complication of this procedure?

- A. Flap dehiscence
- B. Flap necrosis
- C. Flap tearing
- D. Hematoma

ANSWER: A

99. A 50-year-old male patient has come to a dental clinic with complaints of pain and swelling during and after eating food for a long time.

What is the most likely diagnosis in this case?

- A. Sialolithiasis
- B. Sialadinitis
- C. Sialorrhea
- D. Sialosis

ANSWER: A

100. A 45-year-old male patient came to the dental department with a complaint of a firm mass originating from the mandible. The mass measures 6cm in diameter and is in proximity to the inferior alveolar nerve. After clinical assessment, the practitioner decided to do a biopsy.

What is the most likely type of biopsy in this case?

- A. Excisional
- B. Incisional
- C. Aspiration
- D. Cytology

ANSWER: B

101. A 25-year-old male patient came to a dental clinic with a complaint of pain in the lower second molar of one month duration. On radiographic examination, the dentist observed that the last molar on the same side is in the bone with the crown surrounded by well-defined unilocular radiolucency with a sclerotic border. The practitioner diagnosed the lesion as a cyst.

What is the most likely type of cyst in this?

- A. Dentigerous
- B. Periapical
- C. Eruption
- D. Gingival

ANSWER: A

102. A 16-year-old male patient came to a dental clinic with a history of a missing maxillary right canine. The patient denied a history of trauma and extraction of teeth. Radiographic evaluation revealed the absence of the tooth but well-localized radiolucency at the site. The adjacent tooth is vital and slightly displaced.

What is the most likely cyst diagnosed in this case?

- A. Residual
- B. Eruption
- C. Gingival
- D. Primordial

ANSWER: A

103. A 25-year-old female patient came to a dental clinic with a toothache of two days' duration. On examination, there is a grossly decayed tooth; palpation and percussion were positive on tooth 16. In history, there is no known medical illness. IOPA of tooth 15 shows widening of the lamina dura around the apical areas of roots. Extraction is planned under local anesthesia.

What is the baseline laboratory investigation for this patient?

- A. RBS
- B. RFT
- C. CBC
- D. LFT

ANSWER: C

104. A 23-year-old patient came to the dentist with tooth pain in the upper anterior jaw. On examination, there was a caries and root remnant of teeth 11 and 21 seen at the level of the free gingival margin that was left following tooth removal.

Which instrument is most likely important in this procedure?

- A. Mosquito forceps
- B. Straight elevator
- C. Crossbar
- D. Cryer

ANSWER: A

105. A 23-year-old patient came to a dental clinic with a complaint of a toothache on the lower anterior jaw of three days duration. On examination, there was caries and severe mobility of tooth 33. Written informed consent was obtained for extraction under local anesthesia.

What is the most appropriate position of the patient in this scenario?

- A. The mandibular occlusal plane is parallel to the ground
- B. The mandibular occlusal plane is at 45-60 degree
- C. The mandibular occlusal plane is at 90 degree
- D. Mandibular occlusal plane 30 degrees

ANSWER: A

106. A 33-year-old patient came to the hospital with tooth pain in the upper anterior jaw of one month duration. On physical examination, tooth 21 was discolored with pulp necrosis. The patient agreed to undergo extraction.

What is the appropriate final direction force to deliver this tooth?

- A. Apical
- B. Palatal
- C. Labial
- D. Rotational

ANSWER: C

107. A 26-year-old patient came to the dentist with severe tooth pain in the lower posterior jaw of three days duration. The dentist examined and planned for the extraction of tooth 44. During the procedure, there was difficult luxation and minimal mobility of the tooth.

What is the appropriate next step in this patient's management?

- A. Intraoral periapical radiograph
- B. Mucoperiosteal flap raise
- C. Luxation by hammering
- D. Tooth sectioning

ANSWER: D

108. An 18-year-old male patient came to the dental department with a history of difficulty opening his mouth for several years. His medical history is unremarkable. Imaging revealed that reduced temporomandibular joint space and hazy appearance, but normal anatomy of the head of the condyle and glenoid fossa was appreciated.

What is the most likely diagnosis for this case?

- A. Bony ankylosis
- B. Joint dislocation
- C. Fibrous ankylosis
- D. Condylar hypoplasia

ANSWER: C

109. A 15-year-old male patient came to dental clinic after sustaining a sports injury to his dentition, with a duration of seven hours. He had an unremarkable injury to his other body parts. On clinical evaluation of the dentition, the central maxillary incisor has a fracture of the crown confined to enamel and dentin with no root exposure.

What is the classification of this fracture according to the WHO system for disease classification?

- A. Crown infraction
- B. Uncomplicated
- C. Complicated
- D. Craze line

ANSWER: B

110. A 15-year-old male patient came to dental clinic after sustaining of fall down accident and injury to his orofacial region six hours earlier. On clinical evaluation, he had mild soft tissue injury, and on examination of dentition, he showed marked reaction to percussion of maxillary central incisors in both the horizontal and the vertical directions. Otherwise, no abnormal mobility, displacement, or bleeding is observed.

What is the most likely diagnosis for this injury?

- A. Displacement
- B. Concussion
- C. Avulsion
- D. Luxation

ANSWER: B

111. A 50-year-old male patient came to the emergency department of a hospital after sustaining a road traffic accident of 30 minutes duration. A dentist was called to evaluate the dento-alveolar and facial injury for the patient.

What should be considered before any management, including dental evaluation?

- A. Communicate with the attending physician and verify the stability of the patient
- B. Defer any management regardless of stability until he leaves the emergency
- C. Evaluate the patient for any maxillofacial injury immediately
- D. Consult an oral and maxillofacial surgeon immediately

ANSWER: A

112. A 14-year-old patient came to dental department after sustaining a fighting accident of three hours duration. He sustained injury to his mid-face and had a down gaze. The patient could not rotate the left eye inferiorly, whereas the right eye was fully rotated inferiorly.

What is the most likely fracture involved in this case?

- A. Lefort one
- B. Lefort three
- C. Orbital roof
- D. Orbital floor

ANSWER: D

113. A 30-year-old male patient came to the dental clinic after sustaining a bicycle accident of six hours duration. On clinical examination, he had disturbance of occlusion, step-off on the mandible, and lingual hematoma. The practitioner diagnosed the problem as an unfavorable mandibular fracture.

What is the exact nature of the fracture in this case?

- A. Displacement at the fracture site caused by the pull of the muscle
- B. Difficult to reduce by manual manipulation
- C. Invisible fracture line on radiographs
- D. Impossible to manage by any means

ANSWER: A

114. A 20-year-old patient came to a dental clinic with pain in the lower left posterior jaw. On examination, there was deep caries of tooth 38. The pain is exacerbated by cold air and hot drinks. Extraction was planned under local anesthesia by nerve block.

What is an important extra oral landmark in this patient management?

- A. Pterygomandibular raphe
- B. Contralateral cusps of premolars
- C. Contralateral angle of the mouth
- D. Anterior edge of ascending ramus

ANSWER: C

115. A 19-year-old male patient came to the dental department with a complaint of a missing tooth on the left maxillary area. After clinical assessment, a radiograph was taken, and there was an impacted canine on the palatal side for which the practitioner planned disimpaction of the tooth.

What is the most effective technique for profound anesthesia in this case?

- A. Nasopalatine Nerve Block
- B. Greater Palatine Nerve Block
- C. Anterior superior nerve block
- D. Posterior superior nerve block

ANSWER: B

116. A 43-year-old female patient came to the dental department after she had developed a toothache in the right mandibular area. She was sent for X-ray and diagnosed with un-restorable caries, and extraction of the tooth was planned under adequate anesthesia. The anesthesia is effective in depth and duration.

What is the most likely explanation in this case?

- A. The effect of sodium meta-bisulfate
- B. The effect of sodium chloride
- C. The effect of Vasopressor
- D. The effect of sterile water

ANSWER: C

117. A 19-year-old male patient came to a dental clinic with a toothache in the left mandibular area with irreversible pulpitis of the last molar; extraction was planned under local anesthesia. The patient's weight was 50kg, and the dentist administered two percent lidocaine without a vasoconstrictor. The maximum recommended dosage was 4.4 mg/kg for profound anesthesia.

What is the maximum number of cartridges to be used in this case?

- A. 3
- B. 4
- C. 5
- D. 6

ANSWER: C

118. A 44-year-old male patient came to the hospital for the scheduled extraction of the mandibular last molar tooth. During local anesthesia administration, the needle broke and couldn't be readily retrieved.

What is the most appropriate management in this case?

- A. Remain calm and do not panic
- B. Instruct the patient not to move
- C. If the fragment is visible, try to remove it
- D. Do not proceed with an incision or probing

ANSWER: D

119. A 55-year-old male patient came to a dental clinic for the extraction of the maxillary last molar tooth. After local anesthesia administration blocked the posterior superior alveolar nerve, the extraction was done. Two hours after the extraction, the patient developed significant swelling. What is the most appropriate time to start heat therapy in this scenario?
- A. After 6 hours
 - B. After 9 hours
 - C. After 12 hours
 - D. After 24 hours
- ANSWER: D
120. A 10-year-old male child patient was referred from the pediatric hospital to the dental department for the extraction of the mandibular primary canine on the left side. Due to sudden movement on injection of local anesthesia, the lower lip was injured. What is the most appropriate management for this patient?
- A. Irrigate with hydrogen peroxide
 - B. Irrigate with normal saline
 - C. Apply topical anesthetics
 - D. Apply petroleum jelly
- ANSWER: D
121. A 48-year-old female patient came to a dental clinic with a chief complaint of a toothache. On clinical examination, there was a discolored upper central incisor tooth. Root canal treatment was done. The reamers used during the procedure are returned and put in the sterilizer for the next patient. What is the most likely applied sterilization technique in this case?
- A. Ethylene Oxide Gas
 - B. Ultraviolet Rays
 - C. Hot Air Oven
 - D. Infrared
- ANSWER: A
122. A 52-year-old male patient came to the dental department after he developed a toothache in the left maxillary area. History revealed that prosthetic valve replacement is recommended on warfarin treatment. On intraoral examination, there is a root stump of 37. Laboratory investigation, including INR, was sent. What is the most likely acceptable value of INR in this patient care?
- A. 1.5-2.5
 - B. 2.5-3.5
 - C. 3.5-4.5
 - D. 4.5-5.5
- ANSWER: B

123. A 57-year-old male patient came to the dental department after he developed a toothache in the left mandibular area. Extraction was done under local anesthesia for an un-restorable tooth on the second molar. Due to a lack of profound numbness, multiple local anesthesia injection was done, and he developed edema after the extraction.

What is the most effective management in this case?

- A. Post op corticosteroids
- B. Incision and drainage of the fluid
- C. Ice application to a freshly wounded area
- D. Head elevation above the rest of the body

ANSWER: D

124. A 61-year-old female patient went to a dental department for swelling on the right mandibular posterior area due to chronic osteomyelitis. She is a known TB patient and has been on anti-TB for the last five months. Debridement of the area is done safely after proper skin preparation and draping.

What is the chemical of choice in this patient care?

- A. Hexachlorophene
- B. Povidone-iodine
- C. Glutaraldehyde
- D. Chlorhexidine

ANSWER: B

125. A patient came to a dental clinic with a complaint of pain of three days duration. His pain was exacerbated while taking cold water. Upon clinical evaluation, the dentist does not find any decay on the surfaces of the teeth under complaint. And he requested an intra-oral x-ray to rule out interproximal caries.

What is the most appropriate type of X-ray requested in this case?

- A. Occlusal
- B. Bitewing
- C. Periapical
- D. Panoramic

ANSWER: B

126. A dentist was doing the extraction of tooth 16 and removed only the mesio-buccal root. The other roots were pushed to the antrum while trying to remove them with the elevator. The dentist wanted to determine the positions of the roots that were displaced into the antrum during his procedure with occlusal radiography. What type of X-ray is most likely requested in this case?

- A. Upper standard
- B. Upper oblique
- C. Upper anterior
- D. Upper Vertex

ANSWER: B

127. A 20-year-old male patient presented to the dental OPD with a complaint of an avulsed tooth due to a falling down accident of 30 minutes ago. The dentist performed the necessary evaluations and re-implanted the avulsed tooth into the socket and put the patient on a recall visit.

What is the most likely X-ray to be used during the reevaluation?

- A. OPG
- B. IOPA
- C. Occlusal
- D. Bitewing

ANSWER: B

128. A patient came to the dental department with a complaint of bleeding after sustaining a falling accident. Upon clinical evaluation, the dentist found out: bleeding of the gum around 21, and a class-II crown fracture about 21. And he planned to take a periapical x-ray of tooth number 21 by the bisecting angle technique to rule out root fracture of the tooth.

What is the vertical angulation of the X-ray tube in degrees in this case?

- A. +25
- B. +30
- C. +45
- D. +50

ANSWER: D

129. A 17-year-old patient came to a dental clinic with a complaint of pain on his upper right-side teeth. The dentist in charge examined and found out class-I caries on the maxillary right second premolar and first molar teeth. He decided to take a periapical x-ray by the paralleling technique.

What is the most likely position of the image receptor in this case?

- A. Sufficiently posteriorly
- B. Sufficiently anteriorly
- C. Midline of the palate
- D. Lingual sulcus

ANSWER: C

130. A dentist evaluated a 17-year-old patient who came to a dental clinic. Upon examination, the dentist found that missing of all permanent canines. Then he decided to take a panoramic X-ray.

What is the most important patient position in this case?

- (A) Sit comfortably in the chair.
- (B) Upper and lower teeth to be in normal overjet
- (C) The mid-sagittal plane is perpendicular to the floor
- (D) An imaginary Frankfurt plane perpendicular to the floor

ANSWER: C

131. A dentist took a periapical x-ray of the mandibular first molar to rule out apical lesions. He got into the X-ray processing room to complete the image formation with a manual technique.

What is the most important sequence of steps to be followed in this case?

- A. Immerse the film in the fixer, rinse in water, developer, wash with running water, dry, and mount.
- B. Immerse the film in the developer, washing with running water, fixing, rinsing with water, drying, and mounting.
- C. Immerse the film in the fixer, rinse in water, developer, wash with running water, mount, and dry.
- D. Immerse the film in the developer, rinse in water, fix it, wash with running water, dry, and mount.

ANSWER: D

132. A dentist took a bitewing radiography of a patient on the right-side premolars of the jaws. He started processing a film manually. In the dark room, he immersed the film in fixing solution before inserting it into the developer.

What is the most likely feature of the image in the film in this case?

- A. Dark
- B. Light
- C. Blank
- D. Cracked

ANSWER: C

133. A 23-year-old male patient presented to the dental OPD with a complaint of pain in his lower right back jaw of two days duration. On examination, he has a class II cavity on tooth 45, which is tender on percussion. The dentist took IOPA, and after processing the film, the image became too dark.

What is the most appropriate explanation for this?

- A. Diluted developer solution
- B. Too hot a developer solution
- C. Exhausted developer solution
- D. Contaminated developer solution

ANSWER: B

134. A dentist read a periapical x-ray request of the maxillary right central incisor from paper. And, he told the patient to wear a lead apron and thyroid collar for radiation protection. The lead barrier is not available in his clinic.

What is the most likely standing distance of the dentist in feet to be in the zone of maximum radiation safety in this case?

- A. 3
- B. 4
- C. 5
- D. 6

ANSWER: D

135. A dentist working in a dental clinic wants to take multiple X-rays for his patients.

What is the most likely safety measure to be utilized in this case?

- A. He should stand behind a lead barrier
- B. He should stand directly behind the patient
- C. There is minimal radiation risk with dental X-rays
- D. He should stand 3 feet away from the primary beam

ANSWER: A

136. A 40-year-old male patient came to a dental clinic with a complaint of pain in his upper right first molar tooth. In history, he had polydipsia, polyphagia, polyuria, and unexplained weight loss. On physical examination, there was a grossly decayed tooth that mandates extraction.

What is the most likely preoperative investigation in this patient's care?

- A. Thyroid function test
- B. Renal function test
- C. Fructosamine test
- D. Blood sugar test

ANSWER: D

137. A 40-year-old male patient came to a dental OPD with electric shock-like pain. Upon evaluation, the pain was continuous, low intensity, and with no trigger zone.

What is the most likely diagnosis in this patient?

- A. Multiple sclerosis
- B. Alzheimer disease
- C. Parkinson disease
- D. Seizure disorder

ANSWER: A

138. A 37-year-old male patient came to a dental clinic with gum bleeding during brushing. Upon measuring the blood pressure, 145mmHg systolic and 95mmHg diastolic were recorded.

What is the most likely diagnosis in this scenario?

- A. Normal
- B. Prehypertension
- C. Stage I hypertension
- D. Stage II hypertension

ANSWER: C

139. A 45-year-old patient with coronary artery disease came to a dental office with a complaint of swelling on the lower left posterior teeth. In history, she had a recent myocardial infarction. After proper evaluation, there was a grossly decayed mandibular second molar that necessitated extraction.

What is the most likely appropriate time to do the procedure for this case?

- A. Three weeks later
- B. Four weeks later
- C. Five weeks later
- D. Six weeks later

ANSWER: B

140. A 30-year-old female patient came to the dental department for a checkup. In history, she had chronic fatigue and a burning mouth for the past six months. On clinical evaluation, there was angular cheilitis, glossitis, stomatitis, pale oral mucosa, gross calculus deposition on the lingual surface of the lower incisors, and grossly decayed tooth number 38.

What is the most likely diagnosis in this case?

- A. Iron deficiency anemia
- B. Megaloblastic anemia
- C. Sickle cell anemia
- D. Aplastic anemia

ANSWER: A

141. A 35-year-old male patient came to a dental OPD with a complaint of pain in his upper right posterior tooth. The patient was a mouth breather and could not localize the painful tooth. Upon clinical and radiographic evaluation, he had premolar and molar teeth sounded, but the second premolar, first molar, and second molar teeth were painful to vertical percussion.

What is the most likely diagnosis of the case in this scenario?

- A. Chronic apical periodontitis
- B. Acute apical periodontitis
- C. Acute periodontal abscess
- D. Acute maxillary sinusitis

ANSWER: D

142. A 20-year-old male patient came to the dental department for a checkup. In history, he had acute tonsillopharyngitis. The dentist did all the appropriate evaluations and recommended scaling and oral hygiene instruction.

What is the most likely instruction the dentist would recommend in this scenario?

- A. The proper cleaning of toothbrushes
- B. The different toothpastes to be used
- C. The different types of toothbrushes
- D. The methods of brushing teeth

ANSWER: A

143. A 25-year-old female patient came to the dental OPD with a complaint of bleeding after sustaining falling down accident before one hour. Upon evaluation, she had 35 percent blood loss by volume, and she was in shock. The dental surgeon requested investigations and close monitoring of the patient for possible resuscitation.

What is the most likely important parameter for this case?

- A. Respiratory rate
- B. Blood pressure
- C. Pulse pressure
- D. Urine output

ANSWER: D

144. A 20-year-old patient came to a dental OPD with a complaint of bleeding after sustaining a road traffic accident 30 minutes ago. He had a history of bleeding and was in shock.

What is the most likely fluid of choice in this patient management?

- A. Resuscitation with 0.9 percent normal saline
- B. Resuscitation 0.9 percent plasma protein
- C. Resuscitation 0.9 percent blood
- D. Resuscitation 0.9 percent sugar

ANSWER: A

145. A primary hospital had to improve its health care service for patients because of an increase in patients visiting the hospital. The medical director mapped out how to achieve this goal and decided on the necessary steps to accomplish the goal.

What is the most appropriate management function that the medical directors perform in this case?

- A. Staffing
- B. Leading
- C. Planning
- D. Organizing

ANSWER: C

146. A clinical director of a hospital planned to create a new group of experts to reduce repetitive complaints from her employees. She gathered 20 experts from 20 different departments, expecting to solve the problems and to break functional barriers among departments.

What is the most likely type of organizational structure created in this scenario?

- A. Functional
- B. Divisional
- C. Matrix
- D. Team

ANSWER: D

147. In a hospital, a group of experts was assigned to measure and assess the performance of the employees to improve health care delivery for the community. After careful study, the experts have explained that the hospital needs regular monitoring.

What is the use of monitoring for the hospital in this scenario?

- A. To assess the achievement of results
- B. To have selective exercise for research
- C. To provide continuous intervention with early indications
- D. To provide the provision of essential information at periodic intervals.

ANSWER: C

148. A hospital chief executive director is facing resistance from his employees. He couldn't get the best performance from his employees, because he always makes decisions by himself without listening to his employees' ideas and likes to dominate them.

What is the most likely type of Leadership style exhibited by the director in this case?

- A. Participative
- B. Autocratic
- C. Free Rein
- D. General

ANSWER: B

149. A dental doctor joined a hospital with a lot of problems as the medical director. He has been given one year as a goal to eradicate all the problems that the hospital is facing. He must make a lot of decisions to achieve the goal.

What is the most likely correct order to be followed to make effective decisions in this scenario?

- A. Generate options, define the objective, collect information, implement, evaluate, and decide.
- B. Define objective, generate options, decide, collect information, implement, and evaluate
- C. Define objective, collect information, generate options, decide, implement, and evaluate
- D. Collect information, define an objective, implement, evaluate, generate feasible options, and decide.

ANSWER: C

150. A newly opened dental clinic needs a good leader to increase the productivity of the clinic. To achieve this, the clinic hired a dental doctor who can be a leader, not a manager.

What features best describe a leader who increases productivity?

- A. Doing things right
- B. Doing the right things
- C. Focusing on system and structure
- D. Relying on the control of the employee

ANSWER: B

151. The medical director of a primary hospital called the operation room nurse and the surgical unit head to settle a conflict between them about the timing of elective operations on weekends. After having a thorough discussion, they agreed to work harmoniously by minimizing their absolute personal interests.

What conflict resolution technique did they use in this case?

- A. Win-win
- B. Lose-win
- C. Lose-Lose
- D. Cooperativeness

ANSWER: A

152. A patient living in a rural area of Ethiopia visits the nearby primary hospital for dental trauma management. The medical doctor conducted all the necessary examinations and referred the patient to a secondary health care level.

What is the most likely health facility that the patient is referred to in the above case?

- A. Regional hospital
- B. General hospital
- C. Referral hospital
- D. Zonal hospital

ANSWER: B

153. The malaria epidemic emerged in one region in Ethiopia. The regional health bureau organized a team to control the epidemic. And the team is expected to sustain till the control of the epidemic.

What is the most common type of team created in this scenario?

- A. Command team
- B. Reference team
- C. Task force
- D. Committee

ANSWER: C

154. A dental doctor assigned as a manager of a dental clinic noticed a shortage and improper use of materials, which needed purchasing, sorting, and appropriate storage. In the meantime, he tries to classify materials as consumable and non-consumable.

Which of the following materials can be categorized as non-consumable during sorting?

- A. Surgical gloves
- B. Microscope
- C. Syringes
- D. Cotton

ANSWER: B

155. A cross-sectional study was conducted on the prevalence of dental caries among school children aged 14-16 located in the south-west region. The research findings showed, Zone A and Zone D had the highest dental caries prevalence rates of 54.7 percent & 62.1 percent, respectively. In addition, there is an indication that the area is known for suboptimal fluoride content.

What is the most appropriate fluoride application for this scenario?

- A. Fluoride varnishes
- B. Water fluoridation
- C. Topical fluoride application
- D. Dissemination of fluoride-containing toothpaste

ANSWER: B

156. A regional health office is reviewing a project proposal submitted by a local NGO for ethical clearance. The NGO has been working with the community on personal hygiene and sanitation. The current focus of the NGO is to provide drinking water by fluoridation of the main supply used across the region.

What is the most appropriate ethics of water fluoridation to be considered for this case?

- A. The balance of risks and benefits
- B. Benefit is our primary concern over the risk
- C. The role of consent where there is potential harm
- D. The government should have full control over decision-making procedures

ANSWER: A

157. A 25-year-old female patient came to the dental OPD for a follow-up. During her first visit, she had preventive dental care and ultrasonic scaling. However, during the intraoral examination, there was a slight roughness that caught the explorer on the lower molars. And the doctor planned to apply pit and fissure sealants on the lower molar teeth.

Which is the most effective pit and fissure sealant material for this case in terms of prolonged retention?

- A. Resin-based material
- B. Low-viscosity glass-ionomer cements
- C. Polyalkenoic-Based Sealant Material
- D. Bisphenol A-glycidylmethacrylate-based material

ANSWER: A

158. A 22-year-old patient came to the dental OPD with a chief complaint of pain on the right side of the face of two weeks duration. The pain is severe and throbbing, which is associated with a headache. The patient disclosed that he chews gum daily and uses soft drinks with it. On intraoral examinations, there is foul-smelling mouth odor, generalized plaque, and calculus. There is also a class II cavity on teeth numbers 25, 26, 36, and 37.

What is the most likely health determinant to be considered for this patient?

- A. Innate
- B. Lifestyle
- C. Socio-economic
- D. Life circumstances

ANSWER: B

159. A 34-year-old patient came to the dental OPD complaining of pain in his lower front teeth, which is mild, intermittent in nature, initiated on hot and cold exposure, and that is generalized to the lower front tooth area. Upon intraoral examination, the patient has tooth erosion on teeth 31, 32, 41, and 42.

What is the most likely cause of this problem?

- A. Tobacco chewing
- B. Abnormal tooth brushing
- C. Lack of fluoridated water
- D. Excessive use of citrus fruits

ANSWER: D

160. A 65-year-old male patient presented to the dental OPD with a foul-smelling mouth odor for one year. The patient has reported tobacco use ranging up to one pack a day. Upon intraoral examination, there is brownish discoloration in the oral cavity. The dentist realized that it is the smoking habit that caused the increment of keratinization of the filiform papillae and discoloration due to the presence of chromogenic bacteria.

What is the most likely diagnosis for this case?

- A. Oral cancer
- B. Hairy tongue
- C. Tooth staining
- D. Kaposi Sarcoma

ANSWER: B

161. A 20-year-old female patient came to the dental OPD for her follow-up after she had a fixed orthodontic treatment one month ago. She told the doctor that she has difficulty brushing her teeth and is experiencing gum bleeding as she is still getting used to the orthodontic appliances.

What is the most appropriate preventive measure for this case?

- A. Floss
- B. Scaling
- C. Mouth Rinse
- D. Electrical brush

ANSWER: A

162. A dentist working at a regional hospital was asked by the administrator to teach the public about oral hygiene instruction at the triage area of the hospital. He delivered a ten-minute demonstration of the materials used for proper toothbrushing. He stressed that the traditional sticks widely used among the community aren't effective tools to keep oral hygiene and insisted on the use of toothpaste and a toothbrush.

What is the most likely oral health education principle violated by the dentist?

- A. Lack of consideration for limiting factors
- B. Imposing the dentists' ideas on the community
- C. Expecting knowledge to necessarily lead to behavior change
- D. Teaching oral hygiene practice in a value-free and unbiased manner

ANSWER: D

163. A dentist was fired from his job after refusing to accept the manager's new orders. The manager of the private clinic has prepared a bonus for the dentist who orders more investigations and pushes for the costly treatment option. He employed this tactic in order to increase the internal revenue of the clinic and cope with the recent rent increment.

What is the most likely code of ethics employed by the dental surgeon?

- A. Justice
- B. Integrity
- C. Autonomy
- D. Non-maleficence

ANSWER: B

164. A 65-year-old female patient presented to the dental OPD with swelling on the right side of her face of three months duration. After clinical and radiographic examination, the swelling is located on the right mandibular jaw and has irregular boundaries with palpable and tender submandibular lymph nodes. Among the differential diagnoses, the dentist highly suspected the swelling to be a malignant lesion. Therefore, the dentist explained the urgency of seeking further pathologic examinations and appropriate treatment.

What is the most likely code of ethics used by the dentist?

- A. Compassion
- B. Beneficence
- C. Truthfulness
- D. Professionalism

ANSWER: C

165. A 55-year-old male patient has been admitted to the oral and maxillofacial surgery ward with an angular mandibular fracture and has severe, excruciating pain. The dental intern on duty wanted to give the patients analgesics. However, the dental surgeon who saw him at the dental OPD did not write any order to alleviate the pain.

What is the most likely code of ethics breached by the dental surgeon?

- A. Justice
- B. Autonomy
- C. Beneficence
- D. Non-maleficence

ANSWER: C

166. A dental surgeon has appointed a patient for root canal treatment. After the dental assistant prepared the local anaesthetic, dental materials, and instruments, the dental surgeon asked him to swap with the female dental assistant before the patient entered the treatment room and unwrapped her head scarf.

What is the most likely behaviour applied by the dental surgeon?

- A. Care
- B. Respect
- C. Tolerance
- D. Compassion

ANSWER: B

167. A regional hospital has received a cephalometric device as a donation to be used in the dental OPD. The dentist working there has requested a training session before the device starts service to properly operate the device and improve patient care.

What is the most appropriate professional principle used by the dentist?

- A. Fidelity
- B. Veracity
- C. Discernment
- D. Self-reflection

ANSWER: D

168. A dentist encountered an elderly man while entering his private practice. The man explained that he had had severe pain in the left elbow for the past few days but couldn't afford the treatment cost. Therefore, the dentist treated the patient free of charge and appointed him for a week for a follow-up visit.

What is the most likely professional obligation kept by the doctor?

- A. Justice
- B. Respect
- C. Altruism
- D. Compassion

ANSWER: C

169. A dental surgeon went to a regulatory office to renew his professional license after working for the last three years at his private dental clinic. After reviewing his documents, the regulatory office denied his request until he brings a participation certificate of professional training in that time frame.

What is the most likely explanation of the regulatory office?

- A. Certify the credential
- B. Assess the quality of care
- C. Audit patient treatment outcomes
- D. Evaluate competency maintenance

ANSWER: D

170. A dental surgeon enrolled in short-term training on electronic medical record keeping. The training focused on the appropriate recording of patients' demographic data, integrating procedures carried out in the treatment rooms and imaging.

What is the most likely importance of this training?

- A. Enhance clinical skills
- B. Improve patient treatment
- C. Advanced supportive services
- D. Facilitate financial management

ANSWER: C

171. A 35-year-old female patient has come to the dental OPD for a follow-up visit. Her medical record shows that access was done on tooth number 21, and she was appointed to clean and shape for this visit. The dentist wants to take an intraoral periapical x-ray to determine the working length. Then, asks the patient's consent to perform the radiography when she reminded him that she is in her first trimester and has disclosed it on her initial visit.

What is the most appropriate explanation for this case?

- A. Lack of communication among the dentists
- B. Negligence in taking patient history on each visit
- C. Improper documentation of the patient record
- D. Exclusion of the previous procedures during reporting

ANSWER: C

172. A team leader of a private dental clinic has planned an oral health screening and treatment program for elders living in a nursing home. And conducted a meeting for volunteering dentists on the legal implications of proper documentation during care delivery.

What is the most appropriate document to consider in this case?

- A. Past medical history
- B. Medication prescription
- C. Informed consent forms
- D. Specific details of treatment provided

ANSWER: C

173. A quality assurance officer of a regional hospital planned to assess the delivery service in the dental OPD for a two-week duration. The study focuses on the usage of resources, the adherence to clinical guidelines, patient clinicians' communication, and inter-professional collaboration among the OPD team. What is the most appropriate data collection technique for studying?
- A. Interview
 - B. Observation
 - C. Questionnaire
 - D. Focus group discussion
- ANSWER: B
174. A year-five dental student planned to conduct his senior research project on the knowledge, attitude, and practice of oral hygiene among pregnant women coming for prenatal appointments at a regional hospital. According to the hospital records, the average number of women per month who come for prenatal appointments is 432. What is the most appropriate data collection technique for studying?
- A. Interview
 - B. Observation
 - C. Questionnaire
 - D. Focus group discussion
- ANSWER: C
175. A doctor is studying traditional initial sharpening practice among adolescents in District A. His aims are to describe the experience of individuals who went through this practice and understand the group norm. What is the most appropriate sampling technique for this study?
- A. Purposive
 - B. Stratified
 - C. Snowball
 - D. Cluster
- ANSWER: A
176. A doctor submitted a research proposal to the head of the dental department for approval. The study proposal is on the prescription rate of antibiotic drugs by dental surgeons for patients seeking dental treatment. The proposal narrates about the available knowledge from various aspects and perspectives; it develops a rationale for the study with reference to how the study will fill the identified gaps. What is the most appropriate explanation for the stated part of the proposal?
- A. Objective
 - B. Literature review
 - C. Research problem
 - D. Conceptual framework
- ANSWER: C
177. A group of researchers planned to conduct a study on the effect of iatrogenic tooth extraction and the incidence of malocclusion among pediatric patients coming to the dental OPD with rampant caries. The study will follow patients through scheduled dental visits consisting of clinical and radiographic examinations to record growth and development of the dentition in order to identify early signs of malocclusion and administer preventive orthodontic treatment. What is the most likely study design to be applied by the researchers?
- A. Cohort
 - B. Ecological
 - C. Retrospective
 - D. Cross-sectional
- ANSWER: A

178. A dentistry student wants to determine the existence of tooth decay among communities living in the lowland part of a district. Hence, the student wants to be more familiar with the problem as well as different methodological approaches to write his proposal effectively.

Which activity to be taken to develop a research proposal in this case?

- A. Try to refine the title
- B. Consult the expert
- C. Review of the previous articles
- D. Discuss with a classmate

ANSWER: C

179. A doctor planned to assess the association between unhealthy diet intake and obesity. The clinician recruited and followed 200 adults who regularly consume an unhealthy diet, and 200 adults who regularly follow a healthy diet for six months. At the end of six months, 50 of the adults who took an unhealthy diet became obese (BMI greater than 30 kg/m^2), and 30 adults who took a healthy diet became obese (BMI greater than 30 kg/m^2).

What is the most appropriate interpretation for this scenario?

- A. 60 percent of obesity would be averted if unhealthy diet was avoided by adults
- B. 40 percent of obesity would be averted if unhealthy diet was avoided by adults
- C. 66.7 percent of the excess obesity among adults in the healthy diet group would be contributed by the intake of an unhealthy diet
- D. 33.3 percent of the risk of obesity in unhealthy diet groups would be prevented if the unhealthy diet is avoided

ANSWER: A

180. The head of the nursing department is surveying patients' satisfaction with nursing care delivery at the dental OPD. The nurse disseminated standard checklists that assess a degree of satisfaction with communication skills, clinical dexterity, and professionalism of the nurses.

What is the most appropriate analysis to summarize this finding?

- A. Mode
- B. Median
- C. Arithmetic mean
- D. Geometric mean

ANSWER: A

181. A doctor planned to investigate the magnitude of maxillofacial trauma due to motorcycle accidents in the past 10 years in a city.

What is the most appropriate presentation of data for this case?

- A. Pie chart
- B. Histogram
- C. Bar Graph
- D. Line graph

ANSWER: D

182. A 30-year-old male patient came to the dental OPD for a routine check-up. There was no abnormal finding on the intraoral examination except calculus on the lingual surface of the front lower teeth. While consulting about various modalities for oral hygiene practice, the patient asked if powered toothbrushes were superior to manual cleaning for oral health.

What is the most appropriate initial step of evidence-based dental practice for this case?

- A. Converting the patient's inquiry into a clinical question
- B. Assuring the patient about the efficiency of manual brushes
- C. Evaluating the outcomes of powered and manual toothbrushes
- D. Critically appraising the validity and reliability of powered toothbrushes

ANSWER: A

183. A dental practitioner is treating a patient with suspected oral cancer and wants to ensure accurate diagnosis and appropriate management.

What is the most appropriate step to be performed by the dentist?

- A. Consulting with his colleagues to gather information to diagnose and manage the case
- B. Referring a patient to a specialist and following clinical practice guidelines
- C. Rely on his personal experience to diagnose and manage the case
- D. Conduct a case series of similar cases to inform decision-making

ANSWER: B

184. A dental surgeon working at a local hospital was called to the emergency room to examine a patient who had come to the emergency room after sustaining a motor accident. Upon observation, the patient has facial asymmetry with a shift to the left side, suggesting a facial bone fracture. However, the patient was uncooperative and refused to be examined as he claimed not to feel any pain.

What is the most appropriate initial step to be performed by the dentist?

- A. Attempt to examine the patient
- B. Refer to another health care facility
- C. Use a patient-centered approach and recharge the patient
- D. Understand the patient's mental state when expressing his experience

ANSWER: D

185. A mother brings her three-year-old child to the dental OPD for a checkup. Upon clinical examination, the dentist found that the distal surfaces of the maxillary and mandibular deciduous second molars of the child are in the same vertical plane.

What is the most likely assessment?

- A. Distal step
- B. Mesial step
- C. Occlusal plane
- D. Flush terminal plane

ANSWER: D

186. An eight-year-old girl was brought to the dental clinic by her mom for orthodontic treatment. The clinical examination revealed that the girl has midline diastema, flaring of anterior crowns, and spacing on the upper anterior.

What is the most likely assessment?

- A. Permanent dentition
- B. Flush terminal
- C. Leeway space
- D. Ugly duckling

ANSWER: D

187. A patient came to a dental clinic for orthodontic treatment. Intra-oral examinations showed retroclined maxillary central incisors, labially malposed maxillary lateral incisors, a square-shaped maxillary arch, and an excessive overbite. Additionally, the right and left side permanent first molars are in disto-occlusion.

What is the most likely assessment for this patient?

- A. Class I malocclusion
- B. Class II division one malocclusion
- C. Class III subdivision malocclusion
- D. Class II division two malocclusion

ANSWER: D

188. A patient visited the dental clinic with a complaint of difficulty in chewing on the right side. The pertinent intra-oral finding is buccal cusps of two maxillary right posterior teeth occluding lingually to the buccal cusps of the mandibular teeth.

What is the most likely diagnosis for this case?

- A. Buccal non-occlusion
- B. Lingual non-occlusion
- C. Simple posterior crossbite
- D. Complex posterior crossbite

ANSWER: C

189. A 24-year-old male orthodontic patient presented with intraoral features of spacing between the upper and lower anterior teeth, narrow maxillary arch, proclined maxillary and mandibular incisors, and fish-mouse appearance.

What is the most likely diagnosis for this case?

- A. Bimaxillary protrusion
- B. Skeletal anterior open bite
- C. Dental anterior open bite
- D. Complex posterior open bite

ANSWER: C

190. A patient comes to the dental clinic for orthodontic treatment. After careful evaluation, the patient is diagnosed with mild class II malocclusion. The treatment plan is to extract the maxillary first premolars (tooth 14 and 24) and close the extraction space by moving the posteriors forward and the anterior backward with equal distance.

What is the most appropriate anchorage to be used?

- A. Absolute anchorage
- B. Minimum anchorage
- C. Maximum anchorage
- D. Moderate anchorage

ANSWER: D

191. A dentist analyzes the cephalometric x-ray of a 25-year-old patient using Down's analysis. The result indicates decreased facial angle, increased angle of convexity, increased Y-axis (Growth) axis, and increased cant of occlusal plane.

What is the most likely assessment of this patient?

- A. Class I facial pattern
- B. Class II facial pattern
- C. Class III facial pattern
- D. Pseudo-class III facial pattern

ANSWER: B

192. A dentist prepares alginate powder and water in proportion to make an impression of an orthodontic patient. What kind of mixing must the dentist perform?

- A. Zigzag-motion with intermittent rotations (180 degree) of the spatula
- B. Rotation-motion with intermittent rotations (180 degree) of the spatula
- C. Figure-eight motion with intermittent rotations (180 degree) of the spatula
- D. Linear-horizontal motion with intermittent rotations (180 degree) of the spatula

ANSWER: C

193. A cephalometric analysis of a patient presented for orthodontic treatment using the Tweed method showed that FMA is 25 degrees, FMIA is 65 degrees, and IMPA is 90 degrees.

What is the most likely inference that can be drawn from the analysis?

- A. High-angle patient
- B. Well-balanced face
- C. Vertical growth pattern
- D. Horizontal growth pattern

ANSWER: B

194. A dentist assesses the vertical skeletal relationship of the patient's face using the Frankfurt horizontal plane and the lower border of the mandible clinically with two rulers. The two rulers met anterior to the occipital region.

What is the most likely inference that can be made from clinical assessment?

- A. Non-growing face
- B. Normal growing face
- C. Vertical growing face
- D. Horizontal growing face

ANSWER: C

195. A young orthodontic patient comes to the hospital dental clinic for orthodontic treatment. The senior specialist orders the dentist to make a good impression of the lower dental arch after completing the clinical examination.

What is the most appropriate impression tray?

- A. Perforated tray with no flanges
- B. Perforated tray with low flanges
- C. Perforated tray with high flanges
- D. Perforated tray with moderate flanges

ANSWER: C

196. A dentist planned a serial extraction procedure using Nance's method for an eight-year-old child to guide the occlusion.

What is the most appropriate extraction sequence?

- A. Deciduous first molar, deciduous canine, first premolar
- B. Deciduous canine, deciduous first molar, first premolar
- C. Deciduous canine, first premolar, deciduous first molar
- D. Deciduous first molar, first premolar, deciduous canine

ANSWER: D

197. A nine-year-old boy was brought to the dental clinic by his parents for orthodontic treatment. Interview with the child and parents revealed that the child has a habit of breathing through the mouth. The pertinent clinical findings are a long and narrow face, short and flaccid upper lip, anterior marginal gingivitis, increased overjet, and anterior open bite. The water test was positive.

What is the most appropriate appliance to intercept the habit?

- A. Lip bumper
- B. Oral screen
- C. Adhesive tapes
- D. Removable tongue crib

ANSWER: B

198. A dentist wants to perform serial extraction to guide the occlusion for an eight-year-old child who presented with severe tooth size arch length discrepancy. The proposed method is Dewey's method.

What is the most appropriate extraction sequence?

- A. Deciduous canine, deciduous first molar, first premolar
- B. Deciduous first molar, deciduous canine, first premolar
- C. Deciduous first molar, first premolar, deciduous canine
- D. Deciduous canine, first premolar, deciduous first molar

ANSWER: A

199. A clinical evaluation of a seven-year-old boy showed bilateral loss of maxillary deciduous second molars (tooth 55 and 65). The dentist planned space maintenance as the space available is adequate.

What is the most effective space maintainer?

- A. Distal shoe
- B. Lingual arch
- C. Trans palatal arch
- D. Nance palatal holding an arch

ANSWER: D

200. A dentist examines a five-year-old boy who is seeking dental treatment. Premature loss of the left mandibular deciduous second molar (tooth 75) is observed. The boy has not erupted permanent teeth yet.

What is the most appropriate space maintainer?

- A. Distal shoe
- B. Lingual arch
- C. Band and loop
- D. Crown and loop

ANSWER: A