

Short revision

Ear infection

.Mastoiditis

.Mastoiditis is a bacterial infection of the mastoid bone behind the ear. Without treatment it can lead to meningitis and brain abscess.

.a tender swelling behind the ear which pushes the ear forward

.Diagnosis

.Key diagnostic features are:

.☐ High fever

.☐ Tender swelling behind the ear

.Treatment (WHO)

- ☐ Give IM or IV cloxacillin 50mg/kg every 6 h or ceftriaxone 50 mg

- /kg for total of ten days

- ☐ If there is no response to treatment within 48 hours or the child's

- condition deteriorates, consult surgical specialist to consider

2 Acute Otitis Media

Diagnosis

- ☐ Opaque, or perforated tympanic membrane with discharge

.Give oral antibiotics in one of the following regimens:

.☐ First choice: Oral amoxicillin at 40 mg/kg twice a day for at least 5

.days

.☐ If sensitivity for co-trimoxazole is known, give co-trimoxazole

daily for five days

.Ear discharge culture is important

Chronic Otitis Media

- .If pus has been draining from the ear for 2 weeks or longer, the child has a
- .chronic ear infection.

.This is based on a history of pus draining from the ear for more than 2 weeks. On examination, confirm chronic otitis media) by otoscopy

Treatment

- .Keep the ear dry by wicking
- .☐ Instill topical antibiotic (with or without steroids) once daily for 2 weeks.

.Drops containing quinolones (norfloxacin, ciprofloxacin) are more effective than other antibiotic drops and can be applied for 14 days

