

1. Which sign best indicates the presence of coarctation of the aorta?
 - (A) Clubbing of fingers and toes
 - (B) Generalized cyanosis, especially with crying
 - (C) Rapid and irregular apical heartbeat
 - (D) Bounding brachial pulses with weak femoral pulses

- 2. An 18-month-old is admitted to the emergency department with a diagnosis of seizure. Upon assessment, his vital signs are temperature of 40 C, respirations at 26 breaths/minute, pulse at 120 beats/minute, and blood pressure of 90/69 mm Hg. The nurse should:
- (A) give a tepid sponge bath.
- (B) administer phenytoin (Dilantin).
- (C) Give antibiotic
- (D) aggressively hydrate the child with I.V. fluids.

3. Which nursing intervention is most important to perform before administering digoxin (Lanoxin) to a child? ?

- (A) Checking apical pulse for 1 full minute
- (B) Positioning the child with the head slightly elevated
- (C) Counting the child's respiratory rate for 1 full minute
- (D) Calculating the child's urine output

4. An infant is diagnosed with PDA. Which drug may be administered to achieve pharmacologic closure of the defect?

A) digoxin (Lanoxin)

(B) prednisone (Deltasone)

(C) furosemide (Lasix)

(D) indomethacin (Indocin)

5. A child with difficulty breathing and a “barking cough” is displaying signs associated with which condition?

(A) Cystic fibrosis

(B) Asthma

(C) Epiglottitis

(D) Croup

6. Which condition can rapidly obstruct the airway, requiring immediate attention? ?

- (A) Tonsillitis
- (B) Bronchiolitis
- (C) Epiglottitis
- (D) Tuberculosis

7. A child has decreased output of pink-tinged urine, facial edema, and a history of a sore throat “a little while ago.” The nurse anticipates the doctor will be evaluating the child for: ?

- (A) cryptorchidism.
- (B) adverse effects of hemodialysis.
- (C) Hemolytic Uremic Syndrome
- (D) acute poststreptococcal glomerulonephritis.

8. A child is admitted to the pediatric unit with a new diagnosis of nephrotic syndrome. Which set of symptoms would the nurse expect to see? ?

- (A) Periorbital edema, polyuria, proteinuria, and hyperproteinemia
- (B) Hypercholesterolemia, hypoproteinemia, proteinuria, and periorbital edema
- (C) Pedal edema, hypolipidemia, hematuria, and oliguria
- (D) Hyperlipidemia, glycosuria, hyperproteinemia, and generalized edema

9. A 2-month-old male infant is admitted with a diagnosis of pyloric stenosis.

- Due to the projectile vomiting he has had, he's at risk for:
- (A) metabolic acidosis.
- (B) metabolic alkalosis.
- (C) hyperkalemia.
- (D) hypernatremia.

10. A 12-year-old boy is admitted to the pediatric unit with complaints of RLQ abdominal pain and vomiting. When the nurse checks on the child 2 hours later, he states that the pain has stopped. The nurse should suspect that: ?

- (A) he had indigestion, which has been relieved.
- (B) he's afraid of going to surgery.
- (C) his appendix has ruptured.
- (D) he has irritable bowel syndrome.

11. What is Rheumatic fever? ?

- (A) An acquired cardiac disease that typically follows an untreated streptococcal infection such as pharyngitis.
- (B) A disorder characterized by vasculitis in the small vessels and the potential for cardiac aneurysms.
- (C) Pericarditis Inflammation of the sac that surrounds the myocardium and may lead to heart failure.
- (D) Pericarditis Inflammation of the sac that surrounds the myocardium and may lead to heart failure.

- 3. The best home treatment for croup in a 2 year old is:
 - a. antibiotics prescribed by his physician.
 - b. aspirin if there is a fever.
 - c. cool mist humidifier or steam therapy.
 - d. over-the-counter cough syrups and decongestants.

13. A child with known hemophilia A was brought to the emergency room with complaints of nose bleeding and some bruises in the joints. Which of the following should the nurse anticipate to be given to the child?

- A. Oral iron supplement.
- B. Cyclosporine.
- C. Factor X.
- D. Factor VIII.

14. A child was brought to the emergency department with complaints of nausea, vomiting, fruity-scented breath. The resident on duty diagnosed the child with diabetes ketoacidosis. Which of the following should the nurse expect to administer?

- A. Potassium chloride IV infusion.
- B. Dextrose 5% IV infusion.
- C. Ringer's Lactate.
- D. Normal saline IV infusion.

15. Which of the following should the nurse include in the insulin administration instruction for the parents of a child being discharged on insulin?

- A. Insert the needle and aspirate prior to injecting
- B. Inject insulin into the extremity to be exercised to enhance absorption
- C. The muscles in the abdomen and thigh are the easiest to use for self-administration
- D. Clean the site of injection with soap and water and avoid alcohol

16. Nani who is diagnosed with diabetes mellitus type 1 displays symptoms of hypoglycemia; which of the following actions should the nurse instruct the parents?

- A. Give the child honey (simple sugar)
- B. Give the child milk (complex sugar).
- C. Contact the healthcare provider before doing anything.
- D. Give the child nothing by mouth.

17. Nurse Karen is providing postoperative care for Dustin who has cleft palate (CP); she should position the child in which of the following?

- A. In an infant seat
- B. In the supine position
- C. In the prone position
- D. On his side

18. Baby Jonathan was born with cleft lip (CL); Nurse Barbara would be alert that which of the following will most likely be compromised?

- A. GI function
- B. Locomotion
- C. Sucking ability
- D. Respiratory status

19. Nurse Nancy is assessing a child with pyloric stenosis; she is likely to note which of the following?

- A. "Currant jelly" stools
- B. Regurgitation
- C. Steatorrhea
- D. Projectile vomiting

20. Which of the following organisms is responsible for the development of rheumatic fever?

- A. Streptococcal pneumonia
- B. Haemophilus influenza
- C. Group A beta-hemolytic streptococcus
- D. Staphylococcus aureus

21. The nurse is giving instructions to a mother with a child receiving a liquid oral iron supplement. The nurse tells the mother to:

- A. Take it with meals.
- B. Mix it with food.
- C. Mix it with milk.
- D. Administer it using a straw.

- Nurse Angelo admits a child with suspected type 1 DM; which should the nurse ask the parents?
- A. "Does the child complain of headache?"
- B. "How much exercise does the child get?"
- C. "Has the child's number and type of bowel movements changed?"
- D. "Has the child experienced nocturia or bedwetting?"
- E. "How much candy and sweets does your child take daily?"

- The nurse is aware that the following laboratory values support a diagnosis of pyelonephritis?
- A. Myoglobinuria
- B. Ketonuria
- C. Pyuria
- D. Low white blood cell (WBC) count

- The adolescent patient has symptoms of meningitis: nuchal rigidity, fever, vomiting, and lethargy. The nurse knows to prepare for the following test:
 - A. blood culture.
 - B. throat and ear culture.
 - C. CAT scan.
 - D. lumbar puncture.

- Nurse Kim is teaching a group of parents about otitis media. When discussing why children are predisposed to this disorder, the nurse should mention the significance of which anatomical feature?
- A. Eustachian tubes
- B. Nasopharynx
- C. Tympanic membrane
- D. External ear canal

- The nurse is assessing a 9-month-old boy for a well-baby check-up. Which of the following observations would be of most concern?
- A. The baby cannot say “mama” when he wants his mother.
- B. The mother has not given him finger foods.
- C. The child does not sit unsupported.
- D. The baby cries whenever the mother goes out.

27. Parents bring their infant to the clinic, seeking treatment for vomiting and diarrhea that has lasted for 2 days. On assessment, the nurse in charge detects dry mucous membranes and lethargy. What other findings suggest a fluid volume deficit?

- A. A sunken fontanel
- B. Decreased pulse rate
- C. Increased blood pressure
- D. Low urine specific gravity

- The Foley Family is caring for their youngest child, Justin, who is suffering from tetralogy of Fallot. Which of the following are defects associated with this congenital heart condition?
- A. Aorta exits from the right ventricle, pulmonary artery exits from the left ventricle, and two noncommunicating circulations
- B. Ventricular septal defect, overriding aorta, pulmonic stenosis (PS), and right ventricular hypertrophy
- C. Coarctation of aorta, aortic valve stenosis, mitral valve stenosis, and patent ductus arteriosus
- D. Tricuspid valve atresia, atrial septal defect, ventricular septal defect, and hypoplastic right ventricle

28. It is a serious condition in which the bone marrow does not produce enough new blood cells. It may be passed down from the parents or develop sometime during childhood:

- A. Iron deficiency anemia
- B. Sickle cell disease
- C. Aplastic anemia
- D. Thalassemia
- E. Hemophilia

30. Beta-adrenergic agonists such as albuterol are given to Reggie, a child with asthma. Such drugs are administered primarily to do which of the following?

- A. Dilate the bronchioles
- B. Reduce secondary infections
- C. Decrease postnasal drip
- D. Reduce airway inflammation