

- 1. A postpartum patient was in labor for 30 hours and had ruptured membranes for 24 hours. For which of the following would the nurse be alert?
 - A. Endometritis
 - B. Endometriosis
 - C. Salpingitis
 - D. Pelvic thrombophlebitis

- 2. While the postpartum client is receiving heparin for thrombophlebitis, which of the following drugs would the nurse expect to administer if the client develops complications related to heparin therapy?
 - A. Calcium gluconate
 - B. Protamine sulfate
 - C. Methylergonovine (Methergine)
 - D. Nitrofurantoin (Macrochantin)

3. When caring for a 3-day-old neonate who is receiving phototherapy to treat jaundice, the nurse in charge would expect to do which of the following?

- A. Turn the neonate every 6 hours
- B. Encourage the mother to discontinue breastfeeding.
- C. Notify the physician if the skin becomes bronze in color.
- D. Check the vital signs every 2 to 4 hours.

4. A primigravida in active labor is about 9 days post-term. The client desires a bilateral pudendal block anesthesia before delivery. After the nurse explains this type of anesthesia to the client, which of the following locations identified by the client as the area of relief would indicate to the nurse that the teaching was effective?

- A. Back
- B. Abdomen
- C. Fundus
- D. Perineum

5/The nurse is caring for a primigravida at about 2 months and 1-week gestation. After explaining self-care measures for common discomforts of pregnancy, the nurse determines that the client understands the instructions when she says:

- A. "Nausea and vomiting can be decreased if I eat a few crackers before arising."
- B. "If I start to leak colostrum, I should cleanse my nipples with soap and water."
- C. "If I have a vaginal discharge, I should wear nylon underwear."
- D. "Leg cramps can be alleviated if I put an ice pack on the area."

- 6/The nurse plans to instruct the postpartum client about methods to prevent breast engorgement. Which of the following measures would the nurse include in the teaching plan?
- A. Feeding the neonate a maximum of 5 minutes per side on the first day.
- B. Wearing a supportive brassiere with nipple shields.
- C. Breastfeeding the neonate at frequent intervals.
- D. Decreasing fluid intake for the first 24 to 48 hours.

7. When the nurse on duty accidentally bumps the bassinet, the neonate throws out its arms, hands opened, and begins to cry. The nurse interprets this reaction as indicative of which of the following reflexes?

- A. Startle reflex
- B. Babinski reflex
- C. Grasping reflex
- D. Tonic neck reflex

- 8. Which of the following would the nurse in charge do first after observing a 2-cm circle of bright red bleeding on the diaper of a neonate who just had a circumcision?
 - A. Notify the neonate's pediatrician immediately.
 - B. Check the diaper and circumcision again in 30 minutes.
 - C. Secure the diaper tightly to apply pressure on the site.
 - D. Apply gentle pressure to the site with a sterile gauze pad.

9. Which of the following would the nurse most likely expect to find when assessing a pregnant client with abruption placenta?

- A. Excessive vaginal bleeding
- B. Rigid, board-like abdomen
- C. Tetanic uterine contractions
- D. Premature rupture of membranes

- 10. A pregnant patient asks the nurse if she can take castor oil for her constipation. How should the nurse respond?
- A. “Yes, it produces no adverse effect.”
- B. “No, it can initiate premature uterine contractions.”
- C. “No, it can promote sodium retention.”
- D. “No, it can lead to increased absorption of fat-soluble vitamins.”

- 11. A patient in her 14th week of pregnancy has presented with abdominal cramping and vaginal bleeding for the past 8 hours. She has passed several clots. What is the primary nursing diagnosis for this patient?
- A. Knowledge deficit
- B. Fluid volume deficit
- C. Anticipatory grieving
- D. Pain

- 12. A female adult patient is taking a progestin-only oral contraceptive or mini pill. Progestin use may increase the patient's risk for:
 - A. Endometriosis
 - B. Female hypogonadism
 - C. Premenstrual syndrome
 - D. Tubal or ectopic pregnancy

- 13. A patient with pregnancy-induced hypertension probably exhibits which of the following symptoms?
- A. Proteinuria, headaches, vaginal bleeding
- B. Headaches, double vision, vaginal bleeding
- C. Proteinuria, headaches, double vision
- D. Proteinuria, double vision, uterine contractions

- 14. After 3 days of breastfeeding, a postpartum patient reports nipple soreness. To relieve her discomfort, the nurse should suggest that she:
 - A. Apply warm compresses to her nipples just before feeding.
 - B. Lubricate her nipples with expressed milk before feeding.
 - C. Dry her nipples with a soft towel after feedings.
 - D. Apply soap directly to her nipples, and then rinse.

- 15. The nurse is developing a teaching plan for a patient who is 8 weeks pregnant. The nurse should tell the patient that she can expect to feel the fetus move at which time?
- A. Between 10 and 12 weeks' gestation
- B. Between 16 and 20 weeks' gestation.
- C. Between 21 and 23 weeks' gestation.
- D. Between 24 and 26 weeks' gestation.

- 16. Normal lochial findings in the first 24 hours post-delivery include
 - A. Bright red blood
 - B. Large clots or tissue fragments
 - C. A foul odor
 - D. The complete absence of lochia

- 17. Accompanied by her husband, a patient seeks admission to the labor and delivery area. The client states that she is in labor and says she attended the hospital clinic for prenatal care. Which question should the nurse ask her first?
- A. “Do you have any chronic illness?”
- B. “Do you have any allergies?”
- C. “What is your expected due date?”
- D. “Who will be with you during labor?”

- 18. A patient is in the second stage of labor. During this stage, how frequently should the nurse in charge assess her uterine contractions?
- A. Every 5 minutes.
- B. Every 15 minutes.
- C. Every 30 minutes.
- D. Every 60 minutes.

- 19. A patient is in her last trimester of pregnancy. Nurse Vickie should instruct her to notify her primary health care provider immediately if she notices:
 - A. Blurred vision
 - B. Hemorrhoids
 - C. Increased vaginal mucus
 - D. Shortness of breath on exertion

- 20. A client makes a routine visit to the prenatal clinic. Although she is 14 weeks pregnant, the size of her uterus approximates that in an 18- to 20-week pregnancy. Dr. Charles diagnoses gestational trophoblastic disease and orders ultrasonography. The nurse expects ultrasonography to reveal:
 - A. An empty gestational sac.
 - B. Grapelike clusters.
 - C. A severely malformed fetus.
 - D. An extrauterine pregnancy.

- 21. During a prenatal visit at 4 months gestation, a pregnant client asks whether tests can be done to identify fetal abnormalities. Between 18 and 40 weeks gestation, which procedure is used to detect fetal anomalies?
 - A. Amniocentesis
 - B. Chorionic villi sampling
 - C. Fetoscopy
 - D. Ultrasound

22. Which of the following condition is screened by “Pap Smear”?

- A). Ovarian cancer.
- B). Endometrial cancer.
- c). Cervical cancer.
- d). Breast Cancer.

23. First Movements of the baby felt by the mother are known as?

- a) Lightning
- b). Engagement
- c). Quickening
- D). Ballottement

24. Why does the bed rest is essential in pre-eclampsia?

- a).Prevents eclampsia.
- b). Mobilizes tissue fluid circulations and decreases edema.
- c). Prevents premature labor.
- d). It improves blood circulation and decreases edema.

25. Which time is the most difficult to control diabetes during the maternity cycle?

- a). First trimester.
- b). Last trimester.
- c). Labour & delivery.
- d). Puerperium.

26. Which one is the hazard of induction of labor with Pitocin?

- a). Elevation in blood pressure.
- b). Infection.
- c). Rupture of uterus.
- d). Early rupture of membrane.

27. Which is the common ideal female pelvis?

- a). Android.
- b). Gynaecoid.
- c). Platypelloid.
- D). Anthropoid.

- 28. Which hormone is helpful in the growth of Ovarian follicles?
- a). F.S.H.
- b). L.H.
- c). Estrogen.
- d). Progesterone.

29. What is the name of the first secretion from the mother's breast following delivery?

- a). Sequestrum.
- b). Colostrum.
- c). Lochia.
- d). Leucorrhoea.

30. During Pregnancy, deficiency of which of the following leads to neural tube defects?

- A). Folic acid.
- b). Iron.
- c). Calcium.
- d). Zinc.