

Second term sample question

By dr lammii

.1. While assessing the age of a child, he is developmentally normal, is able to sit without support, can transfer objects from one hand to the other, and speaks in a monosyllabic

.babble. What is the probable age of this child?

.A. 6 month B. 9 month C. 4 month D. 12 month

2. In a well-baby clinic, the child you are examining turns her head when her name is called. She is able to say “Mama”. Her mother told you she can also say “Dada.” She could say ‘bye -bye’ to you when she left the room. What is the least consistent age of this child?

.a 11 months.

.b 15 months

.A 6 year old male child has brown urine and healing impetigo lesions. He presents with hypertension, dyspnea, periorbital edema, and hepatomegaly. What is the most likely cause of his problem?

.A. Pyelonephritis

.B. Post-streptococcal glomerulonephritis

.C. IgA nephropathy

.4, A 14-year old boy who had history of watery diarrhea for 4 days now presents with a complaint of recurrent episodes of blood in the urine. Urine analysis revealed proteinuria and hematuria. You suspect that this could be a renal complication following the acute gastroenteritis state. What is the likely etiology of diarrhea for this boy?

.A. E. coli O157:H7

5, Eight months old HIV exposed infant is on follow up at ART clinic. His DNA PCR done at 6 weeks of age was negative. The nurse advised the mother to continue breast feeding and co-trimoxazole prophylaxis. What should be your response if the mother asks you when to stop the co-trimoxazole?

.A. Continue until the age of 5 years

6, A 3-day-old infant born at 32 weeks' gestation and weighing 1700 g has three episodes of apnea, each lasting 20 to 25 s and occurring after a feeding. During these episodes, the heart rate drops from 140 to 100 beats per min, and the child remains motionless; between episodes, however, the child displays normal activity. Blood sugar is 50 mg/dL and serum calcium is normal. The child's apneic periods most likely are

7, A 16-day-old infant presents with fever, irritability, poor feeding, and a bulging fontanelle. Spinal fluid demonstrates gram-positive cocci. The most likely diagnosis is

- .A. *Listeria monocytogenes*
- .B. group A streptococcus
- .C. group B streptococcus
- D. *Streptococcus pneumoniae*

8, A 21-year-old woman is seronegative for rubella. You caution her against exposure to rubella early in pregnancy because it can result in an infant with

- .A. Thrombocytosis
- .B. Wide-set eyes
- .C. Glaucoma
- .D. Tetralogy of Fallot

9, While bathing her 2-year-old son, a mother feels a mass in his abdomen. A thorough medical evaluation of the child reveals aniridia, hypospadias, horseshoe kidney, and hemihypertrophy. The most likely diagnosis for this child is

A. Neuroblastoma

B. Wilms tumor

.10 A neonate is born at 37 weeks. Pregnancy was complicated by intrauterine growth restriction. When examined you note that the infant has microcephaly, cataracts, hepatomegaly, and petechiae. Diagnosis may include:

.A. Cytomegalovirus (CMV)

.B. Rubella

.C. Toxoplasmosis

.11. A 5 year old male presents to your office with a chronic cough, which is productive, increased at night, recurrent, Worse with exercise and with upper respiratory infections. Chest x-ray findings are normal except for mild hyperinflation. Which is the MOST likely diagnosis?

.A. Sinusitis

.B. Asthma

12, You are awakened in the night by your 2-year-old son, who has developed noisy breathing on inspiration, marked retractions of the chest wall, flaring of the nostrils, and has history of drooling of saliva, unable to drink. He has had a mild upper respiratory infection (URI) for 2 days. The most likely diagnosis is

.A. Asthma

14. A 1-day-old healthy infant with a superficial swelling over the right parietotemporal region that does not cross the suture lines.

- A. Intraventricular hemorrhage
- B. Caput succedaneum
- C. Subarachnoid hemorrhage
- D. Cephalohematoma

.15 An infant has the following findings at 5 min of life: pulse, 130 beats/min normal respiration; cyanotic hands and feet; good muscle tone; and a strong cry and grimace. This infant's Apgar score is:

.A. 7

.B. 8

.C. 9

.16, A 30 wk old average-for-gestational-age male infant is born after 24 hr of ruptured membranes and another 10 hr of labor. After birth, he develops grunting, flaring, and retractions, as well as cyanosis. He requires intubation, ventilation, and surfactant therapy. Despite these measures he is still hypoxic and hypotensive. You should now:

.a. .Do a blood culture and await results

17 Parents of a 3 years old child family has been using a neighbor to care for their child while

.the parents work. The neighbor is diagnosed with pulmonary tuberculosis. PPD test of the 3

.years old is negative. Which of the following is indicated for the 3 years old?

.A. Ethambutol chemoprophylaxis

B. Isoniazid chemoprophylaxis